



OPEN ACCESS

EDITED AND REVIEWED BY
Nicholas M. Barnes,
University of Birmingham, United Kingdom

*CORRESPONDENCE

Yazhou Wang
✉ yazhouw@fmmu.edu.cn
Yonghong Liu
✉ liuredred@163.com

†These authors have contributed equally to this work

RECEIVED 07 December 2024
ACCEPTED 03 February 2025
PUBLISHED 17 February 2025

CITATION

Ma T, Wu Y, Chen B, Zhang W, Jin L, Shen C, Wang Y and Liu Y (2025) Corrigendum: D-serine contributes to seizure development via ERK signaling. *Front. Neurosci.* 19:1541189. doi: 10.3389/fnins.2025.1541189

COPYRIGHT

© 2025 Ma, Wu, Chen, Zhang, Jin, Shen, Wang and Liu. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Corrigendum: D-serine contributes to seizure development via ERK signaling

Tie Ma^{1,2†}, Yin Wu^{3†}, Beibei Chen¹, Wenjuan Zhang¹, Lang Jin¹, Chenxi Shen¹, Yazhou Wang^{4*} and Yonghong Liu^{1*}

¹Department of Neurology, Xijing Hospital, Air Force Military Medical University, Xi'an, China,

²Department of Neurology, The Seventh Medical Center of PLA General Hospital, Beijing, China,

³Department of Pharmacy, Xi'an High-tech Hospital, Xi'an, China, ⁴Department of Neurobiology and Institute of Neurosciences, School of Basic Medicine, Air Force Medical University, Xi'an, China

KEYWORDS

d-serine, serine racemase, astrocyte, epilepsy, hippocampus

A Corrigendum on

[D-serine contributes to seizure development via ERK signaling](#)

by Ma, T., Wu, Y., Chen, B., Zhang, W., Jin, L., Shen, C., Wang, Y., and Liu, Y. (2019). *Front. Neurosci.* 13:254. doi: 10.3389/fnins.2019.00254

In the published article, there was an error in [Figure 5A](#) as published. Duplicate images were mistakenly displayed in the inserts 1 and 2 of [Figure 5A](#). The corrected [Figure 5](#) and its caption appear below.

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.




FIGURE 5 Effects of MK801, LaaβH and CBIO on EEG recordings. (A–D) Mean power of EEG recordings in rats treated with LaaβH (A), saline (B), MK801(C) and CBIO (D). (E–H) Representative frequency images of EEG recordings in rats treated with LaaβH (E), saline (F), MK801 (G) and CBIO (H). *N* = 7–9 rats per group. Compared with the saline control, LaaβH could prolong the onset of seizure occurrence and reduce the mean power of the EEG, while CBIO could shorten the onset of seizure induction and increase the mean power of the EEG.