



# Untangling neural networks with dual retrograde transsynaptic viral infection

Shinya Ohara<sup>1</sup>, Ken-ichi Inoue<sup>1</sup>, Menno P. Witter<sup>2</sup> and Toshio Iijima<sup>1\*</sup>

<sup>1</sup> Division of Systems Neuroscience, Tohoku University Graduate School of Life Sciences, Sendai, Japan

<sup>2</sup> Kavli Institute for Systems Neuroscience and Centre for the Biology of Memory, Norwegian University of Science and Technology, Trondheim, Norway

Using recombinant rabies virus (RV), we developed a dual transsynaptic retrograde tracing technique in the rat central nervous system. Two strains of recombinant RV, injected into two separate loci of the brain, were taken up through axon terminals and carried retrogradely and transsynaptically from neuron to neuron. Each viral strain expresses a unique marker in infected neurons. Therefore, neurons that project transsynaptically to two brain loci can be detected by double-labeling. In this review, we will introduce the advantage of dual viral tracing by recombinant RV, and will also address some potential weaknesses of this technique. Although false negative results may arise due to interference between two strains of RV, the ability of the recombinant RV to visualize the morphology of the infected cell and to infect primates in addition to rodents will make this technique a potential tool to provide new insights into the complex organization of brain networks.

**Keywords:** rabies virus, transsynaptic infection, dual retrograde tracing, neural network, rat hippocampus

## Edited by:

Kathleen S. Rockland, RIKEN Brain Science Institute, Japan

## Reviewed by:

Pavel Osten, Max Planck Institute for Medical Research, Germany  
Ian Wickersham, The Salk Institute for Biological Studies, USA

## \* Correspondence:




**Toshio Iijima** is Professor of Systems Neuroscience in the Graduate School of Life Science and Executive Vice President in charge of research at Tohoku University, Japan. He has conducted research in neuroscience on topics ranging from the medial temporal lobe memory system, the cognitive system of whisker representation, to the olfactory system of the brain. He is now working on the development of a second-generation Brain Machine Interface which incorporates cutting edge technology including ultra high-speed optical imaging of brain activity combined with voltage-sensitive FRET.  
t-ijima@mail.tains.tohoku.ac.jp

## BACKGROUND

The central nervous system (CNS) is composed of multiple, interconnected neural circuits. One way to elucidate the complex wiring of the brain is to use tracer substances (dyes or proteins), which are taken up by neurons and carried by axonal transport. In cases where the tracer is transported retrogradely, such as Fast Blue, Diamidino Yellow or Fluorogold, neurons projecting to the injected site will be labeled. Such methods potentially visualize the complete set of converging inputs to a particular brain region. The combination of two retrograde tracers which can be distinguished, for example by different fluorescence wavelength (Kuypers et al., 1980), is referred to as **double retrograde labeling**. This method is useful to assess whether a particular brain region has divergent projections to several regions. Moreover, one can reveal whether the projections originate from single neurons (divergent projection) or from nearby

distinct neurons (parallel projection). These tracing techniques have provided us remarkable insights into the afferent and efferent connections of brain regions.

Since the early 1990's, viruses have also been used to visualize neuronal connections (Kuypers and Ugolini, 1990). Unlike most conventional chemical retrograde tracers, viral retrograde tracers reliably move from one neuron to another neuron mainly via synapses. More importantly, they replicate in infected neurons, resulting in strong amplification of signals. These features enable precise tracing of chains of neurons connected via synapses (**Figure 1**). A variety of neurotropic viruses have been used for such **transsynaptic tracing**, namely, herpes simplex virus type 1 (HSV-1), pseudorabies (PRV), and **rabies virus** (RV). Among these viruses, the Bartha strain of PRV (PRV-Bartha) and the challenge virus standard (CVS) of RV are used often as transsynaptic



### Double retrograde labeling

A tracing method which uses two retrograde tracers in order to detect a neuron with collateralized projections to two different target regions.

### Transsynaptic tracing

A tracing method that can detect synaptically connected neurons (neural circuits) by using tracer substances or viruses that are transported across synapses.

### Rabies virus

Nonsegmented negative-strand RNA-virus which belongs to the genus *Lyssavirus* of the family *Rhabdoviridae*. Rabies virus infects hosts at the periphery and moves to the central nervous system via retrograde axonal and transsynaptic transport. This virus finally causes fatal disease in human and animals.

retrograde tracers (for reviews, see Aston-Jones and Card, 2000; Kelly and Strick, 2000).

To further elucidate the complicated neural networks in the CNS, such as the interaction of multiple neural circuits, a method which combines double retrograde labeling with viral transsynaptic tracing would be useful. The idea of this method, dual viral tracing, is explained in **Figure 2**. Two strains of recombinant viruses are used as transsynaptic tracers, each of which expresses a unique marker protein. With this method, two different neuronal circuits can be labeled distinctly, and a single neuron, from which the two diverging circuits are originating, can be visualized by co-expression of two marker proteins (double labeling). In our original paper (Ohara et al., 2009), we described dual viral tracing in the

rat CNS using recombinant strains of RV. In this article, we will review the reported use of various dual viral tracer approaches, the advantages of dual viral tracing using RV recombinants and point out potential weaknesses and promises of the latter method.

### DUAL TRANSSYNAPTIC TRACING USING PRV RECOMBINANTS

Dual viral tracing has been introduced by Jansen et al. (1995) using recombinant strains of PRV-Bartha. PRV-Bartha is an attenuated strain of PRV which transports along the axon in a retrograde direction. Unlike other  $\alpha$ -herpesviruses (HSV-1 and PRV), PRV-Bartha is less likely to induce cell lysis and therefore will hardly infect anatomically non-connected nearby cells by local spread of the virus. Because of these features, this virus is widely used as a transsynaptic retrograde tracer in rodents. Jansen et al. (1995) used two recombinants of PRV-Bartha which express unique markers [gC viral glycoprotein and  $\beta$ -galactosidase ( $\beta$ -gal)] to address a long-standing issue: whether there are neurons which regulate the functions of two distinct autonomic systems. The two recombinant strains were injected, respectively, into one of two peripheral organs, adrenal gland and stellate ganglion (the major sympathetic source that innervates the heart). The distribution of double labeled neurons in the hypothalamus and brainstem demonstrated the existence of neurons which innervate these two peripheral structures. However, the rate of successful double infection in this first study was very low (8 out of 256 rats) since the virus strains used in this study were nonisogenic strains that differ in their invasive properties. The strain with stronger invasive properties spreads within neural circuits faster than the other strain and prevents the other strain from subsequently infecting the same neuron (Kim et al., 1999). The low rate of successful double infection due to this interference problem (**viral interference**) has been settled partially by using two isogenic PRV recombinants which present similar invasive properties (Cano et al., 2004). Dual viral tracing with these isogenic strains, which express either  $\beta$ -gal or enhanced green fluorescent protein (EGFP) as a marker protein, has been used to identify brainstem neurons that control the simultaneous activation of multiple muscles that are involved in certain actions, such as respiration and emesis (Billig et al., 2000). This method has also been used to demonstrate the existence of neurons which simultaneously coordinate the activations of both motor and autonomic systems (Krout et al., 2003; Kerman et al., 2006).

### Viral interference


An interference mechanism that is established after primary infection, preventing the infected cell from being superinfected by the same or a closely related virus. This inhibition mechanism may occur at various stages of the viral life cycle, such as surface binding, endocytosis, and subsequent replication steps.

### DUAL TRANSSYNAPTIC TRACING USING RV RECOMBINANTS

In the original paper (Ohara et al., 2009), we used recombinant strains of RV instead of the previously used PRV recombinants. Similar to PRV-Bartha, RV transports along the axon exclusively in the retrograde direction and spread to neurons that are synaptically connected (Iwasaki and Clark, 1975; Charlton and Casey, 1979). However, there are several differences in the infection properties in these two viruses. Unlike PRV infection, RV selectively infects neurons, not glia cells. Furthermore, RV does not induce cellular lysis and subsequent local viral spread (Ugolini, 1995). One further prominent characteristic of RV is that it can infect primates in addition to rodents

whereas PRV cannot (Callaway, 2008). These features make RV a powerful tool to reveal the complicated connections within the CNS especially in non-human primates, such as connections of the basal ganglia, cerebellum and cerebral cortex (Kelly and Strick, 2003; Hoshi et al., 2005), or specifically in the visual pathway (Nassi and Callaway, 2006).

We developed a recombinant RV which was derived from a vaccine strain of RV (HEP-Flury) except that the glycoprotein (G) gene was taken from the CVS strain (Inoue et al., 2004). This recombinant virus (rHEP5.0-CVSG) effectively propagated transsynaptically in a retrograde direction. Furthermore, the morphological features of the infected neurons were clearly visu-



alized by using antibodies against the expressed marker protein, such as EGFP or  $\beta$ -gal. We used this recombinant RV and achieved dual transsynaptic tracing in the entorhinal-hippocampal network of the rat. The entorhinal cortex and hippocampal region are known to contribute to encoding, consolidation and retrieval of declarative memories. The afferent and efferent connections within these regions have been well studied (for an overview, see Witter and Amaral, 2004). Briefly, the areas of hippocampus, including the dentate gyrus, areas CA3, CA1 and the entorhinal cortex (EC), exhibit a series of connections that are ideal to test the potential of our recombinant strains of viruses for single and double labeling (Figure 3A). Axons of pyramidal cells in the hippocampal CA3 region project bilaterally to CA1 (Swanson et al., 1980; Laurberg and Sorensen, 1981; Li et al., 1994). In turn, CA1 pyramidal cells project to EC, and with the exception of those in the most dorsal tip of CA1, all neurons project strictly ipsilaterally (van Groen and Wyss, 1990). On the basis of this established anatomy, we hypothesized that a single CA3 cell influences EC bilaterally, with only one CA1 synapse in between. This inferred connectivity, however, has never been demonstrated experimentally. Our new dual viral tracing technique would be perfectly suited to confirm this connection in a straightforward way.


We created two recombinant strains of virus, each expressing one of two distinct marker proteins,  $\beta$ -gal and EGFP (pHEP5.0-CVSG- $\beta$ -gal and pHEP5.0-CVSG-EGFPx2), and injected each of these two strains to the left and right medial entorhinal cortex (MEC) separately. 6 days after injection, which is an approximate time required for the recombinant virus to infect the second-order upstream neurons, many labeled neurons were observed in the dorsal hippocampus. CA1 neurons were predominantly labeled by the recombinant strain that was injected into the ipsilateral MEC, whereas  $\beta$ -gal and EGFP labels intermingled in the bilateral CA3 region. 12.5–21.7% of the labeled CA3 neurons were double-labeled (Figure 3B). In some double-labeled neurons, the level of expression of at least one of the two marker proteins was high enough to visualize the morphological feature of the neuron. The results demonstrate that single neurons in CA3 are connected to MEC bilaterally, with only one synaptic relay at the CA1. In view of the known organization of this neuronal chain, the present finding was expected, but is the first direct demonstration of this transsynaptic connection from individual CA3 neurons to MEC bilaterally. This study also demonstrated the

potential use of this dual viral tracing method using RV recombinants to reveal the complicated circuits in the CNS.

### CAVEATS IN THE USE OF DUAL VIRAL TRACING

Although the dual viral tracing method using RV recombinants is powerful in dissecting multiple circuits, a number of issues should be considered. First, not all cell types or connections can be infected. Viral specificity occurs in case of granule cells in the dentate gyrus, which cannot be infected by RV (Jackson and Reimer, 1989), and CA3 pyramidal cells will not be infected through synapses of Schaffer collaterals by PRV-Bartha (Sik et al., 2006). Second, it is important to assess the possibility of multiple transsynaptic routes between two brain structures. An efficient way is to carry out single viral tracing experiments first and to compare the results with established connectivity. In case of the connections between CA3 and EC, single tracing experiments indicated that a possible connection mediated by the lateral septum (Witter and Amaral, 2004) was an unlikely explanation for the reported transsynaptic labeling in CA3 following injections in EC.

The use of these single tracing experiments also allowed assessing the temporal progression of the two viral strains in the targeted circuit, which is essential to realize efficient double-labeling. As shown in the PRV studies, false negative results may arise in dual viral tracing: neurons which innervate the two injection regions may not be double-labeled due to viral interference. We have demonstrated the effect of this interference on double labeling in an *in vitro* preparation, and showed that the efficiency of double labeling decreases as the interval between the two infections increases, in accordance with previous report using PRV recombinants (Banfield et al., 2003). Thus, for efficient double labeling using viral tracers, two strains of viruses must infect a neuron with only a few hours difference. To achieve this, we must consider the time required for the two recombinant strains to infect the neurons of interest. This difference is determined by the propagation speed of the two strains and by the infection pathways, such as the number of synapses, the length of the axon, and the strength of connections (Lycke and Tsiang, 1987; Ugolini, 1995). Therefore, in order to realize efficient double-labeling the timing of the two injections should be adjusted, taking into account the specifics of each of the two strains such that the two viral strains reach the neurons of interest within a short time period.



### FUTURE PERSPECTIVES

Dual transsynaptic tracing using recombinant RV is a powerful method since RV specifically infects neural circuits in a retrograde manner in a broad range of animals including rodents and primates. Although there still are some problems in the use of this method as described above, recent developments in genetic techniques regarding RV provides the hint of solutions to these problems. Wickersham et al. (2007) reported an elegant technique to label neurons that project to a single, genetically targeted neuron *in vitro* by applying two modifications to RV: (1) deleting the G gene from the genome, and (2) providing the virus with an avian virus enve-

lope (EnvA). This modified virus can only infect specific neurons expressing the EnvA receptor (TVA), and spread transsynaptically to directly connected neurons only if the infected cell is also transfected with G gene. This strategy may be used to restrict the infection pathway of the virus to a certain route and will facilitate the interpretation of the transsynaptic labeling.

Genetic modification may also provide a solution to the problem of false negative results due to viral interference. A previous study in Vesicular stomatitis virus, which belongs to the same family, *Rhabdoviridae*, as RV, has shown that the G protein interferes with the viral uptake of a subsequent

virus (Whitaker-Dowling et al., 1983). Therefore, recombinant RV with low G protein expression may diminish the effect of viral interference, but right now this problem has not been solved. We must further realize that quantitative analysis is difficult in dual viral tracing since the results may underestimate the real number of collateralized neurons, due to interference. On the good side, those neurons that express both marker proteins undoubtedly have anatomical connections to both of the infected target areas. Furthermore, since our recombinant RV can reveal the morphological characteristics of the labeled neurons, the double-labeled neuron can be morphologically typed. The method can also be combined with antibody staining against neural cell markers to classify the cell in more detail. It may also be inter-

esting to combine this method with anterograde tracing, which visualizes inputs from a certain region to the virally-infected neurons. This will allow qualitative and quantitative analyses using confocal- and potentially electron-microscopy. We thus expect that this dual viral tracing with recombinant RV, for which we provided proof of principle in the entorhinal-hippocampal system, will be of use to untangle complex connections in the CNS and to visualize the complex architectures that underlie a variety of brain functions.

## ACKNOWLEDGMENTS

This work was supported by Grant-in-Aid for Scientific Research on Priority Areas (Integrative Brain Research) from the Ministry of Education, Culture, Sports, Science, and Technology of Japan.

## REFERENCES

- Aston-Jones, G., and Card, J. P. (2000). Use of pseudorabies virus to delineate multisynaptic circuits in brain: opportunities and limitations. *J. Neurosci. Methods* 103, 51–61.
- Banfield, B. W., Kaufman, J. D., Randall, J. A., and Pickard, G. E. (2003). Development of pseudorabies virus strains expressing red fluorescent proteins: new tools for multisynaptic labeling applications. *J. Virol.* 77, 10106–10112.
- Billig, I., Foris, J. M., Enquist, L. W., Card, J. P., and Yates, B. J. (2000). Definition of neuronal circuitry controlling the activity of phrenic and abdominal motoneurons in the ferret using recombinant strains of pseudorabies virus. *J. Neurosci.* 20, 7446–7454.
- Callaway, E. M. (2008). Transneuronal circuit tracing with neurotropic viruses. *Curr. Opin. Neurobiol.* 18, 617–623.
- Cano, G., Card, J. P., and Sved, A. F. (2004). Dual viral transneuronal tracing of central autonomic circuits involved in the innervation of the two kidneys in rat. *J. Comp. Neurol.* 471, 462–481.
- Charlton, K. M., and Casey, G. A. (1979). Experimental rabies in skunks: immunofluorescence light and electron microscopic studies. *Lab. Invest.* 41, 36–44.
- Hoshi, E., Tremblay, L., Feger, J., Carras, P. L., and Strick, P. L. (2005). The cerebellum communicates with the basal ganglia. *Nat. Neurosci.* 8, 1491–1493.
- Inoue, K., Ohara, S., Ichijo, H., Kakei, S., and Iijima, T. (2004). Development of rabies virus vectors for neuron-specific and transsynaptic delivery of a foreign gene (Abstract). *Abstr. Soc. Neurosci.* 30, 124.12.
- Iwasaki, Y., and Clark, H. F. (1975). Cell to cell transmission of virus in the central nervous system. II. Experimental rabies in mouse. *Lab. Invest.* 33, 391–399.
- Jackson, A. C., and Reimer, D. L. (1989). Pathogenesis of experimental rabies in mice: an immunohistochemical study. *Acta. Neuropathol.* 78, 159–165.
- Jansen, A. S., Nguyen, X. V., Karpitskiy, V., Mettenleiter, T. C., and Loewy, A. D. (1995). Central command neurons of the sympathetic nervous system: basis of the fight-or-flight response. *Science* 270, 644–646.
- Kelly, R. M., and Strick, P. L. (2000). Rabies as a transneuronal tracer of circuits in the central nervous system. *J. Neurosci. Methods* 103, 63–71.
- Kelly, R. M., and Strick, P. L. (2003). Cerebellar loops with motor cortex and prefrontal cortex of a nonhuman primate. *J. Neurosci.* 23, 8432–8444.
- Kerman, I. A., Akil, H., and Watson, S. J. (2006). Rostral elements of sympatho-motor circuitry: a virally mediated transsynaptic tracing study. *J. Neurosci.* 26, 3423–3433.
- Kim, J. S., Enquist, L. W., and Card, J. P. (1999). Circuit-specific coinfection of neurons in the rat central nervous system with two pseudorabies virus recombinants. *J. Virol.* 73, 9521–9531.
- Krout, K. E., Mettenleiter, T. C., and Loewy, A. D. (2003). Single CNS neurons link both central motor and cardi sympathetic systems: a double-virus tracing study. *Neuroscience* 118, 853–866.
- Kuypers, H. G., Bentivoglio, M., Catsman-Berrevoets, C. E., and Bharos, A. T. (1980). Double retrograde neuronal labeling through divergent axon collaterals, using two fluorescent tracers with the same excitation wavelength which label different features of the cell. *Exp. Brain Res.* 40, 383–392.
- Kuypers, H. G., and Ugolini, G. (1990). Viruses as transneuronal tracers. *Trends Neurosci.* 13, 71–75.
- Laurberg, S., and Sorensen, K. E. (1981). Associational and commissural collaterals of neurons in the hippocampal formation (hilus fasciae dentatae and subfield CA3). *Brain Res.* 212, 287–300.
- Li, X. G., Somogyi, P., Ylinen, A., and Buzsáki, G. (1994). The hippocampal CA3 network: an in vivo intracellular labeling study. *J. Comp. Neurol.* 339, 181–208.
- Lykke, E., and Tsiang, H. (1987). Rabies virus infection of cultured rat sensory neurons. *J. Virol.* 61, 2733–2741.
- Nassi, J. J., and Callaway, E. M. (2006). Multiple circuits relaying primate parallel visual pathways to the middle temporal area. *J. Neurosci.* 26, 12789–12798.
- Ohara, S., Inoue, K., Yamada, M., Yamawaki, T., Koganezawa, N., Tsutsui, K., Witter, M. P., and Iijima, T. (2009). Dual transneuronal tracing in the rat entorhinal-hippocampal circuit by intracerebral injection of recombinant rabies virus vectors. *Front. Neuroanat.* 3, 1. doi: 10.3389/neuro.05.001.2009.
- Sik, A., Cote, A., and Boldogkoi, Z. (2006). Selective spread of neurotropic herpesviruses in the rat hippocampus. *J. Comp. Neurol.* 496, 229–243.
- Swanson, L. W., Sawchenko, P. E., and Cowan, W. M. (1980). Evidence that the commissural, associational and septal projections of the regio inferior of the hippocampus arise from the same neurons. *Brain Res.* 197, 207–212.
- Ugolini, G. (1995). Specificity of rabies virus as a transneuronal tracer of motor networks: transfer from hypoglossal motoneurons to connected second-order and higher order central nervous system cell groups. *J. Comp. Neurol.* 356, 457–480.
- van Groen, T., and Wyss, J. M. (1990). Extrinsic projections from area CA1 of the rat hippocampus: olfactory, cortical, subcortical, and bilateral hippocampal formation projections. *J. Comp. Neurol.* 302, 515–528.
- Whitaker-Dowling, P., Youngner, J. S., Widnell, C. C., and Wilcox, D. K. (1983). Superinfection exclusion by vesicular stomatitis virus. *Virology* 131, 137–143.
- Wickersham, I. R., Lyon, D. C., Barnard, R. J., Mori, T., Finke, S., Conzelmann, K. K., Young, J. A., and Callaway, E. M. (2007). Monosynaptic restriction of transsynaptic tracing from single, genetically targeted neurons. *Neuron* 53, 639–647.
- Witter, M. P., and Amaral, D. G. (2004). Hippocampal Formation: The Rat Nervous System, 3rd edn. Burlington, Academic Press, pp. 635–704.

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 24 July 2009; paper pending published: 13 August 2009; accepted: 20 August 2009; published: 15 December 2009.  
Citation: *Front. Neurosci.* (2009) 3, 3: 334–349. doi: 10.3389/neuro.01.032.2009

Copyright © 2009 Ohara, Inoue, Witter and Iijima. This is an open-access article subject to an exclusive license agreement between the authors and the Frontiers Research Foundation, which permits unrestricted use, distribution, and reproduction in any medium, provided the original authors and source are credited.