



OPEN ACCESS

EDITED BY
Nemat Ali,
King Saud University, Saudi Arabia

REVIEWED BY
Rehan Khan,
Rutgers University, Newark,
United States
Farhat Parween,
National Institutes of Health (NIH),
United States

*CORRESPONDENCE
Norman B. Gaylis
ngaylismd@gmail.com
Yankel Gabet
yankel@tauex.tau.ac.il

†These authors have contributed
equally to this work

SPECIALTY SECTION
This article was submitted to
Nutritional Immunology,
a section of the journal
Frontiers in Nutrition

RECEIVED 01 September 2022
ACCEPTED 03 October 2022
PUBLISHED 25 October 2022

CITATION
Gaylis NB, Kreychman I, Sagliani J,
Mograbi J and Gabet Y (2022) The
results of a unique dietary supplement
(nutraceutical formulation) used
to treat the symptoms of long-haul
COVID.
Front. Nutr. 9:1034169.
doi: 10.3389/fnut.2022.1034169

COPYRIGHT
© 2022 Gaylis, Kreychman, Sagliani,
Mograbi and Gabet. This is an
open-access article distributed under
the terms of the [Creative Commons
Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use,
distribution or reproduction in other
forums is permitted, provided the
original author(s) and the copyright
owner(s) are credited and that the
original publication in this journal is
cited, in accordance with accepted
academic practice. No use, distribution
or reproduction is permitted which
does not comply with these terms.

The results of a unique dietary supplement (nutraceutical formulation) used to treat the symptoms of long-haul COVID

Norman B. Gaylis^{1*†}, Ida Kreychman^{2†}, Joanne Sagliani¹,
Josef Mograbi³ and Yankel Gabet^{2*}

¹The COVID Long Haul Center at Arthritis & Rheumatic Disease Specialties (AARDS), Miami, FL, United States, ²Department of Anatomy and Anthropology, Sackler Faculty of Medicine, Tel Aviv University, Tel Aviv-Yafo, Israel, ³Independent Researcher, Tel Aviv-Yafo, Israel

Long-COVID is a syndrome characterized by debilitating symptoms that persist over 3 months after infection with the SARS-CoV-2 virus. It affects 15 to 33% of COVID-19 recovered patients and has no dedicated treatment. First, we found that β -caryophyllene and pregnenolone have a significant synergistic effect in the resolution of LPS-induced sepsis and inflammation in mice. Then we combined these two compounds with seven others and designed a unique dietary supplement formulation to alleviate long COVID inflammatory and neurological disorders. We performed a one-arm open-labeled study at a single site with 51 eligible patients from 18 states. Each participant recorded the severity level of 12 symptoms (including fatigue, weakness, cardiac and neurological symptoms, shortness of breath, gastrointestinal disorders, ageusia or anosmia, anxiety, joint pain, rash, cough, and insomnia) at baseline, 2- and 4-week time points. On average, all the symptoms were significantly milder after 2 weeks, with further improvement after 4 weeks. Importantly, each symptom was significantly attenuated in 72 to 84% of the participants. There were no significant adverse effects. Our data indicate that the use of this nutraceutical product is a safe and significantly efficient option to reduce multiple symptoms of long COVID.

KEYWORDS

dietary (food) supplements, long-COVID-19, immunology and inflammation, cannabinoids, CB2 agonists

Introduction

Long-haul COVID is characterized by chronic and often debilitating symptoms following acute Coronavirus disease 2019 (COVID-19) caused by the SARS-CoV-2 virus. Long COVID is very challenging to diagnose, treat and categorize as it combines multiple and different symptomatic presentations in sufferers.

The Center for Disease Control (CDC) in June 2022 published a report that 40% of adults in the US reported having COVID-19 in the past, of which nearly 1 in 5 are still having symptoms of long COVID. The CDC defines Long COVID as symptoms that last three or more months after first contracting the virus and did not exist before the COVID-19 infection. Symptoms may last well over 1 year (1). Other studies reported a prevalence ranging from 1 in 8 to 1 in 3 (2, 3), significantly higher than after influenza (2). There appeared to be differences in the prevalence of long COVID between states (2).

Differences in the reported prevalence of long-COVID likely results from the disparity of symptoms and the lack of a unified definition for this condition. Importantly, this condition cannot be ignored as it presents a set of symptoms that often severely impact the quality of life and ability to carry out daily activities. For example, in one survey, 44% percent of patients with Long COVID reported not being able to work at all, compared to their pre-COVID-19 work capacity, and 51% had reduced their working hours (4). Overall, the economic burden may approximate \$50 billion annually in lost salary only (4).

An international cohort of 3,762 participants from 56 countries identified 203 symptoms in 10 organ systems that persisted at least 4 weeks after a confirmed diagnosis of COVID-19 (5). The CDC listed the most common symptoms of COVID-19 in a survey they initiated in April 2022.¹ These included tiredness or fatigue, difficulty thinking, concentrating, forgetfulness, or memory problems (sometimes referred to as "brain fog"), difficulty breathing or shortness of breath, joint or muscle pain, fast-beating or pounding heart (also known as heart palpitations), chest pain, dizziness on standing, menstrual changes, changes to taste/smell, or inability to exercise.

The exact underlying cause of long-COVID remains uncertain, but most reports agree that this condition is associated with a persistent viral infection and long-lasting inflammation (6, 7). Specific neurological symptoms (fatigue, brain fog, anosmia, and ageusia/dysgeusia) in long COVID resemble "sickness behavior," a response of the autonomic nervous system to pro-inflammatory cytokines (8). The long-standing dysautonomia has been proposed to result from sympathetic/parasympathetic imbalance (7, 9).

To the best of our knowledge, no medicine is currently dedicated to treating long-haulers (patients suffering from long-COVID). In the absence of established protocols, each symptom is treated separately with different drugs for different symptoms, even though they all have a common cause. Even when combining the current therapeutic approaches with rehabilitation (10–14), success is minimal, and the treatments are associated with multiple adverse effects.

We thus designed a formula to address the underlying chronic inflammatory status and autonomic imbalance that characterize many long-haulers. The first challenge was to help restore the function of various organ systems; the second challenge was to use only US-recognized dietary supplements to be able to address the unmet need in a very short time.

Tel Aviv University, in collaboration with Arthritis & Rheumatic Disease Specialties (AARDS) based in Miami, FL, USA, formulated a unique oral nutraceutical supplement containing only approved dietary supplements and ingredients "generally recognized as safe" (GRAS) by the United States Food and Drug Administration (FDA). This combination would, by design, be natural, free of any known side effects, and with limited drug interactions. The final formulation relied on new experimental data (presented here), clinical experience in treating long-hauler patients at AARDS with multiple clinical symptoms over 2 years, and scientific literature (Table 1). We selected the ingredients for their unique immuno-modulation properties and activities, reduction in pain, anxiety and depression, potential effect on dysautonomia, anticoagulation activity, as well as the ability to inhibit viral replication. We report here the results of a clinical study aimed at testing this nutraceutical formula as a standalone treatment in a population of COVID long haulers.

We and others have reported that selective CB2 agonists are potent immunomodulators of the innate immune system, both *in vivo* and *in vitro* (15, 16). We showed that CB2 agonists inhibit cytokine expression in LPS-exposed macrophages in cultures and decrease ear edema in a skin inflammation model in mice (16). Following these results, we tested the hypothesis that β -caryophyllene (β CP), a dietary terpene that is also a selective CB2 agonist (17, 18), reduces the exaggerated inflammatory response induced by pathogens.

Another approach to attenuate inflammation and associated tissue damage is using steroids. In contrast to glucocorticosteroids such as dexamethasone, pregnenolone is a steroid hormone precursor with known anti-inflammatory properties in myeloid cells (19), does not induce lymphocyte apoptosis (20) and may even promote thymocyte survival and differentiation (21). Pregnenolone (Preg) and its metabolites suppress the secretion of tumor necrosis factor α and interleukin-6 mediated through TLR2 and TLR4 signaling in macrophages (22). Also relevant to long-haulers, Preg can help with cognitive and neurological issues, partly *via* binding to TRPM3, without causing bone loss (23–26).

In addition to β CP and Preg, we included Dehydroepiandrosterone (DHEA), Bromelain, St. John's Wort extract, Boswellia Serrata gum/resin extract (AKBA), Quercetin, zinc compound, and vitamin D. The rationale for including each compound is summarized in Table 1.

1 cdc.gov

TABLE 1 List of compounds included in our nutraceutical combination and their relevance for COVID long haulers.

Active ingredient	Properties and activities	References
β -caryophyllene (β CP)	Antioxidant, anti-inflammatory, analgesic	(18, 27–29)
Pregnenolone	Modulator of inflammation, regulator of neuroinflammation	(22, 30–32)
Dehydroepiandrosterone (DHEA)	Anti-inflammatory, improves dysfunction of the hypothalamic-pituitary-adrenal (HPA) axis, immune modulator	(33–35)
Bromelain	Anti-inflammatory, antiviral (Anti-SARS-CoV-2), fibrinolytic	(36–39)
St. John's Wort extract	SRI* for treating autonomic dysfunction and impaired balance between the sympathetic and parasympathetic nervous systems	(40, 41)
Boswellia Serrata gum/resin extract (AKBA)	Anti-inflammatory, COVID-19 therapeutic agent, respiratory support	(42–44)
Quercetin	Anti-inflammatory, antiviral, immune modulator, antioxidant	(45–47)
Zinc compound	Essential to preserve natural tissue barriers such as the respiratory epithelium, balanced function of the immune system and the redox system, antiviral	(48, 49)
Vitamin D	Reducing inflammation induced by SARS-CoV-2 infection	(50, 51)

*SRI, serotonin reuptake inhibitor.

Materials and methods

Mouse study

Male wild type 8–10 weeks old mice of inbred strain C57BL/6J-RCC were obtained from Envigo Ltd (Jerusalem, Israel). All experiments were in accordance and with the approval of the institutional animal care and use committee of Tel-Aviv University for these experiments (permit number 01-20-022). The mice were divided into treatment groups 24 h before LPS administration. All treatments were administered intraperitoneally (IP). β -Caryophyllene (β CP) and pregnenolone were injected every 12 h starting 24 h before the LPS injection. A single IP injection of LPS was administered at 25 μ g/gr dose. This study had two control groups as indicated in the experiments below. One that received PBS instead of the LPS injection ("PBS"), and one that received PBS instead of the treatments (CB2 agonist or steroid) before the LPS injection ("LPS"). For humane reasons murine sepsis score (MSS) was

used as a surrogate for survival (52). A low MSS score (0–2) served as a surrogate for survival while mice that reached the score 3 were defined as critically ill and were euthanized.

Statistical analyses for the animal study

For survival experiments in mice, Gehan–Breslow–Wilcoxon test was used for multiple groups comparison. For disease progression we used Dunnett's multiple comparisons test. Differences between groups were considered significant when $p < 0.05$.

Clinical study performed at arthritis and rheumatic disease specialties

The new combination of nutraceuticals including the compounds listed in Table 1 was tested in a one-arm, open-label clinical trial. The amount of each compound in the formulation and the total amount per serving are detailed in Table 2. Participation was voluntary, recruitment was from local patient population as well as social media advertising. All subjects had to have documented evidence of prior COVID infection (PCR or Rapid test) and were having ongoing symptoms (1 or more) for a minimum of 3 months from the time of infection. The existence of any of these symptoms prior to contracting COVID was considered an exclusion. All subjects were required to complete a survey listing and rating their symptoms and all participants were interviewed by the research staff at baseline to verify inclusion criteria.

Following qualification, subjects were provided with the nutraceutical supplement for a 2-week period with dosing instructions (1 serving twice a day with food) after which they were required to repeat the process of documenting any change

TABLE 2 Amount of each compound included in the nutraceutical formulation per serving. The participants were required to take one serving twice a day with food.

Active ingredient	Amount per serving
β -caryophyllene (β CP)	40 mg
Pregnenolone	40 mg
Dehydroepiandrosterone (DHEA)	30 mg
Bromelain (2400 GDU*/g)	416 mg
St. John's Wort extract	150 mg
<i>Boswellia Serrata</i> gum/resin extract (AKBA)	100 mg
Quercetin (<i>Sophora Japonica</i>)	40 mg
Zinc (as Zinc Picolinate)	12 mg
Vitamin D	25 μ g (1000 IU)

*GDU, gelatin digesting unit.

in symptoms relative to baseline and their voluntary decision to continue with treatment for a further 2-week period. At the end of 4 weeks, their change in symptoms from baseline was documented as well as an overall global response.

The total number of patients completing the 4-week evaluation was 51, the age range was 21–73 years of age, female to male ratio was approximately 2 to 1 and the patients originated from 18 US states.

Statistical analysis of the clinical study

We compared the score of each symptom at 2 and 4 weeks to the baseline for each participant using multiple paired *t*-tests. Individual variance was assumed for each symptom and a Benjamini–Krieger–Yekutieli False Rate Discovery Rate (FDR) was used to account for the multiple comparisons. For all comparisons, differences between time points were considered significant when $p < 0.01$.

Results

Animal study

Testing of anti-inflammatory compounds in mice

Here we tested the therapeutic potential of β -caryophyllene and that of pregnenolone in a mouse model of LPS-induced sepsis. β CP (25 mg/kg) and Preg (10 mg/kg) were administered 24 and 12 h before LPS injection (25 mg/kg), and then every 12 h. All the injections were given i.p. A Murine Septic Score (MSS) was recorded every 12 h and survival was defined as MSS < 3. Over the 4-day follow-up, we found a slight beneficial effect of β CP in improving disease progression (improved "well-being"), while Preg alone had no significant effect (Figure 1). When measuring survival, both compounds induced a slight decrease in mortality, but none had a statistically significant effect (Figure 1). Next, we asked whether the combined treatment with both compounds can improve the outcome of the mice to LPS injection. Our results show that combining β CP and Preg had a significant positive effect on both disease progression and survival ($p = 0.026$, Figure 2). Indeed, the combined treatment with β CP and pregnenolone significantly improved the wellbeing by >twofold and the survival from a non-significant effect to a >90% increase (Figure 2). These data demonstrate that β CP and Preg have a synergistic effect in alleviating inflammation as well as disease severity and mortality from sepsis *in vivo*.

Following the demonstration of a synergistic effect of β CP and Preg in the resolution of inflammation, we combined these two compounds with seven additional supplements. We designed the resulting formulation to address the symptoms of

long-COVID, related to persistent viral infection, long-lasting inflammation and neurological disorders.

Clinical trial

Clinical study on COVID long hauler patients

At baseline, 2- and 4-week time points, participants recorded the level of severity for each of the 12 symptoms in addition to their subjective assessment of wellbeing ("global reported"). The symptoms in the survey included fatigue, physical weakness, cardiac (e.g., palpitations) and neurological symptoms (e.g., "brain fog"), shortness of breath, gastrointestinal disorders, loss of smell or taste, anxiety, joint pain, rash or hives, cough and insomnia. All the participants took the recommended daily dose of the nutraceutical for 4 weeks. The surveys and scoring by the participants revealed that the severity score for all the symptoms improved already after 2 weeks and this beneficial effect tended to be even more pronounced after 4 weeks (Figure 3). Notably, the decrease in the severity level of all the symptoms was statistically significant at both the 2- and 4-week time points.



Next, we evaluated the number of symptoms that improved for each participant. The vast majority of the participants (46 out of 51) reported more symptoms that improved than symptoms that worsened. Among the remaining five participants, only one reported feeling a worsening in general wellbeing. We also calculated the percentage of participants who reported an improvement, worsening, or no change for each symptom over the 4-week treatment. For each symptom, we omitted any participants who reported a null severity score ("zero") for this symptom at all time points as these do not denote a lack of effect. Notably, 72 to 84% of the participants reported an improvement for each of the 12 symptoms (Figure 4). When asked about their general wellbeing ("global reported"), 59% reported a noticeable improvement. However, when we calculated the average score of all the symptoms for each participant ("global calculated"), we found that the general profile improved for 88% of the participants.

Safety data

Overall safety data was very good, and no major adverse events were noted. Minor adverse events included three patients with vertigo, one patient with increased anxiety, one patient with a gout attack, one patient with increased joint pain and one patient was reinfected with COVID.

Discussion

The results suggest this unique nutraceutical dietary supplement combination may afford significant symptomatic



benefit to long COVID sufferers. The adverse events were minor and overall safety of the nutraceutical product was confirmed.

As the clinical study results indicate, there were statistically significant improvements in this study population in their overall symptoms after 2 and 4 weeks of treatment. It should be noted that different symptoms improved or worsened in different patients. There were some symptoms such as fatigue and brain fog that appeared to respond more than others, however, no baseline presentations were able to predict individual symptomatic responses.

It is unknown at this time if the duration of benefit will be complete, short term, or long term. Because the follow-up period ended after 4 weeks and treatment was taken until the last day, this study was specifically designed to demonstrate a beneficial effect on the long-COVID symptoms. Further studies are warranted to determine the optimal duration of the treatment and whether symptoms would recur upon cessation of the treatment.

We observed a discrepancy between the sum of improvements for each symptom and the reported global wellbeing. Indeed, calculating the average improvement

of all the symptoms for each individual, 88% of the participants benefited from the treatment; however, only 59% of the participants reported an overall improvement of their wellbeing (Figure 4). A similar discrepancy has been reported by others in post-stroke patients where general wellbeing was poorly associated with changes in executive function and comorbidities (53). In our study, we may speculate that not all participants perceived all symptoms at the same level. For example, a person experiencing an improvement in 4 out of 5 symptoms may still consider no improvement in overall wellbeing if the one unchanged symptom has a severe impact on his or her quality of life.

Our *in vivo* experiment describes for the first time the synergistic effect of two different compounds in the attenuation of systemic inflammation. We could find no report on a putative interaction between β CP and Preg. However, a study showing that Preg may act as an allosteric modulator of CB1 (54), a receptor that shares a 44% homology with CB2 (55), may provide circumstantial evidence to a similar interaction between CB2 activation and Preg. In addition to




FIGURE 3 Survey-based scoring by participants at the intake to the trial (baseline), 2 and 4 weeks while taking the supplement. $N = 51$. Dunnett’s multiple comparisons test, $p < 0.0001$ in 2 and 4 weeks for the effect of the supplement on all the symptoms (vs. baseline). **** $p < 0.006$ paired t-test, at 2 weeks vs. baseline and 4 weeks vs. baseline for each symptom.




FIGURE 4 Percentage of participants reporting an improvement, no change or worsening for each symptom at week 4 relative to baseline. “Global reported” is the score given by each participant on general feeling. “Global calculated” is the average of all the scores for each symptom.

β CP and Preg that have potent anti-inflammatory role, we selected nutraceutical compounds that have been tested and proven effective as anti-viral and/or immunomodulatory agents

(DHEA, Bromelain, AKBA, Quercetin, Zinc, and Vitamin D) or in managing neurological dysfunctions (DHEA, St John Wort, and Zinc, see Table 1). Further studies may be

warranted to elucidate the relative contribution of each of the nine compounds included in this nutraceutical to the various symptoms of Long-COVID.

This product is formulated for adults excluding pregnant and breastfeeding women. It is also contra-indicated in combination with serotonin-related antidepressants medications due to the St. John's Wort extract. When indicated, this new nutraceutical is a safe product to be used in combination with other standard of care therapies prescribed for long COVID. We emphasize that the product is not meant to replace standard recommendations of treatment, vaccinations or other suggested methods of prevention or treatment of COVID-19.

In this study, we compared the treatment outcome after 2 and 4 weeks to the same participant's baseline. While there was no placebo in this clinical study, every patient had tried and failed numerous nutraceutical and pharmaceutical products that were available to them along 3–20 months before the trial with no success. In addition, eligibility included persistence of the symptoms for at least 3 months, which reduces the likelihood that the positive effects are coincidental.

The use of nutraceuticals for specific long-COVID symptoms has been previously suggested (56). Here, we developed a nutraceutical for the treatment of a large array of long-COVID symptoms and we demonstrated statistically significant improvement in all tested endpoints in a relatively short time. Within the limitations of this study, our data indicate that the use of this nutraceutical product is a safe and significantly efficient option to reduce multiple symptoms of long COVID. To the best of our knowledge, our population study represents the largest group of patients to have shown statistically significant symptomatic improvement to a nutraceutical formulation.

Patents

Patent pending, International Patent Application No. PCT/IL2022/050676.

Data availability statement

The original contributions presented in the study are included in the article/supplementary material, further inquiries can be directed to the corresponding authors.

Ethics statement

Ethical review and approval was not required for the study on human participants in accordance with the local legislation and institutional requirements. The patients/participants

provided their written informed consent to participate in this study. The animal study was reviewed and approved by the Tel Aviv University Animal Studies Ethics Committee.

Author contributions

NG, JM, and YG: conceptualization, writing – original draft preparation, and project administration. NG and YG: methodology, resources, supervision, and funding acquisition. NG, IK, JS, and YG: validation and investigation. IK, JS, and YG: formal analysis. JS and IK: data curation. IK and YG: writing – review and editing and visualization. All authors reviewed and approved the final version.

Funding

This work was supported by Israel Science Foundation (ISF) Grant No. 1906/21 (animal studies) to YG and by internal research funding (clinical study) from the AARDS (NG).

Acknowledgments

This work was carried out in partial fulfillment of the requirements for a degree for IK from the Sackler Faculty of Medicine, Tel Aviv University, Tel Aviv-Yafo, Israel.

Conflict of interest

The clinical study has been sponsored in part by NTN Enterprises LLC. Authors NG and JM own stock in NTN Enterprises LLC. Ramot, the technology transfer company of Tel Aviv University also owns stock in this company, and YG was an employee of Tel Aviv University. Authors JM and YG are the inventors of patent PCT/IL2022/050676, which exclusive rights has been acquired by NTN Enterprises LLC.

The remaining authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

References

- Gupta A, Madhavan MV, Sehgal K, Nair N, Mahajan S, Sehrawat TS, et al. Extrapulmonary manifestations of COVID-19. *Nat Med.* (2020) 26:1017–32. doi: 10.1038/s41591-020-0968-3
- Taquet M, Dercon Q, Luciano S, Geddes JR, Husain M, Harrison PJ. Incidence, co-occurrence, and evolution of long-COVID features: a 6-month retrospective cohort study of 273,618 survivors of COVID-19. *PLoS Med.* (2021) 18:e1003773. doi: 10.1371/journal.pmed.1003773
- Ballering AV, van Zon SKR, olde Hartman TC, Rosmalen JGM. Persistence of somatic symptoms after COVID-19 in the Netherlands: an observational cohort study. *Lancet.* (2022) 400:452–61. doi: 10.1016/S0140-6736(22)01214-4
- Cutler DM. The Costs of Long COVID. *JAMA Health Forum.* (2022) 3:e221809. doi: 10.1001/jamahealthforum.2022.1809
- Davis HE, Assaf GS, McCorkell L, Wei H, Low RJ, Re'em Y, et al. Characterizing long COVID in an international cohort: 7 months of symptoms and their impact. *EClinicalMedicine.* (2021) 38:101019. doi: 10.1016/j.eclinm.2021.101019
- Noval Rivas M, Porritt RA, Cheng MH, Bahar I, Arditi M. Multisystem inflammatory syndrome in children and long COVID: the SARS-CoV-2 viral superantigen hypothesis. *Front Immunol.* (2022) 13:941009. doi: 10.3389/fimmu.2022.941009
- Watanabe H, Shima S, Mizutani Y, Ueda A, Ito M. [Long COVID: pathogenesis and therapeutic approach]. *Brain Nerve.* (2022) 74:879–84. doi: 10.11477/mf.1416202142
- Dantzer R, O'Connor JC, Freund GG, Johnson RW, Kelley KW. From inflammation to sickness and depression: when the immune system subjugates the brain. *Nat Rev Neurosci.* (2008) 9:46–56. doi: 10.1038/nrn2297
- Liu LD, Duricka DL. Stellate ganglion block reduces symptoms of long COVID: a case series. *J Neuroimmunol.* (2022) 362:577784. doi: 10.1016/j.jneuroim.2021.577784
- Greenhalgh T, Knight M, A'Court C, Buxton M, Husain L. Management of post-acute covid-19 in primary care. *BMJ.* (2020) 370:m3026. doi: 10.1136/bmj.m3026
- Aiyegbusi OL, Hughes SE, Turner G, Rivera SC, McMullan C, Chandan JS, et al. Symptoms, complications and management of long COVID: a review. *J R Soc Med.* (2021) 114:428–42. doi: 10.1177/01410768211032850
- Schmidt C. COVID-19 long haulers. *Nat Biotechnol.* (2021) 39:908–13. doi: 10.1038/s41587-021-00984-7
- Siso-Almirall A, Brito-Zeron P, Conangla Ferrin L, Kostov B, Moragas Moreno A, Mestres J, et al. Long Covid-19: proposed primary care clinical guidelines for diagnosis and disease management. *Int J Environ Res Public Health.* (2021) 18:4350. doi: 10.3390/ijerph18084350
- Chippa V, Aleem A, Anjum F. *Post Acute Coronavirus (COVID-19) Syndrome.* Treasure Island, FL: StatPearls (2022).
- Nagoor Meeran MF, Sharma C, Goyal SN, Kumar S, Ojha S. CB2 receptor-selective agonists as candidates for targeting infection, inflammation, and immunity in SARS-CoV-2 infections. *Drug Dev Res.* (2021) 82:7–11. doi: 10.1002/ddr.21752
- Raphael-Mizrabi B, Attar-Namdar M, Chourasia M, Cascio MG, Shurki A, Tam J, et al. Osteogenic growth peptide is a potent anti-inflammatory and bone preserving hormone via cannabinoid receptor type 2. *Elife.* (2022) 11:e65834. doi: 10.7554/eLife.65834
- Gertsch J, Leonti M, Raduner S, Racz I, Chen JZ, Xie XQ, et al. Beta-caryophyllene is a dietary cannabinoid. *Proc Natl Acad Sci U.S.A.* (2008) 105:9099–104. doi: 10.1073/pnas.0803601105
- Hashiesh HM, Sharma C, Goyal SN, Sadek B, Jha NK, Kaabi JA, et al. A focused review on CB2 receptor-selective pharmacological properties and therapeutic potential of beta-caryophyllene, a dietary cannabinoid. *Biomed Pharmacother.* (2021) 140:111639. doi: 10.1016/j.biopha.2021.111639
- Henderson E, Weinberg M, Wright WA. Pregnenolone. *J Clin Endocrinol Metab.* (1950) 10:455–74. doi: 10.1210/jcem-10-4-455
- Gold R, Schmied M, Tontsch U, Hartung HP, Wekerle H, Toyka KV, et al. Antigen presentation by astrocytes primes rat T lymphocytes for apoptotic cell death. A model for T-cell apoptosis in vivo. *Brain.* (1996) 119 (Pt 2):651–9. doi: 10.1093/brain/119.2.651
- Vacchio MS, Ashwell JD, King LB. A positive role for thymus-derived steroids in formation of the T-cell repertoire. *Ann N Y Acad Sci.* (1998) 840:317–27. doi: 10.1111/j.1749-6632.1998.tb09571.x
- Murugan S, Jakka P, Namani S, Mujumdar V, Radhakrishnan G. The neurosteroid pregnenolone promotes degradation of key proteins in the innate immune signaling to suppress inflammation. *J Biol Chem.* (2019) 294:4596–607. doi: 10.1074/jbc.RA118.005543
- Sun X, Zhang C, Guo H, Chen J, Tao Y, Wang F, et al. Pregnenolone inhibits osteoclast differentiation and protects against lipopolysaccharide-induced inflammatory bone destruction and ovariectomy-induced bone loss. *Front Pharmacol.* (2020) 11:360. doi: 10.3389/fphar.2020.00360
- Ragozzino FJ, Arnold RA, Fenwick AJ, Riley TP, Lindberg JEM, Peterson B, et al. TRPM3 expression and control of glutamate release from primary vagal afferent neurons. *J Neurophysiol.* (2021) 125:199–210. doi: 10.1152/jn.00229.2020
- Andrabi SS, Kaushik P, Mumtaz SM, Alam MM, Tabassum H, Parvez S. Pregnenolone attenuates the ischemia-induced neurological deficit in the transient middle cerebral artery occlusion model of rats. *ACS Omega.* (2022) 7:19122–30. doi: 10.1021/acsomega.1c07016
- Ramirez S, Haddad-Tovoli R, Radosevic M, Toledo M, Pane A, Alcolea D, et al. Hypothalamic pregnenolone mediates recognition memory in the context of metabolic disorders. *Cell Metab.* (2022) 34:269–284.e9. doi: 10.1016/j.cmet.2021.12.023
- Francomano F, Caruso A, Barbarossa A, Fazio A, La Torre C, Ceramella J, et al. β -Caryophyllene: a sesquiterpene with countless biological properties. *Appl Sci.* (2019) 9:5420.
- Scandiffio R, Geddo F, Cottone E, Querio G, Antoniotti S, Gallo MP, et al. Protective effects of (E)-beta-caryophyllene (BCP) in chronic inflammation. *Nutrients.* (2020) 12:3273. doi: 10.3390/nu12113273
- Jha NK, Sharma C, Hashiesh HM, Arunachalam S, Meeran MN, Javed H, et al. beta-caryophyllene, a natural dietary CB2 receptor selective cannabinoid can be a candidate to target the trinity of infection, immunity, and inflammation in COVID-19. *Front Pharmacol.* (2021) 12:590201. doi: 10.3389/fphar.2021.590201
- Yilmaz C, Karali K, Fodelianaki G, Gravanis A, Chavakis T, Charalampopoulos I, et al. Neurosteroids as regulators of neuroinflammation. *Front Neuroendocrinol.* (2019) 55:100788. doi: 10.1016/j.ynfe.2019.100788
- Naylor JC, Kilts JD, Shampine LJ, Parke GJ, Wagner HR, Szabo ST, et al. Effect of pregnenolone vs placebo on self-reported chronic low back pain among US military veterans: a randomized clinical trial. *JAMA Netw Open.* (2020) 3:e200287. doi: 10.1001/jamanetworkopen.2020.0287
- Pinna G. Sex and COVID-19: a protective role for reproductive steroids. *Trends Endocrinol Metab.* (2021) 32:3–6. doi: 10.1016/j.tem.2020.11.004
- Straub RH, Scholmerich J, Zietz B. Replacement therapy with DHEA plus corticosteroids in patients with chronic inflammatory diseases—substitutes of adrenal and sex hormones. *Z Rheumatol.* (2000) 59(Suppl. 2):II/108–18.
- Oberbeck R, Dahlweid M, Koch R, van Griensven M, Emmendorfer A, Tscherne H, et al. Dehydroepiandrosterone decreases mortality rate and improves cellular immune function during polymicrobial sepsis. *Crit Care Med.* (2001) 29:380–4. doi: 10.1097/00003246-200102000-00029
- Alves VB, Basso PJ, Nardini V, Silva A, Chica JE, Cardoso CR. Dehydroepiandrosterone (DHEA) restrains intestinal inflammation by rendering leukocytes hyporesponsive and balancing colitogenic inflammatory responses. *Immunobiology.* (2016) 221:934–43. doi: 10.1016/j.imbio.2016.05.013
- Pavan R, Jain S, Shradha, Kumar A. Properties and therapeutic application of bromelain: a review. *Biotechnol Res Int.* (2012) 2012:976203. doi: 10.1155/2012/976203
- Rathnavelu V, Alitheen NB, Sohila S, Kanagesan S, Ramesh R. Potential role of bromelain in clinical and therapeutic applications. *Biomed Rep.* (2016) 5:283–8. doi: 10.3892/br.2016.720
- Owoyele BV, Bakare AO, Ologe MO. Bromelain: a review on its potential as a therapy for the management of Covid-19. *Niger J Physiol Sci.* (2020) 35:10–9.
- Sagar S, Rathinavel AK, Lutz WE, Struble LR, Khurana S, Schnaubelt AT, et al. Bromelain inhibits SARS-CoV-2 infection via targeting ACE-2, TMPRSS2, and spike protein. *Clin Transl Med.* (2021) 11:e281. doi: 10.1002/ctm.2.281
- Tiradentes RV, Pires JG, Silva NF, Ramage AG, Santuzzi CH, Futuro Neto HA. Effects of acute administration of selective serotonin reuptake inhibitors on sympathetic nerve activity. *Braz J Med Biol Res.* (2014) 47:554–9. doi: 10.1590/1414-431x20143698
- Dani M, Dirksen A, Taraborrelli P, Torocastro M, Panagopoulos D, Sutton R, et al. Autonomic dysfunction in 'long COVID': rationale, physiology and management strategies. *Clin Med (Lond).* (2021) 21:e63–7. doi: 10.7861/clinmed.2020-0896
- Siddiqui MZ. *Boswellia serrata*, a potential antiinflammatory agent: an overview. *Indian J Pharm Sci.* (2011) 73:255–61. doi: 10.4103/0250-474X.93507

43. Gomaa AA, Mohamed HS, Abd-Ellatif RB, Gomaa MA. Boswellic acids/*Boswellia serrata* extract as a potential COVID-19 therapeutic agent in the elderly. *Inflammopharmacology*. (2021) 29:1033–48. doi: 10.1007/s10787-021-00841-8
44. Gulati K, Verma P, Rai N, Ray A. Chapter 7 – role of nutraceuticals in respiratory and allied diseases. In: Gupta RC, Lall R, Srivastava A editors. *Nutraceuticals (Second Edition)*. Cambridge, MA: Academic Press (2021). p. 101–15.
45. Li Y, Yao J, Han C, Yang J, Chaudhry MT, Wang S, et al. Quercetin, inflammation and immunity. *Nutrients*. (2016) 8:167. doi: 10.3390/nu8030167
46. Agrawal PK, Agrawal C, Blunden G. Quercetin: antiviral significance and possible COVID-19 integrative considerations. *Nat Product Commun*. (2020) 15:1934578X20976293. doi: 10.1177/1934578X20976293
47. Saeedi-Boroujeni A, Mahmoudian-Sani MR. Anti-inflammatory potential of Quercetin in COVID-19 treatment. *J Inflamm (Lond)*. (2021) 18:3. doi: 10.1186/s12950-021-00268-6
48. Read SA, Obeid S, Ahlenstiel C, Ahlenstiel G. The role of zinc in antiviral immunity. *Adv Nutr*. (2019) 10:696–710. doi: 10.1093/advances/nmz013
49. Wessels I, Rolles B, Rink L. The potential impact of zinc supplementation on COVID-19 pathogenesis. *Front Immunol*. (2020) 11:1712. doi: 10.3389/fimmu.2020.01712
50. Ghasemian R, Shamshirian A, Heydari K, Malekan M, Alizadeh-Navaei R, Ebrahimzadeh MA, et al. The role of vitamin D in the age of COVID-19: a systematic review and meta-analysis. *Int J Clin Pract*. (2021) 75:e14675. doi: 10.1111/ijcp.14675
51. Vassiliou AG, Jahaj E, Orfanos SE, Dimopoulou I, Kotanidou A. Vitamin D in infectious complications in critically ill patients with or without COVID-19. *Metabol Open*. (2021) 11:100106. doi: 10.1016/j.metop.2021.100106
52. Mai SHC, Sharma N, Kwong AC, Dwivedi DJ, Khan M, Grin PM, et al. Body temperature and mouse scoring systems as surrogate markers of death in cecal ligation and puncture sepsis. *Intensive Care Med Exp*. (2018) 6:20.
53. Zahuranec DB, Skolarus LE, Feng C, Freedman VA, Burke JF. Activity limitations and subjective well-being after stroke. *Neurology*. (2017) 89:944–50. doi: 10.1212/WNL.0000000000004286
54. Vallée M, Vitiello S, Bellocchio L, Hébert-Chatelain E, Monlezun S, Martin-Garcia E, et al. Pregnenolone can protect the brain from cannabis intoxication. *Science*. (2014) 343:94–8. doi: 10.1126/science.1243985
55. Munro S, Thomas KL, Abu-Shaar M. Molecular characterization of a peripheral receptor for cannabinoids. *Nature*. (1993) 365:61–5. doi: 10.1038/365061a0
56. Tosato M, Ciciarello F, Zazzara MB, Pais C, Savera G, Picca A, et al. Nutraceuticals and dietary supplements for older adults with long COVID-19. *Clin Geriatr Med*. (2022) 38:565–91. doi: 10.1016/j.cger.2022.04.004