Check for updates

OPEN ACCESS

APPROVED BY Frontiers Editorial Office, Frontiers Media SA, Switzerland

*CORRESPONDENCE Rina Wu wrn6956@163.com Junrui Wu junrui Wu@126.com

[†]These authors have contributed equally to this work

SPECIALTY SECTION This article was submitted to Food Chemistry, a section of the journal Frontiers in Nutrition

RECEIVED 27 September 2022 ACCEPTED 29 September 2022 PUBLISHED 13 October 2022

CITATION

Li M, Liu Y, Zhang H, Liu Y, Wang W, You S, Hu X, Song M, Wu R and Wu J (2022) Corrigendum: Anti-cancer potential of polysaccharide extracted from *Polygonatum sibiricum* on HepG2 cells *via* cell cycle arrest and apoptosis. *Front. Nutr.* 9:1054565. doi: 10.3389/fnut.2022.1054565

COPYRIGHT

© 2022 Li, Liu, Zhang, Liu, Wang, You, Hu, Song, Wu and Wu. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Corrigendum: Anti-cancer potential of polysaccharide extracted from *Polygonatum sibiricum* on HepG2 cells *via* cell cycle arrest and apoptosis

Mo Li^{1,2†}, Yumeng Liu^{1†}, Henan Zhang¹, Yanfeng Liu¹, Weiming Wang³, Shengbo You⁴, Xinyu Hu¹, Meijun Song¹, Rina Wu^{1*} and Junrui Wu^{1*}

¹College of Food Science, Shenyang Agricultural University, Liaoning Engineering Research Center of Food Fermentation Technology, Shenyang Key Laboratory of Microbial Fermentation Technology Innovation, Shenyang, China, ²College of Criminal Science and Technology, Criminal Investigation Police University of China, Shenyang, China, ³Heilongjiang Academy of Traditional Chinese Medicine, Harbin, China, ⁴Institute of Crop Germplasm Resources, Shandong Academy of Agricultural Sciences, Jinan, China

KEYWORDS

Polygonatum sibiricum polysaccharide, structural characterization, antitumor activity, HepG2 cells, cell apoptosis

A corrigendum on

Anti-cancer potential of polysaccharide extracted from *Polygonatum* sibiricum on HepG2 cells via cell cycle arrest and apoptosis

by Li, M., Liu, Y., Zhang, H., Liu, Y., Wang, W., You, S., Hu, X., Song, M., Wu, R., and Wu, J. (2022). *Front. Nutr.* 9:938290. doi: 10.3389/fnut.2022.938290

In the published article, there was an error in Figure 5B. The effect of PSP-1 ($200 \,\mu$ g/mL) on MMP in Figure 5B was wrongly used. The corrected Figure 5B and its caption appear below.

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.



The morphological changes of HepG2 cells treated with 0, 100, 200, and 400μ g/mL of PSP-1 for 72 h (**A**), the morphological characterization of HepG2 cells was observed and photographed under a Confocal laser scanning microscope (**A1**), nuclear morphological changes induced by PSP-1 in HepG2 cells after DAPI staining. (**A2**) Effect of PSP-1 on MMP (**B**) Cell cycle progression was assessed using propidium iodide staining detected by fluorescence activated cell sorting. (**C**) The apoptotic rates of the indicated cells induced by PSP-1 at different concentrations for 72 h were detected by ANNexin V/PI double-staining asssy (**D**).