



OPEN ACCESS

EDITED BY

Kotsedi Monyeki,
University of Limpopo, South Africa

REVIEWED BY

Fatemeh Mohammadi-Nasrabadi,
National Nutrition and Food Technology
Research Institute, Iran
Ezequiel Pinto,
University of Algarve, Portugal

*CORRESPONDENCE

Machoene Derrick Sekgala
✉ dsekgala@hsr.ac.za

SPECIALTY SECTION

This article was submitted to
Nutritional Epidemiology,
a section of the journal
Frontiers in Nutrition

RECEIVED 30 November 2022

ACCEPTED 27 January 2023

PUBLISHED 23 February 2023

CITATION

Sekgala MD, Opperman M, Mpahleni B and
Mchiza ZJ-R (2023) Sociodemographic and
lifestyle factors and the risk of metabolic
syndrome in taxi drivers: A focus on street food.
Front. Nutr. 10:1112975.
doi: 10.3389/fnut.2023.1112975

COPYRIGHT

© 2023 Sekgala, Opperman, Mpahleni and
Mchiza. This is an open-access article
distributed under the terms of the [Creative
Commons Attribution License \(CC BY\)](#). The use,
distribution or reproduction in other forums is
permitted, provided the original author(s) and
the copyright owner(s) are credited and that
the original publication in this journal is cited, in
accordance with accepted academic practice.
No use, distribution or reproduction is
permitted which does not comply with these
terms.

Sociodemographic and lifestyle factors and the risk of metabolic syndrome in taxi drivers: A focus on street food

Machoene Derrick Sekgala^{1,2*}, Maretha Opperman³,
Buhle Mpahleni³ and Zandile June-Rose Mchiza^{1,4}

¹School of Public Health, University of the Western Cape, Bellville, South Africa, ²Human and Social Capabilities, Human Sciences Research Council, Cape Town, South Africa, ³Functional Food Research Unit, Department of Biotechnology and Consumer Science, Cape Peninsula University of Technology, Cape Town, South Africa, ⁴Non-Communicable Diseases Research Unit, South African Medical Research Council, Cape Town, South Africa

Background: In South Africa, similar to other populous countries, the taxi industry is an important form of transportation that contributes to the country's development. As a result, minibus taxi driving is an occupation characterized by strenuous activities such as long hours of driving, limited rest, and challenges related to securing passengers, among several others. Consequently, to combat stress, some commercial drivers resort to smoking, overeating unhealthy food sold at transportation interchange areas (i.e., taxi ranks), and participating in sedentary behaviors. Most of these activities are risk factors for metabolic syndrome (MetS).

Aim: Therefore, this study aimed to investigate the sociodemographic and lifestyle factors that predispose South African taxi drivers who work in the Cape Town Metropolitan area to the risk of developing MetS.

Methods: This cross-sectional study used a convenient sampling method that included 185 male minibus taxi drivers aged 20 years or above. The participants were interviewed using a validated questionnaire to gather information regarding their sociodemographic characteristics and lifestyle practices. They also underwent physical and metabolic assessments, and the International Diabetes Federation (IDF) criteria were used to diagnose people with MetS.

Results: Overall, the mean age and driving experience of the taxi drivers were 40.0 years (SD: 10.7) and 9.1 years (SD: 7.4), respectively, with those with MetS being significantly older and having more driving experience than those without. Older participants were 3 and 2.9 times more likely to be diagnosed with MetS than the younger participants. Most taxi drivers (70%) met the IDF diagnostic criteria for MetS. Smokers, those who spent more than 100 ZAR (USD 5.9) and those who spent less than 1.4 MET-minutes per week on physical activity were 1.96, 2.0, and 13.6 times more likely to suffer from MetS than those who were nonsmokers, those who spent less than 100 ZAR and those who spent <1.4 MET-minutes per week on physical activity. Consumption of alcohol and sugar-sweetened beverages (SSBs), as well as takeaway and fried foods, snacks, and sold by the SF vendors, increased the likelihood of developing MetS, abnormal HDL-C, TG, and hypertension, while avoiding takeaway and fried foods decreased this likelihood. Taxi drivers who also avoided consuming fresh fruits had abnormal HDL-C.

Conclusion: These findings have significant public health implications, highlighting the need for South African policymakers to adopt a system-level approach to promote lifestyle changes among taxi drivers within the taxi industry.

This can help reduce the health risks faced by these drivers and improve their overall health profile.

KEYWORDS

street food, metabolic syndrome, male taxi drivers, physical activity, socio-economic status, South Africa, waist circumference

Introduction

Several international epidemiological studies have found the prevalence of metabolic diseases to be high among occupational drivers compared to other professionals, such as industrial and office workers (1–3). For example, the majority of professional drivers are at an increased risk of hypertension, myocardial infarction, and hemorrhagic stroke (4). Furthermore, most drivers are in the habit of eating large main meals and consuming snacks (often oily and fried) and fast-food items from street vendors between trips. In addition, many of them resort to alcohol and smoking to overcome stress. It follows logically that they may have an additional risk of developing metabolic diseases. According to Kurosaka et al. (5), taxi driving is characterized by poor eating habits, ongoing stress from driving, and exposure to various health hazards such as air pollution and a lack of physical activity.

In South Africa, taxi drivers and commuters are major consumers of street food (SF) since it is relatively cheap and easily accessible at taxi ranks and bus stations (6, 7). According to Mchiza et al. (8) and Hill et al. (9), the food sold in the streets of Cape Town and surrounding areas seems to be a public health risk since it is energy-dense and high in saturated fat, trans fats, salt, and sugar. Taxi drivers working in these areas may be at risk of developing metabolic syndrome (MetS) as they have been identified to be among the 38% of individuals who consume SF almost daily (6).

Good health is a basic constitutional right for all South African citizens (10). The Occupational Health and Safety Act (OHSA) Section 12(C) (11) requires medical surveillance for all individuals who have high-risk occupations, such as taxi drivers. Similar to other countries, the taxi industry is an important form of transportation in South Africa, contributing to the country's development (12). However, less focus has been given to this industry to ensure that its workers are in good health. To the best of our knowledge, there has never been any health intervention directed at improving the health condition of taxi drivers in South Africa. Substantiated evidence (13–18) suggests that a healthy lifestyle, including healthy eating and regular physical activity, can help to reduce weight, reduce blood pressure, and improve lipid disorders, including raising high-density lipoprotein cholesterol (HDL-C) and lowering triglycerides (TGs). Moreover, unhealthy eating habits and a sedentary lifestyle are known as modifiable risk factors for MetS among taxi drivers (19).

To our knowledge, to date, there are no data on lifestyle and SF consumption in relation to metabolic syndrome (MetS) among minibus taxi drivers in the Western Cape, South Africa. The current study is the first of its kind in South Africa since it investigated the understudied population of minibus taxi drivers, examining their biochemical parameters, sociodemographic characteristics,

and lifestyle practices, with a particular focus on SF consumption and the association of these factors with MetS and its components. The results of this study provide valuable insights for further public health research in this neglected field. Moreover, it will contribute to developing targeted interventions to curb the escalation of MetS in adult male South Africans, especially those working in long-duration driving business.

Materials and methods

Study participants and sampling size

This cross-sectional study was conducted among 185 professional taxi drivers, who were recruited from taxi ranks in Bellville and Cape Town. They were at least 20 years old. This study used a convenient sampling method, and its aim was not to make generalizations about the entire population but rather to focus on taxi drivers who consume SF. These taxi ranks were chosen because they are the two major transport interchange hubs in the Cape Metropolitan Area in South Africa's Western Cape Province. Some of the data used in this study were used in a previous paper (20). The detailed sample size selection, including the power sampling calculation for the current study, is presented elsewhere (20). The participants of this study were full-time minibus taxi drivers, who had been working in this field for at least 1 year and consumed SF at least three times per week. They donated blood samples that were analyzed in a laboratory to diagnose the presence of MetS. We excluded taxi drivers who had a history of non-communicable diseases (NCDs) such as hypertension, kidney failure, hypo- or hyper- thyroidism, liver diseases, known cardiovascular diseases (CVDs), or diabetes mellitus since their eating habits might have been changed based on the advice given by their health practitioners.

Data collection

Data on sociodemographic and lifestyle practices

A previously validated and structured questionnaire developed and validated for use in South Africans aged 15 years and older, which was successfully used in the first South African National Health and Nutrition Examination Survey (SANHANES-1) (21), was administered by a trained researcher to collect data on sociodemographic characteristics (i.e., age, marital status, race, and education level) and lifestyle practices (i.e., physical activity levels, alcohol consumption, and cigarette smoking) from the taxi drivers *via* face-to-face interviews. Moreover, the duration of

sleep, driving experience, and money spent on purchasing SF were assessed using a validated questionnaire used in the study by Hill et al. (6).

The International Physical Activity Questionnaire (IPAQ) (22) was also used to measure the level of physical activity (PA). The results were then based on the calculated physical activity levels (PAL) using the MET-minutes per week criteria. In this case, a sedentary lifestyle was regarded as $PAL < 1.4$ MET-minutes per week, with low being PAL between 1.4 and 1.69 MET-minutes per week, moderate being PAL between 1.7 and 1.9, and vigorous being $PAL \geq 2$ (23).

Frequency of consuming street food

The SANHANES-1 questionnaire (21) was also used to collect information regarding the frequency of consuming street food (FF). The FF list comprised processed meat (i.e., sausages, polony, and cold cuts, such as Viennas, Frankfurters, Russians, and salami); fast food/takeaway foods, including pizzas, fried chicken, fried fish, and burgers, that were packaged to take home; fried meat and fish dishes (i.e., chips, fried chicken, and fried fish) that were consumed on site; deep-fried snacks (i.e., fries/chips, vetkoek, samoosas, and doughnuts), fresh fruits (i.e., all kinds of fruits, excluding fruit juices and dried fruits), sugar-sweetened beverages (SSBs) (i.e., gas/fizzy and reconstituted cold drinks). Consumption frequency for each food item was measured as “none”, “every day”, “1–3 times per week”, and “4–6 times per week”.

Anthropometric measurements

A nonelastic tape was used to measure the waist circumference (WC) at the narrowest point between the lower rib and the upper iliac crest. A cut-off point of ≥ 94 cm was used to determine abnormal WC levels in men (24).

Blood pressure

After the participant had been seated for 5 min or longer, three blood pressure (BP) readings were taken from the right arm in a sitting position using an electronic Micronta monitoring kit (25). Normal systolic BP (SBP) was regarded as a BP that was ≤ 130 mmHg or a diastolic BP (DBP) that was ≤ 85 mmHg (24).

Biochemical parameters

The fasting blood glucose (FBG) was estimated using the capillary method with a glucometer (OneTouch®). To measure biochemical parameters, a venous fasting blood sample was obtained. The plasma lipid profile was used for MetS analysis. The concentration of triglycerides was assessed using the phosphoglycerides oxidase peroxidase method, while the HDL-C was analyzed using the colorimetric non-precipitation method. The IDF criterion was used to diagnose MetS (26). According to the IDF definition, abdominal obesity (i.e., an abnormal WC reading) and two or more of the other four metabolic risk factors are required to diagnose MetS. The cutoff points for the five MetS risk factors are as follows: WC ≥ 94 cm for men; TG ≥ 1.7 mmol/l; SBP ≥ 130 mmHg or DBP ≥ 85 mmHg; FBG ≥ 5.6 mmol/l; and HDL-C < 1.0 mmol/l.

Statistical analysis

Descriptive statistics were used to describe the basic features such as the categories, distribution, and spread of metabolic status, dietary intake, and lifestyle practices using sociodemographic characteristics. In this case, data were analyzed using the analysis of variance (ANOVA) and the Kruskal–Wallis tests and presented as frequencies, means, medians, and standard deviations, depending on whether they were categorical or continuous. The associations between different variables were analyzed using the Chi-square test. A binary logistic regression analysis was conducted to examine the odds ratios (OR). Multivariate analyses using multiple logistic regression models, which incorporated all risk factors for MetS while adjusting for the effect of possible confounders such as age, employment status, marital status, ethnicity, physical activity, and monthly income, were also applied (AOR). 95% confidence intervals (CIs) that did not overlap and p -values that were less than 0.05 indicated significant differences and associations between variable results. All data were analyzed using the statistical package for social sciences (SPSS version 28.0 for Windows; SPSS Inc., Chicago, IL, USA).

Results

Table 1 presents the sociodemographic characteristics of the study participants based on their MetS status. Overall, the mean age of the participants was 40.0 years (SD: 10.7), with those suffering from MetS being significantly older than those without. There were 10.2% more participants within the age group of 20–39 years. There was a significantly higher prevalence of participants with MetS in the older age group than in the younger age group (61 vs. 39%).

While there were no other significant differences in sociodemographic characteristics in relation to MetS in this cohort, the mean driving experience of the participants was 9.1 years (SD: 7.4). In this case, the participants who presented with MetS had significantly higher driving experience compared to those without. There was also a significantly higher prevalence of participants with MetS who had a driving experience of 8 years or more compared to those with a driving experience of one to seven years (59.7 vs. 40.3%).

Table 2 shows that older participants were 3 and 2.9 times more likely to have MetS than younger participants. While the significant association of MetS with age was unavailable when the data were adjusted for lifestyle practices (i.e., cigarette smoking, alcohol consumption, sleeping duration, physical activity level, and money spent on SF each day), it was available for age after we removed the confounding effects of the other sociodemographic variables explored in the current analysis.

When examining the components of MetS (Table 3), the overall mean values for WC, TG, HDL-C, SBP, DBP, and FBG were 99.1 (SD: 18.3), 1.3 (SD: 1.1), 1.1 (SD: 0.3), 133.4 (SD: 17.2), 84.8 (SD: 13.2), and 6.4 (SD: 3.5), respectively. We also observed that there were many participants with abnormal WC (59.5%), HDL-C (51.4%), and SBP (58.4%). However, there were few participants with abnormal TG (20.5%) and DBP (43.2%). The participants with MetS had significantly higher abnormal WC (64.5 vs. 35.5%), TG (81.6 vs. 18.4%), HDL-C (61.1 vs. 38.9%), SBP (50.9 vs. 49.1%),

TABLE 1 The distribution of sociodemographic characteristics of South African minibus taxi drivers by metabolic syndrome status.

	N = 185	IDF MetS		P value
		No (n = 108)	Yes (n = 77)	
Age_(years)				
M ± SD	40.0 ± 10.7	37.3 ± 10.2	43.7 ± 10.3	<0.001
N(%)				
20-39	102 (55.1)	72 (66.7)	30 (39.0)	
≥40	83 (44.9)	36 (33.3)	47 (61.0)	<0.001
Ethnicity				
N(%)				
Black	146 (78.9)	85 (78.7)	61 (79.2)	
Non-Black	39 (21.1)	23 (21.3)	16 (20.8)	0.932
Level of Education N (%)				
No schooling/primary education	58 (31.4)	35 (32.4)	23 (29.9)	
Some high school/high education	127 (68.6)	73 (67.6)	54 (70.1)	0.714
Marital status				
N (%)				
Single/separated/divorced	97 (52.4)	60 (55.6)	37 (48.1)	
Married/living as married	88 (47.6)	48 (44.4)	40 (51.9)	0.314
Driving experience (years)				
M ± SD	9.1 ± 7.4	7.2 ± 6.1	11.7 ± 8.4	<0.001
N(%)				
1-7	103 (55.1)	72 (66.7)	31 (40.3)	
≥8	82 (44.3)	36 (33.3)	45 (59.7)	<0.001

IDF, International Diabetes Federation; MetS, metabolic syndrome.

DBP (67.5 vs. 32.5%), and FBG (64.1 vs. 35.9%) compared to those without.

Seventy-seven ($n = 77$) study participants met the IDF diagnostic criteria for MetS (i.e., had a clustering of 3 or more metabolic disorders), of which 46 had three (3) risk factors, 25 had four (4) risk factors, and 6 had five (5) risk factors. The distribution is shown in Table 4.

Table 5 presents the lifestyle practices based on the outcomes of MetS. Overall, the participants smoked an average of almost 10 cigarettes (SD: 5.3) a day, slept an average of 6.1 h (SD: 1.1) each day, spent an average of ZARR 92.1 (exchange rate: ZARR 1 = United States Dollar [USD]\$ 17.23) (SD: 36.7) on SF each day, and had an average PAL of 1.42 MET-minutes per week (SD: 0.14). While there were no significant differences regarding the average number of cigarettes smoked by the participants or the average amount of money spent on SF between those who had MetS and those without MetS.

In terms of participant lifestyle distribution based on the MetS status, while there were no significant differences between participants with and without MetS for lifestyle practices such as alcohol consumption and sleeping duration, there were significantly higher number of nonsmokers who were positive for MetS (those who gave an affirmative response for smoking)

and those who were negative for MetS (those who gave a negative response for smoking). There were also significantly more participants with MetS who spent ZARR 100 or more than those who spent less than 100 ZAR (57.1% vs. 42.9%, $p=0.022$). Finally, there were significantly more sedentary participants with MetS compared to those with low and moderate PAL (79.2% vs. 14.3% and 6.5%).

According to Table 6, smokers, those who spent ZARR 100 or more and those who spent <1.4 MET-minute/week were 1.96, 2.0, and 13.6 times significantly more likely to suffer from MetS compared to those who did not smoke, those who spent less than ZARR 100, and those who spent 1.4 or more MET-minute/week. While the increased significant likelihood of MetS for sedentary activity remained, even after removing the confounding effects of sociodemographic characteristics and other lifestyle practices explored in the current study, the likelihood of smoking and the amount spent on SF disappeared. It is also important to note that removing the confounding effects of the other lifestyle facts of the participants resulted in an increased significant likelihood for developing MetS by 2.2 and 2.1 times for those who consumed alcohol and those who slept 7 h or more, respectively.

The frequency of SF consumption in relation to the likelihood of developing MetS and its components was also analyzed and is presented in Supplementary Table 1. Approximately 40.0% of the

TABLE 2 A binary logistic regression analysis to show the association between the sociodemographic characteristics and the MetS status of South African minibus taxi drivers.

	IDF Metabolic syndrome											
				Model 1			Model 2			Model 3		
	Crude	95% CI	<i>P</i> value	AOR	95% CI	<i>P</i> value	AOR	95% CI	<i>P</i> value	AOR	95% CI	<i>P</i> value
Age (years)												
20–39	1			1			1			1		
≥40	3.133	1.706–5.756	<0.001	0.541	0.217–1.351	0.188	0.589	0.225–1.539	0.280	2.955	1.2955–6.969	0.013
Ethnicity												
Black	1			1			1			1		
Non-Black	0.969	0.473–1.987	0.932	0.908	0.356–2.312	0.839	0.991	0.378–2.597	0.986	0.560	0.245–1.280	0.169
Level of Education												
No schooling/primary education	1			1			1			1		
Some high school/high education	1.126	0.598–2.120	0.714	2.880	1.212–6.847	0.017	2.676	1.103–6.506	0.030	2.004	0.940–4.273	0.072
Marital status												
Single/separated/divorced	1			1			1			1		
Married/living as married	1.351	0.752–2.429	0.314	0.557	0.254–1.219	0.143	0.639	0.282–1.450	0.284	0.893	0.449–1.778	0.748

OR, odds ratio; AOR, adjusted odds ratio. Model 1: adjusted for Cigarette smoking, Physical Activity Level and Money spent on Street Food each day; Model 2: adjusted for Cigarettes smoking, Alcohol drinking, sleeping duration, Physical Activity Level and Money spent on Street Food each day; Model 3: adjusted for age ethnicity, level of education, marital status and driving experience.

TABLE 3 The distribution of South African minibus taxi drivers by components of MetS.

	Entire cohort (n = 185)		IDF MetS				P value "between groups"
	Mean ± SD	n (%)	No MetS (77)		With MetS (108)		
			Mean ± SD	n (%)	Mean ± SD	n (%)	
Waist circumference (cm)	99.1 ± 18.3		90.7 (14.5)		110.8(16.7)		<0.001
Normal <94		75 (40.5)		69 (92.0)		6(8.0)	
Abnormal ≥94		110 (59.5)		39 (35.5)		71(64.5)	
Triglycerides (mmol/l)	1.3 ± 1.1		1.0 (0.4)		1.9(1.5)		<0.001
Normal <1.7		147 (79.5)		101 (68.7)		46(31.3)	
Abnormal ≥1.7		38 (20.5)		7 (18.4)		31(81.6)	
HDL-C (mmol/l)	1.1 ± 0.3		1.2 (0.4)		1.0(0.3)		<0.001
Normal ≥1.0		90 (48.6)		71 (78.9)		19(21.1)	
Abnormal <1.0		95 (51.4)		37 (38.9)		58(61.1)	
Systolic blood pressure (mmHg)	133.4 ± 17.2		127 (13.3)		141(18.8)		0.002
Normal <130		77 (41.6)		55 (71.4)		22(28.6)	
Abnormal ≥130		108 (58.4)		53 (49.1)		55(50.9)	
Diastolic blood pressure (mmHg)	84.8 ± 13.2		79 (9.1)		92.7(13.9)		<0.001
Normal <85		105 (56.8)		82 (78.1)		23(21.8)	
Abnormal ≥85		80 (43.2)		26 (32.5)		54(67.5)	
Fasting blood Glucose (mmol/l)	6.4 ± 3.5		5.3 (1.1)		7.9(4.8)		<0.001
Normal <5.5		93 (50.3)		75 (80.6)		18(19.4)	
Abnormal ≥5.5		92 (49.7)		33 (35.9)		59(64.1)	

HDL-C, high density lipoprotein cholesterol; IDF, international diabetes federation; MetS, metabolic syndrome.

TABLE 4 The distribution of South African minibus taxi drivers by the clustering of MetS components.

Number of metabolic disorders	n (%)
0	22 (11.9)
1	40 (21.6)
2	46 (24.9)
3	46 (24.9)
4	25 (13.5)
5	6 (3.2)

Metabolic disorders (abnormal WC, TG, SBP, DBP, HDL-C and FBG).

entire population consumed processed meat (sausages, polony, and cold cuts Viennas, Frankfurters, Russians, and salami) at least 1–3 times a week, with a significantly higher proportion (44.4%) of them experiencing abnormal BP compared to those with normal BP (37.5%). Similar results were also observed for the participants who consumed takeaway foods. Moreover, a higher proportion of participants with MetS and hypertension consumed fried food and snacks (i.e., chips, vetkoek, fried chicken, fried fish) compared to those who did not consume these foods. The daily consumption of deep-fried foods was also associated with an abnormal WC.

As illustrated in [Table 7](#), consuming processed meat daily increases the risk of abnormal HDL-C by 3.7 times, while avoiding processed meat reduces hypertension. Further, avoiding takeout reduced the likelihood of developing MetS by 68.2% and abnormal TG by 25%. Daily takeout meal consumption increased hypertension risk by 3.1 times, even after adjusting for age and sociodemographic characteristics. The daily consumption of fried meat and fish increased the likelihood of developing MetS, abnormal WC, and hypertension by 2.1, 2.2, and 2.3 times, respectively. The association remained unchanged, even after removing the confounding effects of age, ethnicity, money spent on buying these foods, sociodemographic characteristic, and unhealthy lifestyle practices.

Moreover, the consumption of these foods 1–3 times a week increased the likelihood of developing abnormal HDL-C by 2.5 times, and this interaction also remained unchanged, even after removing all the confounding effects. The daily consumption of fried snacks also increased the likelihood of developing MetS, abnormal WC, abnormal HDL-C, and hypertension by 3.8, 1.7, 2.3, and 1.9 times, respectively. The consumption of packaged snacks such as crisps and amazimba (Niknaks Maize Snack) every day also increased the likelihood of abnormal HDL-C by eight times. Moreover, consuming these snacks 1–3 times a week increased the likelihood of developing MetS by 4.1 times. These interactions remained unchanged even after removing

TABLE 5 The distribution of the South African minibus taxi drivers by their lifestyle risk factors (i.e., cigarettes smoking, alcohol consumption, physical activity level, sleep duration and money spent on street food each day) and MetS.

	Entire cohort (n = 185) n (%)	IDF metabolic syndrome		P value n (%)
		No (n = 108) n (%)	Yes (n = 77) n (%)	
Cigarettes smoking				
Yes	80 (43.2)	54 (50.0)	26 (33.8)	0.028
No	105 (56.8)	54 (50.0)	51 (66.2)	
Average number of cigarettes smoked each day				
M ± SD	9.9 ± 5.3	9.3 ± 4.9	11.0 ± 5.9	0.179
1–5	24 (30.0)	17 (31.5)	7 (26.9)	0.793
6–9	7 (8.8)	4 (7.4)	3 (11.5)	
≥10	49 (61.3)	33 (61.1)	16 (61.5)	
Current alcohol drinking				
Yes	100 (54.1)	55 (50.9)	45 (58.4)	0.312
No	85 (45.9)	53 (49.1)	32 (41.6)	
Frequency of alcoholic beverage consumption				
Monthly or less	52 (36.9)	31 (36.9)	24 (37.0)	0.248
2–4 time a month	51 (34.2)	30 (35.7)	21 (32.3)	
2–3 times a week	28 (18.8)	12 (14.3)	16 (24.6)	
4 or more time a week	15 (10.1)	11 (13.1)	4 (6.2)	
Number of alcoholic beverages consumed on a typical day				
1 or 2	9 (9.0)	4 (7.3)	5 (11.1)	0.906
3 or 4	9 (9.0)	4 (7.3)	5 (11.1)	
5 or 6	59 (59.0)	34 (61.8)	25 (55.6)	
7,8 or 9	18 (18.0)	10 (18.2)	8 (17.8)	
10 or more	5 (5.0)	2 (5.5)	2 (4.4)	
Sleeping duration (hours)				
M ± SD	6.1 ± 1.1	6.1 ± 1.0	6.2 ± 1.2	0.624
n (%)				
<6	112 (60.5)	70 (64.8)	42 (54.5)	
≥7	73 (39.5)	38 (35.2)	35 (45.5)	0.159
Money spend on street food each day (ZAR)				
M ± SD	92.1 ± 36.7	88.6 ± 37.7	96.9 ± 34.9	0.056
N (%)				
<R99.00	96 (52.7)	63 (60.0)	33 (42.9)	
≥R100.00	86 (47.5)	42 (40.0)	44 (57.1)	0.022
Physical activity level (MET-minutes per week)				
N ± SD	1.42 ± 0.14	1.35 ± 0.12	1.51 ± 0.10	<0.001
Sedentary PAL				
<1.4	94 (50.8)	33 (30.6)	61 (79.2)	
Low PAL				
1.4–1.69	86 (46.5)	75 (69.4)	11 (14.3)	
Moderate PAL				
1.70–1.99	5 (2.7)	-	5 (6.5)	<0.001
Vigorous PAL				
≥2	-	-	-	

IDF, International Diabetes Federation; ZAR, South African Rand (exchange rate 17.23 United States Dollar); PAL, physical activity level.

TABLE 6 Binary logistic regression analysis to show the association between the lifestyle factors and MetS status of the South African minibus taxi drivers.

	IDF—metabolic syndrome											
	Crude			Model 1			Model 2			Model 3		
	OR	95% CI	<i>P</i> value	AOR	95% CI	<i>P</i> value	AOR	95% CI	<i>P</i> value	AOR	95% CI	<i>P</i> value
Current cigarettes smoking												
No	1	1.072–3.590	0.029	1	0.301–1.063	0.077	1	0.318–1.142	0.120	1	0.270–1.208	0.143
Yes	1.962			0.566			0.602			0.571		
Alcohol consumption												
No	1			1			1			1		
Yes	1.355	0.751–2.444	0.312	1.776	0.935–3.374	0.079	1.706	0.886–3.287	0.110	2.191	1.021–4.699	0.044
Sleeping duration												
<6	1			1			1			1		
≥7	1.535	0.844–2.791	0.160	1.558	0.832–2.915	0.166	1.497	0.786–2.851	0.220	2.107	0.977–4.521	0.057
Money spends on street food each day (ZAR)												
<R99.00	1			1			1			1		
≥R100.00	2.000	1.101–3.633	0.023	1.225	0.619–2.423	0.560	1.400	0.691–2.838	0.350	1.157	0.548–2.441	0.702
PAL (MET—minutes per week)												
Active (low, moderate and vigorous) PAL≥1.4	1	6.388–29.109	<0.001	1	5.449–29.354	<0.001	1	5.769–32.780	<0.001	1	6.769–36.891	<0.001
Sedentary (<1.4)	13.636			12.647			13.751			15.802		

IDF, International Diabetes Federation; ZAR, South African Rand (exchange rate 17.23 United States Dollar); PAL, physical activity level; OR, odds ratio; AOR, adjusted odds ratio = Model 1: adjusted for age and driving experience; Model 2: adjusted for age, ethnicity, level of education, marital status and driving experience; Model 3: adjusted for tobacco smoking, alcohol consumption, sleeping duration, PAL and money spent on street food each day.

TABLE 7 The logistic regression analysis to show the association between the street foods consumed by the South African minibus taxi drivers and their MetS status.

Frequency of food consumption	MetS		Abnormal WC		Abnormal FBG		Abnormal HDL-C		Hypertension		Abnormal triglyceride	
	OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI
Processed meat (sausages, polony, and cold cuts Viennas, Frankfurters, Russians, salami)												
None	0.963	0.416–2.227	1.024	0.477–2.199	0.842	0.390–1.819	1.056	0.477–2.376	0.413*	0.182–0.937	1.165	0.459–2.957
Every day	1.143	0.318–4.109	0.594 [§]	0.208–1.702	0.605 [§]	0.205–1.783	3.667 ^{†*#°@S^}	4.881–15.264	1.353	0.461–3.975	0.879	0.200–3.859
1–3 times last week	0.830	0.366–1.882	1.007	0.481–2.107	0.756	0.357–1.599	0.850	0.388–1.860	0.831	0.390–1.768	0.563	0.210–1.506
4–6 times last week	1		1		1		1		1		1	
Takeaway food (pizza, burgers, chicken and fish parcels)												
None	0.318*	0.115–0.878	1.618	0.266–9.852	0.865	0.163–4.602	1.250	0.173–9.019	0.767	0.075–7.860	0.750 ^{†*#°@S}	1.574–67.602
Every day	1.227	0.263–5.734	1.014	0.448–2.298	1.300	0.583–2.899	0.833	0.338–2.052	3.097 ^{†*#S}	1.123–8.544	3.055	0.680–13.729
1–3 times last week	0.738	0.275–1.981	0.518	0.177–1.514	1.615	0.538–4.853	0.917	0.276–3.040	1.533	0.402–5.841	1.853	0.277–12.389
4–6 times last week	1		1		1		1		1		1	
Fried meat and fish dishes (chips, vetkoek, fried chicken, fried fish)												
None	1.778	0.236–13.405	1.200 [§]	0.471–3.056	1.059	0.406–2.762	1.128	0.393–3.236	1.310	0.465–3.684	2.000	0.533–7.508
Every day	2.051 ^{†*#°@S^}	1.757–5.558	2.223 ^{†*#°@^}	1.007–4.904	1.310	0.595–2.882	1.974	0.851–4.580	2.278*	1.977–5.312	2.082	0.696–6.227
1–3 times last week	1.697	0.447–6.439	1.079	0.850	1.116	0.501–2.486	2.538 ^{†*#°@S^}	1.089–5.918	0.776	0.314–1.916	1.358	0.437–4.228
4–6 times last week	1		1		1		1		1		1	
Fried snacks (vetkoek, samoosas, doughnuts)												
None	1.786	0.550–5.802	0.457	0.191–1.096	1.315	0.551–3.138	1.875	0.678–5.182	0.483	0.180–1.301	0.520	0.139–1.945
Every day	3.772 ^{†*#°@}	1.479–9.616	1.760 ^{†*#°}	0.693–4.470	1.004	0.431–2.342	2.333 ^{†*#}	2.880–6.188	2.194*	1.927–5.191	1.354	0.459–3.998
1–3 times last week	1.875	0.730–4.816	0.656	0.320–1.342	0.994	0.494–2.001	2.035	1.898–4.611	0.842	0.403–1.703	0.780	0.303–2.006
4–6 times last week	1		1		1		1		1		1	
Packaged snacks (chips/crisps, mazimba)												
None	3.036	0.593–15.547	0.606	0.186–1.977	0.643	0.201–2.052	3.354	0.716–18.637	0.381	0.117–1.241	3.056	0.351–26.593
every day	3.125	0.474–20.583	0.273	0.063–1.178	1.111	0.262–4.719	8.000 ^{†*#°@S^}	1.215–52.693	0.889	0.216–3.662	3.300	0.294–37.103
1–3 times last week	4.113 ^{†*#°@^}	1.862–19.626	0.739	0.244–2.237	0.705	0.239–2.086	6.538 [§]	1.373–31.132	0.660	0.226–1.926	3.075	0.379–24.933
4–6 times last week	1		1		1		1		1		1	
Sugar Sweetened beverages (gas/fizzy cold drink and reconstituted)												
None	0.545	0.133–2.236	0.541	0.186–1.577	0.923	0.317–2.685	0.426	0.127–1.427	0.500	0.127–1.965	0.967	0.229–4.087
Every day	1.660*	1.796–3.461	1.164	0.608–2.228	1.620	0.850–3.088	0.833	0.416–1.668	1.691	0.848–3.372	0.767	0.324–1.812
1–3 times last week	1.889	0.772–4.619	0.854	0.383–1.904	0.997	0.444–2.235	1.193	0.502–2.833	1.442	0.610–3.409	1.364	0.507–3.669
4–6 times last week	1		1		1		1		1		1	

FF, food frequencies; OR: *p < 0.05; #Model 1= AOR, p < 0.05 adjusted for age; °Model 2 = AOR, p < 0.05 adjusted for ethnicity; @Model 3 = AOR, p < 0.05 adjusted for money spent on street food each day; §Model 4 = AOR, p < 0.05 adjusted for all socio demographic status, ^Model 5 = AOR, p < 0.05 adjusted for all lifestyle factors.

the confounding effects such as age, ethnicity, money spent on these foods, and sociodemographic status, and unhealthy lifestyle practices. The daily consumption of SSB increased the likelihood of developing MetS by 1.6 times. However, this interaction disappeared after removing the confounding effects such as age, ethnicity, money spent on these foods, sociodemographic characteristics, and unhealthy lifestyle practices.

Discussion

The current study aimed to investigate the risk factors for MetS among the male minibus taxi drivers working in Cape Town and the surrounding areas. The majority of the taxi drivers had abnormal levels of WC, HDL-C, SBP, and FBG. Approximately 70% of the taxi drivers had clusters of three or more of these health issues. These results are corroborated by both national and international literature that show that individuals who are in the long-duration driving occupation, including taxi drivers, have a high likelihood of developing metabolic disorders compared to other professionals such as industrial and office workers (1–3, 27–30). In addition, these studies also identified age, driving duration, and driving experience as factors that accelerate the onset of these metabolic diseases (31–34). More importantly, Hildrum et al. (35) long argued that this condition strongly increases with age, regardless of any algorithm used to measure MetS. Indeed, in the current study, the mean age of the minibus taxi drivers was 40 years, with older participants having more driving experience compared to their younger counterparts. Even though the current analysis did not demonstrate a significant relationship between driving experience and MetS, it showed that old age increased the likelihood of developing MetS by up to 2.95 times.

Moreover, like in the current study, the majority of occupational drivers involved in other studies (28, 36) reported sleeping hours that are less than the recommended 6 h of sleep each day (37). This may be due to these drivers' long and irregular shift hours (28, 31). Unlike the aforementioned international researchers who reported sleep duration and its quality as the determinants of MetS, in the current study, the association between sleeping duration and MetS was significant. However, it is important to note that the majority of the minibus taxi drivers participating in the current research also reported long working hours such that their daily shifts started as early as 5 am most days and sometimes ended after 10 pm. They cite reasons such as the need to secure passengers who start work early and those who knock off late in the evenings from work because of their long working hours.

In the current study, despite no significant differences observed in the number of cigarettes smoked by those who had MetS and those without MetS, on average, the minibus taxi drivers smoked almost 10 cigarettes each day, and smoking increased their likelihood of developing MetS by up to two times. However, this interaction disappeared after we removed the confounding effects, such as sociodemography and other lifestyle practices investigated in the current study. Therefore, this suggests that factors such as age, ethnicity, the number of cigarettes smoked, and so on moderate propensity of smoking and the likelihood of developing

MetS. Additionally, while some of the literature (32, 33) could not establish a relationship between smoking and the likelihood of developing MetS among occupational drivers, Appiah et al. (38) found that non-users of tobacco are less likely to suffer from MetS and its components. Mohebbi et al. (31) also showed that smokers are more likely to suffer from with MetS than nonsmokers. It is also important to explain the differences in the results regarding smokers between the current study and a recent study by Mabetwa et al. (32), which was also conducted for South African taxi drivers. Mabetwa et al.'s (32) study was conducted in the Gauteng province, while the current study was conducted in the Western Cape province. According to Statistics South Africa (39), Cape Town has the highest concentration of male smokers in South Africa. Additionally, it is commonly reported that smokers often smoke in public places, increasing the likelihood of exposure to secondhand smoke for nonsmokers. Therefore, we observed a high prevalence of MetS among nonsmokers in the current study. Moreover, it is important to highlight that the prevalence of smokers in the current study was higher than that of smokers reported in Mabetwa's study (43 vs. 30%).

In the current study, we also found results suggesting that minibus taxi drivers with a sedentary lifestyle had a 13-fold increased risk of developing MetS. This relationship remained strong even after removing the confounding effects such as sociodemography and other lifestyle factors investigated in the current study. This study indicates that physical activity has an independent and significant impact on metabolic health independently, regardless of other social determinants of health. These results are corroborated by substantiated evidence from international studies (40, 41) suggesting a significant negative correlation between physical activity and the likelihood of developing MetS among bus and taxi drivers. Moreover, Chen et al. (1) showed that sedentary occupations, including taxi driving, increase the risk of developing MetS. Several international studies have shown the dose–response relationship between physical activity and metabolic outcomes (13–18). According to Myers et al. (42), most active individuals generally have a low risk of developing metabolic diseases. Additionally, the aforementioned studies found that even meeting the minimal physical activity requirements outlined in the health guidelines (14) (i.e., at least 150 min per week of moderate-intensity activity or 75 min per week of vigorous activities) has significant benefits for reducing metabolic risk. However, we also have to acknowledge that, in our analysis, even though the participants who suffered from MetS had a higher PAL than those without MetS, their activity levels were still within the low PAL range (i.e., they were within the range of 1.4 and 1.7 Met-minutes per week). Thus, the average 1.51 MET-min per week dosage they obtained could not improve their metabolic health. We also must acknowledge that other studies could not find a significant association between physical activity and MetS (32, 38). The reason for this is currently unknown and needs further investigation.

Other interesting results from the current research were that the type and quality of food and beverages consumed by minibus taxi drivers impacted their metabolic health. For instance, when the confounding effects of other lifestyle factors were removed from the current study, alcohol consumption increased the risk of MetS by

up to two times. Even though we did not measure the exact amount of alcohol consumed by the minibus taxi drivers participating in our research, the majority reported that they consumed alcoholic beverages that ranged from 5 to 9 standard drinks most days. This is a cause for concern given that studies by Hernández-Rubio et al. (43) and Fan et al. (44) found that heavy drinking is independently associated with reduced kidney function and metabolic risk factors such as impaired fasting glucose/diabetes mellitus, abdominal obesity, arterial stiffness and plaque buildup, hypertension, and dyslipidemia. In the current analysis, we also found that the consumption of takeaway foods, fried foods, and snacks such as crisps and SSB sold by the SF vendors increased the likelihood of developing MetS, abnormal HDL-C, TG, and hypertension. We also found that avoiding takeaway and fried foods decreased the likelihood of MetS.

International research by Kim and Je (45) corroborates our finding in that individuals with MetS generally consume large quantities of processed meat (such as sausages, polony, and cold cuts such as Viennas, Frankfurters, Russians, and salami). Furthermore, the aforementioned study also found that participants in the highest category of total meat, red meat, and processed meat consumption had an increased risk of developing MetS by approximately 14, 33, and 35%, respectively, compared to those in the lowest consumption category of these foods. A meta-analysis of studies (46, 47) revealed a strong correlation between the consumption of red meat and the likelihood of developing MetS after excluding studies from Asia. For instance, Pan et al. (48) found that even a slight increase in the daily consumption of red and processed meat had a 14% and 32% increase in the likelihood of type 2 diabetes mellitus, respectively. Abete et al. (49) also found high rates of mortality due to metabolic disorders in populations with high consumption of processed meat.

Some potential mechanisms have been explained to indicate the association between processed meat consumption and the likelihood of developing MetS. Among these are the findings that total and saturated fat contained in processed meat increase the risk of MetS through increased body fat centralization, hyperinsulinemia, and hyperglycemia, which are important components of MetS (50). According to Abete et al. (49), the aforementioned mechanism is mediated by nitrosamines. This chemical is toxic to pancreatic cells formed from the nitrates used as preservatives in processed meat. Additionally, these compounds cause insulin resistance.

Moreover, Marku et al. (51) argue that because iron is a strong pro-oxidant, it causes oxidative stress, which can harm tissues such as pancreatic beta cells. Furthermore, the aforementioned researchers argue that high iron levels may inhibit glucose metabolism and reduce pancreatic insulin synthesis and secretion. Based on the literature, we must also acknowledge that high levels of inflammatory mediators, such as C-reactive protein, in people who consume a high amount of red and processed meat could be another reason for the increased risk of MetS. Because C-reactive proteins also increase blood pressure (52), this could explain the association we found in the current research between the consumption of processed meat and hypertension. Griep et al. (53) reported similar results that suggest high consumption of processed meat is positively associated with the risk of hypertension. Another

possible explanation for our findings may be those given by Micha et al. (54), who suggest that the high sodium content of processed meat results in elevated blood pressure.

Our current study further found the association between MetS risk and high consumption of fried food bought from street vendors and consumed on-site (i.e., fries/chips, vetkoek, fried chicken, and fried fish, to be specific). Our results were unsurprising given that the food sold on South African streets, including at the transport interchange areas where we recruited our participants, is not healthy. Additionally, Mchiza et al. (8) and Flores et al. (55) showed that, besides fruits and vegetables, most of the SF sold by vendors are not healthy as they are deep-fried, which is associated with cardiovascular risks. However, we must acknowledge that not all researchers have found associations between fried food and the risk of MetS. For instance, upon investigating a Mediterranean cohort of young Asian adults, Sayon-Orea et al. (56) and Kang and Kim (57) found no association between MetS and the frequency of consuming fried foods. The differences between the results of our study and those of the aforementioned Asian studies could be based on the type of food groups included in the current study and the two Asian studies; among the four groups of fried food included in the Asian studies were fried vegetables, fried fish, and fried seaweed. Therefore, we must always be cognizant of the literature that associates plant foods and fish with preventing metabolic diseases (58, 59). In the current study, on the other hand, the four groups of fried food were deep-fried potato chips (or French fries), vetkoek (a cake of deep-fried dough that is stuffed inside), fried chicken, and fried fish. In this case, fried vegetables and seaweeds impact health differently than fried potatoes and fried starch. Finally, the frying mechanisms in these studies were also different. In Asia, pan frying is mostly preferred, while deep frying is favored in South Africa, and these cooking methods have also been shown to impact health differently (60, 61).

This study's results found a statistically significant association between fast-food consumption and MetS risk. These results are consistent with those of Bahadoran et al. (62), where they showed evidence that regular fast-food consumption has a detrimental effect on general health and can increase the risk of obesity, insulin resistance, and other metabolic abnormalities. Several mechanisms have been proposed to explain the negative effects fast foods have on health outcomes. One such mechanism is that fast foods are energy-dense, thus modulating the weight gain process (63). Indeed, Mchiza et al. (8) showed that most fast foods sold in the streets of South Africa are energy dense and have an energy density that is almost two times the recommended energy for a healthy meal. Moreover, the mean total energy of these meals is estimated to be approximately 158–163 kcal per 100 g of food, with the total fat percent of beef hamburgers, chips, chicken hamburgers, and hot dogs being reported to be about 35.8 ± 10.7 , 35.8 ± 8.7 , 23.0 ± 5.1 , and $34.0 \pm 13.5\%$, respectively, with most of this fat being saturated fat (64).

The current study also showed that consuming packaged snacks (chips/crisps and mazimba) 1–3 times a day was associated with an increased risk of developing MetS. In agreement with this study's results, a significant relationship was also shown between dyslipidemia and the frequency of consumption of hydrogenated

fat, fast foods, cheese puffs, and crisps in both urban and rural areas of Iran (65).

The current analysis also showed that the consumption of SSBs increased the risk of MetS by up to 1.8 times. Consistent with this study's are a few international studies (66, 67) that reported that SSB intake has significant effects on MetS risk. Moreover, a study conducted on 596 young adult South Africans by Seloka et al. (68) also showed that high consumption of SSBs increases the risk of high FBG in men. This is not a surprise since Deshpande et al. (67) have long shown that sweetened beverages disrupt the hormones involved in regulating energy balance and the satiety center within the human limbic system, which may lead to overeating and result in an increase in positive energy balance in the body. Therefore, the results are body weight gain and an increase in WC. It is also important to note that the SSBs that were included in our study consisted of cold drinks and reconstituted gas/fizzy drinks. Overconsumption of fructose and sucrose from these SSBs has been shown to stimulate and initiate lipid production in the liver, resulting in higher serum triglyceride and cholesterol levels, visceral fat accumulation, and plaque buildup (69). Moreover, glucose in SSBs has a higher glycaemic index, which can cause high blood glucose spikes and may lead to glucose intolerance, insulin resistance, and an increase in inflammatory biomarkers (70).

Finally, in the current study, we found that avoiding the consumption of fresh fruits increased the likelihood of developing abnormal HDL-C. Although we could not specify the type, color, and amount of fruit we referred to in our research, we could attribute these significant interactions to the fiber and antioxidants that fresh fruit and vegetables have, which are compounds that have been shown to mitigate metabolic risks (71). Although several epidemiological studies have evaluated the association between fruit and vegetable consumption and the risk of MetS, the results remain controversial. For instance, some studies have emphasized fruits' and vegetables' roles in mitigating metabolic disease risk or eliminating the disease entirely (72–74), while others have shown the opposite or no association at all with disease downregulation (75). However, a meta-analysis of international studies by Tian et al. (76) cleared up the controversy by showing that, when data from these studies were combined, high fruit and vegetable consumers were 13% and 24% less likely to have MetS, respectively. This meta-analysis of 78 studies further investigated the relationship between the consumption of fruits, vegetables, and MetS risk. When these researchers stratified these interactions by continent, the inverse association of fruit and vegetable consumption was observed to be OR: 0.86 (0.77, 0.96) and OR: 0.86 (0.80, 0.92), respectively, with the risk of MetS remaining significant in Asia. Based on these results, they concluded that people should consume more fruits and vegetables to reduce the risk of metabolic diseases. However, we should be cognizant of the amount, type, and quality of fruit and vegetables that bring about this kind of health benefit. Studies by Nguyen et al. (77) and Sharma et al. (78) suggest that plant foods high in fiber, such as brown and white rice, have greater metabolic health benefits. Numerous substantiated pieces of evidence suggest that the consumption of good fatty acids can prevent MetS risk and its components (79–83). Sekgala et al. (81), in their recent research, eloquently indicated that substituting SFA for PUFA significantly decreases the likelihood of elevated BP by 7%.

To end the aforementioned arguments, it is also important to highlight that, unlike many studies conducted in South African populations with financial constraints, the current study was based on a population that could afford to procure food. Hence, the majority spent more than ZARR 100 on buying SF. A hundred ZAR and more a day is way above the recommended amount (ZARR 40 per day) per person recommended as enough budget to spend on healthy food each day. Abraham et al. (84) have long suggested that, on average, for an adult South African man, a healthier diet costs ZARR 17.3 (which is about USD\$ 1) per day. In the current research, the minibus taxi drivers who spent more than ZARR 100 on SF had two times greater risk of MetS than their counterparts who spent ZARR 99 or less. This adverse outcome of MetS could be attributed to the unhealthy food options readily available near transportation hubs. While this relationship was lifestyle and sociodemography dependent, the amount spent on SF was not found to be the mediator/moderator of the type of foods purchased and consumed by the minibus taxi drivers included in the current study.

Limitation

Despite the notable and important results of the current study outlined above, there are a number of limitations to this study that need to be considered. First, the study was cross-sectional. Hence, causal inferences cannot be drawn from this study's results. Second, the results of the current study focused only on South African male taxi drivers. Therefore, they can only be generalized to occupational drivers in the long-duration driving business but not to the general population. Finally, even though most of the major confounders have been taken into account in most of the studies, there is still a chance of unmeasured and residual confounding factors impacting in the current study. The confounders that were taken into account in the current study were also different from those in the other international studies that have been used to corroborate/contrast this study's results. Hence, notable differences were observed.

Conclusion

In the current study, we have shown the significant determinants of MetS and its components among South African minibus taxi drivers who presented with abnormal levels of WC, HDL-C, SBP, and FBG, of whom 70% were diagnosed to have MetS according to the IDF diagnostic criteria. Among these important determinants of MetS, we showed that sociodemographic factors such as age and high experience in taxi driving are significantly associated with MetS risk and its components. Moreover, lifestyle factors such as fewer sleeping hours, smoking many cigarettes each day, alcohol and SSB consumption, spending a lot of money on SF, and being sedentary impacted the minibus taxi drivers' metabolic health. More importantly, the consumption of fried food, processed foods, and commercially packaged snacks like crisps, obtained as takeaways, increased the likelihood of minibus taxi drivers developing MetS and its components. However, avoiding the consumption of takeaway and fried foods reduces the risk of MetS. Finally, avoiding the consumption of fruit increased MetS

risk. These results have significant public health implications, as policymakers need to adopt evidence-based strategies to encourage a healthy lifestyle among South African men, especially minibus taxi drivers.

Data availability statement

The original contributions presented in the study are included in the article/[Supplementary material](#), further inquiries can be directed to the corresponding author.

Ethics statement

The studies involving human participants were reviewed and approved by Ms. Patricia Josias Research Ethics Committee Officer University of the Western Cape. The patients/participants provided their written informed consent to participate in this study.

Author contributions

MS and ZM: conceptualization and funding acquisition. MS: formal data analysis, methodology, and writing—original draft. ZM and MO: supervision, writing, review, and editing. BM: biochemical analysis. All authors contributed to the article and approved its submitted version.

Funding

The study was made possible because of Cochrane South Africa, the South African Medical Research Council (SAMRC), and the Collaboration for Evidence-Based Healthcare and Public Health in Africa (CEBHA+) Scholarship Programme. CEBHA+

References

- Chen CC, Shiu LJ, Li YL, Tung KY, Chan KY, Yeh CJ, Chen SC, et al. Shift work and arteriosclerosis risk in professional bus drivers. *Ann Epidemiol.* (2010) 20:60–6. doi: 10.1016/j.annepidem.2009.07.093
- ĐDindić N, Jovanović J, ĐDindić B, Jovanović M, Pešić M, Jovanović JJ. Work stress related lipid disorders and arterial hypertension in professional drivers: A cross-sectional study. *Vojnosanitetski preglad.* (2013) 70:561–8. doi: 10.2298/VSP1306561D
- Gany FM, Gill PP, Ahmed A, Acharya S, Leng J. “Every disease... man can get can start in this cab”: focus groups to identify south Asian taxi drivers’ knowledge, attitudes and beliefs about cardiovascular disease and its risks. *J Immigrant Minority Health.* (2013) 15:986–92. doi: 10.1007/s10903-012-9682-7
- Bigert C, Gustavsson P, Hallqvist J, Hogstedt C, Lewné M, Plato N, Reuterwall C, Schéele P. Myocardial infarction among professional drivers. *Epidemiology.* (2003) 2003:333–9. doi: 10.1097/01.EDE.0000057141.91012.80
- Kurosaka K, Daida H, Muto T, Watanabe Y, Kawai S, Yamaguchi H. Characteristics of coronary heart disease in Japanese taxi drivers as determined by coronary angiographic analyses. *Ind Health.* (2000) 38:15–23. doi: 10.2486/indhealth.38.15
- Hill J, Mchiza Z, Fourie J, Puoane T, Steyn N. Consumption patterns of street food consumers in Cape Town. *Fam Consum Sci Res J.* (2016) 2016:25–35.

receives funding from the Federal Ministry for Education and Research (Bundesministerium für Bildung und Forschung, BMBF), Germany, through the BMBF funding of Research Networks for Health Innovations in Sub-Saharan Africa (Funding No. 81203621), the Non-communicable Diseases Research Unit (NCD-RU) of the SAMRC, and the Human and Social Capabilities (HSC) division of the Human Science Research Council (HSRC).

Acknowledgments

We thank the taxi drivers who participated in the study. We also thank the following nurses for the blood specimen collection: Sister Ntsiki and Sister Theresa.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher’s note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Supplementary material

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fnut.2023.1112975/full#supplementary-material>

- Steyn NP, Labadarios D. Street foods and fast foods: how much do South Africans of different ethnic groups consume? *Ethn Dis.* (2011) 21:462.
- Mchiza Z, Hill J, Steyn N. Foods currently sold by street food vendors in the Western Cape, South Africa, do not foster good health. *Fast Foods.* (2014) 91–118.
- Hill J, Mchiza Z, Puoane T, Steyn NP. Food sold by street-food vendors in Cape Town and surrounding areas: a focus on food and nutrition knowledge as well as practices related to food preparation of street-food vendors. *J Hunger Environ Nutr.* (2019) 14:401–15. doi: 10.1080/19320248.2018.1434104
- McIntyre D, Gilson L. Putting equity in health back onto the social policy agenda: experience from South Africa. *Soc Sci Med.* (2002) 54:1637–56. doi: 10.1016/S0277-9536(01)00332-X
- Basic Conditions of Employment Act. *Sectoral Determination 7: Domestic Worker Sector.* Available online at: https://www.ilo.org/dyn/legosh/en/f?p=14100:1100:0::NO::P1100_ISO_CODE3,P1100_SUBCODE_CODE,P1100_YEAR:ZAF2013 (accessed November 27, 2022).
- Fouracre PR, Sohail M, Cavill S. A participatory approach to urban transport planning in developing countries. *Transp Plan Technol.* (2006) 29:313–30. doi: 10.1080/03081060600905665

13. Pucci G, Alcidi R, Tap L, Battista F, Mattace-Raso F, Schillaci G. Sex- and gender-related prevalence, cardiovascular risk and therapeutic approach in metabolic syndrome: a review of the literature. *Pharmacol Res.* (2017) 120:34–42. doi: 10.1016/j.phrs.2017.03.008
14. Bull F, Goenka S, Lambert V, Pratt M. Physical activity for the prevention of cardiometabolic disease. *Disease Control Prior.* (2017) 5. doi: 10.1596/978-1-4648-0518-9_ch5
15. Myers J. New American Heart Association/American College of Cardiology guidelines on cardiovascular risk: when will fitness get the recognition it deserves? *Mayo Clin Proc.* 89:722–26. doi: 10.1016/j.mayocp.2014.03.002
16. Franklin BA. Physical activity to combat chronic diseases and escalating health care costs: the unfilled prescription. *Curr Sports Med Rep.* (2008) 7:122–5. doi: 10.1097/01.CSMR.0000319709.18052.e8
17. Berra K, Rippe J, Manson JE. Making physical activity ouselling a priority in clinical practice: the time for action is now. *JAMA.* (2015) 314:2617–8. doi: 10.1001/jama.2015.16244
18. Omura JD, Bellissimo MP, Watson KB, Loustalot F, Fulton JE, Carlson SA. Primary care providers' physical activity ouselling and referral practices and barriers for cardiovascular disease prevention. *Prevent Med.* (2018) 108:115–22. doi: 10.1016/j.jpmed.2017.12.030
19. Yang Y, Fan XS, Tian CH, Zhang W, Li J, Li SQ. Health status, intention to seek health examination, and participation in health education among taxi drivers in Jinan, China. *Iran Red Crescent Med J.* (2014). 16:e13355. doi: 10.5812/ircmj.13355
20. Sekgala MD, Opperman M, Mphahli B, Mchiza ZJ. Anthropometric indices and cut-off points for screening of metabolic syndrome among South African taxi drivers. *Front Nutr.* (2022) 9:974749. doi: 10.3389/fnut.2022.974749
21. Shisana O, Labadarios D, Rehle T, Simbayi L, Zuma K, Dhansay A, et al. *The South African National Health and Nutrition Examination Survey, 2012: SANHANES-1: The Health and Nutritional Status of the Nation.* Cape Town: HSRC Press (2013). Available online at: <http://www.hsrc.ac.za/en/research-outputs/view/6493> (accessed February 11, 2023).
22. Hagströmer M, Oja P, Sjöström M. The International Physical Activity Questionnaire (IPAQ): a study of concurrent and construct validity. *Public Health Nutr.* (2006) 9:755–62.
23. Jess K. *How to Calculate Physical Activity Level.* (2010). Available online at: <https://healthfully.com/calculate-physical-activity-level-7264020.html> (accessed November 27, 2022).
24. Federation ID (IDF). *The IDF Consensus Definition of the Metabolic Syndrome in Children and Adolescents.* Brussels, Belgium: IDF Communication. (2007).
25. Topouchian J, Agnoletti D, Blacher J, Youssef A, Ibanez I, Khabouth J, et al. Validation of four automatic devices for self-measurement of blood pressure according to the international protocol of the European Society of Hypertension. *Vasc Health Risk Manag.* (2011) 7:709. doi: 10.2147/VHRM.S27193
26. Alberti KG, Eckel RH, Grundy SM, Zimmet PZ, Cleeman JI, Donato KA, et al. Harmonizing the metabolic syndrome: a joint interim statement of the international diabetes federation task force on epidemiology and prevention; national heart, lung, and blood institute; American heart association; world heart federation; international atherosclerosis society; and international association for the study of obesity. *Circulation.* (2009) 120:1640–5. doi: 10.1161/CIRCULATIONAHA.109.192644
27. Nasri H, Moazenzadeh M. Coronary artery disease risk factors in drivers versus people in other occupations. *ARYA Atherosclerosis Journal.* (2010). 2(2).
28. Shin SY, Lee CG, Song HS, Kim SH, Lee HS, Jung MS, et al. Cardiovascular disease risk of bus drivers in a city of Korea. *Ann Occup Environ Med.* (2013) 25:1–9. doi: 10.1186/2052-4374-25-34
29. Lakshman A, Manikath N, Rahim A, Anilakumari VP. Prevalence and risk factors of hypertension among male occupational bus drivers in North Kerala, South India: a cross-sectional study. *Int Sch Res Notices.* (2014). 2014:318532. doi: 10.1155/2014/318532
30. Inamasu J, Nakatsukasa M, Tomiyasu K, Mayanagi K, Nishimoto M, Oshima T, et al. Stroke while driving: frequency and association with automobile accidents. *Int J stroke.* (2018) 13:301–7. doi: 10.1177/1747493017728398
31. Mohebbi I, Saadat S, Aghassi M, Shekari M, Matinkhah M, Sehat S. Prevalence of metabolic syndrome in Iranian professional drivers: results from a population based study of 12,138 men. *PLoS ONE.* (2012) 7:e31790. doi: 10.1371/journal.pone.0031790
32. Mabetwa EM, Mokwena KE, Mphahlele PM, Modjadji P. Metabolic syndrome and its components among taxi drivers in the city of Tshwane, South Africa. *Applied Sci.* (2022) 12:1767. doi: 10.3390/app12031767
33. Adedokun AO, Ter Goon D, Owolabi EO, Adeniyi OV, Ajayi AI. On-site evaluation of smoking, alcohol consumption and physical inactivity among commercial taxi drivers in buffalo city metropolitan municipality, South Africa. *Global J Health Sci.* (2019) 11:110. doi: 10.5539/gjhs.v11n2p110
34. Bawa MS, Srivastav M. Study the epidemiological profile of taxi drivers in the background of occupational environment, stress and personality characteristics. *Indian J Occup Environ Med.* (2013) 17:108. doi: 10.4103/0019-5278.130855
35. Hildrum B, Mykletun A, Hole T, Midthjell K, Dahl AA. Age-specific prevalence of the metabolic syndrome defined by the International Diabetes Federation and the National Cholesterol Education Program: the Norwegian HUNT 2 study. *BMC Public Health.* (2007) 7:1–9. doi: 10.1186/1471-2458-7-220
36. Lemke M, Apostolopoulos Y. Health and wellness programs for commercial motor-vehicle drivers: organizational assessment and new research directions. *Workplace Health Saf.* (2015) 63:71–80.
37. Hirshkowitz M, Whiton K, Albert SM, Alessi C, Bruni O, DonCarlos L, et al. National Sleep Foundation's updated sleep duration recommendations. *Sleep health.* (2015) 1:233–43. doi: 10.1177/2165079915569740
38. Appiah CA, Afriyie EO, Hayford FE, Frimpong E. Prevalence and lifestyle-associated risk factors of metabolic syndrome among commercial motor vehicle drivers in a metropolitan city in Ghana. *Pan Afr Med J.* (2020) 36:136. doi: 10.11604/pamj.2020.36.136.16861
39. National Department of Health (NdoH), Stats SA, SAMRC, ICF. *South Africa Demographic and Health Survey.* (2016). South Africa and Rockville Maryland, USA: Pretoria. (2019). Available online at: <https://dhsprogram.com/pubs/pdf/FR337/FR337.pdf>. (accessed November 27, 2022).
40. Rodríguez-Monforte M, Sánchez E, Barrio F, Costa B, Flores-Mateo G. Metabolic syndrome and dietary patterns: a systematic review and meta-analysis of observational studies. *Eur J Nutr.* (2017) 56:925–47. doi: 10.1007/s00394-016-1305-y
41. Chen MS, Chiu CH, Chen SH. Risk assessment of metabolic syndrome prevalence involving sedentary occupations and socioeconomic status. *BMJ Open.* (2021) 11:e042802. doi: 10.1136/bmjopen-2020-042802
42. Myers J, Kokkinos P, Nyelin E. Physical activity, cardiorespiratory fitness, and the metabolic syndrome. *Nutrients.* (2019) 11:1652. doi: 10.3390/nu11071652
43. Hernández-Rubio A, Sanvisens A, Bolao F, Cachón-Suárez I, García-Martín C, Short A, et al. Prevalence and associations of metabolic syndrome in patients with alcohol use disorder. *Sci Rep.* (2022) 12:1–7. doi: 10.1038/s41598-022-06010-3
44. Fan AZ, Russell M, Naimi T, Li Y, Liao Y, Jiles R, et al. Patterns of alcohol consumption and the metabolic syndrome. *J Clin Endocrinol Metab.* (2008) 93:3833–8. doi: 10.1210/jc.2007-2788
45. Kim Y, Je Y. Meat consumption and risk of metabolic syndrome: results from the Korean population and a meta-analysis of observational studies. *Nutrients.* (2018) 10:390.
46. Kim OY, Kwak SY, Kim B, Kim YS, Kim HY, Shin MJ. Selected food consumption mediates the association between education level and metabolic syndrome in Korean adults. *Annals of Nutrition and Metabolism.* (2017) 70:122–31. doi: 10.1159/000470853
47. Baik I, Lee M, Jun NR, Lee JY, Shin C. A healthy dietary pattern consisting of a variety of food choices is inversely associated with the development of metabolic syndrome. *Nutr Res Pract.* (2013) 7:233–41. doi: 10.4162/nrp.2013.7.3.233
48. Pan A, Sun Q, Bernstein AM, Schulze MB, Manson JE, Willett WC, et al. Red meat consumption and risk of type 2 diabetes: 3 cohorts of US adults and an updated meta-analysis. *Am J Clin Nutr.* (2011) 94:1088–96. doi: 10.3945/ajcn.111.018978
49. Abete I, Romaguera D, Vieira AR, de Munain AL, Norat T. Association between total, processed, red and white meat consumption and all-cause, CVD and IHD mortality: a meta-analysis of cohort studies. *Br J Nutrition.* (2014) 112:762–75. doi: 10.1017/S000711451400124X
50. Phillips CM, Kesse-Guyot E, McManus R, Hercberg S, Lairon D, Planells R, et al. High dietary saturated fat intake accentuates obesity risk associated with the fat mass and obesity-associated gene in adults. *J Nutr.* (2012) 142:824–31. doi: 10.3945/jn.111.153460
51. Marku A, Galli A, Marciani P, Dule N, Perego C, Castagna M. Iron metabolism in pancreatic beta-cell function and dysfunction. *Cells.* (2021) 10:2841. doi: 10.3390/cells10112841
52. Ley SH, Sun Q, Willett WC, Eliassen AH, Wu K, Pan A, et al. Associations between red meat intake and biomarkers of inflammation and glucose metabolism in women. *Am J Clin Nutr.* (2014) 99:352–60. doi: 10.3945/ajcn.113.075663
53. Griep LM, Seferidi P, Stamler J, Linda VA, Queenie CH, Tzoulaki I, et al. Relation of unprocessed, processed red meat and poultry consumption to blood pressure in East Asian and Western adults. *J Hypertens.* (2016) 34:1721. doi: 10.1097/HJH.0000000000001008
54. Micha R, Michas G, Mozaffarian D. Unprocessed red and processed meats and risk of coronary artery disease and type 2 diabetes—an updated review of the evidence. *Curr Atheroscler Rep.* (2012) 14:515–24. doi: 10.1007/s11883-012-0282-8
55. Flores M, Meyer L, Jorquera P, Castro P, Saravia C, Galdames C, Orellana S. Consumption of deep-fried food and its association with cardiovascular risk factors among first-year students in a Chilean university. *J Nutr Metab.* (2021). 2021:5591662. doi: 10.1155/2021/5591662
56. Sayon-Orea C, Bes-Rastrollo M, Gea A, Zazpe I, Basterra-Gortari FJ, Martínez-González MA. Reported fried food consumption and the incidence of hypertension in a Mediterranean cohort: the SUN (Seguimiento Universidad de Navarra) project. *Br J Nutr.* (2014) 112:984–91. doi: 10.1017/S0007114514001755
57. Kang Y, Kim J. Association between fried food consumption and hypertension in Korean adults. *Br J Nutr.* (2016) 115:87–94. doi: 10.1017/S000711451500402X

58. Lopes T, Zemlin AE, Hill J, Mchiza ZJ, Peer N, Erasmus RT, et al. Consumption of plant foods and its association with cardiovascular disease risk profile in south africans at high-risk of type 2 diabetes mellitus. *Int J Environ Res Public Health*. (2022) 19:13264. doi: 10.3390/ijerph192013264
59. Song Y, Joung H. A traditional Korean dietary pattern and metabolic syndrome abnormalities. *Nutr Metab Cardiovasc Dis*. (2012) 22:456–62. doi: 10.1016/j.numecd.2010.09.002
60. Van Niekerk C. *Biodiversity of Potato Cultivars as Related to Nutrient Content and Quality*, Doctoral dissertation, University of Pretoria. Available online at: <https://repository.up.ac.za/handle/2263/50821>. (accessed November 30, 2022).
61. Foster R, Williamson CS, Lunn J. Briefing paper. Culinary oils and their health effects. *Nutr Bull*. (2009) 34:4–7. doi: 10.1111/j.1467-3010.2008.01738.x
62. Bahadoran Z, Mirmiran P, Azizi F. Fast food pattern and cardiometabolic disorders: a review of current studies. *Health Promot Persp*. (2015) 5:231. doi: 10.15171/hpp.2015.028
63. Devaraj S, Wang-Polagruto J, Polagruto J, Keen CL, Jialal I. High-fat, energy-dense, fast-food-style breakfast results in an increase in oxidative stress in metabolic syndrome. *Metabolism*. (2008) 57:867–70. doi: 10.1016/j.metabol.2008.02.016
64. Barrado E, Mayo MT, Tesedo A, Romero H. Fat composition of several "fast food". *Nutricion Hospitalaria*. (2008) 23:148–58.
65. Payab M, Kelishadi R, Qorbani M, Motlagh ME, Ranjbar SH, Ardalan G, et al. Association of junk food consumption with high blood pressure and obesity in Iranian children and adolescents: the CASPIAN-IV study. *J Pediatr*. (2015) 91:196–205. doi: 10.1016/j.jpeds.2014.07.006
66. Barrio-Lopez MT, Martinez-Gonzalez MA, Fernandez-Montero A, Beunza JJ, Zazpe I, Bes-Rastrollo M. Prospective study of changes in sugar-sweetened beverage consumption and the incidence of the metabolic syndrome and its components: the SUN cohort. *Br J Nutr*. (2013) 110:1722–31. doi: 10.1017/S0007114513000822
67. Deshpande G, Mapanga RF, Essop MF. Frequent sugar-sweetened beverage consumption and the onset of cardiometabolic diseases: cause for concern? *J Endocr Soc*. (2017) 1:1372–85. doi: 10.1210/js.2017-00262
68. Seloka MA, Matshipi M, Mphlegwana PM, Monyeki KD. The association between the consumption of sugar-sweetened beverages and metabolic syndrome components in young rural adults in South Africa. *Appl Sci*. (2022) 12:3015. doi: 10.3390/app12063015
69. Xi B, Huang Y, Reilly KH, Li S, Zheng R, Barrio-Lopez MT, Martinez-Gonzalez MA, et al. Sugar-sweetened beverages and risk of hypertension and CVD: a dose–response meta-analysis. *Br J Nutr*. (2015) 113:709–17. doi: 10.1017/S0007114514004383
70. Lustig RH, Schmidt LA, Brindis CD. The toxic truth about sugar. *Nature*. (2012) 482:27–9. doi: 10.1038/482027a
71. Becerra-Tomás N, Paz-Graniel I, Tresserra-Rimbau A, Martínez-González MÁ, Barrubés L, Corella D, et al. Fruit consumption and cardiometabolic risk in the PREDIMED-plus study: a cross-sectional analysis. *Nutr Metab Cardiovasc Dis*. (2021) 31:1702–13. doi: 10.1016/j.numecd.2021.02.007
72. Esmailzadeh A, Kimiagar M, Mehrabi Y, Azadbakht L, Hu FB, Willett WC. Fruit and vegetable intakes, C-reactive protein, and the metabolic syndrome. *Am J Clin Nutr*. (2006) 84:1489–97. doi: 10.1093/ajcn/84.6.1489
73. Martins Gregório B, Benchimol De Souza D, Amorim de. Morais Nascimento F, Matta L, Fernandes-Santos C. The potential role of antioxidants in metabolic syndrome. *Curr Pharm Des*. (2016) 22:859–69. doi: 10.2174/1381612822666151209152352
74. Li XT, Liao W, Yu HJ, Liu MW, Yuan S, Tang BW, et al. Combined effects of fruit and vegetables intake and physical activity on the risk of metabolic syndrome among Chinese adults. *PLoS ONE*. (2017) 12:e0188533. doi: 10.1371/journal.pone.0188533
75. Ford ES, Mokdad AH, Giles WH, Brown DW. The metabolic syndrome and antioxidant concentrations: findings from the Third National Health and Nutrition Examination Survey. *Diabetes*. (2003) 52:2346–52. doi: 10.2337/diabetes.52.9.2346
76. Tian Y, Su L, Wang J, Duan X, Jiang X. Fruit and vegetable consumption and risk of the metabolic syndrome: a meta-analysis. *Public Health Nutr*. (2018) 21:756–65. doi: 10.1017/S136898001700310X
77. Nguyen SN, Vien MD, Le TT, Tran TT, Ton NM, Le VV. Effects of enzymatic treatment conditions on dietary fiber content of wheat bran and use of cellulase-treated bran in cookie. *Int J Food Sci Technol*. (2021) 56:4017–25. doi: 10.1111/ijfs.15022
78. Sharma S, Katoch V, Kumar S, Chatterjee S. Functional relationship of vegetable colors and bioactive compounds: implications in human health. *J Nutr Biochem*. (2021) 92:108615. doi: 10.1016/j.jnutbio.2021.108615
79. Krešić G, Koprivnjak O, Lešić T, Jurković M, Sokolić D, Gross-Bošković A, et al. Consumption of canned oily fish as a source of fatty acids. *Rivista italiana delle sostanze grasse*. (2017) 94:239–49.
80. Temple NJ, Steyn NP, Fourie J, De Villiers A. Price and availability of healthy food: a study in rural South Africa. *Nutrition*. (2011) 27:55–8. doi: 10.1016/j.nut.2009.12.004
81. Sekgala MD, Opperman M, Mphaheni B, Mchiza ZJ. Association between macronutrient and fatty acid consumption and metabolic syndrome: a South African taxi driver survey. *Int J Environ Res Public Health*. (2022) 19:15452. doi: 10.3390/ijerph192315452
82. Gillingham LG, Harris-Jan S, Jones PJ. Dietary monounsaturated fatty acids are protective against metabolic syndrome and cardiovascular disease risk factors. *Lipids*. (2011) 46:209–28. doi: 10.1007/s11745-010-3524-y
83. Phillips CM, Goumidi L, Bertrais S, Field MR, McManus R, Hercberg S, et al. Dietary saturated fat, gender and genetic variation at the TCF7L2 locus predict the development of metabolic syndrome. *J Nutr Biochem*. (2012) 23:239–44. doi: 10.1016/j.jnutbio.2010.11.020
84. Abrahams Z, De Villiers A, Steyn NP, Fourie J, Dalais L, Hill J, et al. What's in the lunchbox? Dietary behavior of learners from disadvantaged schools in the Western Cape, South Africa Public Health. *Nutrition*. (2011) 14:1752–8. doi: 10.1017/S1368980011001108