



The Evolving Role of the Oncologic Neurosurgeon: Looking Beyond Extent of Resection in the Modern Era

Stephen G. Bowden and Seunggu Jude Han*

Neurological Surgery, Oregon Health & Science University, Portland, OR, United States

OPEN ACCESS

Edited by:

Gordon Li,
Stanford University, United States

Reviewed by:

Matthew Tate,
Northwestern University, United States
Brad E. Zacharia,
Penn State Milton S. Hershey Medical
Center, United States

*Correspondence:

Seunggu Jude Han
hanse@ohsu.edu

Specialty section:

This article was submitted to
Neuro-Oncology and Neurosurgical
Oncology,
a section of the journal
Frontiers in Oncology

Received: 11 May 2018

Accepted: 06 September 2018

Published: 25 September 2018

Citation:

Bowden SG and Han SJ (2018) The
Evolving Role of the Oncologic
Neurosurgeon: Looking Beyond
Extent of Resection in the Modern Era.
Front. Oncol. 8:406.
doi: 10.3389/fonc.2018.00406

Neurosurgeons have played an essential role in glioma management and research for over a century. While the past twenty years have played witness to many exciting developments in glioma biology, diagnosis, and classification, relatively few novel, effective treatment strategies have been introduced. The role of neurosurgery in glioma management has been clarified, with a large body of evidence in support of maximal safe resection. However, neurosurgeons have also played a critical role in translational research during this period. The development of new MRI technologies has benefited greatly from validation with stereotactically-targeted human tissue. Careful banking of surgically acquired tissue was key to the development of a new classification scheme for glioma. Similarly, we have garnered a considerably deeper understanding of molecular and genetic properties of glioma through analysis of large surgical specimens. As our classification schemes become more sophisticated, incorporating targeted tissue sampling into the development of novel treatment strategies becomes essential. Such ex vivo analysis could be instrumental in determining mechanisms of treatment failure or success. Modern tumor neurosurgeons should consider themselves surgical neuro-oncologists, with engagement in translational research essential to furthering the field and improving outlooks for our patients.

Keywords: glioma, glioma stem cells, cilengitide, tumor treating fields, stereotactic neurosurgery, clinical trials as topic, extent of resection (EOR)

INTRODUCTION

Malignant glioma remains among the most elusive of oncological entities despite increasingly aggressive therapies. Glioma cells' diffuse, microscopic infiltration into the brain parenchyma as well as their histologic, molecular, and genetic heterogeneity facilitates their evasion of any combination of maximal safe surgical resection, radiation, and chemotherapy available. These tumors are thus characterized by inevitable treatment failure and recurrence, with an almost universally dismal prognosis. Due to their aggressive progression and rapid growth, surgical cytoreduction was quickly established as the first line therapy to alleviate mass effect and reduce or prevent neurological decline. The past two decades have played witness to a growing body of evidence in support of maximal safe resection, solidifying the neurosurgeon's role in glioma management in an era when non-surgical treatments are favored for many oncological diseases.

Outcomes remain poor, however, due largely to the fact that resection is routinely limited by nearby eloquent tissue and the consequences of significant neurologic damage or dysfunction to patient outcomes. While the therapeutic benefits of surgery have been affirmed, surgically-acquired tissue continues to play an ever pivotal role in advancing the pursuits of the broader neuro-oncology community. This article serves to review some of the key advancements produced by such multi-disciplinary collaboration. In doing so, we offer a look forward to describe our aims as oncologic neurosurgeons in advancing the field in the modern era.

STANDARD OF CARE

The past two decades have played witness to a growing body of evidence in support of open tumor resection over biopsy in the diagnosis and initial treatment of both low and high grade gliomas. Providing greater amounts of tissue for review facilitates accurate pathologic diagnosis and, subsequently, proper allocation of adjuvant therapies. This has been demonstrated by studies showing diagnostic concordance between stereotactic biopsy and open resection in only 51–79% of cases (1–4). More recently, the survival benefits of increasing extent of resection have been elucidated with a summary of studies provided in a meta-analysis of 41,117 unique patients by Brown et al. Their report, adherent to PRISMA guidelines, demonstrated a clear survival benefit for gross total resection over subtotal resection or biopsy at upfront surgery for glioblastoma (5). Similar findings have been suggested in lower grade gliomas, though the evidence remains limited to smaller studies (6). It should be noted that the survival benefits gained with aggressive resection of glioblastoma appear to be lost if the patient develops a new postoperative neurologic deficit as a result, underscoring the importance of safe resection (7).

Intraoperative mapping is one of many surgical adjuncts to facilitate maximal safe resection that have gained traction in recent years. Stimulation cortical or subcortical mapping has recently been shown to improve both patient outcomes and extent of resection (8). Often used in conjunction with preoperative neuropsychological testing and more sophisticated preoperative imaging modalities such as diffusion tensor imaging and functional MRI, this nuanced approach has led to less severe post-operative neurologic deficits following resection of tumors in both perirolandic and language areas (9, 10). Advancements in these techniques have corroborated more common and tenured surgical adjuncts such as frameless stereotactic guidance (11). With these technological advancements facilitating safe surgery and the diagnostic and therapeutic benefits derived from maximal resection, the role of maximal safe resection has been firmly established in the initial management of glioma.

CHARACTERIZING TUMOR HETEROGENEITY

The purpose of surgical resection in glioma, however, extends well beyond cytoreduction, symptomatic relief, and diagnosis

in the modern era. Recently, studies borne of thoughtful and systematic collection of tissue during surgery have played a critical role in broadening our understanding of glioma's biological complexity. Early studies comparing radiologic and histologic characteristics over the whole tumor volume were essential in introducing both diffusion-weighted and perfusion-weighted sequences to standard tumor imaging (12, 13). More recently, the correlation of stereotactically-targeted tumor samples with MRI characteristics has offered more detailed insight into the histologic and molecular diversity within these tumors as well as validation of interpretations of novel MR technologies (14, 15). Such non-invasive mapping of cellular processes and behavior, with the continued addition of similar tissue-validated techniques, will likely play a key role in the effort to distinguish tumor progression from recurrence or radiation necrosis, which remains one of the greatest challenges in the assessment of treatment response.

Integrated genomic analysis of multiple-site and serial-sampling analyses have similarly revealed a high degree of both spatial and temporal heterogeneity within glioma, thought to be a consequence of serial evolutionary events during tumor growth and progression (16, 17). These findings have promoted the concept of therapy-driven clonal selection and preferential regrowth of resistant cell populations as a potential mechanism of treatment failure. Longitudinal banking of serial samples in the form of primary and paired recurrent tumors have further expounded the effects of adjuvant therapies on tumor progression by identifying routes of mutational evolution (18–20). A more nuanced understanding of chronological hierarchy and these tremendously complex mutational networks is critical to the reliable targeting of new chemical or even genetic therapies. Such impactful translational work is wholly reliant on multi-disciplinary collaboration with carefully planned, thoughtful acquisition of tumor samples during surgery.

IMPROVING DIAGNOSTICS

Several other surgical tissue sampling efforts have contributed to a more sophisticated approach to diagnostics in glioma. Intraoperative fluorescent labeling has been found to facilitate targeted tissue sampling for pathologic review by highlighting areas of greater malignant potential (21) or which may even be predictive of tumor grade (22) in multiple-site sampling studies in non-enhancing, lower grade gliomas. Investigations correlating radiologic features with tissue characteristics has demonstrated the ability to predict tumor grade pre-operatively using dynamic, susceptibility-weighted, contrast-enhanced MRI (23). Perhaps more significantly, work by Choi et al. used surgically acquired tissue to expand on laboratory investigations of isocitrate dehydrogenase (IDH) mutations and their byproduct, 2-hydroxyglutarate (2-HG), to present a novel non-invasive approach to glioma diagnostics. Their initial work demonstrated the efficacy of MR spectroscopy in differentiating between IDH mutated and wild-type tumors (24). A follow-up study incorporated comparisons with histology during multiple phases of treatment and underscored the benefits of such a

reliable, non-invasive approach to diagnostics by showing that 2-HG concentration is related to cellularity and reliably reports the disease state from indolence through post-treatment follow-up (25). This subsequent work is an especially exciting development in light of the unique challenges of post-treatment surveillance.

The ability to non-invasively identify a tumor's molecular status is particularly relevant given the recent shift in classification of glial tumors in the 2016 World Health Organization Classification Scheme (26). These changes were prompted by the discovery that molecular and genetic features of the tumors drive clinical behavior and accurately predict long-term outcomes. Sister studies by Brat et al. and Eckel-Passow et al. analyzed data from large scale, rigorous, expertly-curated tumor banking efforts to illuminate the significance of IDH and telomerase reverse transcriptase (TERT) mutations and 1p/19q co-deletions with respect to patient outcomes (27, 28). Similar studies with continued, prospective banking of tumor tissue acquired during surgery will continue to help drive the translation of laboratory advances to the clinical sphere.

DEVELOPING NOVEL TREATMENTS

Meanwhile, the paucity of enduring novel therapies is attributable in part to the relative lack of rigorous, systematic tissue collection and analysis in advanced clinical trials. The few therapeutic strategies recently tested in large randomized phase III trials have yielded largely disappointing results, while the etiology of treatment failure remains unclear. Retroactive analysis of available tumor tissue has elucidated subsets of patients within “failed” trials who may, in fact, benefit due to certain genetic mutations or molecular characteristics. These follow-up studies have underscored the critical role of tissue analysis in any trial of novel therapy.

The trajectory of anti-vascular endothelial growth factor (VEGF) monoclonal antibody bevacizumab is one such example. Two initial phase II trials showed prolonged progression free survival in recurrent glioblastoma, prompting the US Food and Drug Administration to grant accelerated approval to bevacizumab for recurrent glioblastoma in 2009 (29, 30). However, two subsequent, large randomized trials evaluating the addition of bevacizumab to radiation and temozolomide in the newly diagnosed setting failed to show a similar benefit in overall survival (31, 32). More recently, disappointing phase III trial results have brought its utility in the recurrent setting under scrutiny. However, a subgroup analysis of patient data from the AVAglio trial demonstrated an overall survival benefit in those with a proneural subtype tumor (33). Just as O⁶-methylguanine-DNA-methyltransferase (MGMT) promoter methylation does for temozolomide, the genetic profile of the tumor could be predictive of a clinical response to bevacizumab as well. Such analyses, although performed *post hoc*, have benefited from targeted, prospective collection of tumor tissue at initial surgery. Genetic profiling of newly diagnosed malignant gliomas will need to become routine if this approach is to be utilized widely.

Cilengitide, an inhibitor of aVb3 and aVb5 integrins with antiangiogenic properties, similarly showed initial promise. Preliminary phase I and II trials showed radiological response and/or better outcomes in both recurrent and newly diagnosed glioblastoma, with some suggestion of dependence on MGMT methylation status (34–36). Unfortunately, cilengitide's development was stopped after subsequent trials evaluating cilengitide's efficacy in MGMT methylated (CENTRIC) or unmethylated (CORE) groups failed to reproduce those initial successes (37, 38). Again, a *post hoc* analysis of the tissue available from these trial participants offered clarification for their disappointing results. Improved progression free and overall survival was identified in the CORE cohort (MGMT unmethylated) who had high aVb3 levels in the tumor cells (39). While such context is potentially exciting, interpreting these results on a broader scale is difficult, as tissue was not systematically collected in the prospective trials—less than half of the patients' tumor tissue was available for *post-hoc* analysis. One cannot help but wonder if the outcome would have been different if the enrollment in the phase III trial was enriched for or restricted to tumors with high expression of the aVb3 integrin.

In contrast to the evolution of bevacizumab and cilengitide, the novel approach of tumor treating fields (TTFields), a transcutaneous delivery of low-intensity intermediate frequency alternating electric fields, appears to have taken a different trajectory. A multicenter, randomized phase III trial EF-14, randomized patients with supratentorial glioblastoma without evidence of tumor progression following standard chemoradiation to receive maintenance treatment with TTFields and temozolomide or temozolomide alone (40). The study was terminated early based on results of a planned interim analysis demonstrating benefit in progression free as well as overall survival. Despite the trial's success, TTFields has not been quickly adopted by the neuro-oncology community, with low enthusiasm among clinicians and patients alike (41). Experts have raised concerns regarding the study's open-label design, which lacks a sham-treatment control arm, and the study's generalizability, as the randomization point uniquely occurs after concurrent chemoradiation (42). In addition, the lack of a plausible mechanism of action has also likely contributed to the skepticism. The preclinical data for TTFields is sparse, particularly when applied in conjunction with chemoradiation (41). Even during the early phase studies, the use of robust tissue-based analyses exploring the mechanism of action, such as dose response modeling and multi-site tissue biopsies, may have mitigated some of these concerns and led to wider acceptance by the community. Bevacizumab and cilengitide clearly benefited from retrospective analyses of human tumor tissue to identify susceptible treatment populations, using biomarkers rooted in proposed mechanisms of action. The enthusiasm for TTFields is perhaps tempered by the lack of such studies. Collectively, the evolution of these treatments underscores the value of extensive, targeted tissue sampling in furthering development of new adjunctive therapies.

Targeted tissue acquisition and analysis should continue to play a pivotal role in prospective clinical trial designs.

Comparing pre- and post-treatment tissue profiles may elucidate mechanisms of therapeutic action as well as mechanisms of treatment failures (43, 44). Further, these comparisons provide the opportunity to investigate a number of key fundamental steps along a new therapeutic agent's path to potential clinical success such as: Does the agent successfully traffic to the tumor? If so, how much? And, does the agent then actually have its hypothesized effects at the biochemical or cellular level? Thus, early phase trials should include post-treatment molecular biomarker endpoints when possible. NCT02852655 offers an excellent example of this approach, with both pre- and post-treatment tissue resected and analyzed for the tumor infiltrating T-cell profile as a measure of response to the PD-1 inhibitor pembrolizumab. Should the large phase III trials of PD-1/PD-L1 blockade end up negative, this pilot study will lend valuable information in determining why.

CONCLUSION

Gliomas remain incredibly challenging disease entities due to their inter- and intratumoral heterogeneity and diffuse, microscopic infiltration. Contemporary experience hints that future success in glioma management will hinge on precision medicine, where validated biomarkers and targeted personalized therapies are seamlessly integrated into the treatment paradigm. As our overall approach to the management of gliomas evolves, so should our approach to surgery within this framework. The clinical benefits of maximal resection are clear and

surgery becomes progressively safer with technological advances. However, the benefits of surgery extend well beyond improving patient outcomes in the short term. Instead, the oncological neurosurgeon must now look ahead, recognizing the value of thoughtful acquisition and analysis of tumor tissue, as the potential biomedical benefits of such an approach are profound. Systematic and targeted collection of tumor tissue will validate new surgical adjuncts, such as fluorescent molecules and advanced imaging modalities to improve non-invasive tumor characterization and diagnostics, as well as rigorously test novel therapies with tissue-based analysis. Such prospective sampling will provide a pivot upon which we can expand our understanding of the complex network of oncological processes driving these tumors using novel technologies and treatments. Thus, modern tumor neurosurgeons must consider themselves as surgical neuro-oncologists, and engagement and participation of surgeons into clinical trials, particularly early phase trials, should be encouraged and valued. In turn, we should provide our surgical trainees with the training and tools in neuro-oncology to help them participate and contribute in multidisciplinary collaborative efforts aiming to advance the understanding of brain tumor biology and ultimately improve the outlook of our patients.

AUTHOR CONTRIBUTIONS

All authors listed have made a substantial, direct and intellectual contribution to the work, and approved it for publication.

REFERENCES

- Muragaki Y, Chernov M, Maruyama T, Ochiai T, Taira T, Kubo O, et al. Low-grade glioma on stereotactic biopsy: how often is the diagnosis accurate? *Minim Invasive Neurosurg.* (2008) 51:275–9. doi: 10.1055/s-0028-1082322
- McGirt MJ, Bulsara KR, Cummings TJ, New KC, Little KM, Friedman HS, et al. Prognostic value of magnetic resonance imaging-guided stereotactic biopsy in the evaluation of recurrent malignant astrocytoma compared with a lesion due to radiation effect. *J Neurosurg.* (2003) 98:14–20. doi: 10.3171/jns.2003.98.1.0014
- Woodworth G, McGirt MJ, Samdani A, Garonzik I, Olivi A, Weingart JD. Accuracy of frameless and frame-based image-guided stereotactic brain biopsy in the diagnosis of glioma: comparison of biopsy and open resection specimen. *Neurol Res.* (2005) 27:358–62. doi: 10.1179/016164105X40057
- Jackson RJ, Fuller GN, Abi-Said D, Lang FF, Gokaslan ZL, Shi WM, et al. Limitations of stereotactic biopsy in the initial management of gliomas. *Neuro Oncol.* (2001) 3:193–200. doi: 10.1093/neuonc/3.3.193
- Brown TJ, Brennan MC, Li M, Church EW, Brandmeir NJ, Rakszawski KL, et al. Association of the extent of resection with survival in glioblastoma: a systematic review and meta-analysis. *JAMA Oncol.* (2016) 2:1460–9. doi: 10.1001/jamaoncol.2016.1373
- Duffau H. Long-term outcomes after supratotal resection of diffuse low-grade gliomas: a consecutive series with 11-year follow-up. *Acta Neurochir.* (2016) 158:51–8. doi: 10.1007/s00701-015-2621-3
- Rahman M, Abbatematteo J, De Leo EK, Kubilis PS, Vaziri S, Bova F, et al. The effects of new or worsened postoperative neurological deficits on survival of patients with glioblastoma. *J Neurosurg.* (2017) 127:123–31. doi: 10.3171/2016.7.JNS16396
- De Witt Hamer PC, Robles SG, Zwinderman AH, Duffau H, Berger MS. Impact of intraoperative stimulation brain mapping on glioma surgery outcome: a meta-analysis. *J Clin Oncol.* (2012) 30:2559–65. doi: 10.1200/JCO.2011.38.4818
- Bello L, Gallucci M, Fava M, Carrabba G, Giussani C, Acerbi F, et al. Intraoperative subcortical language tract mapping guides surgical removal of gliomas involving speech areas. *Neurosurgery* (2007) 60:67–80; discussion 80–2. doi: 10.1227/01.NEU.0000249206.58601.DE
- Han SJ, Morshed RA, Troncon I, Jordan KM, Henry RG, Hervey-Jumper SL, et al. Subcortical stimulation mapping of descending motor pathways for perirolandic gliomas: assessment of morbidity and functional outcome in 702 cases. *J Neurosurg.* (2018). doi: 10.3171/2018.3.JNS172494. [Epub ahead of print].
- Hervey-Jumper SL, Berger MS. Maximizing safe resection of low- and high-grade glioma. *J Neurooncol.* (2016) 130:269–82. doi: 10.1007/s11060-016-2110-4
- Sugahara T, Korogi Y, Kochi M, Ikushima I, Shigematu Y, Hirai T, et al. Usefulness of diffusion-weighted MRI with echo-planar technique in the evaluation of cellularity in gliomas. *J Magn Reson Imaging* (1999) 9:53–60. doi: 10.1002/(SICI)1522-2586(199901)9:1<53::AID-JMRI7>3.0.CO;2-2
- Aronen HJ, Gazit IE, Louis DN, Buchbinder BR, Pardo FS, Weisskoff RM, et al. Cerebral blood volume maps of gliomas: comparison with tumor grade and histologic findings. *Radiology* (1994) 191:41–51. doi: 10.1148/radiology.191.1.8134596
- Gill BJ, Pisapia DJ, Malone HR, Goldstein H, Lei L, Sonabend A, et al. MRI-localized biopsies reveal subtype-specific differences in molecular and cellular composition at the margins of glioblastoma. *Proc Natl Acad Sci USA.* (2014) 111:12550–5. doi: 10.1073/pnas.1405839111
- Barajas RF Jr, Phillips JJ, Parvataneni R, Molinaro A, Essock-Burns E, Bourne G, et al. Regional variation in histopathologic features of tumor specimens from treatment-naïve glioblastoma correlates with anatomic and physiologic MR Imaging. *Neuro Oncol.* (2012) 14:942–54. doi: 10.1093/neuonc/nos128

16. Suzuki H, Aoki K, Chiba K, Sato Y, Shiozawa Y, Shiraishi Y, et al. Mutational landscape and clonal architecture in grade II and III gliomas. *Nat Genet.* (2015) 47:458–68. doi: 10.1038/ng.3273
17. Sottoriva A, Spiteri I, Piccirillo SG, Touloumis A, Collins VP, Marioni JC, et al. Intratumor heterogeneity in human glioblastoma reflects cancer evolutionary dynamics. *Proc Natl Acad Sci USA.* (2013) 110:4009–14. doi: 10.1073/pnas.1219747110
18. Kim H, Zheng S, Amini SS, Virk SM, Mikkelsen T, Brat DJ, et al. Whole-genome and multisector exome sequencing of primary and post-treatment glioblastoma reveals patterns of tumor evolution. *Genome Res.* (2015) 25:316–27. doi: 10.1101/gr.180612.114
19. Johnson BE, Mazar T, Hong C, Barnes M, Aihara K, McLean CY, et al. Mutational analysis reveals the origin and therapy-driven evolution of recurrent glioma. *Science* (2014) 343:189–93. doi: 10.1126/science.1239947
20. Wang J, Cazzato E, Ladewig E, Frattini V, Rosenbloom DI, Zairis S, et al. Clonal evolution of glioblastoma under therapy. *Nat Genet.* (2016) 48:768–76. doi: 10.1038/ng.3590
21. Bowden SG, Neira JA, B. Gill JA, Ung TH, Englander ZK, Zanazzi G, et al. Sodium fluorescein facilitates guided sampling of diagnostic tumor tissue in nonenhancing gliomas. *Neurosurgery* (2017) 82:719–27. doi: 10.1093/neuros/nyx271
22. Widhalm G, Kiesel B, Woehrer A, Traub-Weidinger T, Preusser M, Marosi C, et al. 5-Aminolevulinic acid induced fluorescence is a powerful intraoperative marker for precise histopathological grading of gliomas with non-significant contrast-enhancement. *PLoS ONE* (2013) 8:e76988. doi: 10.1371/journal.pone.0076988
23. Law M, Yang S, Babb JS, Knopp EA, Golfinos JG, Zagzag D, et al. Comparison of cerebral blood volume and vascular permeability from dynamic susceptibility contrast-enhanced perfusion MR imaging with glioma grade. *AJNR Am J Neuroradiol.* (2004) 25:746–55.
24. Choi C, Ganji SK, DeBerardinis RJ, Hatanpaa KJ, Rakheja D, Kovacs Z, et al. 2-hydroxyglutarate detection by magnetic resonance spectroscopy in IDH-mutated patients with gliomas. *Nat Med.* (2012) 18:624–9. doi: 10.1038/nm.2682
25. Choi C, Raisanen JM, Ganji SK, Zhang S, McNeil SS, An Z, et al. Prospective longitudinal analysis of 2-hydroxyglutarate magnetic resonance spectroscopy identifies broad clinical utility for the management of patients with IDH-mutant glioma. *J Clin Oncol.* (2016) 34:4030–9. doi: 10.1200/JCO.2016.67.1222
26. Louis DN, Perry A, Reifenberger G, von Deimling A, Figarella-Branger D, Cavenee WK, et al. The 2016 world health organization classification of tumors of the central nervous system: a summary. *Acta Neuropathol.* (2016) 131:803–20. doi: 10.1007/s00401-016-1545-1
27. Cancer Genome Atlas Research, Brat DJ, Verhaak RG, Aldape KD, Yung WK, Salama SR, et al. Comprehensive, integrative genomic analysis of diffuse lower-grade gliomas. *N Engl J Med.* (2015) 372:2481–98. doi: 10.1056/NEJMoa1402121
28. Eckel-Passow JE, Lachance DH, Molinaro AM, Walsh KM, Decker PA, Sicotte H, et al. Glioma groups based on 1p/19q, IDH, and TERT promoter mutations in tumors. *N Engl J Med.* (2015) 372:2499–508. doi: 10.1056/NEJMoa1407279
29. Kreisl TN, Kim L, Moore K, Duic P, Royce C, Stroud I, et al. Phase II trial of single-agent bevacizumab followed by bevacizumab plus irinotecan at tumor progression in recurrent glioblastoma. *J Clin Oncol.* (2009) 27:740–5. doi: 10.1200/JCO.2008.16.3055
30. Vredenburgh JJ, Desjardins A, Herndon JE II, Dowell JM, Reardon DA, Quinn JA, et al. Phase II trial of bevacizumab and irinotecan in recurrent malignant glioma. *Clin Cancer Res.* (2007) 13:1253–9. doi: 10.1158/1078-0432.CCR-06-2309
31. Gilbert MR, Dignam JJ, Armstrong TS, Wefel JS, Blumenthal DT, Vogelbaum MA, et al. A randomized trial of bevacizumab for newly diagnosed glioblastoma. *N Engl J Med.* (2014) 370:699–708. doi: 10.1056/NEJMoa1308573
32. Chinot OL, Wick W, Mason W, Henriksson R, Saran F, Nishikawa R, et al. Bevacizumab plus radiotherapy-temozolomide for newly diagnosed glioblastoma. *N Engl J Med.* (2014) 370:709–22. doi: 10.1056/NEJMoa1308345
33. Sandmann T, Bourgon R, Garcia J, Li C, Cloughesy T, Chinot OL, et al. Patients with proneural glioblastoma may derive overall survival benefit from the addition of bevacizumab to first-line radiotherapy and temozolomide: retrospective analysis of the AVAglio trial. *J Clin Oncol.* (2015) 33:2735–44. doi: 10.1200/JCO.2015.61.5005
34. Reardon DA, Fink KL, Mikkelsen T, Cloughesy TF, O'Neill A, Plotkin S, et al. Randomized phase II study of cilengitide, an integrin-targeting arginine-glycine-aspartic acid peptide, in recurrent glioblastoma multiforme. *J Clin Oncol.* (2008) 26:5610–7. doi: 10.1200/JCO.2008.16.7510
35. Nabors LB, Mikkelsen T, Hegi ME, Ye X, Batchelor T, Lesser G, et al. New approaches to brain tumor therapy central nervous system, a safety run-in and randomized phase 2 study of cilengitide combined with chemoradiation for newly diagnosed glioblastoma (NABTT 0306). *Cancer* (2012) 118:5601–7. doi: 10.1002/cncr.27585
36. Stupp R, Hegi ME, Neyns B, Goldbrunner R, Schlegel U, Clement PM, et al. Phase I/IIa study of cilengitide and temozolomide with concomitant radiotherapy followed by cilengitide and temozolomide maintenance therapy in patients with newly diagnosed glioblastoma. *J Clin Oncol.* (2010) 28:2712–8. doi: 10.1200/JCO.2009.26.6650
37. Stupp R, Hegi ME, Gorlia T, Erridge SC, Perry J, Hong YK, et al. Treatment of, C. Canadian Brain Tumor, and C.s. team, Cilengitide combined with standard treatment for patients with newly diagnosed glioblastoma with methylated MGMT promoter (CENTRIC EORTC 26071–22072 study): a multicentre, randomised, open-label, phase 3 trial. *Lancet Oncol.* (2014) 15:1100–8. doi: 10.1016/S1470-2045(14)70379-1
38. Nabors LB, Fink KL, Mikkelsen T, Grujicic D, Tarnawski R, Nam DH, et al. Two cilengitide regimens in combination with standard treatment for patients with newly diagnosed glioblastoma and unmethylated MGMT gene promoter: results of the open-label, controlled, randomized phase II CORE study. *Neuro Oncol.* (2015) 17:708–17. doi: 10.1093/neuonc/nou356
39. Weller M, Nabors LB, Gorlia T, Leske H, Rushing E, Bady P, et al. Cilengitide in newly diagnosed glioblastoma: biomarker expression and outcome. *Oncotarget* (2016) 7:15018–32. doi: 10.18632/oncotarget.7588
40. Stupp R, Taillibert S, Kanner AA, Kesari S, Steinberg DM, Toms SA, et al. Maintenance therapy with tumor-treating fields plus temozolomide vs. temozolomide alone for glioblastoma: a randomized clinical trial. *JAMA* (2015) 314:2535–43. doi: 10.1001/jama.2015.16669
41. Wick W. TTFIELDS: where does all the skepticism come from? *Neuro Oncol.* (2016) 18:303–5. doi: 10.1093/neuonc/now012
42. Sampson JH. Alternating electric fields for the treatment of glioblastoma. *JAMA* (2015) 314:2511–3. doi: 10.1001/jama.2015.16701
43. Lang FF, Asher A. Prospective clinical trials of brain tumor therapy: the critical role of neurosurgeons. *J Neurooncol.* (2004) 69:151–67. doi: 10.1023/B:NEON.0000041878.93045.f5
44. Lang FF, Gilbert MR, Puduvalli VK, Weinberg J, Levin VA, Yung WK, et al. Toward better early-phase brain tumor clinical trials: a reappraisal of current methods and proposals for future strategies. *Neuro Oncol.* (2002) 4:268–77. doi: 10.1093/neuonc/4.4.268

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Bowden and Han. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.