



RETRACTED: LncRNA BLACAT1 Accelerates Non-small Cell Lung Cancer Through Up-Regulating the Activation of Sonic Hedgehog Pathway

Jiwei Sun^{1†}, Jingzhou Jia^{1†}, Wuying Yuan^{1†}, Shu Liu², Wei Wang³, Lili Ge⁴, Liyue Ge³ and Xiao-Jun Liu^{5*}

¹ Department of Thoracic Surgery, Henan Provincial Chest Hospital, Zhengzhou, China, ² Department of Respiratory, Huai'an Second People's Hospital and the Affiliated Huai'an Hospital of Xuzhou Medical University, Huai'an, China, ³ Department of Oncology, Huai'an Second People's Hospital and the Affiliated Huai'an Hospital of Xuzhou Medical University, Huai'an, China, ⁴ Department of Clinical Laboratory, Huai'an Second People's Hospital and the Affiliated Huai'an Hospital of Xuzhou Medical University, Huai'an, China, ⁵ Outpatient Department of External Injury and Wound, Huai'an Second People's Hospital and the Affiliated Huai'an Hospital of Xuzhou Medical University, Huai'an, China

OPEN ACCESS

Edited by:

Xiaochen Wang,
University of Texas Southwestern
Medical Center, United States

Reviewed by:

Zhicheng Wang,
Jilin University, China
Pranjal Sarma,
University of Cincinnati, United States

*Correspondence:

Xiao-Jun Liu
liuxiaojun202102@163.com

†These authors have contributed
equally to this work

Specialty section:

This article was submitted to
Cancer Genetics,
a section of the journal
Frontiers in Oncology

Received: 02 November 2020

Accepted: 22 February 2021

Published: 14 April 2021

Citation:

Sun J, Jia J, Yuan W, Liu S, Wang W, Ge L, Ge L and Liu X-J (2021) LncRNA BLACAT1 Accelerates Non-small Cell Lung Cancer Through Up-Regulating the Activation of Sonic Hedgehog Pathway. *Front. Oncol.* 11:625253. doi: 10.3389/fonc.2021.625253

Recently, increasing evidence has displayed that lncRNAs can exhibit crucial function in cancer progression, including lung cancer. LncRNA bladder cancer-associated transcript 1 (BLACAT1) is reported to participate in various cancers. The aim of our current study was to investigate the function of BLACAT1 in non-small cell lung cancer progression and study the functional pathway. Here, we reported BLACAT1 was significantly up-regulated in lung cancer tissues in comparison to the adjacent normal tissues, which suggested BLACAT1 might act as an oncogene in lung cancer. Then, A549 and PC9 cells were infected with BLACAT1 overexpression plasmid and shRNA. As shown, we proved up-regulation of BLACAT1 greatly induced the growth of non-small cell lung cancer cells. Reversely, knockdown of BLACAT1 reduced A549 and PC9 cell proliferation, migration and invasion. Sonic hedgehog (shh) signaling is able to exert a significant role in carcinogenesis, including lung cancer. Currently, we proved that up-regulation of BLACAT1 activated shh signaling pathway, via inducing shh, Gli-1 and Smo expression. shh pathway inhibitor GANT-61 reversed the effect of overexpression of BLACAT1 on non-small cell lung cancer. Moreover, we manifested that loss of BLACAT1 remarkably reduced the *in vivo* growth and metastasis of A549 cells via enhancing infiltrating CD3+ T cells. In conclusion, our research revealed a critical role of BLACAT1 in the modulation of non-small cell lung cancer via modulating shh pathway.

Keywords: lung cancer, BLACAT1, shh, Gli-1, Smo

INTRODUCTION

Lung cancer is a common cause of cancer-related death worldwide (1). Particularly, almost 1.4 million people have been diagnosed with lung cancer each year (2, 3). NSCLC accounts for about 80–85% and SCLC accounts for 15–20%, respectively (4). In recent years, although advancing progress has been made in lung cancer, the overall survival rate of lung cancer still remains unfavorable (5). Hence, a wide understanding of the possible mechanisms in non-small cell lung cancer is required.

lncRNAs are non-coding RNAs with a length over 200 nucleotides (6, 7). Increasing lncRNAs are reported to exhibit important roles in many processes, such as cell development, cell differentiation and cell apoptosis (8–10). Dysregulated lncRNAs are reported in multiple types of cancers (11–13). For instance, lncRNA UICLM can promote colorectal cancer through sponging miR-215 and regulate ZEB2 expression (14). GAS5 enhance the killing effect of NK cells in HCC *via* modulating miR-544 and RUNX3 (15). In addition, LINC00673 induces lung cancer development through sponging miR-150-5p (16). The biological mechanisms of lncRNAs in non-small cell lung cancer development remain poorly known.

BLACAT1 is firstly recognized in bladder cancer (17). BLACAT1 is correlated with the malignant types in SCLC (18). In our present research, we reported BLACAT1 was obviously increased in lung cancer. For another, shh signaling pathway has been reported to be physiologically activated during embryogenesis, including in lung development (19). shh signaling is also significantly reactivated in many tumors, such as lung cancer (20). Then, the biological effects of BLACAT1 on non-small cell lung cancer progression were assessed in our research. BLACAT1 could induce lung cancer progression through the activation of shh signaling. Hence, the findings reported in our study might potentiate the use of BLACAT1 interference as a potential strategy to repress non-small cell lung cancer development.

MATERIALS AND METHODS

Clinical Subjects

Twenty paired lung cancer tissues and adjacent non-tumor tissues were collected from patients undergoing surgery at Hua'an Second People's Hospital and the Affiliated Hua'an Hospital of Xuzhou Medical University between 2012 and 2016. Before the operation, these patients received no local or systemic treatment. Meanwhile, 10 lung tissues from health donors were obtained. After tissue samples were harvested, they were snap frozen in liquid nitrogen, and stored at -80°C immediately. The study was approved by the Ethics Committee of Hua'an Second People's Hospital and the Affiliated Hua'an Hospital of Xuzhou Medical University. All the patients signed the informed consent.

Cell Culture

Non-small cell lung cancer cells A549 and PC9 were obtained from the Institute of Biochemistry and Cell Biology of the Chinese Academy of Sciences. Cells were incubated in RPMI-1640 with 10% FBS, 100 U/ml penicillin, and 100 mg/ml streptomycin in humidified environment at 37°C with 5% CO_2 .

Vector Construction and Lentiviral Transfection

The negative controls and shBLACAT1 were provided by Gene Pharma (Shanghai, China). Cells were infected with the lentivirus and harvested 48 h later to carry out further cell function experiments. To obtain the BLACAT1 stable knockdown cell line, cells were infected with LV-shBLACAT1 and LV-NC with 6 $\mu\text{g}/\text{ml}$ polybrene and selected using 10 $\mu\text{g}/\text{ml}$ puromycin.

Then, pcDNA3.1-BLACAT1-OE plasmid and control pcDNA3.1 plasmid (Gene Pharma, Shanghai, China) were transfected into cells by using lipofectamine 3000.

CCK-8 Assay

After cell infection, 3,000 cells were grown into a 96-well-plate. After 1d, 2d, 3d, and 4d, cells were added with CCK-8 (Dojindo Laboratories, Kumamoto, Japan) under the manufacturer's instructions. After 2 h, OD value was tested by the microplate reader (BioTek, Winooski, VT, USA).

Colony Formation Assay

After infection, 400 cells were incubated for 14 days in a 6-well-plate. Then the medium was changed twice a week. Cells were washed twice using PBS, fixed in methanol and then stained by 0.1% crystal violet staining solution for 20 min. Afterwards, cells were observed and photographed.

Cell Apoptosis Analysis

The apoptosis was detected using the Annexin V-FITC/PI kit. Cells were inoculated into a 6-well-plate. Transfected cells were collected using trypsinization and re-suspended in $1 \times$ binding buffer. Afterwards, Annexin V-FITC and PI staining was carried out in the dark for 15 min. Then, apoptotic cells were identified by a FACScan flow cytometer. Subsequently, the data were analyzed using CellQuest software.

Scratch Assay

In brief, cells were seeded in 6-well-plates and incubated for a whole night. After appropriate cell attachment, culture inserts were removed. Then, cells were washed twice using PBS. Afterwards, at 0 and 24 h after scratch would formation, images were obtained using an inverted microscope.

Transwell Assay

To evaluate cell migration and invasion ability, transwell migration and invasion assay were conducted. To conduct migration assay, the experiments were carried out the same as those in the invasion assay except for the Matrigel coating (BD Biosciences, Franklin Lakes, NJ, USA). After 24 h, we removed the filters. Cells adhering to the lower surface were fixed and then stained using 0.1% Crystal Violet. An inverted microscope (Olympus, Wuhan, China) was used to photograph five randomly selected fields in each well.

In situ Hybridization

TMA was de-paraffinized by immersion and rehydrated *via* serial ethanol followed by immersion in DEPC-treated PBS for 5 min (21). The BLACAT1 LNA custom detection probe was utilized to carry out. Then, slides were digested using 20 mg/ml proteinase K. Sections were prehybridized in 50% deionized formamide at 58°C for 60 min and hybridized at 58°C with 0.5 mg/ml DIG-labeled RNA probe. After hybridization, slides were treated with 20 mg/ml RNase A for half an hour and washed with 0.1X SSC at 58°C . Afterwards, hybridized probes were determined using alkaline phosphatase-conjugated anti-DIG antibody.

TABLE 1 | Primers for real-time PCR.

Genes	Forward (5'-3')	Reverse (5'-3')
GAPDH	GTCTCCTCTGACTTCAACAGCG	ACCACCCCTGTTGCTGTAGCCAA
Shh	CCGAGCGATTTAAGGAACCTCACC	AGCGTTCAACTGTCTCTTACACC
Gli1	AGCCTTCAGCAATGCCAGTGAC	GTCAGGACCATGCACTGTCTTG
Smo	TGCTCATCGTGGGAGGCTACTT	ATCTTGCTGGCAGCCTTCTCAC

qRT-PCR

TRIzol was employed to isolate total RNA (Takara, Dalian, China) and the resulting RNA molecules were reverse transcribed into cDNA by utilizing PrimeScript™ RT reagent kit (Takara, Dalian, China). qPCR was conducted using a Taq-Man 2 × Universal PCR Master Mix reagent on a CFX96™ Real-time PCR detection system. Then, primer sequences were displayed in **Table 1**.

Western Blot

Equal proteins were subjected to 10% SDS-PAGE gels. PVDF membranes were used to transfer the protein. Then, the membranes were incubated with Shh, Gli1, Smo, and GAPDH antibodies (Cell Signaling Technology, Danvers, MA, USA) at 4°C for a whole night. Next day, the secondary antibody (Thermo Fisher Scientific, Sunnyvale, CA, USA, 1:2,000) was added. Afterwards, protein bands were visualized by enhanced chemiluminescence (ECL).

Flow Cytometry Analysis of Ki-67

Ki-67 positive cells were assessed using flow cytometry. Ki-67 antibody (BD Biosciences, Franklin Lakes, NJ, USA) was used. Cells were washed twice using staining buffer. About 10⁵ cells were incubated with monoclonal antibodies for 45 min in the dark. Then, data were collected by FACS caliber (Becton Dickinson, NJ, USA).

Confocal Microscopy

Cells were fixed using 4% paraformaldehyde and stained using the indicated primary antibodies (Abcam, Cambridge, USA) at 4°C. Then, secondary antibodies were used for 1 h. Nuclei were stained by DAPI (Beyotime, Shanghai, China) for 3 min. A Nikon A1r confocal microscope was used to obtain immunofluorescence.

Tumor Xenograft Assay

Twelve six-week-old female BALB/C nude mice were obtained from Beijing HFk Bio-Technology. A549 cells down-regulated BLACAT1 were injected into BALB/C mice on the right back. Seven days later, the mice were examined 3 times every week for 1 month. Tumor volume was assessed based on the length and width of tumors. At last, mice were sacrificed *via* cervical dislocation after 17 days. Tumor tissues were extracted and collected for the subsequent researches. Animal protocol was approved by the Ethic Committee on Animal Experimentation of Huai'an Second People's Hospital and the Affiliated Huai'an

Hospital of Xuzhou Medical University. Tumor and brain tissue samples were embedded in paraffin for HE staining and immunohistochemistry staining.

Immunohistochemistry Analysis

Excised tumor tissues were fixed, dehydrated, paraffin-embedded, and then cut into sections. Afterwards, sections were analyzed using antibodies against CD3 (1:200, Abcam, Cambridge, USA) and Ki-67 (1:200, Abcam, Cambridge, USA) was used. Light microscopy (Olympus, Tokyo, Japan) was utilized and the image was quantified using Image-Pro Plus.

Statistical Analysis

Statistical analysis was conducted using SPSS 18.0 and GraphPad Prism 6 software. Chi-square test was carried out to analyze the different distribution of clinical variables. *t*-test or analysis of variance (ANOVA) was carried out to evaluate the difference between different groups. $P < 0.05$ was statistically significant.

RESULTS

BLACAT1 Is Significantly Elevated in Non-small Cell Lung Cancer

We collected lung tissues from healthy donors and lung cancer patients. qRT-PCR analysis displayed BLACAT1 expression in lung tissues was obviously increased in lung cancer patients compared to the healthy donors (**Figure 1A**). Besides, BLACAT1 expression level was up-regulated in a stage dependent manner in **Figure 1B**. In addition, BLACAT1 expression in tumor tissues was overexpressed in **Figure 1C**. Subsequently, RNA hybridization analysis revealed that BLACAT1 expression was strongly stained in three lung cancer patients randomly chosen (**Figure 1D**). These data indicated that BLACAT1 was increased in non-small cell lung cancer.

Increased BLACAT1 Induces the Growth of Non-small Cell Lung Cancer

Then, to study the detailed role of BLACAT1, A549, and PC9 cells were infected with pcDNA3.1-BLACAT1-OE. In **Figure 2A**, we found BLACAT1 expression was significantly enhanced by pcDNA3.1-BLACAT1-OE. Flow cytometry indicated that Ki-67 positive cell ratio was increased in BLACAT1 over-expressed A549 and PC9 cells (**Figure 2B**). Flow cytometry analysis evidenced that cell apoptosis was not altered in BLACAT1 over-expressed A549 and PC9 cells in **Figure 2C**. In **Figures 2D,E**, transwell migration and invasion assay manifested overexpression of BLACAT1 suppressed A549 and PC9 cell migration and invasion. Wound closure of A549 and PC9 cells was triggered by the increased BLACAT1 (**Figure 2F**). These results implied that high expression of BLACAT1 promoted non-small cell lung cancer cell progression.




FIGURE 1 | BLACAT1 is increased in lung cancer tissues. **(A)** BLACAT1 expression in lung tissues from healthy donors and lung cancer patients. **(B)** BLACAT1 expression level in different stages of lung cancer patients. **(C)** BLACAT1 expression in para-tumor and tumor tissues from lung cancer patients. **(D)** RNA hybridization analysis of BLACAT1 expression in para-tumor and tumor tissues from three lung cancer patients. (100 \times). ** $P < 0.01$.

Decreased BLACAT1 Restrains the Progression of Non-small Cell Lung Cancer

Reversely, A549 and PC9 cells were infected with BLACAT1 shRNA. It was manifested that BLACAT1 expression level was greatly reduced by BLACAT1 shRNA *in vitro* in **Figure 3A**. In **Figure 3B**, Ki-67 positive cell percentage was decreased by loss of BLACAT1 in A549 and PC9 cells. CCK-8 assay implied the cell viability was restrained by loss of BLACAT1 expression (**Figure 3C**). Then, transwell migration and invasion indicted A549 and PC9 cell migration and invasion was depressed by BLACAT1 shRNA as demonstrated in **Figures 3D,E**. These implied that loss of BLACAT1 restrained the progression of non-small cell lung cancer.

BLACAT1 Activates Shh Signaling Pathway

To further investigate how BLACAT1 contributed to lung cancer cell progression, the activity of shh pathway was assessed. As displayed in **Figures 4A,B**, western blot and qRT-PCR analysis proved that shh, Gli-1, and Smo expression was induced by BLACAT1 in A549 and PC9 cells. Moreover, confocal analysis indicated Gli-1 positive cells were increased by BLACAT1 overexpression in A549 and PC9 cells (**Figure 4C**). In **Figure 4D**, colony formation assay was carried out and we found that A549 and PC9 cell colony formation ability was increased by BLACAT1. GANT-61 acts as an inhibitor of shh signaling pathway and we proved that GANT-61 was able to reverse



the increased cell colonies as shown in **Figure 4D**. These data suggested shh pathway was involved in lung cancer progression mediated by BLACAT1.


Loss of BLACAT1 Reduces *in vivo* Growth and Metastasis of Non-small Cell Lung Cancer

Furthermore, BLACAT1 was knockdown in A549 cells using BLACAT1 shRNA transfection. Cells were then injected into the BALB/c nude mice on the right back. In **Figure 5A**, we observed that tumor volume was remarkably reduced in BLACAT1-shRNA group time dependently. In **Figure 5B**, tumor weight was increased by BLACAT1-shRNA. Moreover, quantification of infiltrating CD3⁺ T cells was analyzed and CD3⁺ T cells were significantly enriched by silence of BLACAT1 (**Figure 5C**). Flow cytometry analysis indicated that Ki-67 positive cells were decreased by the loss of BLACAT1 (**Figure 5D**). Metastatic brain lesions were reduced in BLACAT1-shRNA mice group as confirmed using H&E staining in **Figure 5E**.

DISCUSSION

The significant data of our study is BLACAT1 demonstrates a crucial role in lung cancer development. We reported that BLACAT1 was significantly up-regulated in human lung cancer tissues. In addition, functional studies indicated that





tumorigenic roles of BLACAT1 in promoting non-small cell lung cancer cell progression was involved in shh signaling pathway. Mechanistically, we found that BLACAT1 activated shh signaling pathway. These results indicated that BLACAT1 acted as a crucial oncogene in non-small cell lung cancer.


Many non-coding transcripts have been reported and lncRNAs have obtained increasing attention due to their various function. Many lncRNAs are abnormally expressed and they can exhibit multiple effects in cancer progression (22–24). As a newly discovered lncRNA, BLACAT1 is to be correlated with poor clinical outcomes in many cancers (25). For instance, lncRNA BLACAT1 can promote glioma cancer progression *via* activating Wnt pathway (26). BLACAT1 induces cervical cancer cell growth through activating Wnt pathway (27). In addition, BLACAT1 is increased in hepatocellular carcinoma and down-regulation of BLACAT1 suppressed cancer development through sponging hsa-miR-485-5p (28). In our work, we indicated a novel biological effect of BLACAT1 in non-small cell lung cancer.

In order to investigate the molecular mechanisms through which BLACAT1 regulated non-small cell lung cancer, shh signaling pathway was concentrated. shh signaling pathway is reactivated in many solid tumors (19, 20). A better understanding of shh signaling can open an avenue for the treatment of various

types of cancers (29). For example, abnormal shh signaling can attenuate lung cancer and enhance chemo-therapy (30). The shh signaling is activated by ligands and they can bind to Ptch. The process of shh ligand binding to Ptch reduces the inhibition of Smo and activates signal transduction through Gli transcription factor. Here, we observed that BLACAT1 significantly induce Shh, Gli1 and Smo expression level in A549 and PC9 cells. shh pathway inhibitor GANT-61 obviously reversed the effect of BLACAT1 on lung cancer growth. The detailed mechanism for lncRNA BLACAT1 in regulating Sonic Hedgehog pathway needs further investigation. We would like to focus on whether there is a target gene of BLACAT1 in lung cancer progression.

In our future study, we should figure out the reason for high BLACAT1 expression in non-small cell lung cancer. In addition, microRNAs have gained widespread attention for their role in diverse cancer, including lung cancer (31). The potential effect of BLACAT1 on its downstream molecular microRNAs is limited. lncRNAs can modulate various biological functions at transcriptional, post-transcriptional level and modulate protein activity (32–34). More researches are warranted to investigate the detailed function of BLACAT1 on non-small cell lung cancer.

In conclusion, our study displayed that BLACAT1 acts as a crucial oncogene in non-small cell lung cancer. BLACAT1 can



promote non-small cell lung cancer cell growth *via* activating shh signaling. Our study suggests a strategy for targeting BLACAT1 as a potential biomarker in non-small cell lung cancer.

DATA AVAILABILITY STATEMENT

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation.

ETHICS STATEMENT

The studies involving human participants were reviewed and approved by Huar'an Second Peoples Hospital and the

Affiliated Huai'an Hospital of Xuzhou Medical University. The patients/participants provided their written informed consent to participate in this study. The animal study was reviewed and approved by Huai'an Second Peoples Hospital and the Affiliated Huai'an Hospital of Xuzhou Medical University.

AUTHOR CONTRIBUTIONS

X-JL designed the research and revised the manuscript. JS, JJ, and WY performed the experiments. JS drafted the manuscript. JJ and WY collected the data and did the analysis. All authors approved the final manuscript.

REFERENCES

- Siegel RL, Miller KD, Jemal A. Cancer statistics, (2019). *CA Cancer J Clin.* (2019) 69:7–34. doi: 10.3322/caac.21551
- Jemal A, Bray F, Center MM, Ferlay J, Ward E, Forman D. Global cancer statistics. *CA Cancer J Clin.* (2011) 61:69–90. doi: 10.3322/caac.20107
- Dela Cruz CS, Tanoue LT, Matthay RA. Lung cancer: epidemiology, etiology, and prevention. *Clin Chest Med.* (2011) 32:605–44. doi: 10.1016/j.ccm.2011.09.001
- Goldstraw P, Ball D, Jett JR, Le Chevalier T, Lim E, Nicholson AG, et al. Non-small-cell lung cancer. *Lancet.* (2011) 378:1727–40. doi: 10.1016/S0140-6736(10)62101-0
- Peters S, Adjei AA, Gridelli C, Reck M, Kerr K, Felip E, et al. Metastatic non-small-cell lung cancer (NSCLC): ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol.* (2012) 23(Suppl 7): vii56–64. doi: 10.1093/annonc/mds226
- Mercer TR, Dinger ME, Mattick JS. Long non-coding RNAs: insights into functions. *Nat Rev Genet.* (2009) 10:155–9. doi: 10.1038/nrg2521
- St Laurent G, Wahlestedt C, Kapranov P. The Landscape of long noncoding RNA classification. *Trends Genet.* (2015) 31:239–51. doi: 10.1016/j.tig.2015.03.007
- Fatica A, Bozzoni I. Long non-coding RNAs: new players in cell differentiation and development. *Nat Rev Genet.* (2014) 15:7–21. doi: 10.1038/nrg3606
- Kretz M, Webster DE, Flockhart RJ, Lee CS, Zehnder A, Lopez-Pajares V, et al. Suppression of progenitor differentiation requires the long noncoding RNA ANCR. *Genes Dev.* (2012) 26:338–43. doi: 10.1101/gad.182121.111
- Kumar MM, Goyal R. LncRNA as a therapeutic target for angiogenesis. *Curr Top Med Chem.* (2017) 17:1750–7. doi: 10.2174/156802661766616116144744
- Bhan A, Soleimani M, Mandal SS. Long noncoding RNA and cancer: a new paradigm. *Cancer Res.* (2017) 77:3965–81. doi: 10.1158/0008-5472.CAN-16-2634
- Schmitt AM, Chang HY. Long noncoding RNAs in cancer pathways. *Cancer Cell.* (2016) 29:452–63. doi: 10.1016/j.ccell.2016.03.010

13. Peng WX, Koirala P, Mo YY. LncRNA-mediated regulation of cell signaling in cancer. *Oncogene*. (2017) 36:5661–7. doi: 10.1038/ncr.2017.184
14. Chen DL, Lu YX, Zhang JX, Wei XL, Wang F, Zeng ZL, et al. Long non-coding RNA UICLM promotes colorectal cancer liver metastasis by acting as a ceRNA for microRNA-215 to regulate ZEB3 expression. *Theranostics*. (2017) 7:4836–49. doi: 10.7150/thno.20942
15. Fang P, Xiang L, Chen W, Li S, Huang S, Li J, et al. LncRNA GAS5 enhanced the killing effect of NK cell on liver cancer through regulating miR-544/RUNX3. *Innate Immun*. (2019) 25:99–109. doi: 10.1177/1753425919827632
16. Lu W, Zhang H, Niu Y, Wu Y, Sun W, Li H, et al. Long non-coding RNA linc00673 regulated non-small cell lung cancer proliferation, migration, invasion and epithelial mesenchymal transition by sponging miR-150-5p. *Mol Cancer*. (2017) 16:118. doi: 10.1186/s12943-017-0716-6
17. He W, Cai Q, Sun F, Zhong G, Wang P, Liu H, et al. linc-UBC1 physically associates with polycomb repressive complex 2 (PRC2) and acts as a negative prognostic factor for lymph node metastasis and survival in bladder cancer. *Biochim Biophys Acta*. (2013) 1832:1528–37. doi: 10.1016/j.bbdis.2013.05.010
18. Chen W, Hang Y, Xu W, Wu J, Chen L, Chen J, et al. BLACAT1 predicts poor prognosis and serves as oncogenic lncRNA in small-cell lung cancer. *J Cell Biochem*. (2019) 120:2540–2546.
19. Giroux-Leprieur E, Costantini A, Ding VW, He B. Hedgehog signaling in lung cancer: from oncogenesis to cancer treatment resistance. *Int J Mol Sci*. (2018) 19:2835. doi: 10.3390/ijms19092835
20. Skoda AM, Simovic D, Karin V, Kardum V, Vranic S, Serman L. The role of the Hedgehog signaling pathway in cancer: a comprehensive review. *Bosn J Basic Med Sci*. (2018) 18:8–20. doi: 10.17305/bjbm.2018.2756
21. Tang H, Deng M, Tang Y, Xie X, Guo J, Kong Y, et al. miR-200b and miR-200c as prognostic factors and mediators of gastric cancer cell progression. *Clin Cancer Res*. (2013) 19:5602–12. doi: 10.1158/1078-0432.CCR-13-1326
22. Schmitz SU, Grote P, Herrmann BG. Mechanisms of long noncoding RNA function in development and disease. *Cell Mol Life Sci*. (2016) 73:2491–509. doi: 10.1007/s00018-016-2174-5
23. Chan JJ, Tay Y. Noncoding RNA:RNA regulatory networks in cancer. *Int J Mol Sci*. (2018) 19:1310. doi: 10.3390/ijms19051310
24. Fang Y, Fullwood MJ. Roles, functions, and mechanisms of long non-coding RNAs in cancer. *Genomics Proteomics Bioinformatics*. (2016) 14:42–54. doi: 10.1016/j.gpb.2015.09.006
25. Lu H, Liu H, Yang X, Ye T, Lv P, Wu X, et al. LncRNA BLACAT1 may serve as a prognostic predictor in cancer: evidence from a meta-analysis. *Biomed Res Int*. (2019) 2019:1275491. doi: 10.1155/2019/1275491
26. Li X, Qi S, Ma D, Fan J, Wang J. Long non-coding RNA BLACAT1 promotes the proliferation and invasion of glioma cells via Wnt/beta-catenin signaling. *Exp Ther Med*. (2019) 17:4703–8. doi: 10.3892/etm.2019.7468
27. Wang CH, Li YH, Tian HL, Bao XX, Wang ZM. Long non-coding RNA BLACAT1 promotes cell proliferation, migration and invasion in cervical cancer through activation of Wnt/beta-catenin signaling pathway. *Eur Rev Med Pharmacol Sci*. (2018) 22:3002–9.
28. Peng Y, Leng W, Duan S, Hong M. Long noncoding RNA BLACAT1 is overexpressed in hepatocellular carcinoma and its downregulation suppressed cancer cell development through endogenously competing against hsa-miR-485-5p. *Biomed Pharmacother*. (2019) 116:109027. doi: 10.1016/j.biopha.2019.109027
29. Carballo GB, Honorato JR, de Lopes GPF, Spohr T. A highlight on Sonic hedgehog pathway. *Cell Commun Signal*. (2018) 16:11. doi: 10.1186/s12964-018-0220-7
30. Du J, Chen W, Yang L, Dai J, Guo J, Wu Y, et al. Disruption of SHH signaling cascade by SBE attenuates lung cancer progression and sensitizes DDP treatment. *Sci Rep*. (2017) 7:1899. doi: 10.1038/s41598-017-02063-x
31. Wu KL, Tsai YM, Lien CT, Kuo PL, Hung AJ. The roles of microRNA in lung cancer. *Int J Mol Sci*. (2019) 20:1611. doi: 10.3390/ijms20071611
32. Wang X, Yin H, Zhang L, Zheng D, Yang Y, Zhang J, et al. The construction and analysis of the aberrant lncRNA-miRNA-mRNA network in non-small cell lung cancer. *J Thorac Dis*. (2019) 11:1772–8. doi: 10.21037/jtd.2019.05.69
33. Ma XY, Ma Y, Zhou H, Zhang H, Sun MJ. Identification of the lncRNA-miRNA-mRNA network associated with gastric cancer via integrated bioinformatics analysis. *Oncol Lett*. (2019) 18:5769–84. doi: 10.3892/ol.2019.10922
34. Sun F, Liang W, Tang K, Hong M, Qian J. Profiling the lncRNA-miRNA-mRNA ceRNA network to reveal potential crosstalk between inflammatory bowel disease and colorectal cancer. *PeerJ*. (2019) 7:e7451. doi: 10.7717/peerj.7451

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2021 Sun, Jia, Yuan, Liu, Wang, Ge, Ge and Liu. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.