



Preclinical Models for Acquired Resistance to Third-Generation EGFR Inhibitors in NSCLC: Functional Studies and Drug Combinations Used to Overcome Resistance

Emna Mahfoudhi¹, Charles Ricordel^{1,2}, Gwendoline Lecuyer¹, Cécile Mouric¹, Hervé Lena^{1,2} and Rémy Pedoux^{1*}

¹ Univ Rennes, Institut Nationale de la Santé et de la Recherche Médicale (INSERM), COSS (Chemistry Oncogenesis Stress Signaling), UMR_S 1242, Centre de Lutte Contre le Cancer (CLOC) Eugène Marquis, Rennes, France, ² Centre Hospitalier Universitaire de Rennes, Service de Pneumologie, Université de Rennes 1, Rennes, France

OPEN ACCESS

Edited by:

Iacopo Petrini,
University of Pisa, Italy

Reviewed by:

Jih-Hsiang Lee,
National Taiwan University, Taiwan
Elena Levantini,
Harvard Medical School, United States

*Correspondence:

Rémy Pedoux
remy.pedoux@univ-rennes1.fr

Specialty section:

This article was submitted to
Thoracic Oncology,
a section of the journal
Frontiers in Oncology

Received: 12 January 2022

Accepted: 02 March 2022

Published: 07 April 2022

Citation:

Mahfoudhi E, Ricordel C, Lecuyer G, Mouric C, Lena H and Pedoux R (2022) Preclinical Models for Acquired Resistance to Third-Generation EGFR Inhibitors in NSCLC: Functional Studies and Drug Combinations Used to Overcome Resistance. *Front. Oncol.* 12:853501. doi: 10.3389/fonc.2022.853501

Epidermal growth factor receptor (EGFR)-tyrosine kinase inhibitors (TKIs) are currently recommended as first-line treatment for advanced non-small-cell lung cancer (NSCLC) with *EGFR*-activating mutations. Third-generation (3rd G) EGFR-TKIs, including osimertinib, offer an effective treatment option for patients with NSCLC resistant 1st and 2nd EGFR-TKIs. However, the efficacy of 3rd G EGFR-TKIs is limited by acquired resistance that has become a growing clinical challenge. Several clinical and preclinical studies are being carried out to better understand the mechanisms of resistance to 3rd G EGFR-TKIs and have revealed various genetic aberrations associated with molecular heterogeneity of cancer cells. Studies focusing on epigenetic events are limited despite several indications of their involvement in the development of resistance. Preclinical models, established in most cases in a similar manner, have shown different prevalence of resistance mechanisms from clinical samples. Clinically identified mechanisms include EGFR mutations that were not identified in preclinical models. Thus, NRAS genetic alterations were not observed in patients but have been described in cell lines resistant to 3rd G EGFR-TKI. Mainly, resistance to 3rd G EGFR-TKI in preclinical models is related to the activation of alternative signaling pathways through tyrosine kinase receptor (TKR) activation or to histological and phenotypic transformations. Yet, preclinical models have provided some insight into the complex network between dominant drivers and associated events that lead to the emergence of resistance and consequently have identified new therapeutic targets. This review provides an overview of preclinical studies developed to investigate the mechanisms of acquired resistance to 3rd G EGFR-TKIs, including osimertinib and rociletinib, across all lines of therapy. In fact, some of the models described were first generated to be resistant to first- and second-generation EGFR-TKIs and often carried the T790M mutation, while others had never been exposed to TKIs. The review further describes the therapeutic opportunities to overcome resistance, based on preclinical studies.

Keywords: preclinical models, 3rd G EGFR-TKI, resistance mechanism, osimertinib, lung cancer

INTRODUCTION

Epidermal growth factor receptor (*EGFR*)-activating mutations in non-small-cell lung cancer (NSCLC) are an important predictor of treatment efficacy with *EGFR* tyrosine kinase inhibitors (TKIs). *EGFR*-TKIs have been shown to prolong the survival of patients with tumors harboring *EGFR*-activating mutations from less than 1 year to approximately 20 to 30 months (1, 2). Despite the initial benefits, almost all tumors develop resistance leading to disease progression. Acquired resistance against the first- and second-generation *EGFR*-TKIs is primarily caused by the development of the secondary *EGFR*^{T790M} mutation (3, 4). Several third-generation *EGFR*-TKIs have been developed to overcome T790M-induced resistance. AZD9291 (osimertinib), CO-1686 (rocicetinib), HM61713 (olmutinib), EGF816 (nazartinib), ASP8273 (naquotinib), lazertinib (YH25448), PF06747775, and AC0010 (avitinib) are third-generation (3rd G) *EGFR*-TKIs, which selectively and irreversibly inhibit *EGFR* with the common activating mutations, exon 19 deletion (Del19) and exon 21 L858R mutation, and the T790M mutation while sparing wild-type *EGFR* (5, 6). Currently, osimertinib is the standard of care for *EGFR*-positive advanced NSCLC with T790M mutation. It has been shown to have remarkably positive results as a first-line treatment for *EGFR*-mutated advanced NSCLC, with a median progression-free survival (PFS) of 18.9 months (7), leading to its approval for first-line treatment of metastatic *EGFR*-mutated NSCLC.

Preclinical modeling and analysis of tumor tissue obtained from patients after disease progression have led to the identification of many mechanisms involved in resistance. Contrary to 1st- and 2nd-generation TKIs, no predominant gatekeeper-resistant gene mutations have been observed (8). In clinical studies, mechanisms responsible for resistance include the emergence of mutations in exon 20 of *EGFR* (e.g., C797S) (9), *MET* and *HER2* amplification, gene fusion, altered cell cycle genes, and *de novo* mutations in *KRAS* (10).

The pattern of resistance mechanisms differs in the reported clinical and preclinical studies. The vast majority of preclinical models developed to date to identify the mechanisms underlying resistance to 3rd G *EGFR*-TKIs have used sensitive cell line models exposed to the drug until resistance emerges. The drug concentrations and exposure duration vary from study to study. However, despite the large number of models available, 40% to 50% of the genetic mechanisms associated with disease progression during osimertinib treatment are still unknown (11). This raises the question of whether continuing to generate preclinical models on a recurrent basis would really help to better decipher the mechanisms involved in resistance acquisition and discover biomarkers of relapse. Resistance to *EGFR*-TKIs therapy is associated with high tumor heterogeneity (12). Such heterogeneity requires tools that mimic the real world for the discovery and evaluation of new therapeutic strategies.

Recent published reviews on resistance mechanisms have focused, in particular, on clinical studies (13–18). Only one review specifically addressed preclinical models, but only those generated from cell lines resistant to first-line osimertinib (19).

In this review, we describe reported preclinical models established to identify mechanisms responsible for or involved in resistance to 3rd G *EGFR*-TKIs and the combinatorial approaches used to circumvent this resistance. We also highlight whether the identified mechanism has been reported in clinical studies. In **Table 1**, the models are listed in the order in which they are cited in the review. Drug doses or the duration of treatment are not listed if not reported in the original articles or references. Models based on modified cell lines generated by transfection, transduction, and/or site-directed mutagenesis are not included in **Table 1**, but are mentioned in the review.

EPIGENETICS IN *EGFR*-TKIS RESISTANCE IN NSCLC

Resistance to *EGFR*-TKIs may be related to the presence of preexisting drug-resistant subclones that will be selected under treatment pressure (52) or to the expansion of persistent cells (with or without an acquired resistance mechanism) after the initial response to targeted therapy (53–55). The question that arises is how, in the absence of a genetic mechanism triggering the development of resistance, persister cells manage to escape *EGFR*-TKI therapy. Previous studies have suggested that entering into a drug-tolerant (DT) persister state is an alternative strategy towards acquiring resistance. It has been reported that small cell populations undergo non-genetic adaptations that allow survival in the presence of the drug from which a fraction of cells can grow into the drug (53, 56). The first attempt to characterize the drug-tolerant cells was reported in 2010 by Sharma et al.; DT persister PC9 cells generated by lethal exposure to erlotinib showed upregulation of histone demethylase KDM5A associated with impaired histone deacetylase (HDAC) activity. Interestingly, the cells returned to spontaneous sensitivity after drug withdrawal (57). Subsequent studies identified other targets related to persister DT cells including IGF1-R (58), and Axl (59), in addition to modulation of apoptosis involving Mcl-1 (60) and Aurora kinases (61). In addition to this, sensitivity to 3rd G *EGFR*-TKIs had been restored in resistant cell lines generated after drug withdrawal (47, 49), suggesting a non-genetic adaptation. Taken together, these studies indicate a potential role for epigenetics in the adaptation persister drug-tolerant cells. Epigenetic modulations are changes that affect cellular phenotype without affecting the DNA sequence. These changes include DNA methylation, post-translational modifications of histones, and small and long non-coding RNA sequences, all of which may be reversible and heritable modifications. While the mechanisms of genetic resistance to 3rd G *EGFR*-TKIs, in particular osimertinib, are widely investigated in clinical and preclinical studies, the epigenetic involvement is not well characterized and remains poorly understood. Nevertheless, some published data show a strong epigenetic involvement in this phenomenon and invite further investigations. First, the methylcytosine dioxygenase TET2 and the methyltransferase DNMT3A appeared in the mutational profile of NSCLC patients on post-osimertinib

TABLE 1 | Preclinical models of resistance to third-generation EGFR-TKIs.

| Model generation method | Cell line | 3rd G TKI | Genetic alteration | Therapy | Method/ approach | References |
|--|-----------------------------------|---|--|--|---|-----------------------|
| On-target EGFR-dependent mechanisms | | | | | | |
| Dose escalation method (0.3 to 1 μ M) for several months | PC9 | Rocilitinib | EGFR amplification | Cetuximab + rocilitinib Afatinib + rocilitinib | FISH, Exome sequencing, DNA qPCR | Nukaga et al. (20) |
| MAPK/PI3K implication | | | | | | |
| - Chronic treatment with osimertinib single dose 160 nM for several months | PC9 and H1975 | Osimertinib WZ4002 | - NRAS gain - NRAS Q61K, NRAS E63K and NRAS G12V/R | Selumetinib (MEK inhibitor) + osimertinib | SnaPshot mutation analysis, targeted and whole exome sequencing | Eberlein et al. (21) |
| -Dose escalation method until 160 nM osimertinib | | | - KRAS gain | | | |
| -Dose escalation method until 1500 nM WZ4002 or osimertinib | | | - MAPK1 gain - CRKL1 gain | | | |
| Dose-escalation exposure (0.01 to 1.0 μ mol/L) for 7.8 months followed by single-cell cloning | Gefetinib resistant PC9 (T790M +) | Naquotinib | NRAS amplification in all sub-clones | Selumetinib/Trametinib (MEK inhibitor) + naquotinib | RNA kinome sequencing, WB, qPCR, and NRAS-GTP pull-down | Ninomiya et al. (22) |
| Escalation dose steps (0.3 to 1 μ M) | PC9 | Osimertinib | KRAS G13D | ND | Whole-exome sequencing (WES) | Nukaga et al. (20) |
| Exposure to increasing concentration (10 nM to 1 μ M) | PC9 | Osimertinib | HRAS G13R with increased MET expression | ND | NGS | Ku et al. (23) |
| Exposure to increasing concentration (10 to 500 nM) followed by cloning | PC9 | Osimertinib | BRAF G469 | Selumetinib/Trametinib + osimertinib | NGS | La Monica et al. (24) |
| Dose escalation method (0.3 to 1 μ M) | H1975 | Osimertinib | Integrin β 1 and phospho-Src upregulation with EMT | Dasatinib/bosutinib (src inhibitor) + osimertinib | WB and Q-PCR | Nukaga et al. (20) |
| MET alterations | | | | | | |
| - PC9 mice xenograft tumors collected after 100 days of rocilitinib administration (150 mg/kg BID) | - | Rocilitinib | MET amplification | Crizotinib + rocilitinib | CAPP-Seq profiling, NGS, RTK array, FISH | Chabon et al. (25) |
| - L858R-positive patient-derived xenograft Exposure to increasing concentrations (10 nM to 500 nM) for approximately 6 months | HCC827 | Osimertinib Cross resist to CO-1686 and erlotinib | MET copy gain | ARQ179/ SGX523 / crizotinib (MET inhibitors) + osimertinib | WB, qPCR | Shi et al. (26) |
| Resistant clones were generated by cloning of Resistant cell populations established from resistant xenograft tumors obtained after a series of continuous drug exposure for 115 days. | H1975 | AC0010 Cross- resist to CO-1686 and to osimertinib | MET upregulation | Crizotinib + AC0010 | RNA-sequencing, WB | Xu et al. (27) |
| Exposure to increasing concentrations (0.01 to 1.0 μ mol/L) during 5.2 months | PC9 | Naquotinib | MET amplification | Crizotinib/SGX523 + naquotinib | Phospho-RTK arrays, WB, FISH | Ninomiya et al. (22) |
| AXL | | | | | | |
| Resistant cells were established from subcutaneous tumors collected from mice treated for 29 days with osimertinib | PC9 | Osimertinib | AXL overexpression | ONO-7475 (AXL inhibitor) + osimertinib | WB | Okura et al. (28) |
| Stepwise escalation up to 3 μ M | H1975 | Osimertinib | STC2 upregulation AXL overexpression | R428 (AXL inhibitor) + osimertinib | WB, qPCR, phospho-RTK array | Liu et al. (29) |
| Exposure to escalating doses (0.001–0.5 μ M) | HCC827 | Osimertinib | GAS6 overexpression AXL overexpression | YD (degrader) + osimertinib | WB, IHC | Kim et al. (30) |
| - Exposure to stepwise escalation (10 nmol/L to 2 μ mol/L) over 6 months | HCC827, HCC4006, | Osimertinib | AXL upregulation AXL upregulation+ EMT+ EGFR copy loss+ ALDHA1 upregulation AXL upregulation + MET amplification | Cabozantinib (TKIs inhibitor including AXL) + osimertinib | WB, NGS, qPCR | Namba et al. (31) |
| - Exposure intermittently to 2 μ mol/L over 6 months | PC-9, H1975 | | | | | |
| Exposure to increasing doses up to 1 μ M for more than 6 months | HCC827 | Osimertinib | AXL upregulation associated with MET amplification | CB469 (dual MET and AXL inhibitor) + osimertinib | Phospho-RTK-array | Yang et al. (32) |

(Continued)

TABLE 1 | Continued

| Model generation method | Cell line | 3rd G TKI | Genetic alteration | Therapy | Method/ approach | References |
|--|---|---|--|--|-------------------------------|--|
| Stepwise dose escalation | Gefitinib-resistant PC9 (T790M+) | Osimertinib | AXL overexpression AXL overexpression with MET activation FGFR1 upregulation | -Foretinib (RTK and AURKB inhibitor) -Barasertib (AURKB-specific inhibitor) -Tozasertib | WB, IHC and Q-PCR | Bertran-Alamillo et al. (33) |
| IGF1-R | | | | | | |
| Exposure to increasing concentrations | Gefitinib-resistant PC9 | WZ4002 | IGF1-R activation with IGFBP3 decreased expression | AG-1024 (IGF1-R inhibitor) or BI836845 (monoclonal anti-IGF1/2 blocking antibody) + WZ4002 | RTK-array | Park et al. (34) |
| Stepwise escalation method from 150 nmol/L to 1 µmol/L over 6 months - Chronic exposure to 1 µmol/L over 3 months | - Gefitinib-resistant PC9 (T790M+) - H1975 | Osimertinib | IGF1R activation | Linsitinib (IGF1R inhibitor) + osimertinib | RTK array | Hayakawa et al. (35) |
| Dose escalation method (0.03 to 1 µmol/L) for several months followed by cloning | PC9 | Osimertinib | IGF1-R activation mediated by IGF2 overexpression | Linsitinib + osimertinib | Phospho-RTK array, ELISA | Manabe et al. (36) |
| EMT and stemness | | | | | | |
| Stepwise escalation (0.1 µM to 1 µM) within 6 months | HCC827 | Osimertinib | Zeb1 upregulation | JMF3086 (HDAC inhibitor) + osimertinib | WB | Weng et al. (37) |
| Stepwise dose escalation (0.03 to 1 µmol/L) followed by limiting dilution | H1975 | Osimertinib | Zeb1 upregulation with miR-200c downregulation | LY2090314 (GSK-3 inhibitor) + osimertinib | WB, miRNA array | Fukuda et al. (38) |
| Stepwise method over 6 months | PC9, HCC827 | Osimertinib | ANKRD1 overexpression with miR-200 family downregulation | Imatinib + osimertinib | WES, cDNA microarray | Takahashi et al. (39) |
| Stepwise-dose escalation (500 nm to 1.5 µM) followed by single-cell dilution | H1975 | Osimertinib | Downregulation of SQSTM1/p62 and up regulation of LC3 | - | WB | Verusingam et al. (40) |
| Mesenchymal-resistant cell line derived from biopsies of NSCLC patients who progressed on 3rd-generation EGFR TKIs | - | EGF816 | Hight expression of FGFR1 and FGF2 | BGJ39 (FGFR1/2/3 inhibitor) with EGF816 (nazartinib) | Whole-genome CRISPR screening | Crystal et al. (41); Raoof et al. (42) |
| Exposure to increasing doses | PC9 | Osimertinib | HES1 upregulation ALDH1A1 upregulation | - | WB | Codony-Servat et al. (43) |
| Apoptosis modulators | | | | | | |
| Gradually increasing concentrations (10 nM to 500 nM) for approximately 6 months | -PC9; Gefitinib-resistant T790M+ PC9; HCC827 | Osimertinib | Bim downregulation with Mcl-1 upregulation | MEK inhibitors (PD0325901; AZD6244; GSK1120212) + osimertinib HDAC inhibitors (SAHA and LBH589) + osimertinib | WB | Shi et al. (44); Zang et al. (45) |
| Escalating dose exposure (20 nM to 5 µM) for 12–16 weeks followed by single-cell cloning for 12–16 weeks | H1975 | AC0010 cross-resist to rociletinib and osimertinib. | BCL-2 upregulation | ABT263 (navitoclax) + AC0010 | RNA sequencing, WB | Xu et al. (27) |
| Stepwise increased concentration (5 µM to 15 µM) over 11 months | H1975 | Osimertinib | BCL-2 upregulation | BCL- 2 inhibitors (ABT263/ABT199) + osimertinib | WB, qPCR | Liu et al. (46) |
| NF-KB | | | | | | |
| Exposure to escalating concentrations up to 1 µM for 8 to 10 months | H1975 | CNX-2006 cross-resist to rociletinib | Overexpression of p105 and of p50 | TPCA-1 + CNX-2006 Bortezomib + of CNX-2006 BEZ-235 + of CNX-2006 | WB, phospho-kinase array | Galvani et al. (47) |
| Gradually increasing concentrations: -from 30 nM to 4 µM, for 10 months -from 200 nM up to 4 µM | Gefitinib-resistant PC9 H1975 | Rociletinib | Overexpression of p50, p65, IKKα/β and Kβα | -Rociletinib + TPCA-1 -Rociletinib + metformin | WB | Pan et al. (48) |

(Continued)

TABLE 1 | Continued

| Model generation method | Cell line | 3rd G TKI | Genetic alteration | Therapy | Method/ approach | References |
|--|---------------------------------|----------------------------|---|------------------------------|-------------------------------|------------------|
| Escalating dose exposure (20 nM to 5 μM) for 12–16 weeks followed by single-cell cloning for 12–16 weeks Other mechanism | H1975 | AC0010 | NFKB1 upregulation | – | RNA sequencing, WB | Xu et al. (27) |
| Stepwise dose escalation (50 nM to 1 μM) | PC9, HCC827, H1975, and HCC4006 | Rociletinib or osimertinib | AURKA activation with TPX2 overexpression | Alisertib + osimertinib | Drug screening, WB | Shah et al. (49) |
| Increased concentrations (5 nM to 1.5 μM) over 22 weeks | H1975 | Osimertinib | Upregulation of CDK4, CDK6 and CCND1 and hyperphosphorylation of Rb | Palbociclib + osimertinib | Cell cycle analysis, qPCR, WB | Qin et al. (50) |
| – | HCC827 | Osimertinib | IRE1α upregulation | STF-083010 (IRE1α inhibitor) | WB | Tang et al. (51) |

therapy (62). Second, HDAC inhibitors have shown synergy with osimertinib in reversing epithelial–mesenchymal transition (EMT)-related resistance linked to stemness, in preclinical models (37, 38). In addition, analysis of circular microRNAs (crmiR) in established osimertinib-resistant cell lines revealed 16,000 differentially expressed crmiRs compared to non-resistant cells (63). MicroRNAs such as the miR-200 family have previously been shown to play a role in acquired resistance to osimertinib (39). Finally, recently, long non-coding RNAs (lncRNAs), CRNDE and DGCR5, have been reported to induce resistance to afatinib and osimertinib *via* downregulation of eIF4A (64).

Notably, a recent report showed that the emergence of EGFR inhibitor resistance in NSCLC may also be nonheritable and attributed to stochastic variations (65).

PRECLINICAL MODELS FOR ACQUIRED RESISTANCE TO 3RD G EGFR-TKIs

EGFR-Dependent Mechanisms

The mechanisms of on-target EGFR resistance consist of genetic alterations in EGFR occurring during progression under 3rd G EGFR-TKIs. In clinical studies, EGFR-dependent resistance is related to additional somatic EGFR mutations and to gene amplification (9, 27). EGFR point mutations occur in the kinase domain and affect the osimertinib covalent binding residue (C797S/G, exon 20), the EGFR solvent-front (G796S/R, exon 20), the hinge region (L792H/F), and residues inducing steric interaction (L718Q/V, G719C/S/A and G724S, exon 18) (62, 66, 67).

In preclinical models, EGFR amplification was reported in an established rociletinib-resistant cell line (Table 1) and sensitivity to rociletinib was restored by cetuximab, a specific anti-EGFR monoclonal antibody, or by afatinib (20). Somatic EGFR mutations, however, have not been identified, which could be explained by the efficacy of 3rd G EGFR-TKIs in inhibiting EGFR protein. Thus, to better understand the involvement of EGFR aberrations in the induction of resistance, studies have

been limited to ectopic overexpression of the wild-type (25) or mutated protein or to site-directed mutagenesis replicating mutations described in relapsed patients. The C797S mutation, engineered with a deletion within exon 19 in Ba/F3 cells, conferred significant resistance against osimertinib compared to other EGFR variants such as L718V, L792F/H, and G724S. However, when associated with L858R, C797S/G and L718Q/V conferred comparable resistance, which was greater than in a Del19 background. L792F/H, in contrast, induced significantly less resistance with L858R (68). This indicates that the initial activating mutation may play a role in the potency of resistance to osimertinib. Consistent with this finding, it was reported in the FLAURA study that patients with the L858R mutation have a worse prognosis than those with Del19 (7). Importantly, as observed in the clinic (69–72), earlier-generation EGFR-TKIs may be effective against osimertinib resistance; this may depend not only on the position of the mutation but also on its allelic context. Afatinib inhibited EGFR phosphorylation and cell growth in osimertinib-resistant Ba/F3 cells that exogenously express the G724S mutation, alone or with Del19 (67). The S724 variant induced conformational changes that are incompatible with EGFR-TKIs 3rd G and 1st G binding but not with 2nd G (67). Ba/F3 cells expressing EGFR^{L858R/C797S} by N-ethyl-N-nitrosourea mutagenesis were found to be sensitive to gefitinib (73). A recent study showed that sensitivity and response to EGFR-TKIs are also heterogeneous within the same EGFR exon and proposes a new classification rather based on the structure function of the mutation to determine potential future therapeutic approaches (74).

In clinical studies, emergence of the EGFR^{C797X} mutation is the most common mechanism of resistance to EGFR-dependent osimertinib regardless of treatment line. It was detected in 15% of patients progressing to second-line osimertinib therapy (75) and in only 7% of disease progression when osimertinib is administered in first-line therapy (76). C797S occurred more frequently in association with Del19 than with L858R mutation (77, 78). Otherwise, C797S was observed in less than 3% of cases in rociletinib-resistant patients (25) and was not observed in patients who progressed after AC0010 treatment (79), suggesting that the resistance mechanism might be drug dependent.

EGFR-Independent Mechanisms

Resistance to osimertinib mediated by EGFR-independent mechanisms can be acquired through activation of alternative bypass pathways, aberrant downstream signaling or histologic transformation. **Figure 1** illustrates the involved pathways described in the review.

MAPK and PI3K Pathways Implication

MAPK and AKT are common downstream modulators of receptor tyrosine kinases (RTKs). Activation of the MAPK pathway *via* ERK activation is a common feature of nearly all preclinical models of resistance to 3rd G EGFR-TKIs. However, alterations of other upstream effectors have also been identified as a driving event in the occurrence of resistance. Copy number gains of MAPK1, CRKL, NRAS, and KRAS as well as single mutations in NRAS (G12V/R, Q61K and E63K) were identified in resistant populations established from PC9 and H1975 cells after chronic exposure to osimertinib or WZ4002 (**Table 1**). Combination with MEK inhibitor Selumetinib prevented the emergence of resistant PC9 cells after 34 days at which time resistant cells appeared in the presence of osimertinib alone, and delayed resistance in H1975 from 17 days in the presence of osimertinib alone to 40 days in the presence of selumetinib. Interestingly, combination therapy induced regression in osimertinib-resistant tumors in transgenic mice carrying EGFR^{L858R/T790M} (21). NRAS amplification was also reported, in a subsequent study, in naquotinib-resistant PC9 cells

harboring the T790M mutation (**Table 1**). Either selumetinib or trametinib, a second MEK inhibitor, resensitized naquotinib-resistant cells (22). Single KRAS mutations have also been described in resistance to 3rd G EGFR-TKIs. First, to assess the relevance of KRAS^{G12S} identified in a patient with acquired osimertinib resistance, Ortiz-Cuaran et al. showed that exogenous expression of KRAS^{G12S} in PC9 and HCC827 cells reduced sensitivity to osimertinib and rociletinib, indicating that expression of an activated KRAS allele is sufficient to drive resistance to 3rd-generation TKI (10). In a second study, KRAS^{G13D} mutation was reported as a potential resistance mechanism in osimertinib-resistant PC9 cells (20) (**Table 1**). Another variant, HRAS^{G13R}, in association with increased MET expression was reported in osimertinib-resistant PC9 cells (23) (**Table 1**). Moreover, BRAF^{G469A} mutation occurred in osimertinib-resistant PC9 clones (**Table 1**). Similar to the models mentioned, dual therapy using selumetinib or trametinib with osimertinib was effective in overcoming resistance and enhancing cell death in mutated cells (24). To our knowledge, no NRAS alteration has been reported in clinical studies. Alterations in KRAS, however, are widely reported in clinical data. The KRAS^{G12S} mutation was identified in the lymph node biopsy, collected after relapse of osimertinib treatment, but not in plasma sample. Interestingly, EGFR^{C797S} variant, which in turn appeared after relapse, had been found in the patient's plasma but not in the lymph node biopsy, indicating that different resistance mechanisms may develop at the same




FIGURE 1 | Schematic representation of an overview of pathways implicated in resistance emergence to third-generation EGFR-TKIs. Mechanisms of resistance to third-generation EGFR-TKIs include aberrant activation of receptor tyrosine kinases (MET, AXL, IGF1-R, FGFR, and EGFR) and/or the downstream pathways (PI3K/AKT, RAS/MAPK, and NFκB) and histological transformation. RTK activation is due to overexpression of the protein with or without copy number gain, through its transactivation involving transcription factors (e.g., Jun and FOXA1) or consequently to the overexpression of its specific ligand (e.g., IGF2 and GAS6). Activation of the downstream cascades can also be due to somatic mutations (e.g., RAS, RAF, and PI3K). Histological transformations consist of EMT and EMT-related stemness features including downregulation of E-cadherin and miR200 family, upregulation of Vimentin, Zeb1 and ANKRD1, enrichment in CD44^{high}/CD24^{low} and ALDH1^{high} populations, HES1 overexpression, and autophagy activity. Resistance also required apoptosis modulation through Bim degradation and Bcl-1 upregulation. Non-classified resistance mechanisms include activation of AURKA and its coactivator TPX2 and upregulation of CDK4/6 and IRE1. Green color indicates activation or overexpression; red indicates down-regulation. P: phosphorylation. Star: point mutation. This figure was created using the free version of the Biorender website.

time (10). In another study, the KRAS^{G12D} mutation detected in patient's plasma relapsing on first-line osimertinib therapy was associated with the CTNNB1^{S37F} mutation. Notably, the initial Del19 has not been detected in the post-therapy plasma sample (11). Clearance of the Del19 subpopulation may be due to different sources of the pre- and post-therapy samples or to selection of EGFR^{WT} cells during the development of resistance. The KRAS^{Q61R} variant has also been reported in the occurrence of osimertinib resistance (78). KRAS G12A, Q61H, and A146T variants were found in patients treated with rociletinib that were not detected in their pre-treatment plasma specimens (25). KRAS amplification has also been involved in osimertinib resistance. Molecular profiling of patients who relapsed on osimertinib therapy showed one case of KRAS/MDM2/CDK4 co-amplification (78). Finally, the BRAF^{V600E} mutation has been reported as a mechanism of resistance to osimertinib treatment, both alone (80) and in combination with MET amplification (81). The variant was also identified in a liquid biopsy sample from a patient undergoing treatment with ASP8273, a Japanese 3rd G EGFR-TKI (82). Combination therapies with 3rd G EGFR-TKIs and MEK inhibitors have been developed for lung cancers with EGFR mutations (NCT02143466).

As with ERK, maintained AKT activation is shown in the majority of preclinical models resistant to 3rd G TKIs and its restoration to normal status often required dual therapy. Shigenari et al. reported Src-AKT pathway activation, through Integrin β 1 overexpression, as a resistance mechanism in established osimertinib- and rociletinib-resistant H1975 cells (**Table 1**). Either dasatinib or bosutinib, both Src inhibitors, suppressed Src phosphorylation and restored sensitivity to 3rd G EGFR-TKIs. Finally, to understand if the co-occurring of PIK3CA^{G106V} with CTNNB1^{S37F} mutations observed in a patient that had progressed on rociletinib treatment had a role in acquired resistance, HCC827 cells were engineered with single or both mutations. While PIK3CA^{G106V} expression promoted invasion and migration, CTNNB1^{S37F} activated the wnt/beta-catenin pathway, promoted cellular invasion and suppressed apoptosis in response to rociletinib. Authors suggested a non-redundant cooperation of CTNNB1 with PI3CA alterations to promote tumor metastasis or limit EGFR inhibitor response (83). β -catenin has been shown to play an important role in acquired resistance to EGFR-TKIs and EMT in NSCLC cells (84, 85).

In patients, PIK3CA mutations, including E542K, E545K, and E418K, are frequently observed in resistance to 3rd G EGFR-TKIs (25, 62, 78, 83). They accounted for 7% of the resistance mechanisms to first-line osimertinib therapy (76).

ALTERNATIVE RECEPTOR TYROSINE KINASE ACTIVATION

MET

Activation of C-mesenchymal-epithelial transition factor (c-MET) appears to be a common resistance mechanism to third-generation EGFR-TKIs, and has been found to be associated with

resistance to osimertinib, to rociletinib (10), and to AC0010 (86). In preclinical studies, MET activation is due to gene copy gain or protein overexpression. No MET point mutations have been reported as observed in patients. In an preclinical, *in vivo*, model, rociletinib-resistant tumors were collected from mice bearing PC9 tumors (**Table 1**). Genomic and biochemical analysis revealed MET amplification and activation as the only mechanism of acquired resistance. The combination of rociletinib with crizotinib, a kinase inhibitor with multiple targets including MET, reduced significantly the viability of cell lines derived from rociletinib-resistant tumors. Dual therapy effectively decreased growth on mice of patient-derived xenograft tumors harboring L858R and MET copy gain (25). In a second original model, established by Wanhong et al., AC0010-resistant H1975 cells were generated in two steps first, *in vivo*, and then after selection, *in vitro* (**Table 1**). The derived cells were cross-resistant to EGFR-TKIs of other generations and showed upregulation of MET. Consistent with the above mentioned, double inhibition of EGFR and MET with AC0010 and crizotinib, respectively, prevented colony formation and suppressed MET activation in resistant cell lines and reduced significantly tumor growth in xenograft, compared with single therapy (86). MET activation has also been reported in cell line-based models generated by exposure to progressively increasing concentrations of 3rd G EGFR-TKI. Increased MET copy number was observed in osimertinib-resistant HCC827 cells (**Table 1**), which were also cross-resistant to rociletinib and the 1st-generation EGFR-TKI, erlotinib. MET inhibitors such as crizotinib, ARQ179, or SGX523 sensitized osimertinib-resistant cells (26). In a second model, increased MET phosphorylation was reported to be responsible for acquired resistance in naquotinib-resistant PC9 cells (**Table 1**). Interestingly, dual therapy by naquotinib with crizotinib or SGX-523 had a limited effect on bulk resistant cells, while drastically reducing the proliferation of monoclonal resistant cells, suggesting that heterogeneity may underlie the resistance to a specific TKI target. Notably, the one clone that overexpressed MET protein showed an increase in MET copy number, which was not observed in the resistant parental cells arguing for clonal evolution in the development of resistance. In addition, naquotinib administrated with crizotinib induced robust regression in mice bearing monoclonal resistant cell tumors without apparent cytotoxicity (22).

In clinical studies, MET copy number gains are the most common alternative bypass mechanisms for osimertinib resistance, regardless of treatment line (76, 78). In a recent report, MET amplification was found in 66% ($n = 6/9$) of patients treated with first-line osimertinib (12). Besides osimertinib, MET amplification has also been reported in tumor biopsies from patients with lung adenocarcinoma who developed resistance to rociletinib (10). Point mutations, such as MET P97Q, I865F, and H1094Y, have also been identified in patients with lung cancer progressing on osimertinib (62, 77). The efficacy of MET inhibitors alone or in combined therapy with 3rd G EGFR-TKIs has been reported in clinical studies (87, 88). Recently, the feasibility of combining osimertinib with

savolitinib, a potent and selective MET inhibitor, has been tested in clinical trials (89).

AXL

In NSCLC, it has been reported that anaxelekto (AXL) plays a role in resistance to many anti-cancer drugs including EGFR inhibitors (90). Indeed, Tanguichi et al. showed that PC9 cells do not have basal AXL activity as it is the case for MET and HER3 and that AXL phosphorylation appears shortly (4 h) after exposure to osimertinib and increases throughout the exposure period. They also observed a concomitant increase in MET, HER3, and EGFR phosphorylation, suggesting that AXL activation may accelerate the emergence of tolerant cells (59). In addition, drug-tolerant cells isolated from PC9, HCC4011, or H1975, 9 days after high-dose osimertinib treatment (3 $\mu\text{mol/L}$) expressed a higher level of AXL than parental cells and were highly sensitive to the AXL inhibitor, ONO-7475, in contrast to parental cells (28). Moreover, initial combined treatment with osimertinib and ONO-7475 had more effective effect on tumor regression PC9-derived xenograft when used as the initial treatment than as an alternative therapy once resistance to osimertinib developed (28). Since PC9 cells are enriched in AXL, this indicates that AXL expression level may be a predictor of response to osimertinib. Moreover, primary PE2988 cells, established from the pleural effusion of a patient who developed resistance to osimertinib showed high level of total and phosphorylated AXL and of stanniocalcin (STC2) and responded to the combination of AXL inhibitors and osimertinib (29). STC2 is involved in EGFR-TKIs resistance and was found upregulated in established gefitinib-resistant PC9 and osimertinib-resistant H1975 cells (**Table 1**). Indeed, exogenous overexpression of STC2 activated AXL and increased c-jun level and phosphorylation. In fact, c-jun forms with c-Fos the transcription factor, activation protein-1 (AP-1), which binds to AXL promoter (91), suggesting the involvement of the STC2-JUN-AXL axis in EGFR-TKI resistance (29, 92). Another mechanism of resistance to osimertinib and gefitinib, involving AXL, was described by Kim et al. in osimertinib-resistant HCC827 cells (**Table 1**). The model showed increased expression level of GAS6, a ligand of AXL, and prolonged protein degradation rates in parallel with AXL overexpression. Thus, YD-mediated AXL degradation synergized with osimertinib to restore osimertinib sensitivity *in vitro* and *in vivo* (30). AXL activation was identified as the mechanism of resistance in different established osimertinib-resistant cell lines, despite different EGFR mutational profiles (**Table 1**). It has been observed alone, with an EMT phenotype, or with MET amplification (31–33). Cabozantinib, a multiple TKI including AXL (31), or CB469, a dual inhibitor of MET and AXL, with osimertinib (32) overcame resistance. Notably, resistant clones generated with AXL upregulation lost the T790M subpopulation and some the Del19 population, indicating that clonal evolution leads to heterogeneity in resistance mechanisms. In testing a library of drugs, foretinib, a type II inhibitor targeting a panel of RTKs including MET and AXL, showed the lowest IC₅₀ in resistant cell lines, which were T790M-negative (33), indicating

that clonal heterogeneity is very likely to impair the efficacy of targeted therapy.

In clinical data, high AXL expression was associated with low RR to osimertinib in patients with EGFR-mutated NSCLC (59). Furthermore, PFS and ORR were inversely correlated with AXL mRNA expression in patients with EGFR-mutated NSCLC (93). Results of phase I clinical trials to assess the safety and tolerability of DS-1205c, a specific AXL inhibitor, when combined with osimertinib in metastatic or unresectable subjects with EGFR-mutant NSCLC (NCT03255083) are not yet published.

IGF1-R

Insulin-like growth factor receptor (IGF-R) activation is involved in EGFR-TKIs resistance in NSCLC cell lines (94) and patients (95). Regarding its implication in resistance to 3rd G EGFR-TKIs, it was shown that drug-tolerant cells obtained 72 h after osimertinib treatment expressed elevated levels of total and phosphorylated IGF1-R without changes in the expression of its ligands, IGF1 and IGF2. This activation in the presence of osimertinib was due to epigenetic activation of its own transcription, mediated by the transcription factor FOX1. Moreover, osimertinib treatment enhanced the association of IGF1-R with its adaptor proteins Gab1 and IRS1, thereby promoting cell survival (58). Indeed, WZ4002-resistant PC9 cells (**Table 1**) showed activated IGF1-R associated with IGF1R3 downregulation. Chemical inhibition of IGF1-R with AG-1024 or the blocking monoclonal anti-IGF1/2 antibody, BI836845, restored sensitivity to WZ4002, *in vitro*, and in xenograft mice (34). Loss of IGF1R3 was shown to induce activation of IGF1-R signaling and enhance resistance to WZ4002 in gefitinib-resistance cell line. Reciprocally, addition of recombinant IGF1R3 was sufficient to restore sensitivity to the 3rd G EGFR-TKI (96). In osimertinib-resistant cell line models, IGF1-R activation was observed in the presence (35) or absence of T790M (**Table 1**). Indeed, a protein phosphorylation array performed in PC9 osimertinib cells (**Table 1**) detected high activity of the “p-Y-IRS1 p-IRS2 bind PI3K” pathway, which is involved in IGF1-R signaling. Interestingly, the pathway was not activated in gefitinib- or erlotinib-resistant PC9, suggesting a mechanism specific to third-generation EGFR inhibitors. In contrast to the above models, the resistant cells did not show IGF1R3 downregulation but did show increased IGF2 expression. The IGF1-R inhibitor linsitinib overcame osimertinib resistance in resistant cell lines and in the patient-derived KOLK43 cells [established from pleural effusion of a erlotinib- and osimertinib-resistant patient with high IGF1-R phosphorylation (36)]. Increased phosphorylation of IGF1-R was observed, by immunohistochemistry, in the tumor sample of an EGFR-mutated NSCLC patient who acquired resistance to osimertinib (35).

HISTOLOGICAL TRANSFORMATION

Histological and phenotypic transformations in preclinical models of resistance to 3rd-generation EGFR-TKIs correspond

mainly to epithelial–mesenchymal transition (EMT) and to EMT-related stemness. In contrast to clinical data, transformation into small cell lung cancer has not been reported in preclinical models to date. Features of EMT, including decreased E-cadherin and increased vimentin expression, were reported in the generated osimertinib-resistant cell lines. The phenotype was observed in association with upregulation of the zinc finger transcription factor ZEB1 and the formation of spheroids, a feature of stemness (**Table 1**). Reversal of EMT by dual HDAC and the 3-hydroxy-3-methylglutaryl coenzyme A reductase inhibitor, JMF3086, successfully restored sensitivity to osimertinib (37). In fact, ZEB1 could recruit HDAC1 or DNMT1 to the E-cadherin promoter leading to E-cadherin silencing and EMT induction (97). A similar model (**Table 1**) had shown, in addition, a decrease in microRNA-200c expression (38). In fact, the EMT process is governed by a mutually inhibitory miR-200/ZEB feedback loop (98). Glycogen synthase kinase-3-beta (GSK-3 β) inhibitor (LY2090314) that emerged in drug screening with significant inhibition of resistant cell growth, in combination with osimertinib, bypassed resistance by suppressing AKT signaling and restoring apoptosis in resistant cells (38). GSK-3 β inhibition has been shown to decrease mesenchymal markers and to reduce the associated properties of cancer stem cells (CSC) in aggressive breast cancer (99). A third preclinical model of osimertinib-resistant cells (**Table 1**) reported that the EMT phenotype and downregulation of the miR-200 family were associated with overexpression of Ankyrin Repeat Domain1 (ANKRD1) (39). Indeed, when upregulated, ZEB1 forms a transactivation complex of ANKRD1 with YAP and JUN (100). Imatinib, by inhibiting ANKRD1 and ZEB1, restored apoptosis in resistant cells by increasing levels of Bcl-2 and cleaved PARP (39). Recently, established osimertinib-resistant H1975 clones (**Table 1**) had exhibited EMT characteristics and autophagy activity by downregulating of p62 and upregulation of LC3 (40). Notably, autophagy has been shown to play an important role in promoting cancer metastasis, and inhibition of autophagy might be an effective treatment strategy for malignant cancer (101). Interestingly, whole-genome CRISPR screening in a resistant mesenchymal cell line established from biopsies of NSCLC patients who progressed on 3rd G EGFR-TKIs (41) identified FGFR1 as the top sensitization target of EGF816-resistant cells. Dual EGFR/FGFR inhibition by combining EGF816 with BGJ398, a selective FGFR1–3 inhibitor, induced mesenchymal cell death but had no effect on patient-derived epithelial cell lines (42). In accordance with this, *in vitro* analysis demonstrated that FGF2 supplementation conferred resistance to osimertinib in EGFR mutant NSCLC cells. The same study reported FGFR amplification in patients after progression on osimertinib (102).

Anticancer drug resistance and EMT have been associated with CSCs. However, there are no suitable CSC markers for NSCLC-associated drug resistance and EMT. Upregulation of aldehyde dehydrogenase ALDH1A1, a widely used cancer stem cell marker, was observed in osimertinib-resistant HCC827 cells

with EMT features and MET amplification (31) (**Table 1**). EGFR-TKIs, including osimertinib, have been shown to induce enrichment of ALDH positive subpopulations in EGFR-mutated NSCLC models (103, 104), suggesting that specific dual targeting could overcome this adverse effect. Furthermore, osimertinib-resistant PC9 clones (**Table 1**) showed ALDH1A1 or Hairy and enhancer of split homolog-1 (HES1) overexpression (43). HES1 is a transcriptional factor that plays a critical role in gaining and retaining stemness capacity (105). Clinical studies showed HES1 protein levels increased during relapse and were negatively correlated with PFS in EGFR-mutated patients treated with TKIs including osimertinib (43, 106).

APOPTOSIS MODULATORS

Bcl2-like 11 (BIM) has emerged as a key modulator of EGFR-TKI induced apoptosis. Low levels of *BIM* expression in primary tumors are reported to be associated with shorter PFS in patients treated with EGFR-TKI (107). In preclinical studies, osimertinib-resistant PC9 and HCC827 cells (**Table 1**) showed Bim downregulation and Mcl-1 upregulation in association with ERK activation (44). Bim and Mcl-1 are known to be regulated by ERK (108, 109). MEK inhibitors such as PD0325901, AZD6244, or GSK1120212 suppressed phosphorylation of ERK, Bim, and Mcl-1 in cell lines and effectively decreased the growth of osimertinib-resistant xenografts (44). Alternatively, HDAC inhibitors (SAHA and LBH589) plus osimertinib induced significant growth inhibition of osimertinib-resistant cells and xenografts through Bim stabilization (45). Furthermore, a drug screen performed in gefitinib-resistant cells in which WZ4002 failed to restore Bim expression identified ABT-263 (navitoclax), a dual inhibitor of BCL-XL and BCL-2 at the head of compounds that achieve maximal growth inhibition in combination with WZ4002, suggesting a role for BCL-2 in the occurrence of resistance against 3rd G EGFR-TKIs (55). Indeed, RNA sequencing of AC0010-resistant H1975 cells generated *in vitro* (**Table 1**) revealed an overexpression of BCL-2 (8.6-fold compared to parental cells). Dual therapy with ABT263 and AC0010 enhanced apoptosis in resistant cells and reduced colony formation (86). In another model of osimertinib-resistant H1975 cells (**Table 1**) with BCL-2 upregulation, ABT263 as well as ABT199 (BCL-2 inhibitor) synergized with osimertinib to overcome resistance through downregulation of p21 or downregulation of SQSTM1 and Survivin (46). Clinical trials studying oral combination therapy with navitoclax and osimertinib in advanced EGFR-mutant NSCLC with prior TKI treatment have reported an ORR of up to 100% and a median PFS of 16.8 months. However, thrombocytopenia and lymphopenia were the most common adverse events (37%) observed in the study (110). Finally, it was found that C-FLIP knockdown restored osimertinib-induced apoptosis in resistant cells (**Table 1**), suggesting that C-FLIP depletion may be an effective strategy to overcome osimertinib resistance in NSCLC

(44, 111). Moreover, silencing of C-FLIP had sensitized EGFR-mutant NSCLC to erlotinib and, conversely, its overexpression rescued EGFR-mutant lung cancer cells from erlotinib treatment, presumably through modulation of NF- κ B activity (112).

NF-KB PATHWAY

Enhanced nuclear factor binding near the κ light chain gene in B cells (NF- κ B) signaling activity has been implicated as a possible mechanism of resistance to EGFR-TKIs since patients with EGFR mutations who had developed resistance to erlotinib showed low expression of the NF- κ B inhibitor, I κ B α (112). Activation of NF- κ B by overexpression of NF- κ B1 (p50) and its precursor (p105), without altering the expression level of p65, has been reported as a mechanism of acquired resistance against CNX-2006, a prototype for rociletinib, in resistant H1975 cells (Table 1). Notably, resistant cells showed a variety of differences compared to parental cells, but the involvement of NF- κ B was the most studied. Bortezomib, TPCA-1, or BEZ-235, all inhibitors of the NF- κ B pathway, synergized with CNX-2006 to inhibit cell growth, but with different efficiencies (47). A similar phenotype was observed in established rociletinib-resistant cells (Table 1) that expressed higher levels of p50, p65, phospho-IKK α / β , and phospho-KB α proteins than the parental cells. As in the previous model, combination treatment of rociletinib with TPCA-1 or with Metformin, which is known to inhibit the NF- κ B activity (113), overcame resistance (48). More recently, NF- κ B1 was identified among the top ten upregulated gene in AC0010-resistant H1975 generated *in vitro* (Table 1), but no further investigations have been conducted to understand the mechanism involved in acquired resistance (86).

Others

AURKA

Aurora kinase A (AURKA) is a serine/threonine kinase that plays a key role during cell division particularly in the process of chromosome segregation (114). The Aurora kinase inhibitors, barasertib and VX680, were identified at the top of the list of drugs that synergized with osimertinib or rociletinib to reduce the growth of generated resistant cell lines, respectively (Table 1). Mechanistically, phosphorylation of AURKA was associated with increasing TPX2 protein level following abolition of its CDH1-dependent degradation due to CDH1 sequestration in the cytosol (49). Moreover, barasertib and tozasertib, a second AURK inhibitor, showed a significant antiproliferative effect on osimertinib-resistant cells with no observed difference in AURK expression level (33). Recently, the importance of AURK inhibition in enhancing BIM- and PUMA-mediated apoptosis upon EGFR-TKI therapy in EGFR-mutated lung cancer cells has been described (61). TPX2 expression was significantly increased in tumor tissue samples obtained from patients with advanced EGFR-mutant NSCLC after erlotinib treatment failure compared with results from pre-treatment samples (49).

A Phase 1/1b clinical trial of AURKA inhibitor, Alisertib, with osimertinib in EGFR-mutant stage IV metastatic lung cancer is currently recruiting participants (NCT04085315).

CDK4/6

Upregulation of CDK4 and CDK6 together with hyperphosphorylation of Rb have been reported as a mechanism of resistance to osimertinib in H1975-resistant cells (Table 1). The combination of palbociclib, a selective and potent inhibitor of CDK4/6, with osimertinib overcomes the resistance (50). Acquired alterations in cell cycle genes, including amplification of CDK4/6, CCND1, CCND2, and CCNE1, account for 10% of the acquired resistance mechanisms detected in patients who relapsed after first-line treatment with osimertinib (76).

IRE1

Zheng-Hai Tang et al. suggested an increase in Inositol requiring enzyme 1 α (IRE1 α) expression as a mechanism of resistance to osimertinib in resistant HCC827 established *in vitro* (Table 1). Indeed, IRE1 α knockdown or STF-083010, an inhibitor of IRE1 α , reduces cell viability in resistant cells (51).

CONCLUSION AND PERSPECTIVES

In preclinical studies, resistance to 3rd G EGFR-TKIs is mainly due to genetic alterations that increase activity of receptor tyrosine kinases (such as MET, IGF1-R, and AXL) and downstream signaling cascades (such as RAS/MAPK and PI3K/AKT). Histological transformations are limited to EMT and EMT-related stemness. Multiple mechanisms of resistance could be observed in the same population highlighting the heterogeneity of the process, which may be explained in part by clonal evolution, and suggesting that combination therapies will be required to overcome acquired resistance. At this stage, we cannot conclude which of the mechanisms identified in the *in vitro* or *in vivo* models are the closest to what is described in the clinic because of limited data on *in vivo* studies. In general, the developed models do not really reflect the diversity of mechanisms observed in the clinic. Somatic alterations in EGFR, HER2 amplification, and gene fusions (e.g., ALK, RET, and BRAF fusions) are not identified as mechanisms of resistance to 3rd G EGFR-TKIs in the preclinical models. Nevertheless, the models described highlight the utility of early dual therapy and provide insights into possible combination therapies to optimize treatment lines. They also allow the identification of potential biomarkers in pre-existing resistant cells that will emerge under selective pressures, hence the need to develop new relevant preclinical models. NSCLC organoids derived from primary patient tumors or patient-derived xenograft tumors have been shown to maintain the histologic and tumorigenic properties of the parental cancer cells and reflect the drug responses of the parental tumor (115). Such models as well as murine models of patient-derived xenograft and syngeneic lung cancer may be suitable to further investigate resistance mechanisms that are not identified *in vitro*, such as

EGFR mutations, or small cell or squamous cell transformation, while preserving the authenticity of the tumor. They could also allow the anticipation of investigations on new-generation therapeutic strategies.

AUTHOR CONTRIBUTIONS

RP conceived and supervised the project direction. EM and GL inquired and collected the literature. EM read the literature, wrote the manuscript, and prepared the figure and table. CR and

CM revised the manuscript. RP and HL read and approved the final draft. All authors contributed to the article and approved the submitted version.

FUNDING

This work was funded by post-doctoral fellowship from Association pour la recherche contre le cancer (ARC), Ligue contre le cancer Grand Ouest, Institut National de la Santé et de la Recherche Médicale (INSERM).

REFERENCES

- Mok TS, Wu Y-L, Thongprasert S, Yang C-H, Chu D-T, Saijo N, et al. Gefitinib or Carboplatin–Paclitaxel in Pulmonary Adenocarcinoma. *N Engl J Med* (2009) 361:947–57. doi: 10.1056/NEJMoa0810699
- Sequist LV, Yang JC-H, Yamamoto N, O'Byrne K, Hirsh V, Mok T, et al. Phase III Study of Afatinib or Cisplatin Plus Pemetrexed in Patients With Metastatic Lung Adenocarcinoma With EGFR Mutations. *J Clin Oncol* (2013) 31:3327–34. doi: 10.1200/JCO.2012.44.2806
- Kobayashi S, Boggon TJ, Dayaram T, Jänne PA, Kocher O, Meyerson M, et al. EGFR Mutation and Resistance of Non-Small-Cell Lung Cancer to Gefitinib. *N Engl J Med* (2005) 352:786–92. doi: 10.1056/NEJMoa044238
- Kosaka T, Yatabe Y, Endoh H, Yoshida K, Hida T, Tsuboi M, et al. Analysis of Epidermal Growth Factor Receptor Gene Mutation in Patients With Non-Small Cell Lung Cancer and Acquired Resistance to Gefitinib. *Clin Cancer Res* (2006) 12:5764–9. doi: 10.1158/1078-0432.CCR-06-0714
- Pirker R. Third-Generation Epidermal Growth Factor Receptor Tyrosine Kinase Inhibitors in Advanced Nonsmall Cell Lung Cancer. *Curr Opin Oncol* (2016) 28:115–21. doi: 10.1097/CCO.0000000000000260
- Barnes TA, O'Kane GM, Vincent MD, Leigh NB. Third-Generation Tyrosine Kinase Inhibitors Targeting Epidermal Growth Factor Receptor Mutations in Non-Small Cell Lung Cancer. *Front Oncol* (2017) 7:113:113. doi: 10.3389/fonc.2017.00113
- Soria J-C, Ohe Y, Vansteenkiste J, Reungwetwattana T, Chewaskulyong B, Lee KH, et al. Osimertinib in Untreated EGFR -Mutated Advanced Non-Small-Cell Lung Cancer. *N Engl J Med* (2018) 378:113–25. doi: 10.1056/NEJMoa1713137
- Zhang Q, Zhang X-C, Yang J-J, Yang Z-F, Bai Y, Su J, et al. EGFR L792H and G796R: Two Novel Mutations Mediating Resistance to the Third-Generation EGFR Tyrosine Kinase Inhibitor Osimertinib. *J Thorac Oncol* (2018) 13:1415–21. doi: 10.1016/j.jtho.2018.05.024
- Thress KS, Pawletz CP, Felip E, Cho BC, Stetson D, Dougherty B, et al. Acquired EGFR C797S Mutation Mediates Resistance to AZD9291 in Non-Small Cell Lung Cancer Harboring EGFR T790M. *Nat Med* (2015) 21:560–2. doi: 10.1038/nm.3854
- Ortiz-Cuaran S, Scheffler M, Plenker D, Dahmen L, Scheel AH, Fernandez-Cuesta L, et al. Heterogeneous Mechanisms of Primary and Acquired Resistance to Third-Generation EGFR Inhibitors. *Clin Cancer Res* (2016) 22:4837–47. doi: 10.1158/1078-0432.CCR-15-1915
- Ramalingam SS, Yang JC-H, Lee CK, Kurata T, Kim D-W, John T, et al. Osimertinib As First-Line Treatment of EGFR Mutation–Positive Advanced Non-Small-Cell Lung Cancer. *JCO* (2018) 36:841–9. doi: 10.1200/JCO.2017.74.7576
- Roper N, Brown A-L, Wei JS, Pack S, Trindade C, Kim C, et al. Clonal Evolution and Heterogeneity of Osimertinib Acquired Resistance Mechanisms in EGFR Mutant Lung Cancer. *Cell Rep Med* (2020) 1:100007. doi: 10.1016/j.xcrm.2020.100007
- Leonetti A, Sharma S, Minari R, Perego P, Giovannetti E, Tiseo M. Resistance Mechanisms to Osimertinib in EGFR-Mutated Non-Small Cell Lung Cancer. *Br J Cancer* (2019) 121:725–37. doi: 10.1038/s41416-019-0573-8
- Schmid S, Li J-J, Leigh NB. Mechanisms of Osimertinib Resistance and Emerging Treatment Options. *Lung Cancer* (2020) 147:123–9. doi: 10.1016/j.lungcan.2020.07.014
- Lazzari C, Gregorc V, Karachaliou N, Rosell R, Santarpia M. Mechanisms of Resistance to Osimertinib. *J Thorac Dis* (2020) 12:2851–8. doi: 10.21037/jtd.2019.08.30
- Piper-Vallillo AJ, Sequist LV, Piotrowska Z. Emerging Treatment Paradigms for EGFR-Mutant Lung Cancers Progressing on Osimertinib: A Review. *JCO* (2020) 38:2926–36. doi: 10.1200/JCO.19.03123
- Passaro A, Jänne PA, Mok T, Peters S. Overcoming Therapy Resistance in EGFR-Mutant Lung Cancer. *Nat Cancer* (2021) 2:377–91. doi: 10.1038/s43018-021-00195-8
- Ricordel C, Friboulet L, Facchinetti F, Soria J-C. Molecular Mechanisms of Acquired Resistance to Third-Generation EGFR-TKIs in EGFR T790M-Mutant Lung Cancer. *Ann Oncol* (2018) 29:i28–37. doi: 10.1093/annonc/mdx705
- Ohara S, Suda K, Mitsudomi T. Cell Line Models for Acquired Resistance to First-Line Osimertinib in Lung Cancers—Applications and Limitations. *Cells* (2021) 10:354. doi: 10.3390/cells10020354
- Nukaga S, Yasuda H, Tsuchihara K, Hamamoto J, Masuzawa K, Kawada I, et al. Amplification of EGFR Wild-Type Alleles in Non-Small Cell Lung Cancer Cells Confers Acquired Resistance to Mutation-Selective EGFR Tyrosine Kinase Inhibitors. *Cancer Res* (2017) 77:2078–89. doi: 10.1158/0008-5472.CAN-16-2359
- Eberlein CA, Stetson D, Markovets AA, Al-Kadhimi KJ, Lai Z, Fisher PR, et al. Acquired Resistance to the Mutant-Selective EGFR Inhibitor AZD9291 Is Associated With Increased Dependence on RAS Signaling in Preclinical Models. *Cancer Res* (2015) 75:2489–500. doi: 10.1158/0008-5472.CAN-14-3167
- Ninomiya K, Ohashi K, Makimoto G, Tomida S, Higo H, Kayatani H, et al. MET or NRAS Amplification is an Acquired Resistance Mechanism to the Third-Generation EGFR Inhibitor Naquotinib. *Sci Rep* (2018) 8:1955. doi: 10.1038/s41598-018-20326-z
- Ku BM, Choi MK, Sun J-M, Lee S-H, Ahn JS, Park K, et al. Acquired Resistance to AZD9291 as an Upfront Treatment Is Dependent on ERK Signaling in a Preclinical Model. *PLoS One* (2018) 13:e0194730. doi: 10.1371/journal.pone.0194730
- La Monica S, Minari R, Cretella D, Bonelli M, Fumarola C, Cavazzoni A, et al. Acquired BRAF G469A Mutation as a Resistance Mechanism to First-Line Osimertinib Treatment in NSCLC Cell Lines Harboring an EGFR Exon 19 Deletion. *Targ Oncol* (2019) 14:619–26. doi: 10.1007/s11523-019-00669-x
- Chabon JJ, Simmons AD, Lovejoy AF, Esfahani MS, Newman AM, Haringsma HJ, et al. Circulating Tumour DNA Profiling Reveals Heterogeneity of EGFR Inhibitor Resistance Mechanisms in Lung Cancer Patients. *Nat Commun* (2016) 7:11815. doi: 10.1038/ncomms11815
- Shi P, Oh Y-T, Zhang G, Yao W, Yue P, Li Y, et al. Met Gene Amplification and Protein Hyperactivation Is a Mechanism of Resistance to Both First and Third Generation EGFR Inhibitors in Lung Cancer Treatment. *Cancer Lett* (2016) 380:494–504. doi: 10.1016/j.canlet.2016.07.021
- Xu H, Shen J, Xiang J, Li H, Li B, Zhang T, et al. Characterization of Acquired Receptor Tyrosine–Kinase Fusions as Mechanisms of Resistance to

- EGFR Tyrosine-Kinase Inhibitors. *CMAR* (2019) 11:6343–51. doi: 10.2147/CMAR.S197337
28. Okura N, Nishioka N, Yamada T, Taniguchi H, Tanimura K, Katayama Y, et al. ONO-7475, a Novel AXL Inhibitor, Suppresses the Adaptive Resistance to Initial EGFR-TKI Treatment in *EGFR*-Mutated Non-Small Cell Lung Cancer. *Clin Cancer Res* (2020) 26:2244–56. doi: 10.1158/1078-0432.CCR-19-2321
 29. Liu Y, Tsai M, Wu S, Chang T, Tsai T, Gow C, et al. Acquired Resistance to EGFR Tyrosine Kinase Inhibitors Is Mediated by the Reactivation of STC2/JUN/AXL Signaling in Lung Cancer. *Int J Cancer* (2019) 145:1609–24. doi: 10.1002/ijc.32487
 30. Kim D, Bach D-H, Fan Y-H, Luu T-T, Hong J-Y, Park HJ, et al. AXL Degradation in Combination With EGFR-TKI can Delay and Overcome Acquired Resistance in Human Non-Small Cell Lung Cancer Cells. *Cell Death Dis* (2019) 10:361. doi: 10.1038/s41419-019-1601-6
 31. Namba K, Shien K, Takahashi Y, Torigoe H, Sato H, Yoshioka T, et al. Activation of AXL as a Preclinical Acquired Resistance Mechanism Against Osimertinib Treatment in *EGFR*-Mutant Non-Small Cell Lung Cancer Cells. *Mol Cancer Res* (2019) 17:499–507. doi: 10.1158/1541-7786.MCR-18-0628
 32. Yang Y-M, Jang Y, Lee SH, Kang B, Lim SM. AXL/MET Dual Inhibitor, CB469, has Activity in Non-Small Cell Lung Cancer With Acquired Resistance to EGFR TKI With AXL or MET Activation. *Lung Cancer* (2020) 146:70–7. doi: 10.1016/j.lungcan.2020.05.031
 33. Bertran-Alamillo J, Cattán V, Schoumacher M, Codony-Servat J, Giménez-Capitán A, Cantero F, et al. AURKB as a Target in Non-Small Cell Lung Cancer With Acquired Resistance to Anti-EGFR Therapy. *Nat Commun* (2019) 10:1812. doi: 10.1038/s41467-019-09734-5
 34. Park JH, Choi YJ, Kim SY, Lee J-E, Sung KJ, Park S, et al. Activation of the IGF1R Pathway Potentially Mediates Acquired Resistance to Mutant-Selective 3rd-Generation EGF Receptor Tyrosine Kinase Inhibitors in Advanced Non-Small Cell Lung Cancer. *Oncotarget* (2016) 7:22005–15. doi: 10.18632/oncotarget.8013
 35. Hayakawa D, Takahashi F, Mitsuishi Y, Tajima K, Hidayat M, Winardi W, et al. Activation of Insulin-Like Growth Factor-1 Receptor Confers Acquired Resistance to Osimertinib in Non-Small Cell Lung Cancer With *EGFR* T790M Mutation. *Thorac Cancer* (2020) 11:140–9. doi: 10.1111/1759-7714.13255
 36. Manabe T, Yasuda H, Terai H, Kagiwada H, Hamamoto J, Ebisudani T, et al. IGF2 Autocrine-Mediated IGF1R Activation Is a Clinically Relevant Mechanism of Osimertinib Resistance in Lung Cancer. *Mol Cancer Res* (2020) 18:549–59. doi: 10.1158/1541-7786.MCR-19-0956
 37. Weng C-H, Chen L-Y, Lin Y-C, Shih J-Y, Lin Y-C, Tseng R-Y, et al. Epithelial-Mesenchymal Transition (EMT) Beyond EGFR Mutations Per Se Is a Common Mechanism for Acquired Resistance to EGFR TKI. *Oncogene* (2019) 38:455–68. doi: 10.1038/s41388-018-0454-2
 38. Fukuda K, Takeuchi S, Arai S, Kita K, Tanimoto A, Nishiyama A, et al. Glycogen Synthase Kinase-3 Inhibition Overcomes Epithelial-Mesenchymal Transition-Associated Resistance to Osimertinib in *EGFR*-Mutant Lung Cancer. *Cancer Sci* (2020) 111:2374–84. doi: 10.1111/cas.14454
 39. Takahashi A, Seike M, Chiba M, Takahashi S, Nakamichi S, Matsumoto M, et al. Ankyrin Repeat Domain 1 Overexpression Is Associated With Common Resistance to Afatinib and Osimertinib in *EGFR*-Mutant Lung Cancer. *Sci Rep* (2018) 8:14896. doi: 10.1038/s41598-018-33190-8
 40. Verusingam ND, Chen Y-C, Lin H-F, Liu C-Y, Lee M-C, Lu K-H, et al. Generation of Osimertinib-Resistant Cells From Epidermal Growth Factor Receptor L858R/T790M Mutant Non-Small Cell Lung Carcinoma Cell Line. *J Chin Med Assoc* (2021) 84:248–54. doi: 10.1097/JCMA.0000000000000438
 41. Crystal AS, Shaw AT, Sequist LV, Friboulet L, Niederst MJ, Lockerman EL, et al. Patient-Derived Models of Acquired Resistance can Identify Effective Drug Combinations for Cancer. *Science* (2014) 346:1480–6. doi: 10.1126/science.1254721
 42. Raouf S, Mulford IJ, Frisco-Cabanos H, Nangia V, Timonina D, Labrot E, et al. Targeting FGFR Overcomes EMT-Mediated Resistance in *EGFR* Mutant Non-Small Cell Lung Cancer. *Oncogene* (2019) 38:6399–413. doi: 10.1038/s41388-019-0887-2
 43. Codony-Servat J, Codony-Servat C, Cardona AF, Giménez-Capitán A, Drozdowskyj A, Berenguer J, et al. Cancer Stem Cell Biomarkers in *EGFR*-Mutation-Positive Non-Small-Cell Lung Cancer. *Clin Lung Cancer* (2019) 20:167–77. doi: 10.1016/j.clcc.2019.02.005
 44. Shi P, Oh Y-T, Deng L, Zhang G, Qian G, Zhang S, et al. Overcoming Acquired Resistance to AZD9291, A Third-Generation EGFR Inhibitor, Through Modulation of MEK/ERK-Dependent Bim and Mcl-1 Degradation. *Clin Cancer Res* (2017) 23:6567–79. doi: 10.1158/1078-0432.CCR-17-1574
 45. Zang H, Qian G, Zong D, Fan S, Owonikoko TK, Ramalingam SS, et al. Overcoming Acquired Resistance of Epidermal Growth Factor Receptor-Mutant Non-Small Cell Lung Cancer Cells to Osimertinib by Combining Osimertinib With the Histone Deacetylase Inhibitor Panobinostat (LBH589). *Cancer* (2020) 126:2024–33. doi: 10.1002/cncr.32744
 46. Liu Z, Gao W. Synergistic Effects of Bcl-2 Inhibitors With AZD9291 on Overcoming the Acquired Resistance of AZD9291 in H1975 Cells. *Arch Toxicol* (2020) 94:3125–36. doi: 10.1007/s00204-020-02816-0
 47. Galvani E, Sun J, Leon LG, Sciarillo R, Narayan RS, Tjin Tham Sjin R, et al. NF- κ B Drives Acquired Resistance to a Novel Mutant-Selective EGFR Inhibitor. *Oncotarget* (2015) 6:42717–32. doi: 10.18632/oncotarget.3956
 48. Pan Y-H, Lin C-Y, Lu C-H, Li L, Wang Y-B, Chen H-Y, et al. Metformin Synergistically Enhances the Antitumor Activity of the Third-Generation EGFR-TKI CO-1686 in Lung Cancer Cells Through Suppressing NF- κ B Signaling. *Pan Et al. Clin Respir J* (2018) 12:2642–52. doi: 10.1111/crj.12970
 49. Shah KN, Bhatt R, Rotow J, Rohrberg J, Olivias V, Wang VE, et al. Aurora Kinase A Drives the Evolution of Resistance to Third-Generation EGFR Inhibitors in Lung Cancer. *Nat Med* (2019) 25:111–8. doi: 10.1038/s41591-018-0264-7
 50. Qin Q, Li X, Liang X, Zeng L, Wang J, Sun L, et al. CDK4/6 Inhibitor Palbociclib Overcomes Acquired Resistance to Third-Generation EGFR Inhibitor Osimertinib in Non-Small Cell Lung Cancer (NSCLC). *Thorac Cancer* (2020) 11:2389–97. doi: 10.1111/1759-7714.13521
 51. Tang Z, Su M-X, Guo X, Jiang X-M, Jia L, Chen X, et al. Increased Expression of IRE1 α Associates With the Resistant Mechanism of Osimertinib (AZD9291)-Resistant Non-Small Cell Lung Cancer HCC827/OSIR Cells. *ACAMC* (2018) 18:550–5. doi: 10.2174/1871520617666170719155517
 52. Wang J, Wang B, Chu H, Yao Y. Intrinsic Resistance to EGFR Tyrosine Kinase Inhibitors in Advanced Non-Small-Cell Lung Cancer With Activating EGFR Mutations. *Oncol Targets Ther* (2016) 9:3711–26. doi: 10.2147/OTT.S106399
 53. Ramirez M, Rajaram S, Steininger RJ, Osipchuk D, Roth MA, Morinishi LS, et al. Diverse Drug-Resistance Mechanisms can Emerge From Drug-Tolerant Cancer Persister Cells. *Nat Commun* (2016) 7:10690. doi: 10.1038/ncomms10690
 54. Oxnard GR. The Cellular Origins of Drug Resistance in Cancer. *Nat Med* (2016) 22:232–4. doi: 10.1038/nm.4058
 55. Hata AN, Niederst MJ, Archibald HL, Gomez-Caraballo M, Siddiqui FM, Mulvey HE, et al. Tumor Cells can Follow Distinct Evolutionary Paths to Become Resistant to Epidermal Growth Factor Receptor Inhibition. *Nat Med* (2016) 22:262–9. doi: 10.1038/nm.4040
 56. De Conti G, Dias MH, Bernards R. Fighting Drug Resistance Through the Targeting of Drug-Tolerant Persister Cells. *Cancers* (2021) 13:1118. doi: 10.3390/cancers13051118
 57. Sharma SV, Lee DY, Li B, Quinlan MP, Takahashi F, Maheswaran S, et al. A Chromatin-Mediated Reversible Drug-Tolerant State in Cancer Cell Subpopulations. *Cell* (2010) 141:69–80. doi: 10.1016/j.cell.2010.02.027
 58. Wang R, Yamada T, Kita K, Taniguchi H, Arai S, Fukuda K, et al. Transient IGF-1R Inhibition Combined With Osimertinib Eradicates AXL-Low Expressing EGFR Mutated Lung Cancer. *Nat Commun* (2020) 11:4607. doi: 10.1038/s41467-020-18442-4
 59. Taniguchi H, Yamada T, Wang R, Tanimura K, Adachi Y, Nishiyama A, et al. AXL Confers Intrinsic Resistance to Osimertinib and Advances the Emergence of Tolerant Cells. *Nat Commun* (2019) 10:259. doi: 10.1038/s41467-018-08074-0
 60. Song K-A, Hosono Y, Turner C, Jacob S, Lochmann TL, Murakami Y, et al. Increased Synthesis of MCL-1 Protein Underlies Initial Survival of *EGFR*-Mutant Lung Cancer to EGFR Inhibitors and Provides a Novel Drug Target. *Clin Lung Cancer* (2018) 24:5658–72. doi: 10.1158/1078-0432.CCR-18-0304
 61. Tanaka K, Yu HA, Yang S, Han S, Selcuklu SD, Kim K, et al. Targeting Aurora B Kinase Prevents and Overcomes Resistance to EGFR Inhibitors in

- Lung Cancer by Enhancing BIM- and PUMA-Mediated Apoptosis. *Cancer Cell* (2021) 39:1245–61.e6. doi: 10.1016/j.ccell.2021.07.006
62. Yang Z, Yang N, Ou Q, Xiang Y, Jiang T, Wu X, et al. Investigating Novel Resistance Mechanisms to Third-Generation EGFR Tyrosine Kinase Inhibitor Osimertinib in Non-Small Cell Lung Cancer Patients. *Clin Cancer Res* (2018) 24:3097–107. doi: 10.1158/1078-0432.CCR-17-2310
 63. Chen T, Luo J, Gu Y, Huang J, Luo Q, Yang Y. Comprehensive Analysis of Circular RNA Profiling in AZD9291-Resistant Non-Small Cell Lung Cancer Cell Lines. *Thorac Cancer* (2019) 10:930–41. doi: 10.1111/1759-7714.13032
 64. Takahashi S, Noro R, Seike M, Zeng C, Matsumoto M, Yoshikawa A, et al. Long Non-Coding RNA CRNDE Is Involved in Resistance to EGFR Tyrosine Kinase Inhibitor in EGFR-Mutant Lung Cancer via Eif4a3/MUC1/EGFR Signaling. *IJMS* (2021) 22:4005. doi: 10.3390/ijms22084005
 65. Hayford CE, Tyson DR, Robbins CJ, Frick PL, Quaranta V, Harris LA. An *In Vitro* Model of Tumor Heterogeneity Resolves Genetic, Epigenetic, and Stochastic Sources of Cell State Variability. *PLoS Biol* (2021) 19:e3000797. doi: 10.1371/journal.pbio.3000797
 66. Klempner S, Mehta P, Schrock A, Ali S, Ou S-HI. Cis-Oriented Solvent-Front EGFR G796S Mutation in Tissue and ctDNA in a Patient Progressing on Osimertinib: A Case Report and Review of the Literature. *LCTT* (2017) 8:241–7. doi: 10.2147/LCTT.S147129
 67. Fassunke J, Müller F, Keul M, Michels S, Dammert MA, Schmitt A, et al. Overcoming EGFRG724S-Mediated Osimertinib Resistance Through Unique Binding Characteristics of Second-Generation EGFR Inhibitors. *Nat Commun* (2018) 9:4655. doi: 10.1038/s41467-018-07078-0
 68. Nishino M, Suda K, Kobayashi Y, Ohara S, Fujino T, Koga T, et al. Effects of Secondary EGFR Mutations on Resistance Against Upfront Osimertinib in Cells With EGFR-Activating Mutations *In Vitro*. *Lung Cancer* (2018) 126:149–55. doi: 10.1016/j.lungcan.2018.10.026
 69. Peled N, Roisman LC, Miron B, Pfeffer R, Lanman RB, Ilouze M, et al. Subclonal Therapy by Two EGFR TKIs Guided by Sequential Plasma Cell-Free DNA in EGFR-Mutated Lung Cancer. *J Thorac Oncol* (2017) 12:e81–4. doi: 10.1016/j.jtho.2017.02.023
 70. Niederst MJ, Hu H, Mulvey HE, Lockerman EL, Garcia AR, Piotrowska Z, et al. The Allelic Context of the C797S Mutation Acquired Upon Treatment With Third-Generation EGFR Inhibitors Impacts Sensitivity to Subsequent Treatment Strategies. *Clin Cancer Res* (2015) 21:3924–33. doi: 10.1158/1078-0432.CCR-15-0560
 71. Wang Z, Yang J-J, Huang J, Ye J-Y, Zhang X-C, Tu H-Y, et al. Lung Adenocarcinoma Harboring EGFR T790M and *In Trans* C797S Responds to Combination Therapy of First- and Third-Generation EGFR TKIs and Shifts Allelic Configuration at Resistance. *J Thorac Oncol* (2017) 12:1723–7. doi: 10.1016/j.jtho.2017.06.017
 72. Arulananda S, Do H, Musafir A, Mitchell P, Dobrovic A, John T. Combination Osimertinib and Gefitinib in C797S and T790M EGFR-Mutated Non-Small Cell Lung Cancer. *J Thorac Oncol* (2017) 12:1728–32. doi: 10.1016/j.jtho.2017.08.006
 73. Rangachari D, To C, Shpilsky JE, VanderLaan PA, Kobayashi SS, Mushajiang M, et al. EGFR-Mutated Lung Cancers Resistant to Osimertinib Through EGFR C797S Respond to First-Generation Reversible EGFR Inhibitors But Eventually Acquire EGFR T790M/C797S in Preclinical Models and Clinical Samples. *J Thorac Oncol* (2019) 14:1995–2002. doi: 10.1016/j.jtho.2019.07.016
 74. Robichaux JP, Le X, Vijayan RSK, Hicks JK, Heeke S, Elamin YY, et al. Structure-Based Classification Predicts Drug Response in EGFR-Mutant NSCLC. *Nature* (2021) 597:732–7. doi: 10.1038/s41586-021-03898-1
 75. Papadimitrakopoulou VA, Wu Y-L, Han J-Y, Ahn M-J, Ramalingam SS, John T, et al. Analysis of Resistance Mechanisms to Osimertinib in Patients With EGFR T790M Advanced NSCLC From the AURA3 Study. *Ann Oncol* (2018) 29:viii741. doi: 10.1093/annonc/mdy424.064
 76. Ramalingam SS, Cheng Y, Zhou C, Ohe Y, Imamura F, Cho BC, et al. Mechanisms of Acquired Resistance to First-Line Osimertinib: Preliminary Data From the Phase III FLAURA Study. *Ann Oncol* (2018) 29:viii740. doi: 10.1093/annonc/mdy424.063
 77. Schoenfeld AJ, Chan JM, Kubota D, Sato H, Rizvi H, Daneshbod Y, et al. Tumor Analyses Reveal Squamous Transformation and Off-Target Alterations As Early Resistance Mechanisms to First-Line Osimertinib in EGFR -Mutant Lung Cancer. *Clin Cancer Res* (2020) 26:2654–63. doi: 10.1158/1078-0432.CCR-19-3563
 78. Le X, Puri S, Negrao MV, Nilsson MB, Robichaux J, Boyle T, et al. Landscape of EGFR-Dependent and -Independent Resistance Mechanisms to Osimertinib and Continuation Therapy Beyond Progression in EGFR -Mutant NSCLC. *Clin Cancer Res* (2018) 24:6195–203. doi: 10.1158/1078-0432.CCR-18-1542
 79. Ma Y, Zheng X, Zhao H, Fang W, Zhang Y, Ge J, et al. First-In-Human Phase I Study of AC0010, a Mutant-Selective EGFR Inhibitor in Non-Small Cell Lung Cancer: Safety, Efficacy, and Potential Mechanism of Resistance. *J Thorac Oncol* (2018) 13:968–77. doi: 10.1016/j.jtho.2018.03.025
 80. Ho C-C, Liao W-Y, Lin C-A, Shih J-Y, Yu C-J, Yang J-C-H. Acquired BRAF V600E Mutation as Resistant Mechanism After Treatment With Osimertinib. *J Thorac Oncol* (2017) 12:567–72. doi: 10.1016/j.jtho.2016.11.2231
 81. Minari R, Bordi P, La Monica S, Squadrilli A, Leonetti A, Bottarelli L, et al. Concurrent Acquired BRAF V600E Mutation and MET Amplification as Resistance Mechanism of First-Line Osimertinib Treatment in a Patient With EGFR-Mutated NSCLC. *J Thorac Oncol* (2018) 13:e89–91. doi: 10.1016/j.jtho.2018.03.013
 82. Bearz A, De Carlo E, Doliana R, Schiappacassi M. Acquired BRAF V600E Mutation as Resistant Mechanism After Treatment With Third-Generation EGFR Tyrosine Kinase Inhibitor. *J Thorac Oncol* (2017) 12:e181–2. doi: 10.1016/j.jtho.2017.07.017
 83. Blakely CM, Watkins TBK, Wu W, Gini B, Chabon JJ, McCoach CE, et al. Evolution and Clinical Impact of Co-Occurring Genetic Alterations in Advanced-Stage EGFR-Mutant Lung Cancers. *Nat Genet* (2017) 49:1693–704. doi: 10.1038/ng.3990
 84. Fang X, Gu P, Zhou C, Liang A, Ren S, Liu F, et al. β -Catenin Overexpression Is Associated With Gefitinib Resistance in Non-Small Cell Lung Cancer Cells. *Pulm Pharmacol Ther* (2014) 28:41–8. doi: 10.1016/j.pupt.2013.05.005
 85. Togashi Y, Hayashi H, Terashima M, de Velasco MA, Sakai K, Fujita Y, et al. Inhibition of β -Catenin Enhances the Anticancer Effect of Irreversible EGFR-TKI in EGFR-Mutated Non-Small-Cell Lung Cancer With a T790M Mutation. *J Thorac Oncol* (2015) 10:93–101. doi: 10.1097/JTO.0000000000000353
 86. Xu W, Tang W, Li T, Zhang X, Sun Y. Overcoming Resistance to AC0010, a Third Generation of EGFR Inhibitor, by Targeting C-MET and BCL-2. *Neoplasia* (2019) 21:41–51. doi: 10.1016/j.neo.2018.11.004
 87. Ou S-HI, Cui J, Schrock AB, Goldberg ME, Zhu VW, Albacker L, et al. Emergence of Novel and Dominant Acquired EGFR Solvent-Front Mutations at Gly796 (G796S/R) Together With C797S/R and L792F/H Mutations in One EGFR (L858R/T790M) NSCLC Patient Who Progressed on Osimertinib. *Lung Cancer* (2017) 108:228–31. doi: 10.1016/j.lungcan.2017.04.003
 88. Sequist LV, Han J-Y, Ahn M-J, Cho BC, Yu H, Kim S-W, et al. Osimertinib Plus Savolitinib in Patients With EGFR Mutation-Positive, MET-Amplified, Non-Small-Cell Lung Cancer After Progression on EGFR Tyrosine Kinase Inhibitors: Interim Results From a Multicentre, Open-Label, Phase 1b Study. *Lancet Oncol* (2020) 21:373–86. doi: 10.1016/S1470-2045(19)30785-5
 89. Oxnard GR, Yang J-C-H, Yu H, Kim S-W, Saka H, Horn L, et al. TATTON: A Multi-Arm, Phase Ib Trial of Osimertinib Combined With Selumetinib, Savolitinib, or Durvalumab in EGFR-Mutant Lung Cancer. *Ann Oncol* (2020) 31:507–16. doi: 10.1016/j.annonc.2020.01.013
 90. Zhang Z, Lee JC, Lin L, Olivas V, Au V, LaFramboise T, et al. Activation of the AXL Kinase Causes Resistance to EGFR-Targeted Therapy in Lung Cancer. *Nat Genet* (2012) 44:852–60. doi: 10.1038/ng.2330
 91. Mudduluru G, Leupold JH, Stroebel P, Allgayer H. PMA Up-Regulates the Transcription of Axl by AP-1 Transcription Factor Binding to TRE Sequences via the MAPK Cascade in Leukaemia Cells. *Biol Cell* (2011) 103:21–33. doi: 10.1042/BC20100094
 92. Liu Y-N, Chang T-H, Tsai M-F, Wu S-G, Tsai T-H, Chen H-Y, et al. IL-8 Confers Resistance to EGFR Inhibitors by Inducing Stem Cell Properties in Lung Cancer. *Oncotarget* (2015) 6:10415–31. doi: 10.18632/oncotarget.3389
 93. Karachaliou N, Chaib I, Cardona AF, Berenguer J, Bracht JWP, Yang J, et al. Common Co-Activation of AXL and CDCP1 in EGFR-Mutation-Positive Non-Small Cell Lung Cancer Associated With Poor Prognosis. *EBioMedicine* (2018) 29:112–27. doi: 10.1016/j.ebiom.2018.02.001

94. Peled N, Wynnes MW, Ikeda N, Ohira T, Yoshida K, Qian J, et al. Insulin-Like Growth Factor-1 Receptor (IGF-1R) as a Biomarker for Resistance to the Tyrosine Kinase Inhibitor Gefitinib in Non-Small Cell Lung Cancer. *Cell Oncol (Dordr)* (2013) 36:277–88. doi: 10.1007/s13402-013-0133-9
95. Yeo CD, Park KH, Park CK, Lee SH, Kim SJ, Yoon HK, et al. Expression of Insulin-Like Growth Factor 1 Receptor (IGF-1R) Predicts Poor Responses to Epidermal Growth Factor Receptor (EGFR) Tyrosine Kinase Inhibitors in non-Small Cell Lung Cancer Patients Harboring Activating EGFR Mutations. *Lung Cancer* (2015) 87:311–7. doi: 10.1016/j.lungcan.2015.01.004
96. Cortot AB, Repellin CE, Shimamura T, Capelletti M, Zejnullahu K, Ercan D, et al. Resistance to Irreversible EGF Receptor Tyrosine Kinase Inhibitors Through a Multistep Mechanism Involving the IGF1R Pathway. *Cancer Res* (2013) 73:834–43. doi: 10.1158/0008-5472.CAN-12-2066
97. Aghdassi A, Sandler M, Guenther A, Mayerle J, Behn C-O, Heidecke C-D, et al. Recruitment of Histone Deacetylases HDAC1 and HDAC2 by the Transcriptional Repressor ZEB1 Downregulates E-Cadherin Expression in Pancreatic Cancer. *Gut* (2012) 61:439–48. doi: 10.1136/gutjnl-2011-300060
98. Brabletz S, Brabletz T. The ZEB/miR-200 Feedback Loop—A Motor of Cellular Plasticity in Development and Cancer? *EMBO Rep* (2010) 11:670–7. doi: 10.1038/embor.2010.117
99. Vijay GV, Zhao N, Den Hollander P, Toneff MJ, Joseph R, Pietila M, et al. Gsk3 β Regulates Epithelial-Mesenchymal Transition and Cancer Stem Cell Properties in Triple-Negative Breast Cancer. *Breast Cancer Res* (2019) 21:37. doi: 10.1186/s13058-019-1125-0
100. Feldker N, Ferrazzi F, Schuhwerk H, Widholz SA, Guenther K, Frisch I, et al. Genome-Wide Cooperation of EMT Transcription Factor ZEB 1 With YAP and AP -1 in Breast Cancer. *EMBO J* (2020) 39(17):e103209. doi: 10.15252/embj.2019103209
101. DeNardo DG, Barreto JB, Andreu P, Vasquez L, Tawfik D, Kolhatkar N, et al. CD4(+) T Cells Regulate Pulmonary Metastasis of Mammary Carcinomas by Enhancing Protumor Properties of Macrophages. *Cancer Cell* (2009) 16:91–102. doi: 10.1016/j.ccr.2009.06.018
102. Kim TM, Song A, Kim D-W, Kim S, Ahn Y-O, Keam B, et al. Mechanisms of Acquired Resistance to AZD9291. *J Thorac Oncol* (2015) 10:1736–44. doi: 10.1097/JTO.0000000000000688
103. Arasada RR, Amann JM, Rahman MA, Huppert SS, Carbone DP. EGFR Blockade Enriches for Lung Cancer Stem-like Cells Through Notch3-Dependent Signaling. *Cancer Res* (2014) 74:5572–84. doi: 10.1158/0008-5472.CAN-13-3724
104. Codony-Servat C, Codony-Servat J, Karachaliou N, Molina MA, Chaib I, Ramirez JL, et al. Activation of Signal Transducer and Activator of Transcription 3 (STAT3) Signaling in EGFR Mutant non-Small-Cell Lung Cancer (NSCLC). *Oncotarget* (2017) 8:47305–16. doi: 10.18632/oncotarget.17625
105. Goto N, Ueo T, Fukuda A, Kawada K, Sakai Y, Miyoshi H, et al. Distinct Roles of HES1 in Normal Stem Cells and Tumor Stem-Like Cells of the Intestine. *Cancer Res* (2017) 77:3442–54. doi: 10.1158/0008-5472.CAN-16-3192
106. Bousquet Mur E, Bernardo S, Papon L, Mancini M, Fabbriozio E, Goussard M, et al. Notch Inhibition Overcomes Resistance to Tyrosine Kinase Inhibitors in EGFR-Driven Lung Adenocarcinoma. *J Clin Invest* (2019) 130:612–24. doi: 10.1172/JCI126896
107. Faber AC, Corcoran RB, Ebi H, Sequist LV, Waltman BA, Chung E, et al. BIM Expression in Treatment-Naïve Cancers Predicts Responsiveness to Kinase Inhibitors. *Cancer Discov* (2011) 1:352–65. doi: 10.1158/2159-8290.CD-11-0106
108. Ley R, Balmanno K, Hadfield K, Weston C, Cook SJ. Activation of the ERK1/2 Signaling Pathway Promotes Phosphorylation and Proteasome-Dependent Degradation of the BH3-Only Protein, Bim. *J Biol Chem* (2003) 278:18811–6. doi: 10.1074/jbc.M301010200
109. Domina AM, Vrana JA, Gregory MA, Hann SR, Craig RW. MCL1 is Phosphorylated in the PEST Region and Stabilized Upon ERK Activation in Viable Cells, and at Additional Sites With Cytotoxic Okadaic Acid or Taxol. *Oncogene* (2004) 23:5301–15. doi: 10.1038/sj.onc.1207692
110. Bertino EM, Gentzler RD, Clifford S, Kolesar J, Muzikansky A, Haura EB, et al. Phase IB Study of Osimertinib in Combination With Navitoclax in EGFR -Mutant NSCLC Following Resistance to Initial EGFR Therapy (ETCTN 9903). *Clin Cancer Res* (2021) 27:1604–11. doi: 10.1158/1078-0432.CCR-20-4084
111. Shi P, Zhang S, Zhu L, Qian G, Ren H, Ramalingam SS, et al. The Third-Generation EGFR Inhibitor, Osimertinib, Promotes C-FLIP Degradation, Enhancing Apoptosis Including TRAIL-Induced Apoptosis in NSCLC Cells With Activating EGFR Mutations. *Trans Oncol* (2019) 12:705–13. doi: 10.1016/j.tranon.2019.02.006
112. Bivona TG, Hieronymus H, Parker J, Chang K, Taron M, Rosell R, et al. FAS and NF- κ B Signalling Modulate Dependence of Lung Cancers on Mutant EGFR. *Nature* (2011) 471:523–6. doi: 10.1038/nature09870
113. Moiseeva O, Deschênes-Simard X, St-Germain E, Igelmann S, Huot G, Cadar AE, et al. Metformin Inhibits the Senescence-Associated Secretory Phenotype by Interfering With IKK / NF - κ B Activation. *Aging Cell* (2013) 12:489–98. doi: 10.1111/acer.12075
114. Tang A, Gao K, Chu L, Zhang R, Yang J, Zheng J. Aurora Kinases: Novel Therapy Targets in Cancers. *Oncotarget* (2017) 8:23937–54. doi: 10.18632/oncotarget.14893
115. Shi R, Radulovich N, Ng C, Liu N, Notsuda H, Cabanero M, et al. Organoid Cultures as Preclinical Models of Non-Small Cell Lung Cancer. *Clin Cancer Res* (2020) 26:1162–74. doi: 10.1158/1078-0432.CCR-19-1376

Conflict of Interest: RP received a grant from AstraZeneca to conduct a project in the laboratory but it was not used to fund this review.

The remaining authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2022 Mahfoudhi, Ricordel, Lecuyer, Mouric, Lena and Pedoux. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.