



The Likelihood of Preventing Respiratory Exacerbations in Children and Adolescents with either Chronic Suppurative Lung Disease or Bronchiectasis

Kerry-Ann F O'Grady^{1*} and Keith Grimwood²

¹Institute of Health and Biomedical Innovation, Queensland University of Technology, South Brisbane, QLD, Australia,

²Menzies Health Research Institute Queensland, Griffith University, Gold Coast Health, Southport, QLD, Australia

OPEN ACCESS

Edited by:

Francesca Santamaria,
University of Naples Federico II, Italy

Reviewed by:

Nicola Ullmann,
Bambino Gesù Ospedale Pediatrico
(IRCCS), Italy
Paolo Bottau,
Imola Hospital, Italy
Bülent Taner Karadağ,
Marmara University, Turkey

*Correspondence:

Kerry-Ann F O'Grady
kerryann.ogradey@qut.edu.au

Specialty section:

This article was submitted to
Pediatric Pulmonology,
a section of the journal
Frontiers in Pediatrics

Received: 13 December 2016

Accepted: 08 March 2017

Published: 24 March 2017

Citation:

O'Grady KF and Grimwood K (2017)
The Likelihood of Preventing
Respiratory Exacerbations in Children
and Adolescents with either Chronic
Suppurative Lung Disease or
Bronchiectasis.
Front. Pediatr. 5:58.
doi: 10.3389/fped.2017.00058

Chronic suppurative lung disease (CSLD) and bronchiectasis in children and adolescents are important causes of respiratory morbidity and reduced quality of life (QoL), also leading to subsequent premature death during adulthood. Acute respiratory exacerbations in pediatric CSLD and bronchiectasis are important markers of disease control clinically, given that they impact upon QoL and increase health-care-associated costs and can adversely affect future lung functioning. Preventing exacerbations in this population is, therefore, likely to have significant individual, familial, societal, and health-sector benefits. In this review, we focus on therapeutic interventions, such as drugs (antibiotics, mucolytics, hyperosmolar agents, bronchodilators, corticosteroids, non-steroidal anti-inflammatory agents), vaccines and physiotherapy, and care-planning, such as post-hospitalization management and health promotion strategies, including exercise, diet, and reducing exposure to environmental toxicants. The review identified a conspicuous lack of moderate or high-quality evidence for preventing respiratory exacerbations in children and adolescents with CSLD or bronchiectasis. Given the short- and long-term impact of exacerbations upon individuals, their families, and society as a whole, large studies addressing interventions at the primary and tertiary prevention phases are required. This research must include children and adolescents in both developing and developed countries and address long-term health outcomes.

Keywords: chronic suppurative lung disease, bronchiectasis, respiratory exacerbations, children, prevention

Abbreviations: AUC, area under the curve; CF, cystic fibrosis; cHRCT, chest high-resolution computed tomography; CI, confidence interval; COPD, chronic obstructive pulmonary disease; CRP, C-reactive protein; CSLD, chronic suppurative lung disease; FEV₁%, forced expiratory volume in one-second percentage; NPV, negative predictive value; NSAIDs, non-steroidal anti-inflammatory drugs; OR, odds ratio; OTC, over-the-counter; PBB, protracted bacterial bronchitis; PPV, positive predictive value; QoL, quality of life; RCT, randomized controlled trial; rhDNase, recombinant human DNase; RR, relative risk; TSANZ, Thoracic Society of Australia and New Zealand; US, United States.

INTRODUCTION

Chronic (>4 weeks duration) wet cough in children implies increased airway secretions and lower airway infection (1). It is associated with often unrecognized morbidity and impaired quality of life (QoL) scores (2, 3). Within Australia, the most common cause of chronic wet cough in children is protracted bacterial bronchitis (PBB), followed by bronchiectasis (4). These two entities are at the opposite ends of a spectrum of overlapping conditions characterized by chronic wet cough, airway neutrophilia, and high bacterial densities in bronchoalveolar lavage (BAL) fluid cultures (5). They also share a common risk factor of impaired airway clearance facilitating microbial colonization of the lower airways (6), while their lower airway microbiota are dominated by strains of non-typeable *Haemophilus influenzae* (NTHi), *Streptococcus pneumoniae*, and *Moraxella catarrhalis* (1, 7–15).

Protracted bacterial bronchitis is defined as an illness with an isolated wet cough >4 weeks duration that resolves with a prolonged 2–4 weeks of appropriate oral antibiotic treatment, usually amoxicillin–clavulanate, and where other pointers suggestive of an alternative cause of cough are absent (16). It is a relatively newly defined diagnosis, but likely represents illnesses previously termed as chronic bronchitis, persistent endobronchial infection, or pre-bronchiectasis (16, 17).

By contrast, bronchiectasis has many underlying causes and is diagnosed when there is a chronic wet cough, which has a variable response to even prolonged courses of oral antibiotics, frequent respiratory exacerbations, and radiographic demonstration of irreversible dilatation of one or more bronchi (18). These children often have other accompanying respiratory symptoms and signs too, including growth failure, exertional dyspnea, digital clubbing, chest wall deformity, hyperinflation, and adventitial sounds on chest auscultation. The most common etiologies of bronchiectasis in children include post-infectious causes following recurrent and/or severe lower respiratory infections in early life, cystic fibrosis (CF), primary ciliary dyskinesia, immunodeficiency, retained foreign body, and chronic aspiration (12, 18).

Between the diagnostic spectrum of PBB and bronchiectasis are children with chronic suppurative lung disease (CSLD) who have the clinical symptoms and signs suggestive of bronchiectasis but lack the chest high-resolution computed tomography (cHRCT) scan evidence to support a diagnosis of bronchiectasis, such as an increased broncho-arterial ratio and absence of bronchial tapering while progressing toward the lung periphery (18, 19). Some of those categorized as having CSLD live in remote Indigenous communities in Australia and Alaska where undergoing cHRCT scans is difficult logistically and may be delayed until late in adolescence. Consequently, some of the younger children with CSLD may have undiagnosed bronchiectasis. In light of this and as management is the same for both diagnostic entities (19), this review focuses upon preventing respiratory exacerbations in children with either CSLD or bronchiectasis.

While published data are currently limited, recurrent episodes of PBB can also occur. In a study of 106 children with PBB followed for 2 years (20), 43.5% had >3 episodes per year and bronchiectasis was diagnosed subsequently in 8.1% of study

participants (20). *H. influenzae* infection in the lower airways and recurrent episodes of PBB were independently associated with a future diagnosis of bronchiectasis [odds ratio (OR) 7.60, 95% confidence interval (CI) 1.53, 37.79 and OR 11.48, 95% CI 2.33, 56.50, respectively] (20). Nevertheless, children with recurrent PBB are not included in this review as the point where they progress to CSLD or bronchiectasis is yet to be determined, especially if they respond promptly to antibiotics and do not have additional symptoms or signs of respiratory disease. While CF is an important cause of chronic wet cough and bronchiectasis, children with CF are also excluded here given differences in the type of disease and concerns that the findings of studies in this patient population should not be extrapolated to those with bronchiectasis unrelated to CF (21).

ACUTE PULMONARY EXACERBATIONS

Acute pulmonary exacerbations are associated with disease severity (7), and although there are no direct data for CSLD or bronchiectasis in children, exacerbations are likely to contribute to a decline in lung function. Determinants of accelerated lung function decline in adults with bronchiectasis are frequency of hospitalized exacerbations, increased systemic inflammatory markers, and infection with *Pseudomonas aeruginosa* (22). Among other factors, increased mortality risk is associated with degree of lung function impairment (23). In children, studies from Australia and the United Kingdom reported that children with normal lung function at the time bronchiectasis was diagnosed maintained normal lung function 5 years later (24–26). Those children with poor lung function at diagnosis, although substantially improved, were still likely to have poor lung function 5 years afterward with the only significant predictor of pulmonary decline being the frequency of hospitalized exacerbations (25). With each hospitalized exacerbation, the forced expiratory volume in 1-s percentage (FEV₁%) predicted decreased by 1.95% adjusted for time (25). Thus, interventions that minimize exacerbation frequencies are likely to be important in the overall management of bronchiectasis (19). In addition, preventing exacerbations will potentially reduce the associated economic and social costs. In the United States, between 1993 and 2006, the median cost for adult inpatient care was US\$7,827 (27). Hospitalization rates increased significantly over the study time period with an annual percentage increase of 2.4% for males and 3.0% for females (27). At a New Zealand Hospital in 2004, half of the service's pediatric bronchiectasis patients required at least one hospital admission per year (range 1–10) for exacerbations (28). The mean length of stay was 7 days (range 1–25). Eight percent of these children had regular three to four monthly admissions for 14 days to maintain respiratory status at a mean cost in 2004 of US\$5,492 (not including theater time or costs for inserting peripherally inserted central catheter vascular lines) for each hospitalization (28).

DEFINITION

As with the paucity of data on CSLD and bronchiectasis in children in general (29), there is a dearth of literature defining an acute

exacerbation. In adults, acute exacerbations of chronic obstructive pulmonary disease (COPD) are characterized by worsening dyspnea and increasing sputum volume and/or purulence (30). In adult bronchiectasis, exacerbations have similar features to that of COPD with increased cough frequency and sputum volume or purulence and are often associated with culturing respiratory bacterial pathogens in sputum (31). In research settings, particularly studies with exacerbations as the primary outcome, the definition may also include the need for hospitalization and intravenous antibiotics (32).

Clinical trials in children are currently using (33–36) similar definitions to that used in adults, including variations in symptoms of increased cough, dyspnea, increased sputum volume or color intensity, new chest examination or radiographic findings, deterioration in FEV₁% predicted >10%, or hemoptysis. Not all definitions have, however, included the need for hospitalization or intravenous antibiotics.

In a study of 30 children with 115 exacerbations of bronchiectasis (25), increased cough frequency (88%) and change in cough character (67%) were the most frequent symptoms reported. Fever (28%), increased sputum volume (42%) and purulence (35%), and deteriorating chest auscultatory findings (58%) were also common, while changes in spirometry values compared to the stable state were not (25). Further work was undertaken from this cohort's data (81 exacerbations included) (37) to develop a standardized definition of an exacerbation in children with bronchiectasis. Wet cough and cough severity (score ≥ 2) over 72 h were the best predictors of an exacerbation [receiver operating characteristic area under the curve of 0.85 (95% CI 0.79, 0.92) and 0.84 (95% CI 0.77, 0.91), respectively] (37). Sputum color, dyspnea, hemoptysis, chest signs, and chest pain were considered minor criteria and the addition of serum C-reactive protein (CRP), amyloid-A, and interleukin-6 improved the specificity and positive predictive value of the definition (37). The authors subsequently classified symptoms and laboratory measures into three sets of criteria (major, minor, and laboratory—**Box 1**) and then devised three options that could be used to define an exacerbation in bronchiectasis (**Box 1**). A limitation of the study is that there was a lack of an objective gold standard for diagnosing an exacerbation in bronchiectasis and hence the criteria were assessed against a pediatric pulmonologist's definition that included change in symptoms and the need for additional treatment beyond the stable state. Further studies validating the definition in larger cohorts are now awaited.

ETIOLOGY

The exact cause of exacerbations in young children with CSLD or bronchiectasis is not well understood, and it is unclear whether these are new infections, a resurgence of chronic infection, or a combination of both (31, 38). Part of the problem is that despite their regular use in clinical practice, upper airway secretions collected by throat or cough swabs do not reliably predict organisms within the lower airways (9), especially as the potential pathogens of interest (NTHi, *S. pneumoniae*, and *M. catarrhalis*) are also found commonly in the upper airway spaces of healthy children

BOX 1 | Proposed criteria for defining a pulmonary exacerbation in children with bronchiectasis (37).

- (I) Major criteria
 - o Significant frequency of cough (median cough score ≥ 2) over 72 h.
 - o Wet cough for 72 h.
- (II) Minor criteria
 - o Sputum color ≥ 3 on BronkoTest™.
 - o Parent/child perceived breathlessness.
 - o Chest pain.
 - o Auscultatory crackles.
 - o Wheeze.
 - o Hypoxia (oxygen saturation $\leq 93\%$ by pulse oximetry).
- (III) Laboratory criteria
 - o CRP > 3 mg/L on high sensitive testing.
 - o Serum interleukin-6 > 2 ng/L.
 - o Serum amyloid-A > 5 mg/L.
 - o Raised peripheral blood neutrophil % (age appropriate).

The three combinations considered the best to define an exacerbation:

- (Option-A) *One major PLUS any one laboratory criteria* positive [sensitivity 63%, specificity 94%, AUC 0.784, $p < 0.001$; positive predictive value (PPV) 91%, negative predictive value (NPV) 71.6%], OR
- (Option-B) *Two major criteria* positive (sensitivity 92.6%, specificity 75.3%, AUC 0.84, $p < 0.001$; PPV 79%, NPV 91%), OR
- (Option-C) *One major PLUS any two minor criteria* positive (sensitivity 95%, specificity 75%, AUC 0.84, $p < 0.001$; PPV 78.5%, NPV 93.75%).

(39). Repeated bronchoscopies with multi-lobar BAL to collect lower airway specimens before, during, and following acute exacerbations in children too young to reliably expectorate sputum would be ideal, but impractical given their invasive nature, including the need for repeated sedation or general anesthesia. Hence the data required to confidently assign causality in research and patient care are limited.

Haemophilus influenzae, *S. pneumoniae*, and *M. catarrhalis* are frequently isolated at high densities ($\geq 10^4$ colony-forming units/mL) from BAL specimens collected from children and adults during an acute exacerbation, but mixed infections are also common (9, 13), and few studies have searched systematically for viruses. *P. aeruginosa* and non-tuberculous mycobacteria are uncommon in children with bronchiectasis and, when present, raises the possibility of undiagnosed CF while also being associated with more severe underlying bronchiectasis and comorbidities (13). A small study of 69 children with bronchiectasis that included 900 child-months of follow-up (40) identified at least one respiratory virus in nasopharyngeal aspirates in 48% of exacerbations, most commonly human rhinovirus (54% of virus-positive events). Children with virus-positive exacerbations were more likely to require hospitalization (59 vs 32.5%; $p = 0.02$), have fever (OR 3.1, 95% CI 1.2, 11.1), hypoxia (OR 25.5, 95% CI 2.0, 322.6), chest signs (OR 3.3, 95% CI 1.1, 10.2), and raised CRP (OR 4.7, 95% CI 1.7, 13.1), when compared with virus-negative exacerbations (40).

In a cross-sectional study of 245 children diagnosed with PBB or mild bronchiectasis undergoing bronchoscopy and BAL for clinical indication (median age 30 months) (41), a standard respiratory panel for viruses using polymerase chain reaction assays

was performed on all specimens. Human adenovirus (HAdV) was the most common virus detected, being identified in 40 children; influenza virus was detected in 3 children, parainfluenza virus in 12, respiratory syncytial virus in 11, and human metapneumovirus in 5 children (41). HAdV detection was more common in the young age groups ($p = 0.001$) and was positively associated with each of the three major bacterial pathogens (41). However, this association disappeared after adjustment for age and, given both the cross-sectional nature of the study with a lack of controls, a causal association between detecting these viruses and clinical illness at the time of bronchoscopy cannot be determined.

Similarly, there are very few studies examining the risk factors for exacerbations of CSLD or bronchiectasis. A study of 93 Indigenous children from Alaska and Australia with CSLD or bronchiectasis (42) reported 74% of children experienced >2 exacerbations over a 3-year period. In this study, the factors associated with recurrent episodes were young age (<3 years), hospitalization for an acute exacerbation in the first-year of life, and pneumonia or hospitalization for an acute exacerbation in the year preceding enrollment (42). Exacerbations are also more frequent in severe bronchiectasis, and one study of 111 children reported that intensive medical intervention reduced the annual exacerbation rate by 56%; however, children still experienced a mean of 2.9 episodes per year (43).

PREVENTION

Preventing exacerbations and reducing severity, are important goals in managing children with CSLD or bronchiectasis in order to maintain lung health and enhance their QoL (19). The cornerstone of CSLD management is a combination of airway clearance techniques and antibiotic therapy, with or without other therapies such as anti-inflammatory agents and bronchodilators (31, 38, 44). However, moderate or high-quality studies in children for either CSLD or bronchiectasis are limited (45), including those addressing broader approaches such as vaccines, health promotion, and chronic disease management strategies.

Pharmacological Agents

Antibiotics

A Cochrane review of antibiotic efficacy for preventing recurrent lower respiratory tract infections in high-risk children aged <12 years (46) identified a lack of evidence supporting prophylactic antibiotics and emphasized the need for high-quality trials to be conducted. That review included three studies involving children infected with human immunodeficiency virus, four with CF, and one each with children with sickle cell disease, cancer, and low birth weight neonates in a pediatric intensive care unit with underlying respiratory disorders (46). The only studies that addressed acute respiratory exacerbations were three of the CF studies; one (82 children) used ciprofloxacin in conjunction with colistin (polymyxin E) (47) and two (total of 312 children) used azithromycin (48, 49). The review excluded five studies [four in children with CF (50–53) and one involving children with either CSLD or bronchiectasis (54)] on the grounds that the children were already infected (51–54) or where infection could not be

discounted (50). Given the underlying nature of these endobronchial disorders, where infection has a central pathogenic role, the justification for excluding the five studies is unconvincing, while the basis for being currently infected by unidentified organisms is also unclear.

In the Australian study excluded from the review (54), 89 children with either CSLD or bronchiectasis were randomized to receive either azithromycin (30 mg/kg once-a-week) or placebo for up to 24 months. Children receiving azithromycin had significantly lower exacerbation rates (incidence rate ratio 0.50; 95% CI 0.35, 0.71) than the control group (54). However, children in the azithromycin group also developed significantly higher carriage of azithromycin-resistant bacteria (19 of 41, 46%) than those receiving placebo (4 of 37, 11%; $p = 0.002$) (54). Acquiring macrolide-resistant organisms was more prevalent in those who were poorly adherent (76%) than those complying with their therapy (52%; OR 2.94, 95% CI 1.23, 7.14) and post-intervention *Staphylococcus aureus* strains remained resistant to macrolides (55). These findings support a role for directly observed therapy in individuals and communities where adherence may be suboptimal, which will then help to ensure the benefits of long-term antibiotics, while minimizing potential harm from acquiring antibiotic-resistant organisms. A dose–response relationship between azithromycin use and an increase in macrolide-resistant strains of *S. pneumoniae* and *S. aureus* in the nasopharynx was also observed in a prospective study of 79 remote Australian children with CSLD or bronchiectasis conducted between 2004 and 2008 (56). These findings are consistent with those of the 2015 Cochrane review (57), which reported a threefold increase in antibiotic-resistant bacteria.

The 2015 Cochrane review of the efficacy of long-term antibiotics (≥ 4 weeks duration) in children and adults with bronchiectasis (57), identified only three studies (33, 58, 59) that included children (total 148 participants), and one did not report exacerbations (58). Of the remaining two, one used oral azithromycin and is described above (54), and the second used either oral roxithromycin or placebo in 25 children for 12 weeks (59). This latter study failed to demonstrate a statistically significant difference in exacerbation rates, likely a result of the small sample size and short duration of follow-up, although the point estimate of effect provided a weak signal toward a reduction (OR 0.16, 95% CI 0.01, 3.60) (59). The study did, however, report significantly lower sputum purulence scores in the antibiotic group compared with those receiving placebo (1.39 ± 0.6 vs 2.17 ± 0.72 ; $p < 0.01$) (59). With respect to short-term antibiotics (<4 weeks duration), a further Cochrane review (60) identified no studies in children and, therefore, concluded there was no evidence to support this approach for reducing the severity and frequency of exacerbations in children with bronchiectasis.

Expectorants, Mucolytics, and Mucokinetics

Retained mucus in the lower airways promotes bacterial growth, persistent airway inflammation, and bronchial wall injury (61). Hence reducing mucus retention is an important aspect of airway clearance techniques. Mucoactive agents include expectorants, mucolytics, and mucokinetic agents.

Expectorants aim to increase the volume of airway water or secretion in order to increase the effectiveness of cough (62) and include both over-the-counter (OTC) cough medications and inhaled hyperosmolar saline and mannitol. There is no good evidence for the use of OTC cough medications and expectorants in children (63), and concerns about adverse events, including some reports of infant deaths (64), led to several countries recommending OTC cough medications should not be given to young children. Inhaled hyperosmolar agents, including hyperosmolar saline and mannitol, potentially alter the physical properties of mucus by increasing water in the airway lumen and disrupting mucin networks (65). While there is evidence to support the role of hyperosmolar agents in children with CF (62), there are insufficient data to support their use in children with CSLD or bronchiectasis (45).

Mucolytics make mucus less viscous so that it can be more easily expectorated, and they are available in both oral and inhaled forms. This group of drugs includes *N*-acetylcysteine, ambroxol (3), sobrerol, carbocysteine, sobrerol, letosteine, cithiolone, iodinated glycerol, *N*-isobutyrylcysteine, myrtol and erdosteine, and recombinant human DNase (RhDNase) (62). The exact mechanism of action of most mucolytics is unclear, except for RhDNase that targets neutrophils at the site of infection in the lungs (66, 67). There have been no pediatric trials of mucolytics in children with CSLD or bronchiectasis (66). Moreover, RhDNase is not recommended in either children or adults with bronchiectasis given a study in adults reporting its negative effects on lung function and exacerbation rates (68).

Mucokinetics increase the effectiveness of cough, either by increasing expiratory cough airflow or by removing secretions from the airway walls; aerosol surfactant is one of this class of medications (62). However, as with the other pharmacologic airway clearance therapies described above, there is no high-level evidence to support their use in children with CSLD or bronchiectasis (69).

Bronchodilators

Bronchiectasis may be associated with an obstructive ventilatory defect, which may worsen during an exacerbation (70). Although patients with bronchiectasis may also have asthma (71) and the presence of bronchiectasis may worsen asthma exacerbations (72), small adult studies suggest the airflow limitation is not readily reversible (73). There are, however, scarce data on the prevalence of asthma in children with CSLD or bronchiectasis, although there are suggestions that children with either CSLD or bronchiectasis have been misdiagnosed with asthma previously (19). There are no trials of bronchodilators in children with CSLD or bronchiectasis in the absence of a confirmed diagnosis of asthma, and their use in this group of children is not recommended currently (74).

Corticosteroids

Airway inflammation is a key characteristic of both CSLD and bronchiectasis where the resultant symptoms can be similar or falsely attributed to asthma (75). While the data are limited, the presence of asthma-like symptoms in the presence of

bronchiectasis has been associated with accelerated lung function decline (76–78). Inhaled corticosteroids are commonly used to control asthma symptoms, including preventing exacerbations. This class of drugs demonstrates broad anti-inflammatory actions, which can occur either rapidly or over a period of hours or days (79). Side effects are, however, not uncommon, and there are data to suggest children with persistent asthma receiving daily high-dose inhaled corticosteroids have an increased risk of linear growth faltering over a 12-month treatment period (80). Furthermore, for this strategy to be effective treatment, adherence is especially important (81) and, while data are limited in children with CSLD or bronchiectasis, non-adherence to inhaled corticosteroids regimens in adults with bronchiectasis has been documented (81, 82), and is known to be problematic in children with asthma, particularly adolescents. A Cochrane review of inhaled corticosteroids in children and adults with bronchiectasis identified no pediatric studies, and hence there is no available evidence to support their use (75). However, a small study of inhaled corticosteroid withdrawal over a 12-week period in 27 children with bronchiectasis (83) reported a significant increase in bronchial hyper-reactivity after ceasing the drug (37.0 vs 46.8%, $p = 0.016$) and a decrease in neutrophilic apoptosis in induced sputum (42.8 vs 20.2%, $p = 0.03$), but no change in sputum inflammatory markers or in the number of infectious exacerbations (83). Larger studies are required to confirm these findings and to determine whether they are of any clinical significance. By contrast, the role of systemic corticosteroids in children or adults with CSLD or bronchiectasis has not been examined (84, 85).

Non-Steroidal Anti-inflammatory Agents

Non-steroidal anti-inflammatory drugs (NSAIDs) are a class of medications that act as non-selective inhibitors of the enzyme cyclo-oxygenase and induce analgesic, antipyretic, and anti-inflammatory effects. Both oral and inhaled NSAIDs are available. A recently updated Cochrane review of inhaled NSAIDs in children and adults with bronchiectasis identified no relevant trials of NSAID efficacy in children and concluded there was no evidence to either support or refute their use in bronchiectasis (86). Another review of the efficacy of oral NSAIDs published in 2007 identified no trials in children or adults and could, therefore, make no recommendations regarding their use (87).

Vaccines

With respect to pathogens associated with respiratory illnesses, pediatric vaccines currently exist for influenza, *S. pneumoniae* (23 valent polysaccharide vaccine and 7, 10, and 13 valent pneumococcal conjugate vaccines), *Bordetella pertussis*, *H. influenzae* type b, measles, and varicella. The 10-valent pneumococcal vaccine uses Protein D, an outer membrane protein derivative from NTHi strains, as the conjugate. There is currently no evidence for any of these vaccines in preventing acute exacerbations of either CSLD or bronchiectasis in children (88), primarily because the trials have not been done. A randomized controlled trial (RCT) of the 10-valent pneumococcal Protein D conjugate vaccine in children with CSLD or bronchiectasis using acute

exacerbations as the primary endpoint has concluded recently in Australia (36), although the results were not yet published at the time of this review. The need for a safe and effective vaccine targeting NTHi for children is being recognized increasingly as a priority preventive and therapeutic intervention (89–91). Despite the lack of evidence for preventing exacerbations, children with chronic lung diseases are at high risk for severe infections from most of these organisms (92–94), and disease management should include ensuring children receive on-time and age-appropriate immunizations in all countries where these vaccines are recommended through national immunization programs.

Non-Pharmacological Airway Clearance Techniques

Non-pharmacological airway clearance techniques are strategies that aim to clear the airway of secretions, improve gas exchange, and hence reduce the risk of infection in the lower airways (95). The core components involve postural drainage, percussion, vibration of the chest wall, and coughing (96), and specific techniques include active cycle of breathing techniques, forced expiration techniques, autogenic drainage, postural drainage, oscillating positive expiratory pressure, high frequency chest wall oscillation, and exercise and/or pulmonary rehabilitation (96). While a Cochrane review found these techniques have clinically important effects on health-related QoL scores in adults (97), it identified only one study in children with bronchiectasis (98). This study involved only nine children with bronchiectasis and the intervention was oscillatory positive expiratory pressure administered three times a day with lung function measured after a 3-month period. The study did not include exacerbations as an outcome. Despite the lack of trials, non-pharmacological airway clearance techniques are recommended in managing children with CSLD or bronchiectasis (71, 74, 99) given the need to reduce mucus retention in the lower airways. The efficacy of these treatments with respect to preventing exacerbations is unknown.

Chronic Disease Management Plans

The complexity of managing patients with chronic diseases necessitates a multi-disciplinary approach that involves shared decision-making and includes the patient and their carers/family in developing treatment plans (100, 101). Personalized chronic disease management plans involve joint goal setting and agreement on actions necessary to achieve those goals (102). A systematic review of 19 studies involving 10,856 adults with a range of chronic health conditions concluded that modest improvements can be made through personalized management plans in physical and psychological parameters, as well as in patient capacity to self-manage their conditions (102). There appears to be no published studies of the efficacy of chronic disease management plans in preventing exacerbations of CSLD or bronchiectasis in children. A systematic review of action plans for adults with COPD that included limited self-management education identified improvements in the recognition of, and initiation of treatment for, acute exacerbations, but there was no evidence for reduced health-care

utilization or improved health-related QoL (103). A review of shared care of people with chronic diseases between primary and specialty health services concluded there was insufficient evidence for the benefits of that approach other than improved prescribing (104).

Health Promotion Strategies

Optimizing healthy growth and development is important for all children; however, additional efforts are required for children with CSLD or bronchiectasis given the links between respiratory infections and key factors, such as nutrition, exposure to environmental toxicants (particularly tobacco smoke), and healthy weight.

Optimize Nutritional Status

Nutritional deficiencies in children may lead to poor lung function, increased susceptibility to infections, and a greater likelihood of acute illnesses in childhood and chronic illness in adulthood (105). Micronutrient deficiencies, particularly of vitamins A, C, and D, folic acid, and the trace elements, zinc and iron, have been linked to many childhood infectious diseases, including respiratory infections (105). There is increasing interest in the role of vitamin D in respiratory diseases given its modulatory and regulatory role in inflammation and immunity (106). There are some data suggesting adults with chronic lung diseases are frequently deficient (107) and that deficiency is associated with disease severity (108). However, the studies supporting a role for vitamin D in protection against acute and chronic respiratory infections are associative and come from observational studies. A recent systematic review and meta-analysis of seven RCT of vitamin D supplementation in 6,503 children aged <18 years found no evidence to support their use for preventing or treating acute respiratory infections in healthy children (109). Interestingly, in this meta-analysis, vitamin D supplementation reduced the risk of asthma exacerbations by 74% [relative risk (RR) 0.26, 95% CI 0.11, 0.59], although this result needs to be interpreted cautiously as it involved two small studies of 430 and 48 subjects, respectively, and where significant heterogeneity was observed. Thus larger RCT are needed to determine whether vitamin D supplementation can reduce the risk of acute asthma exacerbations and if it should also be studied in children with other chronic lung diseases, including CSLD and bronchiectasis.

In addition, high intake of fructose-containing beverages such as soft drinks and fruit juice has been associated with increased prevalence of asthma in children (110, 111) and with chronic bronchitis in adults (112). Currently, there are no data on the role of micronutrient deficiencies, micronutrient supplementation, and high fructose intake in children with either CSLD or bronchiectasis. However, promoting a healthy diet is important for all children. A study of 141 Polish children with recurrent bronchitis found higher levels of body fat and muscle mass deficiency compared to reference ranges for healthy children (113). Whether this result reflects cause or effect is unknown, but while there are no efficacy trials of weight reduction in preventing exacerbations of pediatric CSLD or bronchiectasis, there are some data, albeit

limited, to suggest an effect in adults with asthma (114). Given the global rise of high calorie diets and associated obesity, and the complex socioeconomic circumstances where that has occurred, reducing calorie intake in children requires a multi-sectorial, population-based approach that includes strong government policy (115).

Reduce Exposure to Tobacco Smoke

Active and passive exposure to tobacco smoke is well-known as important risk factors for respiratory diseases. Children of parents who smoke are at approximately twice the risk of being hospitalized for respiratory illnesses (116). Hence avoiding exposure is not only beneficial to lung health but reduces the risk of the multitude of other adverse health outcomes associated with smoking. While knowledge around the effects of a child's passive exposure to tobacco smoke is improving in some settings (117), challenges remain in improving community and smoker perceptions of the risk to children, including among families of children with chronic lung diseases (117–119). Several strategies have been studied to reduce household tobacco smoke exposure in children, including motivational interviewing (with or without education) (120), maternal and child health nurse interventions (121), and family-centered interventions (122); however, the effectiveness of many strategies is modest at best (123).

Efforts are required to ensure children with CSLD or bronchiectasis do not take up smoking, particularly as they enter the vulnerable adolescent and young adult years (124). This must include efforts to prevent the use of e-cigarettes, which are becoming increasingly popular among adolescents despite emerging evidence to suggest adverse health effects (125). In a study of 2,086 North American adolescents of e-cigarette use and reported chronic cough, sputum production, or bronchitis symptoms in the previous 12 months (126), the risk of symptoms was almost twofold higher among past users (OR 1.85, 95% CI 1.37, 2.49), compared to never users, and by 2.02-fold (95% CI 1.42, 2.88) among current users. The risk also increased with frequency of current use (126).

Physical Activity

Promoting physical activity in children and adolescents should be standard practice for health-care providers given its well-known benefits to overall health and well-being. However, chronic illness can impact on activity levels in children, potentially affecting disease progression (127, 128). People with advanced lung disease experience static and dynamic hyperinflation due to airflow limitation, leading to reduced tidal volumes, increased dead space ventilation, and exertional dyspnea making exercise difficult (127). Physical activity provides higher health-related QoL scores (129) and greater aerobic capacity (130).

A study of an 8-week exercise training program in 55 adults with bronchiectasis who were subsequently followed for 12 months (131) reported fewer exacerbations in the intervention group compared to controls (RR 0.69) (95% CI 0.49, 0.98) as well as fewer exacerbations that required antibiotics. There are no data on the patterns of physical activity in children with CSLD or bronchiectasis, nor are there data on the effectiveness of physical activities in reducing exacerbations in these children.

CONCLUSION

Chronic suppurative lung disease and bronchiectasis in children and adolescents are serious illnesses with significant adverse impacts on the child, their family, the health-care sector, and broader community. Despite this, there has been scant attention paid to the spectrum of disease, its management, and prevention in comparison to CF and chronic lung diseases in adults. Exacerbations of CSLD and bronchiectasis in children are important predictors of lung function decline and are associated with significant morbidity. Preventing exacerbations should, therefore, be the cornerstone of long-term disease management. However, there are only limited data on preventing acute respiratory exacerbations in children with CSLD or bronchiectasis. Hence management is currently informed largely by clinical opinion and extrapolating data from adults and/or children with CF and asthma. Unfortunately, such an approach is not without risk, where treatments highly successful in CF patients such as inhaled rhDNase, tobramycin, aztreonam, and colistin have provided disappointing results in adults with bronchiectasis (132). A standardized, validated definition of an acute exacerbation in children with CSLD and bronchiectasis is needed urgently as an outcome measure in clinical trials and to help guide clinical practice. In a similar vein, studies to determine when recurrent episodes of PBB should be classified and managed as CSLD are also required. Substantially more, and larger studies, are needed to determine the efficacy and effectiveness of both pharmacological and non-pharmacological interventions to prevent exacerbations. These studies must also include health-related QoL assessments and cost-effectiveness of the interventions. Given the burden of disease is likely to be highest in low- and middle-income countries, measures to prevent exacerbations must be translated to low socioeconomic and disadvantaged community and health-care settings that do not have access to the drugs and technology available in high-income nations.

For the clinician, the lack of robust well-designed, intervention-based RCT to guide prevention of acute exacerbations of CSLD and bronchiectasis in children and adolescents means that guidelines are more consensus than evidence-based. We follow the Thoracic Society of Australia and New Zealand (TSANZ) guidelines (19, 74) where children and adolescents are seen three times monthly and where we recommend they practice regular airway clearance techniques in consultation with a pediatric respiratory physiotherapist, encourage physical activity, monitor growth and optimize nutrition, counsel against active and passive tobacco smoke exposure, promote avoidance of other environmental inhaled toxicants, and ensure they receive their vaccinations according to the National Immunization Program, including the pneumococcal and annual seasonal influenza vaccines. At the same time, co-morbidities, such as underlying immunodeficiency or chronic aspiration, are often managed with the aid of other specialty services. Children and adolescents who are frequent exacerbators (≥ 3 exacerbations and/or ≥ 2 respiratory hospitalizations in the previous 12 months) are prescribed low-dose oral azithromycin for a trial period of 12–24 months according to the TSANZ guidelines. Selected patients are offered inhaled hypertonic saline as a mucoactive agent and/or an inhaled antibiotic, usually tobramycin, if exacerbations continue,

especially if they have a chronic *P. aeruginosa* infection of their lower airways. These agents are trialed separately and if tolerated are continued long term. Finally, bronchodilators and inhaled corticosteroids are administered only to those with an established diagnosis of asthma and are not used routinely for other patients with CSLD or bronchiectasis.

AUTHOR CONTRIBUTIONS

KO devised the scope of the manuscript, performed the literature reviews, and wrote the first and final draft. KG made a substantial

contribution to the content and writing of the manuscript. Both the authors approved the final version.

FUNDING

KO is supported by a National Health and Medical Research Council (NHMRC) Career Development Fellowship (1045157) and a Queensland Government Smart Futures Fellowship. This work was also supported by a NHMRC Centre for Research Excellence for Lung Health in Aboriginal and Torres Strait Islanders grant (1040830).

REFERENCES

1. Wurzel DF, Marchant JM, Clark JE, Masters IB, Yerkovich ST, Upham JW, et al. Wet cough in children: infective and inflammatory characteristics in broncho-alveolar lavage fluid. *Pediatr Pulmonol* (2013) 49:561–8. doi:10.1002/ppul.22792
2. Marchant JM, Newcombe PA, Juniper EF, Sheffield JK, Stathis SL, Chang AB. What is the burden of chronic cough for families? *Chest* (2008) 134(2):303–9. doi:10.1378/chest.07-2236
3. Newcombe PA, Sheffield JK, Petsky HL, Marchant JM, Willis C, Chang AB. A child chronic cough-specific quality of life measure: development and validation. *Thorax* (2016) 71(8):695–700. doi:10.1136/thoraxjnl-2015-207473
4. Chang AB, Robertson CF, van Asperen PP, Glasgow NJ, Masters IB, Teoh L, et al. A cough algorithm for chronic cough in children: a multicenter, randomized controlled study. *Pediatrics* (2013) 131(5):e1576–83. doi:10.1542/peds.2012-3318
5. Goyal V, Grimwood K, Marchant JM, Masters IB, Chang AB. Paediatric chronic suppurative lung disease: clinical characteristics and outcomes. *Eur J Pediatr* (2016) 175(8):1077–84. doi:10.1007/s00431-016-2743-5
6. Chang AB, Redding GJ, Everard ML. Chronic wet cough: protracted bronchitis, chronic suppurative lung disease and bronchiectasis. *Pediatr Pulmonol* (2008) 43(6):519–31. doi:10.1002/ppul.20821
7. Wang Y, Hao C, Chi F, Yu X, Sun H, Huang L, et al. Clinical characteristics of protracted bacterial bronchitis in Chinese infants. *Sci Rep* (2015) 5:13731. doi:10.1038/srep13731
8. Hare KM, Binks MJ, Grimwood K, Chang AB, Leach AJ, Smith-Vaughan H. Culture and PCR detection of *Haemophilus influenzae* and *Haemophilus haemolyticus* in Australian indigenous children with bronchiectasis. *J Clin Microbiol* (2012) 50(7):2444–5. doi:10.1128/JCM.00566-12
9. Hare KM, Grimwood K, Leach AJ, Smith-Vaughan H, Torzillo PJ, Morris PS, et al. Respiratory bacterial pathogens in the nasopharynx and lower airways of Australian indigenous children with bronchiectasis. *J Pediatr* (2010) 157(6):1001–5. doi:10.1016/j.jpeds.2010.06.002
10. Narang R, Bakewell K, Peach J, Clayton S, Samuels M, Alexander J, et al. Bacterial distribution in the lungs of children with protracted bacterial bronchitis. *PLoS One* (2014) 9(9):e108523. doi:10.1371/journal.pone.0108523
11. Priftis KN, Litt D, Mangani S, Anthracopoulos MB, Thickett K, Tzanakaki G, et al. *Streptococcus pneumoniae* and non-typable *Haemophilus influenzae* causing bacterial bronchitis in children and the impact of vaccination. *Chest* (2012) 143:152–7. doi:10.1378/chest.12-0623
12. Grimwood K. Airway microbiology and host defences in paediatric non-CF bronchiectasis. *Paediatr Respir Rev* (2011) 12(2):111–8. doi:10.1016/j.prrv.2010.10.009
13. Kapur N, Grimwood K, Masters IB, Morris PS, Chang AB. Lower airway microbiology and cellularity in children with newly diagnosed non-CF bronchiectasis. *Pediatr Pulmonol* (2012) 47(3):300–7. doi:10.1002/ppul.21550
14. Das L, Kovesi TA. Bronchiectasis in children from Qikiqtani (Baffin) region, Nunavut, Canada. *Ann Am Thorac Soc* (2015) 12(1):96–100. doi:10.1513/AnnalsATS.201406-257OC
15. van der Gast CJ, Cuthbertson L, Rogers GB, Pope C, Marsh RL, Redding GJ, et al. Three clinically distinct chronic pediatric airway infections share a common core microbiota. *Ann Am Thorac Soc* (2014) 11(7):1039–48. doi:10.1513/AnnalsATS.201312-456OC
16. Chang AB, Upham JW, Masters IB, Redding GR, Gibson PG, Marchant JM, et al. Protracted bacterial bronchitis: the last decade and the road ahead. *Pediatr Pulmonol* (2016) 51(3):225–42. doi:10.1002/ppul.23351
17. Kantar A. Update on pediatric cough. *Lung* (2016) 194:9–14. doi:10.1007/s00408-015-9815-6
18. Goyal V, Grimwood K, Marchant J, Masters IB, Chang AB. Pediatric bronchiectasis: no longer an orphan disease. *Pediatr Pulmonol* (2016) 51(5):450–69. doi:10.1002/ppul.23380
19. Chang AB, Bell SC, Byrnes CA, Grimwood K, Holmes PW, King PT, et al. Chronic suppurative lung disease and bronchiectasis in children and adults in Australia and New Zealand. *Med J Aust* (2010) 193(6):356–65.
20. Wurzel DF, Marchant JM, Yerkovich ST, Upham JW, Petsky H, Smith-Vaughan H, et al. Protracted bacterial bronchitis in children: natural history and risk factors for bronchiectasis. *Chest* (2016) 150(5):1101–8. doi:10.1016/j.chest.2016.06.030
21. ElMaraachli W, Conrad DJ, Wang AC. Using cystic fibrosis therapies for non-cystic fibrosis bronchiectasis. *Clin Chest Med* (2016) 37(1):139–46. doi:10.1016/j.ccm.2015.11.005
22. Martinez-Garcia MA, Soler-Cataluna JJ, Perpina-Tordera M, Roman-Sanchez P, Soriano J. Factors associated with lung function decline in adult patients with stable non-cystic fibrosis bronchiectasis. *Chest* (2007) 132(5):1565–72. doi:10.1378/chest.07-0490
23. Loebinger MR, Wells AU, Hansell DM, Chinyanganya N, Devaraj A, Meister M, et al. Mortality in bronchiectasis: a long-term study assessing the factors influencing survival. *Eur Respir J* (2009) 34(4):843–9. doi:10.1183/09031936.00003709
24. Bastardo CM, Sonnappa S, Stanojevic S, Navarro A, Lopez PM, Jaffe A, et al. Non-cystic fibrosis bronchiectasis in childhood: longitudinal growth and lung function. *Thorax* (2009) 64(3):246–51. doi:10.1136/thx.2008.100958
25. Kapur N, Masters IB, Chang AB. Exacerbations in noncystic fibrosis bronchiectasis: clinical features and investigations. *Respir Med* (2009) 103(11):1681–7. doi:10.1016/j.rmed.2009.05.007
26. Kapur N, Masters IB, Chang AB. Longitudinal growth and lung function in pediatric non-cystic fibrosis bronchiectasis: what influences lung function stability? *Chest* (2010) 138(1):158–64. doi:10.1378/chest.09-2932
27. Seitz AE, Olivier KN, Steiner CA, Montes de Oca R, Holland SM, Prevots DR. Trends and burden of bronchiectasis-associated hospitalizations in the United States, 1993–2006. *Chest* (2010) 138(4):944–9. doi:10.1378/chest.10-0099
28. Edwards EA, Twiss J, Byrnes CA. Treatment of paediatric non-cystic fibrosis bronchiectasis. *Expert Opin Pharmacother* (2004) 5(7):1471–84. doi:10.1517/14656566.5.7.1471
29. Flight WG, Jones AM. Cystic fibrosis, primary ciliary dyskinesia and non-cystic fibrosis bronchiectasis: update 2008–11. *Thorax* (2012) 67(7):645–9. doi:10.1136/thoraxjnl-2011-200467
30. Stoller JK. Acute exacerbations of chronic obstructive pulmonary disease. *N Engl J Med* (2002) 346:988–94. doi:10.1056/NEJMc012477
31. European Respiratory Society. Bronchiectasis. *European Lung White Book: Respiratory Health and Disease in Europe*. Sheffield: European Respiratory Society (2013). Available from: <http://www.erswhitebook.org>
32. Grillo L, Irving S, Hansell DM, Nair A, Annan B, Ward S, et al. The reproducibility and responsiveness of the lung clearance index in bronchiectasis. *Eur Respir J* (2015) 46:1645–53. doi:10.1183/13993003.00152-2015

33. Valery PC, Morris PS, Grimwood K, Torzillo PJ, Byrnes CA, Masters IB, et al. Azithromycin for Indigenous children with bronchiectasis: study protocol for a multi-centre randomized controlled trial. *BMC Pediatr* (2012) 12:122. doi:10.1186/1471-2431-12-122
34. Chang AB, Grimwood K, Robertson CF, Wilson AC, van Asperen PP, O'Grady KA, et al. Antibiotics for bronchiectasis exacerbations in children: rationale and study protocol for a randomised placebo-controlled trial. *Trials* (2012) 13:156. doi:10.1186/1745-6215-13-156
35. Chang AB, Grimwood K, Wilson AC, van Asperen PP, Byrnes CA, O'Grady KA, et al. Bronchiectasis exacerbation study on azithromycin and amoxicillin-clavulanate for respiratory exacerbations in children (BEST-2): study protocol for a randomized controlled trial. *Trials* (2013) 14:53. doi:10.1186/1745-6215-14-53
36. O'Grady KA, Grimwood K, Cripps A, Mulholland EK, Morris P, Torzillo PJ, et al. Does a 10-valent pneumococcal-*Haemophilus influenzae* protein D conjugate vaccine prevent respiratory exacerbations in children with recurrent protracted bacterial bronchitis, chronic suppurative lung disease and bronchiectasis: protocol for a randomised controlled trial. *Trials* (2013) 14:282. doi:10.1186/1745-6215-14-282
37. Kapur N, Masters IB, Morris PS, Galligan J, Ware R, Chang AB. Defining pulmonary exacerbation in children with non-cystic fibrosis bronchiectasis. *Pediatr Pulmonol* (2012) 47(1):68–75. doi:10.1002/ppul.21518
38. Chang AB, Bilton D. Exacerbations in cystic fibrosis: 4 – non-cystic fibrosis bronchiectasis. *Thorax* (2008) 63(3):269–76. doi:10.1136/thx.2006.060913
39. Jourdain S, Smeesters PR, Denis O, Dramaix M, Sputael V, Malaviole X, et al. Differences in nasopharyngeal bacterial carriage in preschool children from different socio-economic origins. *Clin Microbiol Infect* (2011) 17(6):907–14. doi:10.1111/j.1469-0691.2010.03410.x
40. Kapur N, Mackay IM, Sloots TP, Masters IB, Chang AB. Respiratory viruses in exacerbations of non-cystic fibrosis bronchiectasis in children. *Arch Dis Child* (2014) 99(8):749–53. doi:10.1136/archdischild-2013-305147
41. Wurzel DF, Mackay IM, Marchant JM, Wang CY, Yerkovich ST, Upham JW, et al. Adenovirus species C is associated with chronic suppurative lung diseases in children. *Clin Infect Dis* (2014) 59(1):34–40. doi:10.1093/cid/ciu225
42. Redding GJ, Singleton RJ, Valery PC, Williams H, Grimwood K, Morris PS, et al. Respiratory exacerbations in indigenous children from two countries with non-cystic fibrosis chronic suppurative lung disease/bronchiectasis. *Chest* (2014) 146(3):762–74. doi:10.1378/chest.14-0126
43. Karadag B, Karakoc F, Ersu R, Kut A, Bakac S, Dagli E. Non-cystic-fibrosis bronchiectasis in children: a persisting problem in developing countries. *Respiration* (2005) 72(3):233–8. doi:10.1159/000085362
44. Chang AB, Grimwood K, Maguire G, King PT, Morris PS, Torzillo PJ. Management of bronchiectasis and chronic suppurative lung disease in indigenous children and adults from rural and remote Australian communities. *Med J Aust* (2008) 189(7):386–93.
45. Welsh EJ, Evans DJ, Fowler SJ, Spencer S. Interventions for bronchiectasis: an overview of Cochrane systematic reviews. *Cochrane Database Syst Rev* (2015) (7):CD010337. doi:10.1002/14651858.CD010337.pub2
46. Onakpoya IJ, Hayward G, Heneghan CJ. Antibiotics for preventing lower respiratory tract infections in high-risk children aged 12 years and under. *Cochrane Database Syst Rev* (2015) (9):CD011530. doi:10.1002/14651858.CD011530.pub2
47. Tramper-Stranders GA, Wolfs TE, van Haren Noman S, van Aalderen WM, Nagelkerke AF, Nuijsink M, et al. Controlled trial of cycled antibiotic prophylaxis to prevent initial *Pseudomonas aeruginosa* infection in children with cystic fibrosis. *Thorax* (2010) 65(10):915–20. doi:10.1136/thx.2009.126128
48. Clement A, Tamalet A, Leroux E, Ravilly S, Fauroux B, Jais JP. Long term effects of azithromycin in patients with cystic fibrosis: a double blind, placebo controlled trial. *Thorax* (2006) 61(10):895–902. doi:10.1136/thx.2005.057950
49. Saiman L, Anstead M, Mayer-Hamblett N, Lands LC, Kloster M, Hocesvar-Trnka J, et al. Effect of azithromycin on pulmonary function in patients with cystic fibrosis uninfected with *Pseudomonas aeruginosa*: a randomized controlled trial. *JAMA* (2010) 303(17):1707–15. doi:10.1001/jama.2010.563
50. Dogru D, Dalgic F, Kiper N, Ozcelik U, Yalcin E, Aslan AT, et al. Long-term clarithromycin in cystic fibrosis: effects on inflammatory markers in BAL and clinical status. *Turk J Pediatr* (2009) 51(5):416–23.
51. Equi A, Balfour-Lynn IM, Bush A, Rosenthal M. Long term azithromycin in children with cystic fibrosis: a randomised, placebo-controlled crossover trial. *Lancet* (2002) 360(9338):978–84. doi:10.1016/S0140-6736(02)11081-6
52. Murphy TD, Anbar RD, Lester LA, Nasr SZ, Nickerson B, VanDevanter DR, et al. Treatment with tobramycin solution for inhalation reduces hospitalizations in young CF subjects with mild lung disease. *Pediatr Pulmonol* (2004) 38(4):314–20. doi:10.1002/ppul.20097
53. Stutman HR, Lieberman JM, Nussbaum E, Marks MI. Antibiotic prophylaxis in infants and young children with cystic fibrosis: a randomized controlled trial. *J Pediatr* (2002) 140(3):299–305. doi:10.1067/mpd.2002.121930
54. Valery PC, Morris PS, Byrnes CA, Grimwood K, Torzillo PJ, Bauert PA, et al. Long-term azithromycin for indigenous children with non-cystic-fibrosis bronchiectasis or chronic suppurative lung disease (Bronchiectasis Intervention Study): a multicentre, double-blind, randomised controlled trial. *Lancet Respir Med* (2013) 1(8):610–20. doi:10.1016/S2213-2600(13)70185-1
55. Hare KM, Grimwood K, Chang AB, Chatfield MD, Valery PC, Leach AJ, et al. Nasopharyngeal carriage and macrolide resistance in indigenous children with bronchiectasis randomized to long-term azithromycin or placebo. *Eur J Clin Microbiol Infect Dis* (2015) 34(11):2275–85. doi:10.1007/s10096-015-2480-0
56. Hare KM, Singleton RJ, Grimwood K, Valery PC, Cheng AC, Morris PS, et al. Longitudinal nasopharyngeal carriage and antibiotic resistance of respiratory bacteria in indigenous Australian and Alaska native children with bronchiectasis. *PLoS One* (2013) 8(8):e70478. doi:10.1371/journal.pone.0070478
57. Hnin K, Nguyen C, Carson KV, Evans DJ, Greenstone M, Smith BJ. Prolonged antibiotics for non-cystic fibrosis bronchiectasis in children and adults. *Cochrane Database Syst Rev* (2015) (8):CD001392. doi:10.1002/14651858.CD001392.pub3
58. Yalcin E, Kiper N, Ozcelik U, Dogru D, Firat P, Sahin A, et al. Effects of clarithromycin on inflammatory parameters and clinical conditions in children with bronchiectasis. *J Clin Pharm Ther* (2006) 31(1):49–55. doi:10.1111/j.1365-2710.2006.00708.x
59. Koh YY, Lee MH, Sun YH, Sung KW, Chae JH. Effect of roxithromycin on airway responsiveness in children with bronchiectasis: a double-blind, placebo-controlled study. *Eur Respir J* (1997) 10(5):994–9. doi:10.1183/09031936.97.10050994
60. Wurzel D, Marchant JM, Yerkovich ST, Upham JW, Masters IB, Chang AB. Short courses of antibiotics for children and adults with bronchiectasis. *Cochrane Database Syst Rev* (2011) (6):CD008695. doi:10.1002/14651858.CD008695.pub2
61. Rubin BK. Mucolytics, expectorants, and mucokinetic medications. *Respir Care* (2007) 52(7):859–65.
62. Nair GB, Ilowite JS. Pharmacologic agents for mucus clearance in bronchiectasis. *Clin Chest Med* (2012) 33(2):363–70. doi:10.1016/j.ccm.2012.02.008
63. Smith SM, Schroeder K, Fahey T. Over-the-counter (OTC) medications for acute cough in children and adults in community settings. *Cochrane Database Syst Rev* (2014) (11):CD001831. doi:10.1002/14651858.CD001831.pub5
64. Hampton LM, Nguyen DB, Edwards JR, Budnitz DS. Cough and cold medication adverse events after market withdrawal and labeling revision. *Pediatrics* (2013) 132(6):1047–54. doi:10.1542/peds.2013-2236
65. Daviskas E, Anderson SD. Hyperosmolar agents and clearance of mucus in the diseased airway. *J Aerosol Med* (2006) 19(1):100–9. doi:10.1089/jam.2006.19.100
66. Wilkinson M, Sugumar K, Milan SJ, Hart A, Crockett A, Crossingham I. Mucolytics for bronchiectasis. *Cochrane Database Syst Rev* (2014) (5):CD001289. doi:10.1002/14651858.CD001289.pub2
67. Pressler T. Review of recombinant human deoxyribonuclease (rhDNase) in the management of patients with cystic fibrosis. *Biologics* (2008) 2(4):611–7. doi:10.2147/BTT.S3052
68. O'Donnell AE, Barker AE, Ilowite JS, Fick RB. Treatment of idiopathic bronchiectasis with aerosolized recombinant human DNase I. rhDNase Study Group. *Chest* (1998) 113(5):1329–34. doi:10.1378/chest.113.5.1329
69. Strickland SL, Rubin BK, Haas CF, Volsko TA, Drescher GS, O'Malley CA. AARC clinical practice guideline: effectiveness of pharmacologic airway clearance therapies in hospitalized patients. *Respir Care* (2015) 60(7):1071–7. doi:10.4187/respcare.04165
70. AL-Shirawi N, AL-Jahdali HH, Al Shimemeri A. Pathogenesis, etiology and treatment of bronchiectasis. *Ann Thorac Med* (2006) 1(1):41–51. doi:10.4103/1817-1737.25870
71. Pasteur MC, Bilton D, Hill AT, British Thoracic Society Non-CF Bronchiectasis Guideline Group. British Thoracic Society guideline for non-CF bronchiectasis. *Thorax* (2010) 65(7):577. doi:10.1136/thx.2010.142778

72. Amalakuhan B, Maselli DJ, Martinez-Garcia MA. Update in bronchiectasis 2014. *Am J Respir Crit Care Med* (2015) 192(10):1155–61. doi:10.1164/rccm.201505-0926UP
73. Murphy MB, Reen DJ, Fitzgerald MX. Atopy, immunological changes, and respiratory function in bronchiectasis. *Thorax* (1984) 39(3):179–84. doi:10.1136/thx.39.3.179
74. Chang AB, Bell SC, Torzillo PJ, King PT, Maguire GP, Byrnes CA, et al. Chronic suppurative lung disease and bronchiectasis in children and adults in Australia and New Zealand Thoracic Society of Australia and New Zealand guidelines. *Med J Aust* (2015) 202(3):130. doi:10.5694/mjac14.00287
75. Kapur N, Bell S, Kolbe J, Chang AB. Inhaled steroids for bronchiectasis. *Cochrane Database Syst Rev* (2009) (1):CD000996. doi:10.1002/14651858.CD000996.pub2
76. Keistinen T, Saynajakangas O, Tuuponen T, Kivela SL. Bronchiectasis: an orphan disease with a poorly-understood prognosis. *Eur Respir J* (1997) 10(12):2784–7. doi:10.1183/09031936.97.10122784
77. Saynajakangas O, Keistinen T, Tuuponen T, Kivela SL. Links between hospital diagnoses of bronchiectasis and asthma. *Allergy* (1997) 52(11):1120–2. doi:10.1111/j.1398-9995.1997.tb00186.x
78. Field CE. Bronchiectasis. Third report on a follow-up study of medical and surgical cases from childhood. *Arch Dis Child* (1969) 44(237):551–61. doi:10.1136/adc.44.237.551
79. Hossny E, Rosario N, Lee BW, Singh M, El-Ghoneimy D, Soh JY, et al. The use of inhaled corticosteroids in pediatric asthma: update. *World Allergy Organ J* (2016) 9:26. doi:10.1186/s40413-016-0117-0
80. Ohn M, Jaffe A, Selvadurai H. Persistent growth effects of inhaled corticosteroids. *J Paediatr Child Health* (2016) 52(10):964–6. doi:10.1111/jpc.13356
81. McCullough AR, Tunney MM, Quittner AL, Elborn JS, Bradley JM, Hughes CM. Treatment adherence and health outcomes in patients with bronchiectasis. *BMC Pulm Med* (2014) 14:107. doi:10.1186/1471-2466-14-107
82. McCullough AR, Tunney MM, Elborn JS, Bradley JM, Hughes CM. 'All illness is personal to that individual': a qualitative study of patients' perspectives on treatment adherence in bronchiectasis. *Health Expect* (2015) 18(6):2477–88. doi:10.1111/hex.12217
83. Guran T, Ersu R, Karadag B, Karakoc F, Demirel GY, Hekim N, et al. Withdrawal of inhaled steroids in children with non-cystic fibrosis bronchiectasis. *J Clin Pharm Ther* (2008) 33(6):603–11. doi:10.1111/j.1365-2710.2008.00951.x
84. ten Hacken NH, Wijkstra PJ, Kerstjens HA. Treatment of bronchiectasis in adults. *BMJ* (2007) 335(7629):1089–93. doi:10.1136/bmj.39384.657118.80
85. Lasserson T, Holt K, Greenstone M. Oral steroids for bronchiectasis (stable and acute exacerbations). *Cochrane Database Syst Rev* (2001) (4):CD002162. doi:10.1002/14651858.CD002162
86. Pizzutto SJ, Upham JW, Yerkovich ST, Chang AB. Inhaled non-steroid anti-inflammatories for children and adults with bronchiectasis. *Cochrane Database Syst Rev* (2016) (1):CD007525. doi:10.1002/14651858.CD007525.pub3
87. Kapur N, Chang AB. Oral non steroid anti-inflammatories for children and adults with bronchiectasis. *Cochrane Database Syst Rev* (2007) (4):CD006427. doi:10.1002/14651858.CD006427.pub2
88. O'Grady KA, Chang AB, Grimwood K. Vaccines for children and adults with chronic lung disease: efficacy against acute exacerbations. *Expert Rev Respir Med* (2014) 8(1):43–55. doi:10.1586/17476348.2014.852960
89. Murphy TE. Immunity to nontypeable *Haemophilus influenzae*: elucidating protective responses. *Am J Respir Crit Care Med* (2003) 167(4):486–7. doi:10.1164/rccm.2211009
90. Murphy TE. Vaccines for nontypeable *Haemophilus influenzae*: the future is now. *Clin Vaccine Immunol* (2015) 22(5):459–66. doi:10.1128/CVI.00089-15
91. Pizzutto SJ, Yerkovich ST, Upham JW, Hales BJ, Thomas WR, Chang AB. Improving immunity to *Haemophilus influenzae* in children with chronic suppurative lung disease. *Vaccine* (2015) 33(2):321–6. doi:10.1016/j.vaccine.2014.11.024
92. Aigbogun NW, Hawker JL, Stewart A. Interventions to increase influenza vaccination rates in children with high-risk conditions – a systematic review. *Vaccine* (2015) 33(6):759–70. doi:10.1016/j.vaccine.2014.12.013
93. Pelton SI, Weycker D, Farkouh RA, Stratton DR, Shea KM, Edelsberg J. Risk of pneumococcal disease in children with chronic medical conditions in the era of pneumococcal conjugate vaccine. *Clin Infect Dis* (2014) 59(5):615–23. doi:10.1093/cid/ciu348
94. Montella S, De Stefano S, Sperli F, Barbarano F, Santamaria F. Increased risk of chronic suppurative lung disease after measles or pertussis in non-vaccinated children. *Vaccine* (2007) 25(3):402–3. doi:10.1016/j.vaccine.2006.09.045
95. Main E, Grillo L, Rand S. Airway clearance strategies in cystic fibrosis and non-cystic fibrosis bronchiectasis. *Semin Respir Crit Care Med* (2015) 36(2):251–66. doi:10.1055/s-0035-1546820
96. Snijders D, Fernandez Dominguez B, Calgario S, Bertozzi I, Escribano Montaner A, Perilongo G, et al. Mucociliary clearance techniques for treating non-cystic fibrosis bronchiectasis: is there evidence? *Int J Immunopathol Pharmacol* (2015) 28(2):150–9. doi:10.1177/0394632015584724
97. Lee AL, Burge AT, Holland AE. Airway clearance techniques for bronchiectasis. *Cochrane Database Syst Rev* (2015) (11):CD008351. doi:10.1002/14651858.CD008351.pub3
98. Kurz M, Ahrens P, Zielen S, Hofmann D. Objective and subjective results of a comparative study in children with chronic lung diseases [Physiotherapie mit dem VRP1-Gerat: Objektive und subjektive Ergebnisse einer vergleichenden Studie bei Kindern mit chronischen Lungenerkrankungen]. *Prev Rehabil* (1997) 9(3):105–9.
99. Chang AB, Oppenheimer JJ, Weinberger M, Rubin BK, Irwin RS. Children with chronic wet or productive cough – treatment and investigations; a systematic review. *Chest* (2016) 149(1):120–42. doi:10.1378/chest.15-2065
100. van Dongen JJ, van Bokhoven MA, Daniels R, van der Weijden T, Emonts WW, Beurskens A. Developing interprofessional care plans in chronic care: a scoping review. *BMC Fam Pract* (2016) 17(1):137. doi:10.1186/s12875-016-0535-7
101. van Dongen JJ, Lenzen SA, van Bokhoven MA, Daniels R, van der Weijden T, Beurskens A. Interprofessional collaboration regarding patients' care plans in primary care: a focus group study into influential factors. *BMC Fam Pract* (2016) 17:58. doi:10.1186/s12875-016-0456-5
102. Coulter A, Entwistle VA, Eccles A, Ryan S, Shepperd S, Perera R. Personalised care planning for adults with chronic or long-term health conditions. *Cochrane Database Syst Rev* (2015) (3):CD010523. doi:10.1002/14651858.CD010523.pub2
103. Walters JA, Turnock AC, Walters EH, Wood-Baker R. Action plans with limited patient education only for exacerbations of chronic obstructive pulmonary disease. *Cochrane Database Syst Rev* (2010) (5):CD005074. doi:10.1002/14651858.CD005074.pub3
104. Smith SM, Allwright S, O'Dowd T. Effectiveness of shared care across the interface between primary and specialty care in chronic disease management. *Cochrane Database Syst Rev* (2007) (3):CD004910. doi:10.1002/14651858.CD004910.pub2
105. Karim T, Muhit M, Khandaker G. Interventions to prevent respiratory diseases – nutrition and the developing world. *Paediatr Respir Rev* (2017) 22:31–7. doi:10.1016/j.prrv.2016.09.003
106. Solidoro P, Bellocchia M, Facchini F. The immunobiological and clinical role of vitamin D in obstructive lung diseases. *Minerva Med* (2016) 107(3 Suppl 1):12–9.
107. Zdrenghea MT, Makrinioti H, Bagacean C, Bush A, Johnston SL, Stanciu LA. Vitamin D modulation of innate immune responses to respiratory viral infections. *Rev Med Virol* (2016) 27:R1909. doi:10.1002/rmv.1909
108. Zhu B, Zhu B, Xiao C, Zheng Z. Vitamin D deficiency is associated with the severity of COPD: a systematic review and meta-analysis. *Int J Chron Obstruct Pulmon Dis* (2015) 10:1907–16. doi:10.2147/COPD.S89763
109. Xiao L, Xing C, Yang Z, Xu S, Wang M, Du H, et al. Vitamin D supplementation for the prevention of childhood acute respiratory infections: a systematic review of randomised controlled trials. *Br J Nutr* (2015) 114(7):1026–34. doi:10.1017/S000711451500207X
110. DeChristopher LR, Uribarri J, Tucker KL. The link between soda intake and asthma: science points to the high-fructose corn syrup, not the preservatives: a commentary. *Nutr Diabetes* (2016) 6(11):e234. doi:10.1038/ntd.2016.46
111. DeChristopher LR, Uribarri J, Tucker KL. Intake of high-fructose corn syrup sweetened soft drinks, fruit drinks and apple juice is associated with prevalent arthritis in US adults, aged 20-30 years. *Nutr Diabetes* (2016) 6:e199. doi:10.1038/ntd.2016.7
112. DeChristopher LR, Uribarri J, Tucker KL. Intake of high fructose corn syrup sweetened soft drinks is associated with prevalent chronic bronchitis in U.S. Adults, ages 20-55 y. *Nutr J* (2015) 14:107. doi:10.1186/s12937-015-0097-x

113. Umlawska W, Lipowicz A. Growth, nutritional status, and pulmonary function in children with chronic recurrent bronchitis. *Adv Exp Med Biol* (2016) 885:1–9. doi:10.1007/5584_2015_192
114. Stenius-Aarniala B, Poussa T, Kvarnstrom J, Gronlund EL, Ylikahri M, Mustajoki P. Immediate and long term effects of weight reduction in obese people with asthma: randomised controlled study. *BMJ* (2000) 320(7238):827–32. doi:10.1136/bmj.320.7238.827
115. Popkin BM, Hawkes C. Sweetening of the global diet, particularly beverages: patterns, trends, and policy responses. *Lancet Diabetes Endocrinol* (2016) 4(2):174–86. doi:10.1016/S2213-8587(15)00419-2
116. Li JS, Peat JK, Xuan W, Berry G. Meta-analysis on the association between environmental tobacco smoke (ETS) exposure and the prevalence of lower respiratory tract infection in early childhood. *Pediatr Pulmonol* (1999) 27(1):5–13. doi:10.1002/(SICI)1099-0496(199901)27:1<5::AID-PPUL3>3.0.CO;2-5
117. Lonergan BJ, Meaney S, Perry JJ, Comber H, Power B, Bradley C, et al. Smokers still underestimate the risks posed by secondhand smoke: a repeated cross-sectional study. *Nicotine Tob Res* (2014) 16(8):1121–8. doi:10.1093/ntr/ntu046
118. Borrelli B, McQuaid EL, Wagener TL, Hammond SK. Children with asthma versus healthy children: differences in secondhand smoke exposure and caregiver perceived risk. *Nicotine Tob Res* (2014) 16(5):554–61. doi:10.1093/ntr/ntt180
119. Fedele DA, Tooley E, Busch A, McQuaid EL, Hammond SK, Borrelli B. Comparison of secondhand smoke exposure in minority and nonminority children with asthma. *Health Psychol* (2016) 35(2):115–22. doi:10.1037/hea0000220
120. Eakin MN, Rand CS, Borrelli B, Bilderback A, Hovell M, Riekert KA. Effectiveness of motivational interviewing to reduce head start children's secondhand smoke exposure. a randomized clinical trial. *Am J Respir Crit Care Med* (2014) 189(12):1530–7. doi:10.1164/rccm.201404-0618OC
121. Daly JB, Freund M, Burrows S, Considine R, Bowman JA, Wiggers JH. A cluster randomised controlled trial of a brief child health nurse intervention to reduce infant secondhand smoke exposure. *Matern Child Health J* (2017) 21:108–17. doi:10.1007/s10995-016-2099-5
122. Walker N, Johnston V, Glover M, Bullen C, Trenholme A, Chang A, et al. Effect of a family-centered, secondhand smoke intervention to reduce respiratory illness in indigenous infants in Australia and New Zealand: a randomized controlled trial. *Nicotine Tob Res* (2015) 17(1):48–57. doi:10.1093/ntr/ntu128
123. Priest N, Roseby R, Waters E, Polnay A, Campbell R, Spencer N, et al. Family and carer smoking control programmes for reducing children's exposure to environmental tobacco smoke. *Cochrane Database Syst Rev* (2008) (4):CD001746. doi:10.1002/14651858.CD001746.pub2
124. Das JK, Salam RA, Arshad A, Finkelstein Y, Bhutta ZA. Interventions for adolescent substance abuse: an overview of systematic reviews. *J Adolesc Health* (2016) 59(4S):S61–75. doi:10.1016/j.jadohealth.2016.06.021
125. Dunbar MS, Tucker JS, Ewing BA, Pedersen ER, Miles JN, Shih RA, et al. Frequency of E-cigarette use, health status, and risk and protective health behaviors in adolescents. *J Addict Med* (2017) 11:55–62. doi:10.1097/ADM.0000000000000272
126. McConnell R, Barrington-Trimis JL, Wang K, Urman R, Hong H, Unger J, et al. Electronic-cigarette use and respiratory symptoms in adolescents. *Am J Respir Crit Care Med* (2016). doi:10.1164/rccm.201604-0804OC
127. Burtin C, Hebestreit H. Rehabilitation in patients with chronic respiratory disease other than chronic obstructive pulmonary disease: exercise and physical activity interventions in cystic fibrosis and non-cystic fibrosis bronchiectasis. *Respiration* (2015) 89(3):181–9. doi:10.1159/000375170
128. LeBlanc CM, Lands LC. Can I play? *Pediatr Ann* (2014) 43(12):e316–24. doi:10.3928/00904481-20141124-11
129. Saunders TJ, Gray CE, Poitras VJ, Chaput JB, Janssen I, Katzmarzyk PT, et al. Combinations of physical activity, sedentary behaviour and sleep: relationships with health in children and youth. *Med Sci Sports Exerc* (2016) 48(5 Suppl 1):912. doi:10.1249/01.mss.0000487732.62636.11
130. Bai Y, Saint-Maurice PF, Welk GJ, Russell DW, Allums-Featherston K, Candelaria N. The longitudinal impact of NFL PLAY 60 programming on youth aerobic capacity and BMI. *Am J Prev Med* (2017) 52:311–23. doi:10.1016/j.amepre.2016.10.009
131. Lee AL, Hill CJ, Cecins N, Jenkins S, McDonald CF, Burge AT, et al. The short and long term effects of exercise training in non-cystic fibrosis bronchiectasis – a randomised controlled trial. *Respir Res* (2014) 15:44. doi:10.1186/1465-9921-15-44
132. Grimwood K, Bell SC, Chang AB. Antimicrobial treatment of non-cystic fibrosis bronchiectasis. *Expert Rev Anti Infect Ther* (2014) 12(10):1277–96. doi:10.1586/14787210.2014.952282

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2017 O'Grady and Grimwood. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.