



## OPEN ACCESS

## EDITED BY

Ruth Heying,  
University Hospital Leuven, Belgium

## REVIEWED BY

Ozgur Kasapcopur,  
Istanbul University-Cerrahpasa, Türkiye

## \*CORRESPONDENCE

Ji-whan Han  
✉ harji59@gmail.com

RECEIVED 20 January 2025

ACCEPTED 26 February 2025

PUBLISHED 11 March 2025

## CITATION

Kim BJ, Choi A, Kim S-i and Han J-w (2025)  
Corrigendum: The incidence of periungual  
desquamation and thrombocytosis in  
Kawasaki disease and the importance of  
systematic observation in the subacute phase.  
Front. Pediatr. 13:1564036.  
doi: 10.3389/fped.2025.1564036

## COPYRIGHT

© 2025 Kim, Choi, Kim and Han. This is an  
open-access article distributed under the  
terms of the [Creative Commons Attribution  
License \(CC BY\)](#). The use, distribution or  
reproduction in other forums is permitted,  
provided the original author(s) and the  
copyright owner(s) are credited and that the  
original publication in this journal is cited, in  
accordance with accepted academic practice.  
No use, distribution or reproduction is  
permitted which does not comply with  
these terms.

# Corrigendum: The incidence of periungual desquamation and thrombocytosis in Kawasaki disease and the importance of systematic observation in the subacute phase

Beom Joon Kim<sup>1</sup> , Arum Choi<sup>2</sup> , Suk-il Kim<sup>3</sup> and  
Ji-whan Han <sup>1\*</sup>

<sup>1</sup>Department of Pediatrics, College of Medicine, Catholic University of Korea, Seoul, Republic of Korea, <sup>2</sup>Department of Biomedical Systems Informatics, Yonsei University College of Medicine, Seoul, Republic of Korea, <sup>3</sup>Department of Preventive Medicine and Public Health, College of Medicine, Catholic University of Korea, Seoul, Republic of Korea

## KEYWORDS

desquamation, fever, incomplete Kawasaki Disease, intravenous immunoglobulin, Kawasaki Disease, Kawasaki-like Disease, subacute phase, thrombocytosis

## A Corrigendum on

The incidence of periungual desquamation and thrombocytosis in Kawasaki disease and the importance of systematic observation in the subacute phase

By Kim BJ, Choi A, Kim S-i and Han J-w (2024). Front. Pediatr. 12. doi: 10.3389/fped.2024.1384015

## Error in table

In the published article, there was an error in [Table 4](#), “Proportion of Patients Diagnosed with Incomplete KD Among All KD Cases during the Acute and Subacute Phase.” The total number for this group is currently listed as 521, but it should be 819. The corrected [Table 4](#) and its caption appear below.

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

TABLE 4 Proportion of patients diagnosed with incomplete KD among all KD cases during the acute and subacute phases.

Variable	Group I ( <i>n</i> = 60)	Group II ( <i>n</i> = 819)	<i>P</i> value
Incomplete KD during Acute Phase (without desquamation)	15 (25.0)	291 (35.5)	0.130
Incomplete KD during subacute phase (with desquamation)	8 (13.3)	268 (34.9)	0.001

Data are expressed as number (%). KD, Kawasaki disease.

## Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.