



Corrigendum: Ethnopharmacological Approaches for Therapy of Jaundice: Part II. Highly Used Plant Species from Acanthaceae, Euphorbiaceae, Asteraceae, Combretaceae, and Fabaceae Families

Devesh Tewari^{1†}, Andrei Mocan^{2,3†}, Emil D. Parvanov⁴, Archana N. Sah¹, Seyed M. Nabavi⁵, Lukasz Huminiecki⁶, Zheng Feei Ma^{7,8}, Yeong Yeh Lee⁷, Jaroslaw O. Horbańczuk⁶ and Atanas G. Atanasov^{6,9,10*}

¹ Department of Pharmaceutical Sciences, Faculty of Technology, Kumaun University, Nainital, India, ² Department of Pharmaceutical Botany, "Iuliu Hatieganu" University of Medicine and Pharmacy, Cluj-Napoca, Romania, ³ ICHAT and Institute for Life Sciences, University of Agricultural Sciences and Veterinary Medicine, Cluj-Napoca, Romania, ⁴ Division BIOCEV, Institute of Molecular Genetics, Academy of Sciences of the Czech Republic, Prague, Czechia, ⁵ Applied Biotechnology Research Center, Baqiyatallah University of Medical Sciences, Tehran, Iran, ⁶ Institute of Genetics and Animal Breeding of the Polish Academy of Sciences, Jastrzebiec, Poland, ⁷ School of Medical Sciences, Universiti Sains Malaysia, Kota Bharu, Malaysia, ⁸ Department of Public Health, Xi'an Jiaotong-Liverpool University, Suzhou, China, ⁹ Department of Pharmacognosy, University of Vienna, Vienna, Austria, ¹⁰ Department of Vascular Biology and Thrombosis Research, Centre for Physiology and Pharmacology, Medical University of Vienna, Vienna, Austria

OPEN ACCESS

Edited and reviewed by:

Jianbo Xiao,
University of Macau, China

*Correspondence:

Atanas G. Atanasov
a.atanasov.mailbox@gmail.com

[†]These authors have contributed
equally to this work.

Specialty section:

This article was submitted to
Ethnopharmacology,
a section of the journal
Frontiers in Pharmacology

Received: 19 August 2017

Accepted: 15 September 2017

Published: 06 October 2017

Citation:

Tewari D, Mocan A, Parvanov ED, Sah AN, Nabavi SM, Huminiecki L, Ma ZF, Lee YY, Horbańczuk JO and Atanasov AG (2017) Corrigendum: Ethnopharmacological Approaches for Therapy of Jaundice: Part II. Highly Used Plant Species from Acanthaceae, Euphorbiaceae, Asteraceae, Combretaceae, and Fabaceae Families. *Front. Pharmacol.* 8:690. doi: 10.3389/fphar.2017.00690

Keywords: jaundice, bilirubin, oxidative stress, traditional use, phytoconstituents, serum enzymes, alkaline phosphatase

A corrigendum on

Ethnopharmacological Approaches for Therapy of Jaundice: Part II. Highly Used Plant Species from Acanthaceae, Euphorbiaceae, Asteraceae, Combretaceae, and Fabaceae Families by Tewari, D., Mocan, A., Parvanov, E. D., Sah, A. N., Nabavi, S. M., Huminiecki, L., et al. (2017). *Front. Pharmacol.* 8:519. doi: 10.3389/fphar.2017.00519

In the original article, there was a mistake in the legend for **Figure 4** as published (the spelling of isosilbin was incorrect). The correct legend appears below.

In the original article, there was a mistake in **Figure 4** as published (CH₃ group was missing in the Silybin structure). The corrected **Figure 4** appears below.

The authors apologize for these errors and state that this does not change the scientific conclusions of the article in any way.

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

The handling Editor declared a past co-authorship with the author SN.

Copyright © 2017 Tewari, Mocan, Parvanov, Sah, Nabavi, Huminiecki, Ma, Lee, Horbańczuk and Atanasov. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

