



Molecular Insights Into SARS COV-2 Interaction With Cardiovascular Disease: Role of RAAS and MAPK Signaling

OPEN ACCESS

Zena Wehbe^{1†}, Safaa Hammoud^{2†}, Nadia Soudani³, Hassan Zaraket^{3*}, Ahmed El-Yazbi^{4,5*} and Ali H. Eid^{4,6*}

Edited by:

Ismail Laher,
University of British Columbia,
Canada

Reviewed by:

Urs T. Ruegg,
Université de Genève,
Switzerland
Christian Aalkjaer,
Aarhus University, Denmark
Michael A. Hill,
University of Missouri, United States
Chris R. Triggler,
Weill Cornell Medicine-Qatar, Qatar

*Correspondence:

Hassan Zaraket
hz34@aub.edu.lb
Ahmed El-Yazbi
ae88@aub.edu.lb
Ali H. Eid
ae81@aub.edu.lb

[†]These authors have contributed
equally to this work

Specialty section:

This article was submitted to
Cardiovascular and Smooth
Muscle Pharmacology,
a section of the journal
Frontiers in Pharmacology

Received: 01 April 2020

Accepted: 21 May 2020

Published: 03 June 2020

Citation:

Wehbe Z, Hammoud S, Soudani N,
Zaraket H, El-Yazbi A and Eid AH
(2020) Molecular Insights Into
SARS COV-2 Interaction With
Cardiovascular Disease: Role of
RAAS and MAPK Signaling.
Front. Pharmacol. 11:836.
doi: 10.3389/fphar.2020.00836

¹ Department of Biology, American University of Beirut, Beirut, Lebanon, ² Department of Pharmacology and Therapeutics, Beirut Arab University, Beirut, Lebanon, ³ Department of Experimental Pathology, Immunology and Microbiology, American University of Beirut, Beirut, Lebanon, ⁴ Department of Pharmacology and Toxicology, American University of Beirut, Beirut, Lebanon, ⁵ Department of Pharmacology and Toxicology, Faculty of Pharmacy, Alexandria University, Alexandria, Egypt, ⁶ Department of Biomedical Sciences, College of Health, Qatar University, Doha, Qatar

In December 2019, reports of viral pneumonia came out of Wuhan city in Hubei province in China. In early 2020, the causative agent was identified as a novel coronavirus (CoV) sharing some sequence similarity with SARS-CoV that caused the severe acute respiratory syndrome outbreak in 2002. The new virus, named SARS-CoV-2, is highly contagious and spread rapidly across the globe causing a pandemic of what became known as coronavirus infectious disease 2019 (COVID-19). Early observations indicated that cardiovascular disease (CVD) patients are at higher risk of progression to severe respiratory manifestations of COVID-19 including acute respiratory distress syndrome. Moreover, further observations demonstrated that SARS-CoV-2 infection can induce *de novo* cardiac and vascular damage in previously healthy individuals. Here, we offer an overview of the proposed molecular pathways shared by the pathogenesis of CVD and SARS-CoV infections in order to provide a mechanistic framework for the observed interrelation. We examine the crosstalk between the renin-angiotensin-aldosterone system and mitogen activated kinase pathways that potentially links cardiovascular predisposition and/or outcome to SARS-CoV-2 infection. Finally, we summarize the possible effect of currently available drugs with known cardiovascular benefit on these pathways and speculate on their potential utility in mitigating cardiovascular risk and morbidity in COVID-19 patients.

Keywords: severe COVID-19, cardiovascular burden, signaling pathways, RAAS, MAPK signaling

INTRODUCTION

Coronaviruses (CoVs) are single-stranded, positive-sense RNA viruses (26–32 kb) that belong to the *Coronaviridae* family (Su et al., 2016). Common human coronaviruses include HCoV-NL63, -229E, -OC43, and -HKU1 and are usually associated with mild acute respiratory illnesses or “common cold.” In December 2019, clusters of pneumonia cases of unknown etiology were reported in Wuhan

City, Hubei Province in China (Huang et al., 2020). Within a few weeks, scientists determined that these mysterious pneumonia cases were caused by a novel coronavirus (CoV) that shares around 79.5% sequence similarity with the SARS-CoV and 96.2% with bat-CoV RaTG13 (Lu et al., 2020; Zhou P. et al., 2020). Therefore, the virus was named SARS-CoV-2 and the disease COVID-19 stands for coronavirus infectious disease 2019. On March 11, 2020, the World Health Organization announced COVID-19 as a pandemic and requested all countries to scale up their emergency response mechanisms (Who, 2020b).

SARS-CoV2 is highly contagious with an average incubation period of 5–6 days (range 1–14) (Who, 2020a). It can be transmitted by droplets generated during coughing or sneezing, close contact, and touching contaminated surfaces (Prevention, 2020). The typical symptoms of COVID-19 are fever, cough, fatigue, and shortness of breath (Who, 2020a). About 15% of SARS-CoV2 positive cases become severe-to-critical with the following complications: pneumonia, acute respiratory distress syndrome, arrhythmia, septic shock, and/or multiple organ dysfunction/failure (Who, 2020a; Zheng et al., 2020).

A relationship between COVID-19 and cardiovascular disease (CVD) is becoming increasingly evident. Indeed, patients with CVD are at a higher risk of developing severe COVID-19 complications, and viral infection might, itself, induce cardiovascular injury. In this review, we provide an overview of the possible pathways that are common to SARS-CoV infections and CVD that might underlie this mutually reinforcing relationship. We also examine the possible modifying effect of some of the available therapies that could confer a protective effect or mitigate the severity of the disease complications.

CARDIOVASCULAR INVOLVEMENT IN COVID-19: A TWO-WAY ROAD

Increased Severity of COVID-19 in Patients With CVD

CVD is a risk factor for the progression of severe disease following lower respiratory tract infection by SARS-CoV or MERS-CoV (Oudit et al., 2009; Alhagbani, 2016). However, many factors complicate the accurate identification of prevalence of CVD in infected patients. Nevertheless, association between existing CVD and increased risk of progression to severe COVID-19 complications was suggested (Liu et al., 2020; Rodriguez-Morales et al., 2020; Zhou F. et al., 2020). The incidence of these comorbidities was higher in patients requiring intensive care admission (ICU) than non-ICU patients. Other studies show that the incidence of CVD in COVID-19 patients is relatively high (Guan et al., 2020; Yang et al., 2020). Moreover, the risk of developing acute respiratory distress or in-hospital death was shown to increase by at least 2-fold for hypertension and over 20-fold for coronary artery disease (Wu et al., 2020; Zhou F. et al., 2020). A report on the incidence of CVDs in the general population in China showed a 23.2% incidence rate for hypertension (Ma et al., 2020). This is

close to the rates at which these comorbidities appear among COVID-19 patients. As such, it is likely that CVD patients have an increased severity of COVID-19 complications rather than an increased vulnerability to infection. Indeed, this was shown to be the case in several recent studies (Li et al., 2020; Mehra et al., 2020). Therefore, it is important to identify if and which signalling pathways in CVD may augment SARS-CoV-2 pathogenesis.

SARS-CoV-2-Induced Myocardial Damage

Accumulating case studies document acute cardiac manifestations in COVID-19 patients, who were previously healthy (Inciardi et al., 2020). Typically, most studies define acute myocardial injury by elevated cardiac troponin I levels (Bansal, 2020; Huang et al., 2020; Zhou F. et al., 2020). High levels of troponin or creatine kinase were noted in a significant fraction of patients diagnosed with COVID-19 (Huang et al., 2020; Wang et al., 2020). Moreover, among patients who were without previous CVD but died following SARS-CoV-2 infection, 11.8% had high levels of troponin indicating myocardial injury and cardiac arrest (Wang et al., 2020). Another study showed that 27.8% of COVID-19 patients had myocardial injury resulting in cardiac dysfunction and arrhythmias (Guo et al., 2020). This study highlighted the impact of myocardial injury showing that patients with high troponin levels and no previous history of CVD had a mortality rate of 37.5%, whereas even those with underlying CVD and normal troponin levels demonstrated a better prognosis (13% mortality). Troponin levels were correlated with increases in C-reactive proteins suggesting a tight link between myocardial injury and inflammatory pathogenesis (Guo et al., 2020). Systemic inflammation is intimately related to reduced coronary blood flow and decreased oxygen supply to the heart even in the absence of other risk factors of coronary artery disease (Recio-Mayoral et al., 2009). Yet, myocardial damage induced by direct viral entry through binding to angiotensin converting enzyme-2 (ACE2) on cardiac cells could not be eliminated either (Oudit et al., 2009). Patients who recovered from infections by previous SARS-CoVs reported disruption in blood lipids and blood pressure after 12 years of recovery (Wu et al., 2017). Given the structural similarity between SARS-CoV and SARS-CoV-2, a similar profile could be expected.

SARS-CoV-2-Induced Endothelial Damage and Intravascular Coagulation

Viral infections are known to activate coagulation cascade, a mechanism thought to be protective as to limit viral spread (Milbrandt et al., 2009; Antoniak and Mackman, 2014). Yet, excessive coagulation could eventually lead to disseminated intravascular coagulation. Contextually, D-dimer and fibrin/fibrinogen degradation products were both higher in COVID-19 patients than controls, with the majority of non-survivals having disseminated intravascular coagulation during their hospital stay (Guan et al., 2020; Han et al., 2020; Tang et al., 2020; Zhou F. et al., 2020). The mechanism behind this activated and accelerated coagulation in COVID-19 patients could be an

inflammatory-immunological stimulation likely caused by vascular endothelial damage. Significantly, ACE2 is localized on endothelial cells (Gallagher et al., 2008), and SARS-CoV-2 has recently been shown to cause endothelial cell infections across vascular beds in COVID-19 patients (Varga et al., 2020). On the other hand, endothelial dysfunction and the activation of the clotting cascade are common occurrences in CVD (Lowe and Rumley, 2014; Widmer and Lerman, 2014). As such, a thorough understanding of the signaling pathways common to the pathogenesis of CVD and SARS-CoV-2 infection is necessary to identify crucial sites of crosstalk and direct future investigation of potential therapeutic interventions mitigating both COVID-19 complications in CVD patients as well as short- and long-term viral-induced cardiovascular impairment.

MOLECULAR PATHWAYS IMPLICATED IN THE CROSSTALK BETWEEN CVD AND SARS-COV-2 PATHOGENESIS

ACE2, the Renin-Angiotensin-Aldosterone System, and MAPK Pathways

CVD conditions are typically associated with increased RAAS activity, favoring increased AngII levels (Nehme and Zibara, 2017b). In this regard, AT₁ receptor activation facilitates various intracellular pathways involved in cardiac and vascular remodeling, endothelial dysfunction, and atherosclerosis (Muslin, 2008; Nehme and Zibara, 2017a). AngII-mediated AT₁ receptor stimulation activates protein kinases such as mitogen-activated protein kinases (MAPK) (Muslin, 2008; Kawai et al., 2017). On the other hand, Ang(1-7), produced by ACE2-catalyzed cleavage of AngII, is essential in repressing MAPK cascades (Zhang et al., 2014) and decreasing inflammation by keeping a well-balanced RAAS activity, between ACE/AngII pathway and ACE2/Ang(1-7) (Nehme and Zibara, 2017b). While the cardiovascular outcomes of ACE2 activity and Ang(1-7) production have been described in reasonable detail, it remains uncertain as to the nature of the receptor mediating these effects. Early studies reported that mitochondrial assembly 1 receptors (MasR) mediated the AngII/AT₁R antagonistic effect of Ang(1-7) including PI3k/Akt activation of nitric oxide synthase and NO production, vasodilation, and anti-fibrosis effects (Azushima et al., 2020). Yet, recent studies showed that Ang(1-7)-mediated AT₂R activation might contribute to the protective effect *via* activation of phosphotyrosine phosphatase and suppression of MAPK activity (Azushima et al., 2020). Moreover, other studies implicated other Mas-related G protein-coupled receptors as mediators of Ang(1-7) effects in the cardiovascular system (Tetzner et al., 2016). Nevertheless, the RAAS and MAPK pathways interaction has many consequences on CVD (summarized in **Table 1** and **Figure 1**) and is implicated in several steps of SARS-CoV-2 pathogenesis. As such, it is important to highlight potential points of hijack by SARS-

CoV-2, which are potentially augmented in CVD or could induce *de novo* cardiovascular injury. These points are summarized in **Figure 2**.

ACE2 and Viral Binding to Host Cells: Impact on CVD and Myocardial Injury

Current knowledge suggests that the membrane bound spikes of coronaviruses, known as S glycoproteins, are solely responsible for enabling host cell attachment and entry (Fehr and Perlman, 2015). SARS-CoV-2 gains host cell entry *via* ACE2 expressed on pulmonary alveolar epithelial cells (Letko et al., 2020). Recently, it has been suggested that two S glycoproteins can bind to a single ACE2 homodimer (Yan et al., 2020). Binding occurs *via* direct interaction of the ACE2 peptidase domain and the ACE2-binding domain on the S1 subunit of the S protein. ACE2 is also expressed on the surface of epithelial cells in lungs, heart, kidneys, and intestine, facilitating viral entry to these tissues and hence might underlie the multi-organ dysfunction seen with COVID-19 (Zhang et al., 2020). However, for proper entry into the cell, the S protein must be primed by host cell proteases at a second cleavage site on the membrane embedded S2 subunit (Bosch et al., 2003; Lu et al., 2015; Coutard et al., 2020). Specifically, cleavage at the S2' cleavage site exposes S2 domains, like the fusion peptide and internal fusion peptide, and enables them to partition into the membrane and facilitate entry (Lu et al., 2015). A recent study showed that compared to the proteases required for viral entry, ACE2 has a lower expression level in different epithelial cell types indicating that it is likely to be the rate limiting step in viral entry (Sungnak et al., 2020).

An early study on human ventricular tissue from heart failure patients showed an increased expression and activity of ACE2 (Zisman et al., 2003). Correspondingly, a recent study showed that heart failure patients with COVID-19 had increased levels of ACE2 placing them at higher risk of severity of symptoms (Chen et al., 2020). This might result from increased viral entry due to a higher ACE2 expression level. However, it may also be due to pathways triggered by the virus that can exacerbate CVD symptoms. Interestingly, the same study indicated that cardiac pericytes demonstrated a much higher level of ACE2 expression than cardiac myocytes raising the possibility that the initial cardiac injury might be microvascular. Previously, it was demonstrated that the endothelial cell expression of ACE2 provides a possible route of entry to SARS-CoV once it enters the blood (Ding et al., 2003). The same may be expected of SARS-CoV-2. Viral entry and inflammation could upregulate tissue factor expression on the cell surface, promoting the blood coagulation cascade (Witkowski et al., 2016). Fibrinoid necrosis and thrombotic events will therefore follow in microvessels leading to disseminated intravascular coagulation. Thus, endothelial injury exacerbates the immuno-inflammatory condition by inducing coagulation abnormalities (Vallet and Wiel, 2001). As such, supply of soluble ACE units could, in principle, reduce the infectious burden and rescue the tissues/organs from the

TABLE 1 | Interactions and examples of cardiovascular consequences of RAAS imbalance and MAPK pathway activation

Contribution of RAAS imbalance to CVD and interaction with MAPK pathways:	- \uparrow AngII \rightarrow \uparrow mechanisms involving growth factor receptors (e.g., PDGF and TGF- β receptors) \rightarrow \uparrow ERK activation cardiac and vascular hypertrophy and fibrosis	(Schellings et al., 2006; Ehanire et al., 2015)
\uparrow AngII/AT ₁ R pathway \rightarrow \uparrow MAPK signaling \rightarrow \uparrow signs of CVD	- \uparrow AngII activity in CVD \rightarrow \downarrow cell surface expression of ACE2 in an AT ₁ /p38 MAPK-dependent pathway leading to protein excision and shedding	(Patel et al., 2014; Xu et al., 2017)
\uparrow ACE2/Ang(1-7) pathway \rightarrow \downarrow MAPK signaling \rightarrow \downarrow inflammation \rightarrow amelioration of CVD	-ACE2 deletion \rightarrow \uparrow NADPH oxidase activity, superoxide generation, MAPK signaling and inflammatory cytokine production in the aorta of mice receiving AngII	(Jin et al., 2012)
Contribution of MAPK signaling to CVD	- \uparrow ACE2/Ang(1-7) pathway \rightarrow \downarrow AngII-induced inflammation in cardiac tissue and in hypothalamic cardiovascular centers, cardiac remodeling, and hypertension	(Grobe et al., 2007; Sriramula et al., 2011)
	-ACE2 overexpression \rightarrow \downarrow TNF- α , IL-1 β , and IL-6 in and \uparrow anti-inflammatory cytokine IL-10 in autoimmune myocarditis	(Sukumaran et al., 2011)
	-Erk1/2, JNK, and p38 MAPK activation \rightarrow \uparrow cardiomyopathic remodeling	(Wang, 2007)
	-Pressure overload \rightarrow \uparrow Ras/c-RAF/MKK1/ERK1/2 pathway \rightarrow cardiac hypertrophy, \uparrow cardiomyocyte size, diastolic dysfunction, and myofibril disarray	(Hunter et al., 1995)
	-ERK1/2 \rightarrow \uparrow c-Fos \rightarrow \uparrow GLUT1 transporter expression in hypertrophic and ischemic heart	(Babu et al., 2000) (Santalucia et al., 2003) (Shao and Tian, 2015)
	-MAPK pathway activation \rightarrow progressive endothelial dysfunction	(Huang et al., 2012)
	-Oxidized LDL \rightarrow \uparrow MAPK pathway activation \rightarrow \uparrow GM-CSF \rightarrow \uparrow macrophage infiltration of atherosclerotic plaques	(Proctor et al., 2007; Muslin, 2008)
	- \uparrow MAPK pathway activation \rightarrow \uparrow interleukins and TNF α \rightarrow \uparrow atherosclerotic lesion progression	(Bachstetter and Van Eldik, 2010; De Souza et al., 2014)
	-TNF α \rightarrow \uparrow ERK1/2 activity \rightarrow \uparrow MMP9 \rightarrow \uparrow fibrotic lesion formation	(Cheng et al., 2012; Zhang et al., 2017)
	-Viral infection \rightarrow \uparrow IL-1, TNF α , IL-6 \rightarrow \uparrow p38 MAPK and JNK \rightarrow \uparrow viral cardiomyopathy	(Monsuez et al., 2007; Wang et al., 2014; Yue-Chun et al., 2016; Niu et al., 2017)

PDGF, platelet-derived growth factor; TGF, transforming growth factor; TNF, tumor necrosis factor; MKK, MAPK kinase; GLUT, glucose transporter; LDL, low-density lipoprotein; MMP, matrix metalloproteinase.

detrimental outcome of SARS-CoV-2 infection. Indeed, a recent study showed that treatment with clinical grade human recombinant ACE2 reduced viral recovery from infected cell culture systems (Monteil et al., 2020). The same study also showed that soluble ACE2 treatment inhibited SARS-CoV-2 infection in blood vessel and kidney organoids.

On the other hand, binding of SARS-Cov-2 to ACE2 could eventually deplete ACE2 levels. Indeed, prior studies on the earlier SARS-CoV strain showed that pulmonary and myocardial infections were associated with marked reduction of ACE2 expression (Kuba et al., 2005; Oudit et al., 2009). This would inhibit the protective effects of the ACE2/Ang(1-7) pathway, offsetting the RAAS balance and causing potential exacerbation of respiratory symptoms and cardiovascular complications (Sun et al., 2020). In fact, ACE2 downregulation induced by viral spike glycoprotein injection led to increased pulmonary failure *in vivo* (Kuba et al., 2005). Therefore, ACE2 plays two contradictory roles in patients with cardiovascular/cardiometabolic diseases inflicted with COVID-19. On one hand, ACE2 serves as a gate for SARS-COV-2 infection and on another it protects against CVD as well as the increased production of inflammatory cytokines.

MAPK, Viral Entry, and the Hijacked Proteases

Non-endosomal SARS-CoV-2 uptake is mostly associated with the transmembrane serine protease TMPRSS2, in contrast to the less frequently used endosomal cysteine proteases cathepsin B and L (CatB/L) (Hoffmann et al., 2020). Interestingly, a unique furin-like protease recognition sequence at the S1/S2 cleavage site in SARS-CoV-2 is absent from the other members of the coronavirus family (Coutard et al., 2020). Furin is a serine protease and considered a proprotein convertase, which transforms protein precursors into active forms upon cleavage. Its function extends to a variety of proteins including growth hormones, cytokines, and surface glycoproteins of viruses (Coutard et al., 2020). It was previously established in another class of viruses that the presence of furin cleavage sites is associated with more pathogenicity as compared to cleavage by trypsin proteases (Kido et al., 2012). Incidentally, atherosclerotic patients demonstrate elevated levels of furin, particularly in foam cells of plaques (Zhao et al., 2018). Furthermore, this protease is regulated by ERK pathways under inflammatory conditions (Ventura et al., 2017). In fact, ERK1 is essential for maintaining activity of furin and has been implicated as a




FIGURE 1 | The RAAS balance between AngII/AT₁-R and ACE2/Ang(1-7)/Ang(1-7)R axes. The former is enhanced in CVD conditions triggering various intracellular pathways via MAPK signaling, which ultimately leads to cardiac and vascular remodeling, endothelial dysfunction, and atherosclerosis. ACE2, on the other hand, decreases inflammation by countering the effect of the ACE/AngII/AT₁-R axis. ACE2/Ang(1-7)/Ang(1-7)R signaling pathway not only ameliorates cellular proliferation, hypertrophy, oxidative stress, and vascular fibrosis but also reduces the activation of the downstream MAPK cascades. AT₁-R activation triggers p38 MPAK-dependent ACE2 excision and shedding leading to a reduced cell surface expression. As demonstrated in other coronaviruses, SARS-CoV-2 replication might be enhanced as a result of increased MAPK activity downstream of AngII in CVD patients.




FIGURE 2 | The MAPK pathway can be induced by several triggers in various cells within the context of CVDs. For example, pressure overload can cause hypertrophy in cardiomyocytes via the ERK1/2/GLUT pathway. Oxidized LDL triggers macrophage proliferation via ERK1/2/GM-CSF, which contributes to the development of atherosclerosis. All cells involved in atherosclerosis can release cytokines in an ERK1/2 dependent manner, which ultimately propel plaque and clot growth. Cytokines can subsequently enhance MMP9 production, which leads to plaque rupture and thrombosis. Meanwhile, coronaviruses have been shown to involve p38 MAPK, JNK, and MKK1/ERK1/2 pathways for viral pathogenesis. MKK1/ERK1/2 pathway also upregulates the protease furin, which is implicated in SARS-CoV-2 entry due to the unique furin-like S1/S2 cleavage site. It is worth examining if inhibition of MKK1/ERK1/2 mitigates production of SARS-CoV-2 viral progeny. On the other hand, SARS-CoV-2 can also amplify production of cytokines, which can worsen existing CVDs.

potential therapeutic target to combat the pro-inflammatory effects of TGF- β . Because ERK1 activity is heightened in CVD patients, it is likely that furin activity is also increased. Therefore, it is important to consider the role of elevated furin expression/function in the virulence of SARS-CoV-2.

MAPK and Viral Pathogenesis

Not only is the Raf/MEK1/2/ERK1/2 pathway heavily implicated in CVDs (**Table 1**), but it is also indispensable for effective viral replication (Li et al., 2019). Specifically, SARS-CoV spike and nucleocapsid proteins were reported to trigger ERK1/2 phosphorylation with subsequent induction of pro-inflammatory pathways including increased cyclooxygenase-2 expression and IL-8 release (Chang et al., 2004; Mizutani et al., 2004a; Yan et al., 2006; Liu et al., 2007). Significantly, while some report no change in SARS-CoV infected cell death after ERK1/2 inhibition (Mizutani et al., 2004a), inhibition of the MEK1/2/ERK1/2 pathway has been shown to significantly impair coronavirus replication in mice (Cai et al., 2007). Similarly, JNK activity increased in cells exposed to SARS-CoV spike and nucleocapsid proteins (Mizutani et al., 2004b; Surjit et al., 2004; Liu et al., 2007). Interestingly, JNK inhibition precluded the development of persistent SARS-CoV infections (Mizutani et al., 2005). Furthermore, SARS-CoV infection was shown to activate p38 MAPK and the downstream signaling possibly leading to cell death (Mizutani et al., 2004b; Surjit et al., 2004; Kopecky-Bromberg et al., 2006). Contextually, p38 MAPK inhibition reduced human coronavirus HCoV-229E viral replication in human lung epithelial cells (Kono et al., 2008). Moreover, regardless of the cell type or strain of the virus, MEK1/2 inhibition diminishes the production of viral progeny (Cai et al., 2007). As previously discussed, many MAPK pathways are already significantly upregulated in CVD possibly downstream of the AngII/AT₁ pathway activation. The chronic nature and gradual development timeframe of these diseases might argue that the crosstalk among increased MAPK signaling cascades and SARS-CoV-2 pathogenesis leads to increased infection severity and complications in patients with established CVD rather than being involved in viral-triggered cardiovascular involvement. Nevertheless, viral-induced myocardial injury and endothelial involvement are thought to include a strong inflammatory component as well as an offset of the RAAS balance away from ACE2/Ang(1-7) arm, both of which interacts closely with MAPK pathways (**Table 1**).

PHARMACOLOGICAL INTERVENTIONS MODIFYING PATHWAYS RELEVANT TO VIRAL INFECTION

Accumulating evidence shows that cardiovascular risk reduction, especially on the long term, involves modification of RAAS and MAPK signalling pathways, regardless of the particular intervention applied. For instance, exercise training reduces

p38 MAPK and ERK activities decreasing the incidence of heart failure (Nagata and Hattori, 2011) and ameliorates the AngII/Ang(1-7) imbalance in hypertensive rats (Ren et al., 2016). Similarly, dietary restriction reduced ERK activation (Xie et al., 2007; Castello et al., 2011) and decreased serum ACE activity (Harp et al., 2002). In this section, we provide an overview of the activity of some of the available therapeutic agents or classes with cardiovascular benefit. We focus on drug classes that can potentially modulate ACE2 and/or MAPK signalling pathways, hence decreasing cardiovascular risk on long-term use. We will then speculate on the possible impact of these agents to reduce risk of progression to severe COVID-19 in CVD patients. As will be seen with some of these agents, clinical and research interest in their use for the mitigation of acute complications of SARS-CoV-2 infections has also emerged.

ACEIs/ARBs

Antihypertensive drugs such as angiotensin converting enzyme inhibitors (ACEI) and AngII receptor blockers (ARBs) gained interest in the context of treating patients with COVID-19. Several studies reported conflicting findings regarding the effects of ACEI/ARBs on ACE2 levels. Some of which reported an increase in ACE2 levels that could be explained based on inhibition of AT₁ receptor-mediated ACE2 shedding (Patel et al., 2014; Xu et al., 2017), whereas others argued for a role for higher levels of AngII detected after treatment with ARBs, a case of higher substrate availability leading to an increase in expression of the linked enzyme (Dickstein et al., 1994; Gavras and Gavras, 1999; Ferrario et al., 2005; Soler et al., 2009; Sukumaran et al., 2011; Furuhashi et al., 2015; Esler and Esler, 2020). Yet, other studies have documented no effect of ARBs in this regard (Campbell et al., 2004; Burchill et al., 2012). While it could be postulated that treatment with ACEI/ARBs increasing ACE2 levels might increase risk of SARS-COV-2 infection severity, recent studies showed that continuation of these drugs in COVID-19 patients was not associated with such risk (Mancia et al., 2020; Mehra et al., 2020; Reynolds et al., 2020). On the other hand, ACE2 internalization was shown to be dependent on heterodimerization with AT₁-R and subsequent activation of AT₁-R with AngII (Deshotels et al., 2014). Hence, even if these drugs increase ACE2 expression, they could confer a protective effect *via* suppression of ACE2-mediated viral internalization (Deshotels et al., 2014) by blocking AT₁-R in case of ARBs or reducing the availability of AngII in case of ACEIs. Moreover, reduced induction of MAPK pathways triggered downstream of AT₁-R might contribute to a less severe viral pathogenesis. Along these lines, a study showed that ACE2 knockout exacerbated severe acute respiratory failure following acid aspiration in mice that was rescued by ARB or recombinant ACE2 treatment (Imai et al., 2005). In this context, elevated systemic AngII in CVD patients could underlie higher risk of infection and further complications as discussed previously. Nevertheless, the studies that reported lack of increased risk with ACEI/ARB use were not designed to detect potential benefit; thus, more focused prospective studies are required to answer this question. On

this basis, losartan, an ARB, is now being incorporated in two clinical studies investigating possible reduction of lung injury in patients with COVID-19 (clinicaltrials.gov/ct2/show/NCT04311177, clinicaltrials.gov/ct2/show/NCT04312009).

Statins

Statins are HMG-CoA reductase inhibitors used for treatment of CVD associated with dyslipidemia. Aside from their hypocholesterolemic effect, statins have a proven anti-inflammatory effect. They have been proposed as an adjunctive therapy in influenza virus infection (Arabi et al., 2020). Simvastatin improved survival in patients with hyper-inflammatory acute respiratory distress syndrome (Calfee et al., 2018). Specifically, atorvastatin attenuated NF- κ B signaling within 24 h of MERS-CoV infection (Yuan, 2015). Similarly, atorvastatin showed an anti-inflammatory effect that was dose- and time-dependent effectively attenuating NF- κ B ultimately leading to decreased synthesis of inflammatory cytokines and chemokines (Chansrichavala et al., 2009). Significantly, statins, *via* activating Akt signaling, enhance endothelial junction integrity and reduce plasma leakage and acute lung injury (Fedson, 2016). Moreover, rosuvastatin reduced AngII production, AT1 receptor expression, and Erk activation and upregulated ACE2 in rats following vascular injury (Li et al., 2013). Taken together, these observations suggest that CVD patients undergoing statin treatment might benefit from a mitigated basal and SARS-CoV-2-induced inflammatory reaction and hence a reduced risk of progression to severe COVID-19.

SGLT-2 Inhibitors

Sodium-glucose cotransporter type 2 (SGLT-2) inhibitors have been approved to reduce major cardiovascular events in patients with diabetes though direct mechanisms not necessarily related to their hypoglycemic effects (Kaplan et al., 2018; Scheen, 2018). SGLT-2 inhibitors were found to induce a modest increase in plasma levels of aldosterone and AngII, but still within the low range typical in diabetes, as well as an increase in ACE2, ACE, and angiotensinogen (Kawanami et al., 2017). When administered with ACEI or ARBs, the effect of SGLT-2 inhibitors will shunt the RAAS activation towards ACE2 (Kawanami et al., 2017), thus significantly increasing the beneficial effects of Ag(1-7) leading a vasodilatory, anti-inflammatory, and antioxidative effect in the heart and kidneys. More research is needed to evaluate the possible beneficial use of SGLT-2 inhibitors with ARBs or ACEIs on the cardiovascular and metabolic events in COVID-19.

Aldosterone Receptor Antagonists

Eplerenone, an aldosterone receptor antagonist, is recommended for the treatment of hypertension, left ventricular hypertrophy, and congestive heart failure (Jewell et al., 2006). Patients treated with eplerenone had lower morbidity and mortality due to CVD,

a protective effect thought to be independent on blood pressure. Aldosterone induces cell growth and differentiation through activation of Erk and downstream signaling including NF- κ B, suggesting a protective effect of eplerenone *via* inhibition of Erk (Kobayashi et al., 2006). In addition, aldosterone receptor blockade improves cardiac function and remodeling *via* increasing eNOS expression and Akt-mediated phosphorylation (Kobayashi et al., 2005). This raises the possibility for a beneficial effect of eplerenone or other aldosterone receptor antagonists in reducing risk of severe COVID-19 infection in CVD patients or to mitigate the cardiovascular burden of SARS-CoV-2 infection.

Tocilizumab

Interest in the use of tocilizumab, a monoclonal antibody against IL-6, in COVID-19 patients has emerged especially in those at risk of exaggerated inflammatory response (Luo et al., 2020). In the latter study, about two-thirds of the patients with elevated serum IL-6 levels had co-morbid CVD. Treatment with tocilizumab, originally approved for treatment of rheumatoid arthritis, demonstrated several cardiovascular effects. Tocilizumab led to an increased total cholesterol, low density lipoprotein, and high-density lipoprotein levels increasing the risk of dyslipidemia (Singh et al., 2010). Yet other studies reported a strong reduction in the incidence of CVD with tocilizumab treatment despite the change in lipid profile (Rao et al., 2015). It could be argued that high IL-6 levels induce cardiac fibrosis and increase myocardial infarction and CVDs, a pathway linked to aldosterone activity (Chou et al., 2018), thus justifying a positive cardiovascular impact of tocilizumab treatment. As such, more studies are needed to clarify the possible modulatory effect exerted by IL-6 antagonism in COVID-19 patients, and its impact on the intersecting pathways leading to respiratory and cardiovascular complications.

CONCLUSIONS AND FUTURE DIRECTIONS

Whereas knowledge of the exact mechanisms underlying the pathogenesis and the sequelae of SARS-CoV-2 infection is still emerging, the available evidence could be perceived to form a framework implicating RAAS and MAPK pathway imbalances in the observed association between COVID-19 and cardiovascular dysfunction. On the one hand, predominance of the AngII/AT₁ receptor arm of RAAS as well as increased MAPK activity in CVD could enhance viral pathogenesis and thus predispose the patients for increased COVID-19 severity. While on the other hand, viral activity including cell entry, viral replication, and induction of inflammatory response might also trigger the same pathway imbalances inducing or exacerbating cardiovascular damage. Despite the presumed efficacy of some of the available therapeutic tools in interrupting this vicious cycle, significant investigation is needed to direct evidence-based clinical use.

Moreover, a large window of opportunity exists for the identification and design of selective tools to interfere with these pathways in COVID-19 patients with CVD risk.

AUTHOR CONTRIBUTIONS

AE-Y and AE developed the idea and the review framework. ZW, SH, HZ, and NS wrote the first draft of the manuscript. All

authors contributed to corrections and adjustment of subsequent iterations of the manuscript. All authors approve and agree with the content.

FUNDING

This paper was supported by an AUM-FM MPP grant number 320148 to AE-Y and grant number 320133 to AE.

REFERENCES

- Alhighbani, T. (2016). Acute myocarditis associated with novel Middle east respiratory syndrome coronavirus. *Ann. Saudi Med.* 36, 78–80. doi: 10.5144/0256-4947.2016.78
- Antoniak, S., and Mackman, N. (2014). Multiple roles of the coagulation protease cascade during virus infection. *Blood* 123, 2605–2613. doi: 10.1182/blood-2013-09-526277
- Arabi, Y. M., Fowler, R., and Hayden, F. G. (2020). Critical care management of adults with community-acquired severe respiratory viral infection. *Intensive Care Med.* 46, 315–328. doi: 10.1007/s00134-020-05943-5
- Azushima, K., Morisawa, N., Tamura, K., and Nishiyama, A. (2020). Recent Research Advances in Renin-Angiotensin-Aldosterone System Receptors. *Curr. Hypertens. Rep.* 22, 22. doi: 10.1007/s11906-020-1028-6
- Babu, G. J., Lalli, M. J., Sussman, M. A., Sadoshima, J., and Periasamy, M. (2000). Phosphorylation of elk-1 by MEK/ERK pathway is necessary for c-fos gene activation during cardiac myocyte hypertrophy. *J. Mol. Cell Cardiol.* 32, 1447–1457. doi: 10.1006/jmcc.2000.1185
- Bachstetter, A. D., and Van Eldik, L. J. (2010). The p38 MAP Kinase Family as Regulators of Proinflammatory Cytokine Production in Degenerative Diseases of the CNS. *Aging Dis.* 1, 199–211.
- Bansal, M. (2020). Cardiovascular disease and COVID-19. *Diabetes Metab. Syndr.* 14, 247–250. doi: 10.1016/j.dsx.2020.03.013
- Bosch, B. J., Van Der Zee, R., De Haan, C. A., and Rottier, P. J. (2003). The coronavirus spike protein is a class I virus fusion protein: structural and functional characterization of the fusion core complex. *J. Virol* 77, 8801–8811. doi: 10.1128/JVI.77.16.8801-8811.2003
- Burchill, L. J., Velkoska, E., Dean, R. G., Griggs, K., Patel, S. K., and Burrell, L. M. (2012). Combination renin-angiotensin system blockade and angiotensin-converting enzyme 2 in experimental myocardial infarction: implications for future therapeutic directions. *Clin. Sci. (Lond)* 123, 649–658. doi: 10.1042/CS20120162
- Cai, Y., Liu, Y., and Zhang, X. (2007). Suppression of coronavirus replication by inhibition of the MEK signaling pathway. *J. Virol* 81, 446–456. doi: 10.1128/JVI.01705-06
- Calfee, C. S., Delucchi, K. L., Sinha, P., Matthay, M. A., Hackett, J., Shankar-Hari, M., et al. (2018). Acute respiratory distress syndrome subphenotypes and differential response to simvastatin: secondary analysis of a randomised controlled trial. *Lancet Respir. Med.* 6, 691–698. doi: 10.1016/S2213-2600(18)30177-2
- Campbell, D. J., Zeitz, C. J., Esler, M. D., and Horowitz, J. D. (2004). Evidence against a major role for angiotensin converting enzyme-related carboxypeptidase (ACE2) in angiotensin peptide metabolism in the human coronary circulation. *J. Hypertens.* 22, 1971–1976. doi: 10.1097/00004872-200410000-00020
- Castello, L., Maina, M., Testa, G., Cavallini, G., Biasi, F., Donati, A., et al. (2011). Alternate-day fasting reverses the age-associated hypertrophy phenotype in rat heart by influencing the ERK and PI3K signaling pathways. *Mech. Ageing Dev.* 132, 305–314. doi: 10.1016/j.mad.2011.06.006
- Chang, Y. J., Liu, C. Y., Chiang, B. L., Chao, Y. C., and Chen, C. C. (2004). Induction of IL-8 release in lung cells via activator protein-1 by recombinant baculovirus displaying severe acute respiratory syndrome-coronavirus spike proteins: identification of two functional regions. *J. Immunol.* 173, 7602–7614. doi: 10.4049/jimmunol.173.12.7602
- Chansrichavala, P., Chantharaksri, U., Sritara, P., and Chaiyaroj, S. C. (2009). Atorvastatin attenuates TLR4-mediated NF-kappaB activation in a MyD88-dependent pathway. *Asian Pac J. Allergy Immunol.* 27, 49–57.
- Chen, L., Li, X., Chen, M., Feng, Y., and Xiong, C. (2020). The ACE2 expression in human heart indicates new potential mechanism of heart injury among patients infected with SARS-CoV-2. *Cardiovasc. Res.* 116, 1097–1100. doi: 10.1093/cvr/cvaa078
- Cheng, C. Y., Hsieh, H. L., Hsiao, L. D., and Yang, C. M. (2012). PI3-K/Akt/JNK/NF- κ B is essential for MMP-9 expression and outgrowth in human limbal epithelial cells on intact amniotic membrane. *Stem Cell Res.* 9, 9–23. doi: 10.1016/j.scr.2012.02.005
- Chou, C. H., Hung, C. S., Liao, C. W., Wei, L. H., Chen, C. W., Shun, C. T., et al. (2018). IL-6 trans-signalling contributes to aldosterone-induced cardiac fibrosis. *Cardiovasc. Res.* 114, 690–702. doi: 10.1093/cvr/cvy013
- Coutard, B., Valle, C., De Lamballerie, X., Canard, B., Seidah, N. G., and Decroly, E. (2020). The spike glycoprotein of the new coronavirus 2019-nCoV contains a furin-like cleavage site absent in CoV of the same clade. *Antiviral Res.* 176, 104742. doi: 10.1016/j.antiviral.2020.104742
- De Souza, A. P., Vale, V. L., Silva, M., Araújo, I. B., Trindade, S. C., De Moura-Costa, L. F., et al. (2014). MAPK involvement in cytokine production in response to *Corynebacterium pseudotuberculosis* infection. *BMC Microbiol* 14, 230. doi: 10.1186/s12866-014-0230-6
- Deshotel, M. R., Xia, H., Sriramula, S., Lazartigues, E., and Filipeanu, C. M. (2014). Angiotensin II mediates angiotensin converting enzyme type 2 internalization and degradation through an angiotensin II type I receptor-dependent mechanism. *Hypertension* 64, 1368–1375. doi: 10.1161/HYPERTENSIONAHA.114.03743
- Dickstein, K., Gottlieb, S., Fleck, E., Kostis, J., Levine, B., Dekock, M., et al. (1994). Hemodynamic and neurohumoral effects of the angiotensin II antagonist losartan in patients with heart failure. *J. Hypertens. Suppl.* 12, S31–S35.
- Ding, Y., Wang, H., Shen, H., Li, Z., Geng, J., Han, H., et al. (2003). The clinical pathology of severe acute respiratory syndrome (SARS): a report from China. *J. Pathol.* 200, 282–289. doi: 10.1002/path.1440
- Ehanire, T., Ren, L., Bond, J., Medina, M., Li, G., Bashirov, L., et al. (2015). Angiotensin II stimulates canonical TGF- β signaling pathway through angiotensin type 1 receptor to induce granulation tissue contraction. *J. Mol. Med. (Berl)* 93, 289–302. doi: 10.1007/s00109-014-1211-9
- Esler, M., and Esler, D. (2020). Can angiotensin receptor-blocking drugs perhaps be harmful in the COVID-19 pandemic? *J. Hypertens.* 38, 781–782. doi: 10.1097/HJH.0000000000002450
- Fedson, D. S. (2016). Treating the host response to emerging virus diseases: lessons learned from sepsis, pneumonia, influenza and Ebola. *Ann. Transl. Med.* 4, 421. doi: 10.21037/atm.2016.11.03
- Fehr, A. R., and Perlman, S. (2015). Coronaviruses: an overview of their replication and pathogenesis. *Methods Mol. Biol.* 1282, 1–23. doi: 10.1007/978-1-4939-2438-7_1
- Ferrario, C. M., Jessup, J., Chappell, M. C., Averill, D. B., Brosnihan, K. B., Tallant, E. A., et al. (2005). Effect of angiotensin-converting enzyme inhibition and angiotensin II receptor blockers on cardiac angiotensin-converting enzyme 2. *Circulation* 111, 2605–2610. doi: 10.1161/CIRCULATIONAHA.104.510461
- Furuhashi, M., Moniwa, N., Mita, T., Fuseya, T., Ishimura, S., Ohno, K., et al. (2015). Urinary angiotensin-converting enzyme 2 in hypertensive patients may be increased by olmesartan, an angiotensin II receptor blocker. *Am. J. Hypertens.* 28, 15–21. doi: 10.1093/ajh/hpu086

- Gallagher, P. E., Ferrario, C. M., and Tallant, E. A. (2008). Regulation of ACE2 in cardiac myocytes and fibroblasts. *Am. J. Physiol. Heart Circ. Physiol.* 295, H2373–H2379. doi: 10.1152/ajpheart.00426.2008
- Gavras, I., and Gavras, H. (1999). Effects of eprosartan versus enalapril in hypertensive patients on the renin-angiotensin-aldosterone system and safety parameters: results from a 26-week, double-blind, multicentre study. Eprosartan Multinational Study Group. *Curr. Med. Res. Opin.* 15, 15–24. doi: 10.1185/03007999909115169
- Grobe, J. L., Mecca, A. P., Lingis, M., Shenoy, V., Bolton, T. A., Machado, J. M., et al. (2007). Prevention of angiotensin II-induced cardiac remodeling by angiotensin-(1-7). *Am. J. Physiol. Heart Circ. Physiol.* 292, H736–H742. doi: 10.1152/ajpheart.00937.2006
- Guan, W. J., Ni, Z. Y., Hu, Y., Liang, W. H., Ou, C. Q., He, J. X., et al. (2020). Clinical Characteristics of Coronavirus Disease 2019 in China. *N Engl. J. Med.* 382, 1708–1720. doi: 10.1056/NEJMoa2002032
- Guo, T., Fan, Y., Chen, M., Wu, X., Zhang, L., He, T., et al. (2020). Cardiovascular Implications of Fatal Outcomes of Patients With Coronavirus Disease 2019 (COVID-19). *JAMA Cardiol.* doi: 10.1001/jamacardio.2020.1017
- Han, H., Yang, L., Liu, R., Liu, F., Wu, K. L., Li, J., et al. (2020). Prominent changes in blood coagulation of patients with SARS-CoV-2 infection. *Clin. Chem. Lab. Med.* doi: 10.1515/cclm-2020-0188
- Harp, J. B., Henry, S. A., and Digirolamo, M. (2002). Dietary Weight Loss Decreases Serum Angiotensin-Converting Enzyme Activity in Obese Adults. *Obes Res.* 10, 985–990. doi: 10.1038/oby.2002.134
- Hoffmann, M., Kleine-Weber, H., Schroeder, S., Krüger, N., Herrler, T., Erichsen, S., et al. (2020). SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. *Cell* 181, 271–280.e278. doi: 10.1016/j.cell.2020.02.052
- Huang, A., Yang, Y.-M., Yan, C., Kaley, G., Hintze, T. H., and Sun, D. (2012). Altered MAPK signaling in progressive deterioration of endothelial function in diabetic mice. *Diabetes* 61, 3181–3188. doi: 10.2337/db12-0559
- Huang, C., Wang, Y., Li, X., Ren, L., Zhao, J., Hu, Y., et al. (2020). Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 395, 497–506. doi: 10.1016/S0140-6736(20)30183-5
- Hunter, J. J., Tanaka, N., Rockman, H. A., Ross, J.Jr., and Chien, K. R. (1995). Ventricular expression of a MLC-2v-ras fusion gene induces cardiac hypertrophy and selective diastolic dysfunction in transgenic mice. *J. Biol. Chem.* 270, 23173–23178. doi: 10.1074/jbc.270.39.23173
- Imai, Y., Kuba, K., Rao, S., Huan, Y., Guo, F., Guan, B., et al. (2005). Angiotensin-converting enzyme 2 protects from severe acute lung failure. *Nature* 436, 112–116. doi: 10.1038/nature03712
- Inciardi, R. M., Lupi, L., Zaccone, G., Italia, L., Raffo, M., Tomasoni, D., et al. (2020). Cardiac involvement in a patient with coronavirus disease 2019 (COVID-19). *JAMA Cardiol.*
- Jewell, C., Watson, L. E., Mock, J., and Dostal, D. E. (2006). Aldosterone receptor antagonists and cardiovascular disease: Do we need a change of the guard? *Cardiovasc. Hematol. Agents Med. Chem.* 4, 129–153. doi: 10.2174/187152506776369935
- Jin, H. Y., Song, B., Oudit, G. Y., Davidge, S. T., Yu, H. M., Jiang, Y. Y., et al. (2012). ACE2 deficiency enhances angiotensin II-mediated aortic profilin-1 expression, inflammation and peroxynitrite production. *PLoS One* 7, e38502. doi: 10.1371/journal.pone.0038502
- Kaplan, A., Abidi, E., El-Yazbi, A., Eid, A., Booz, G. W., and Zouein, F. A. (2018). Direct cardiovascular impact of SGLT2 inhibitors: mechanisms and effects. *Heart Fail Rev.* 23, 419–437. doi: 10.1007/s10741-017-9665-9
- Kawai, T., Forrester, S. J., O'Brien, S., Baggett, A., Rizzo, V., and Eguchi, S. (2017). AT1 receptor signaling pathways in the cardiovascular system. *Pharmacol. Res.* 125, 4–13. doi: 10.1016/j.phrs.2017.05.008
- Kawanami, D., Matoba, K., Takeda, Y., Nagai, Y., Akamine, T., Yokota, T., et al. (2017). SGLT2 Inhibitors as a Therapeutic Option for Diabetic Nephropathy. *Int. J. Mol. Sci.* 18, 1083. doi: 10.3390/ijms18051083
- Kido, H., Okumura, Y., Takahashi, E., Pan, H. Y., Wang, S., Yao, D., et al. (2012). Role of host cellular proteases in the pathogenesis of influenza and influenza-induced multiple organ failure. *Biochim. Biophys. Acta* 1824, 186–194. doi: 10.1016/j.bbapap.2011.07.001
- Kobayashi, N., Hara, K., Tojo, A., Onozato, M. L., Honda, T., Yoshida, K., et al. (2005). Eplerenone shows renoprotective effect by reducing LOX-1-mediated adhesion molecule, PKCepsilon-MAPK-p90RSK, and Rho-kinase pathway. *Hypertension* 45, 538–544. doi: 10.1161/01.HYP.0000157408.43807.5a
- Kobayashi, N., Yoshida, K., Nakano, S., Ohno, T., Honda, T., Tsubokou, Y., et al. (2006). Cardioprotective mechanisms of eplerenone on cardiac performance and remodeling in failing rat hearts. *Hypertension* 47, 671–679. doi: 10.1161/01.HYP.0000203148.42892.7a
- Kono, M., Tatsumi, K., Imai, A. M., Saito, K., Kuriyama, T., and Shirasawa, H. (2008). Inhibition of human coronavirus 229E infection in human epithelial lung cells (L132) by chloroquine: Involvement of p38 MAPK and ERK. *Antiviral Res.* 77, 150–152. doi: 10.1016/j.antiviral.2007.10.011
- Kopecky-Bromberg, S. A., Martinez-Sobrido, L., and Palese, P. (2006). 7a protein of severe acute respiratory syndrome coronavirus inhibits cellular protein synthesis and activates p38 mitogen-activated protein kinase. *J. Virol* 80, 785–793. doi: 10.1128/JVI.80.2.785-793.2006
- Kuba, K., Imai, Y., Rao, S., Gao, H., Guo, F., Guan, B., et al. (2005). A crucial role of angiotensin converting enzyme 2 (ACE2) in SARS coronavirus-induced lung injury. *Nat. Med.* 11, 875–879. doi: 10.1038/nm1267
- Letko, M., Marzi, A., and Munster, V. (2020). Functional assessment of cell entry and receptor usage for SARS-CoV-2 and other lineage B betacoronaviruses. *Nat. Microbiol* 5, 562–569. doi: 10.1038/s41564-020-0688-y
- Li, Y.-H., Wang, Q.-X., Zhou, J.-W., Chu, X.-M., Man, Y.-L., Liu, P., et al. (2013). Effects of rosuvastatin on expression of angiotensin-converting enzyme 2 after vascular balloon injury in rats. *J. Geriatr. Cardiol.* 10, 151–158. doi: 10.3969/j.issn.1671-5411.2013.02.009
- Li, C. C., Wang, X. J., and Wang, H. R. (2019). Repurposing host-based therapeutics to control coronavirus and influenza virus. *Drug Discovery Today* 24, 726–736. doi: 10.1016/j.drudis.2019.01.018
- Li, B., Yang, J., Zhao, F., Zhi, L., Wang, X., Liu, L., et al. (2020). Prevalence and impact of cardiovascular metabolic diseases on COVID-19 in China. *Clin. Res. Cardiol.* 109, 531–538. doi: 10.1007/s00392-020-01626-9
- Liu, M., Yang, Y., Gu, C., Yue, Y., Wu, K. K., Wu, J., et al. (2007). Spike protein of SARS-CoV stimulates cyclooxygenase-2 expression via both calcium-dependent and calcium-independent protein kinase C pathways. *FASEB J.* 21, 1586–1596. doi: 10.1096/fj.06-6589.com
- Liu, K., Fang, Y. Y., Deng, Y., Liu, W., Wang, M. F., Ma, J. P., et al. (2020). Clinical characteristics of novel coronavirus cases in tertiary hospitals in Hubei Province. *Chin Med. J. (Engl)* 133, 1025–1031. doi: 10.1097/CM9.0000000000000744
- Lowe, G., and Rumley, A. (2014). The relevance of coagulation in cardiovascular disease: what do the biomarkers tell us? *Thromb. Haemost.* 112, 860–867. doi: 10.1160/th14-03-0199
- Lu, G., Wang, Q., and Gao, G. F. (2015). Bat-to-human: spike features determining 'host jump' of coronaviruses SARS-CoV, MERS-CoV, and beyond. *Trends Microbiol* 23, 468–478. doi: 10.1016/j.tim.2015.06.003
- Lu, R., Zhao, X., Li, J., Niu, P., Yang, B., Wu, H., et al. (2020). Genomic characterisation and epidemiology of 2019 novel coronavirus: implications for virus origins and receptor binding. *Lancet* 395, 565–574. doi: 10.1016/S0140-6736(20)30251-8
- Luo, P., Liu, Y., Qiu, L., Liu, X., Liu, D., and Li, J. (2020). Tocilizumab treatment in COVID-19: A single center experience. *J. Med. Virol.* doi: 10.1002/jmv.25801
- Ma, L. Y., Chen, W. W., Gao, R. L., Liu, L. S., Zhu, M. L., Wang, Y. J., et al. (2020). China cardiovascular diseases report 2018: an updated summary. *J. Geriatr. Cardiol.* 17, 1–8. doi: 10.11909/j.issn.1671-5411.2020.01.001
- Mancia, G., Rea, F., Ludergnani, M., Apolone, G., and Corrao, G. (2020). Renin-Angiotensin-Aldosterone System Blockers and the Risk of Covid-19. *N Engl. J. Med.* doi: 10.1056/NEJMoa2006923
- Mehra, M. R., Desai, S. S., Kuy, S., Henry, T. D., and Patel, A. N. (2020). Cardiovascular Disease, Drug Therapy, and Mortality in Covid-19. *N Engl. J. Med.* doi: 10.1056/NEJMoa2007621
- Milbrandt, E. B., Reade, M. C., Lee, M., Shook, S. L., Angus, D. C., Kong, L., et al. (2009). Prevalence and significance of coagulation abnormalities in community-acquired pneumonia. *Mol. Med.* 15, 438–445. doi: 10.2119/molmed.2009.00091
- Mizutani, T., Fukushi, S., Murakami, M., Hirano, T., Saijo, M., Kurane, I., et al. (2004a). Tyrosine dephosphorylation of STAT3 in SARS coronavirus-infected Vero E6 cells. *FEBS Lett.* 577, 187–192. doi: 10.1016/j.febslet.2004.10.005
- Mizutani, T., Fukushi, S., Saijo, M., Kurane, I., and Morikawa, S. (2004b). Phosphorylation of p38 MAPK and its downstream targets in SARS

- coronavirus-infected cells. *Biochem. Biophys. Res. Commun.* 319, 1228–1234. doi: 10.1016/j.bbrc.2004.05.107
- Mizutani, T., Fukushi, S., Saijo, M., Kurane, I., and Morikawa, S. (2005). JNK and PI3k/Akt signaling pathways are required for establishing persistent SARS-CoV infection in Vero E6 cells. *Biochim. Biophys. Acta* 1741, 4–10. doi: 10.1016/j.bbdis.2005.04.004
- Monsuez, J.-J., Escaut, L., Teicher, E., Charniot, J.-C., and Vittecoq, D. (2007). Cytokines in HIV-associated cardiomyopathy. *Int. J. Cardiol.* 120, 150–157. doi: 10.1016/j.ijcard.2006.11.143
- Monteil, V., Kwon, H., Prado, P., Hagelkrüys, A., Wimmer, R. A., Stahl, M., et al. (2020). Inhibition of SARS-CoV-2 Infections in Engineered Human Tissues Using Clinical-Grade Soluble Human ACE2. *Cell* 181, 905–913. doi: 10.1016/j.cell.2020.04.004
- Muslin, A. J. (2008). MAPK signalling in cardiovascular health and disease: molecular mechanisms and therapeutic targets. *Clin. Sci. (Lond)* 115, 203–218. doi: 10.1042/CS20070430
- Nagata, K., and Hattori, T. (2011). Cardioprotective mechanisms of lifestyle modifications and pharmacotherapies on cardiac remodeling and dysfunction in hypertensive heart disease: an overview. *Nagoya J. Med. Sci.* 73, 91–105.
- Nehme, A., and Zibara, K. (2017a). Cellular distribution and interaction between extended renin-angiotensin-aldosterone system pathways in atherosclerosis. *Atherosclerosis* 263, 334–342. doi: 10.1016/j.atherosclerosis.2017.05.029
- Nehme, A., and Zibara, K. (2017b). Efficiency and specificity of RAAS inhibitors in cardiovascular diseases: how to achieve better end-organ protection? *Hypertens. Res.* 40, 903–909. doi: 10.1038/hr.2017.65
- Niu, L., Li, C., Wang, Z., Xu, H., and An, X. (2017). Effects of the MAPK pathway and the expression of CAR in a murine model of viral myocarditis. *Exp. Ther. Med.* 13, 230–234. doi: 10.3892/etm.2016.3909
- Oudit, G. Y., Kassiri, Z., Jiang, C., Liu, P. P., Poutanen, S. M., Penninger, J. M., et al. (2009). SARS-coronavirus modulation of myocardial ACE2 expression and inflammation in patients with SARS. *Eur. J. Clin. Invest.* 39, 618–625. doi: 10.1111/j.1365-2362.2009.02153.x
- Patel, V. B., Clarke, N., Wang, Z., Fan, D., Parajuli, N., Basu, R., et al. (2014). Angiotensin II induced proteolytic cleavage of myocardial ACE2 is mediated by TACE/ADAM-17: a positive feedback mechanism in the RAS. *J. Mol. Cell Cardiol.* 66, 167–176. doi: 10.1016/j.yjmcc.2013.11.017
- Prevention, C.C.F.D.C.A. (2020). *Coronavirus (COVID-19)*, [Online]. [Accessed March 29, 2020].
- Proctor, B. M., Ren, J., Chen, Z., Schneider, J. G., Coleman, T., Lupu, T. S., et al. (2007). Grb2 is required for atherosclerotic lesion formation. *Arterioscler Thromb. Vasc. Biol.* 27 (6), 1361–1367. doi: 10.1161/ATVBAHA.106.134007
- Rao, V. U., Pavlov, A., Klearman, M., Musselman, D., Giles, J. T., Bathon, J. M., et al. (2015). An evaluation of risk factors for major adverse cardiovascular events during tocilizumab therapy. *Arthritis Rheumatol* 67, 372–380. doi: 10.1002/art.38920
- Recio-Mayoral, A., Mason, J. C., Kaski, J. C., Rubens, M. B., Harari, O. A., and Camici, P. G. (2009). Chronic inflammation and coronary microvascular dysfunction in patients without risk factors for coronary artery disease. *Eur. Heart J.* 30, 1837–1843. doi: 10.1093/eurheartj/ehp205
- Ren, C.-Z., Yang, Y.-H., Sun, J.-C., Wu, Z.-T., Zhang, R.-W., Shen, D., et al. (2016). Exercise Training Improves the Altered Renin-Angiotensin System in the Rostral Ventrolateral Medulla of Hypertensive Rats. *Oxid. Med. Cell Longev* 2016, 7413963–7413963. doi: 10.1155/2016/7413963
- Reynolds, H. R., Adhikari, S., Pulgarin, C., Troxel, A. B., Itrurra, E., Johnson, S. B., et al. (2020). Renin–Angiotensin–Aldosterone System Inhibitors and Risk of Covid-19. *N Engl. J. Med.* doi: 10.1056/NEJMoa2008975
- Rodriguez-Morales, A. J., Cardona-Ospina, J. A., Gutierrez-Ocampo, E., Villamizar-Pena, R., Holguin-Rivera, Y., Escalera-Antezana, J. P., et al. (2020). Clinical, laboratory and imaging features of COVID-19: A systematic review and meta-analysis. *Travel Med. Infect. Dis.* 101623. doi: 10.1016/j.tmaid.2020.101623
- Santalucia, T., Christmann, M., Yacoub, M. H., and Brand, N. J. (2003). Hypertrophic agonists induce the binding of c-Fos to an AP-1 site in cardiac myocytes: implications for the expression of GLUT1. *Cardiovasc. Res.* 59, 639–648. doi: 10.1016/S0008-6363(03)00472-3
- Scheen, A. J. (2018). Cardiovascular outcome studies in type 2 diabetes: Comparison between SGLT2 inhibitors and GLP-1 receptor agonists. *Diabetes Res. Clin. Pract.* 143, 88–100. doi: 10.1016/j.diabres.2018.06.008
- Schellings, M. W., Baumann, M., Van Leeuwen, R. E., Duisters, R. F., Janssen, S. H., Schroen, B., et al. (2006). Imatinib attenuates end-organ damage in hypertensive homozygous TGR(mRen2)27 rats. *Hypertension* 47, 467–474. doi: 10.1161/01.HYP.0000202487.68969.f7
- Shao, D., and Tian, R. (2015). Glucose Transporters in Cardiac Metabolism and Hypertrophy. *Compr. Physiol.* 6, 331–351. doi: 10.1002/cphy.c150016
- Singh, J. A., Beg, S., and Lopez-Olivo, M. A. (2010). Tocilizumab for rheumatoid arthritis. *Cochrane Database Syst. Rev.* 38 (1), Cd008331. doi: 10.1002/14651858.CD008331
- Soler, M. J., Ye, M., Wysocki, J., William, J., Lloveras, J., and Batlle, D. (2009). Localization of ACE2 in the renal vasculature: amplification by angiotensin II type 1 receptor blockade using telmisartan. *Am. J. Physiol. Renal Physiol.* 296, F398–F405. doi: 10.1152/ajprenal.90488.2008
- Sriramula, S., Cardinale, J. P., Lazartigues, E., and Francis, J. (2011). ACE2 overexpression in the paraventricular nucleus attenuates angiotensin II-induced hypertension. *Cardiovasc. Res.* 92, 401–408. doi: 10.1093/cvr/cvr242
- Su, S., Wong, G., Shi, W., Liu, J., Lai, A. C. K., Zhou, J., et al. (2016). Epidemiology, Genetic Recombination, and Pathogenesis of Coronaviruses. *Trends Microbiol* 24, 490–502. doi: 10.1016/j.tim.2016.03.003
- Sukumaran, V., Veeraveedu, P. T., Gurusamy, N., Yamaguchi, K., Lakshmanan, A. P., Ma, M., et al. (2011). Cardioprotective effects of telmisartan against heart failure in rats induced by experimental autoimmune myocarditis through the modulation of angiotensin-converting enzyme-2/angiotensin 1-7/mas receptor axis. *Int. J. Biol. Sci.* 7, 1077–1092. doi: 10.7150/ijbs.7.1077
- Sun, M. L., Yang, J. M., Sun, Y. P., and Su, G. H. (2020). [Inhibitors of RAS Might Be a Good Choice for the Therapy of COVID-19 Pneumonia]. *Zhonghua Jie He He Hu Xi Za Zhi* 43, 219–222. doi: 10.3760/cma.j.issn.1001-0939.2020.03.016
- Sungnak, W., Huang, N., Bécavin, C., Berg, M., Queen, R., Litvinukova, M., et al. (2020). SARS-CoV-2 entry factors are highly expressed in nasal epithelial cells together with innate immune genes. *Nat. Med.* 26, 681–687. doi: 10.1038/s41591-020-0868-6
- Surjit, M., Liu, B., Jameel, S., Chow, V. T., and Lal, S. K. (2004). The SARS coronavirus nucleocapsid protein induces actin reorganization and apoptosis in COS-1 cells in the absence of growth factors. *Biochem. J.* 383, 13–18. doi: 10.1042/BJ20040984
- Tang, N., Li, D., Wang, X., and Sun, Z. (2020). Abnormal coagulation parameters are associated with poor prognosis in patients with novel coronavirus pneumonia. *J. Thromb. Haemost.* 18 (4), 844–847. doi: 10.1111/jth.14768
- Tetzner, A., Gebolys, K., Meinert, C., Klein, S., Uhlich, A., Trebicka, J., et al. (2016). G-Protein-Coupled Receptor mRgD Is a Receptor for Angiotensin-(1-7) Involving Adenylyl Cyclase, cAMP, and Phosphokinase A. *Hypertension* 68, 185–194. doi: 10.1161/HYPERTENSIONAHA.116.07572
- Vallet, B., and Wiel, E. (2001). Endothelial cell dysfunction and coagulation. *Crit. Care Med.* 29, S36–S41. doi: 10.1097/00003246-200107001-00015
- Varga, Z., Flammer, A. J., Steiger, P., Haberecker, M., Andermatt, R., Zinkernagel, A. S., et al. (2020). Endothelial cell infection and endotheliitis in COVID-19. *Lancet* 395, 1417–1418. doi: 10.1016/S0140-6736(20)30937-5
- Ventura, E., Weller, M., and Burghardt, I. (2017). Cutting Edge: ERK1 Mediates the Autocrine Positive Feedback Loop of TGF- β and Furin in Glioma-Initiating Cells. *J. Immunol.* 198, 4569–4574. doi: 10.4049/jimmunol.1601176
- Wang, D., Chen, Y., Jiang, J., Zhou, A., Pan, L., Chen, Q., et al. (2014). Carvedilol has stronger anti-inflammation and anti-virus effects than metoprolol in murine model with coxsackievirus B3-induced viral myocarditis. *Gene* 547, 195–201. doi: 10.1016/j.gene.2014.06.003
- Wang, D., Hu, B., Hu, C., Zhu, F., Liu, X., Zhang, J., et al. (2020). Clinical Characteristics of 138 Hospitalized Patients With 2019 Novel Coronavirus-Infected Pneumonia in Wuhan, China. *JAMA.* 323 (11), 1061–1069. doi: 10.1001/jama.2020.1585
- Wang, Y. (2007). Mitogen-activated protein kinases in heart development and diseases. *Circulation* 116, 1413–1423. doi: 10.1161/CIRCULATIONAHA.106.679589
- Who (2020a). *Report of the WHO-China Joint Mission on Coronavirus Disease 2019 (COVID-19)* [Online], [Accessed March 29, 2020].
- Who (2020b). *WHO Director-General's opening remarks at the media briefing on COVID-19 - 16 March 2020* [Online], [Accessed March 29, 2020].

- Widmer, R. J., and Lerman, A. (2014). Endothelial dysfunction and cardiovascular disease. *Glob Cardiol. Sci. Pract.* 2014, 291–308. doi: 10.5339/gcsp.2014.43
- Witkowski, M., Landmesser, U., and Rauch, U. (2016). Tissue factor as a link between inflammation and coagulation. *Trends Cardiovasc. Med.* 26, 297–303. doi: 10.1016/j.tcm.2015.12.001
- Wu, Q., Zhou, L., Sun, X., Yan, Z., Hu, C., Wu, J., et al. (2017). Altered Lipid Metabolism in Recovered SARS Patients Twelve Years after Infection. *Sci. Rep.* 7, 9110. doi: 10.1038/s41598-017-09536-z
- Wu, C., Chen, X., Cai, Y., Xia, J., Zhou, X., Xu, S., et al. (2020). Risk Factors Associated With Acute Respiratory Distress Syndrome and Death in Patients With Coronavirus Disease 2019 Pneumonia in Wuhan, China. *JAMA Intern Med.* doi: 10.1001/jamainternmed.2020.0994
- Xie, L., Jiang, Y., Ouyang, P., Chen, J., Doan, H., Herndon, B., et al. (2007). Effects of Dietary Calorie Restriction or Exercise on the PI3K and Ras Signaling Pathways in the Skin of Mice. *J. Biol. Chem.* 282, 28025–28035. doi: 10.1074/jbc.M604857200
- Xu, J., Sriramula, S., Xia, H., Moreno-Walton, L., Culicchia, F., Domenig, O., et al. (2017). Clinical Relevance and Role of Neuronal AT1 Receptors in ADAM17-Mediated ACE2 Shedding in Neurogenic Hypertension. *Circ. Res.* 121, 43–55. doi: 10.1161/CIRCRESAHA.116.310509
- Yan, X., Hao, Q., Mu, Y., Timani, K. A., Ye, L., Zhu, Y., et al. (2006). Nucleocapsid protein of SARS-CoV activates the expression of cyclooxygenase-2 by binding directly to regulatory elements for nuclear factor-kappa B and CCAAT/enhancer binding protein. *Int. J. Biochem. Cell Biol.* 38, 1417–1428. doi: 10.1016/j.biocel.2006.02.003
- Yan, R., Zhang, Y., Li, Y., Xia, L., Guo, Y., and Zhou, Q. (2020). Structural basis for the recognition of SARS-CoV-2 by full-length human ACE2. *Science* 367, 1444–1448. doi: 10.1126/science.abb2762
- Yang, X., Yu, Y., Xu, J., Shu, H., Xia, J., Liu, H., et al. (2020). Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centered, retrospective, observational study. *Lancet Respir. Med.* 8, 475–481. doi: 10.1016/S2213-2600(20)30079-5
- Yuan, S. (2015). Statins May Decrease the Fatality Rate of Middle East Respiratory Syndrome Infection. *mBio* 6, e01120–e01120. doi: 10.1128/mBio.01120-15
- Yue-Chun, L., Guang-Yi, C., Li-Sha, G., Chao, X., Xinqiao, T., Cong, L., et al. (2016). The Protective Effects of Ivabradine in Preventing Progression from Viral Myocarditis to Dilated Cardiomyopathy. *Front. Pharmacol.* 7, 408. doi: 10.3389/fphar.2016.00408
- Zhang, Z., Chen, L., Zhong, J., Gao, P., and Oudit, G. Y. (2014). ACE2/Ang-(1-7) signaling and vascular remodeling. *Sci. China Life Sci.* 57, 802–808. doi: 10.1007/s11427-014-4693-3
- Zhang, P., Wu, C., Huang, X. H., Shen, C. L., Li, L., Zhang, W., et al. (2017). Aspirin suppresses TNF- α -induced MMP-9 expression via NF- κ B and MAPK signaling pathways in RAW264.7 cells. *Exp. Ther. Med.* 14, 5597–5604. doi: 10.3892/etm.2017.5252
- Zhang, H., Penninger, J. M., Li, Y., Zhong, N., and Slutsky, A. S. (2020). Angiotensin-converting enzyme 2 (ACE2) as a SARS-CoV-2 receptor: molecular mechanisms and potential therapeutic target. *Intensive Care Med.* 46, 586–590. doi: 10.1007/s00134-020-05985-9
- Zhao, G., Yang, W., Wu, J., Chen, B., Yang, X., Chen, J., et al. (2018). Influence of a Coronary Artery Disease-Associated Genetic Variant on FURIN Expression and Effect of Furin on Macrophage Behavior. *Arterioscler. Thromb. Vasc. Biol.* 38, 1837–1844. doi: 10.1161/ATVBAHA.118.311030
- Zheng, Y.-Y., Ma, Y.-T., Zhang, J.-Y., and Xie, X. (2020). COVID-19 and the cardiovascular system. *Nat. Rev. Cardiol.* 17, 259–260. doi: 10.1038/s41569-020-0360-5
- Zhou, F., Yu, T., Du, R., Fan, G., Liu, Y., Liu, Z., et al. (2020). Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* 395, 1054–1062. doi: 10.1016/S0140-6736(20)30566-3
- Zhou, P., Yang, X. L., Wang, X. G., Hu, B., Zhang, L., Zhang, W., et al. (2020). A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature* 579, 270–273. doi: 10.1038/s41586-020-2012-7
- Zisman, L. S., Keller, R. S., Weaver, B., Lin, Q., Speth, R., Bristow, M. R., et al. (2003). Increased angiotensin-(1-7)-forming activity in failing human heart ventricles: evidence for upregulation of the angiotensin-converting enzyme Homologue ACE2. *Circulation* 108, 1707–1712. doi: 10.1161/01.CIR.0000094734.67990.99

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Wehbe, Hammoud, Soudani, Zaraket, El-Yazbi and Eid. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.