



Corrigendum: Tubeimoside 1 Acts as a Chemotherapeutic Synergist via Stimulating Macropinocytosis

OPEN ACCESS

Approved by:

Frontiers Editorial Office,
Frontiers Media SA, Switzerland

*Correspondence:

Xuegang Sun
sxg_smu@126.com

†These authors have contributed
equally to this work

Specialty section:

This article was submitted to
Ethnopharmacology,
a section of the journal
Frontiers in Pharmacology

Received: 25 June 2021

Accepted: 28 June 2021

Published: 12 July 2021

Citation:

Gong X, Sun R, Gao Z, Han W, Liu Y,
Zhao L, Jing L, Yao X and Sun X (2021)
Corrigendum: Tubeimoside 1 Acts as a
Chemotherapeutic Synergist via
Stimulating Macropinocytosis.
Front. Pharmacol. 12:730775.
doi: 10.3389/fphar.2021.730775

Xianling Gong^{1,2†}, Ruibo Sun^{1†}, Zhuowei Gao^{1,3†}, Weili Han⁴, Yawei Liu⁵, Liang Zhao⁶,
Linlin Jing⁷, Xueqing Yao⁷ and Xuegang Sun^{1*}

¹The Key Laboratory of Molecular Biology, State Administration of Traditional Chinese Medicine, School of Traditional Chinese Medicine, Southern Medical University, Guangzhou, China, ²School of Pharmacy, Guangdong Medical University, Dongguan, China, ³Shunde Hospital, Southern Medical University, Foshan, China, ⁴School of Public Health, Guangzhou, China, ⁵Nanfeng Hospital, Southern Medical University, Guangzhou, China, ⁶School of Basic Medical Sciences, Southern Medical University, Guangzhou, China, ⁷Traditional Chinese Medicine Integrated Hospital, Southern Medical University, Guangzhou, China

Keywords: tubeimoside 1, chemotherapeutic synergist, macropinocytosis, methuosis, endocytosis

A Corrigendum on

Tubeimoside 1 Acts as a Chemotherapeutic Synergist via Stimulating Macropinocytosis
by Gong, X., Sun, R., Gao, Z., Han, W., Liu, Y., Zhao, L., Jing, L., Yao, X., and Sun X. (2018). *Front. Pharmacol.* 9:1044. doi:10.3389/fphar.2018.01044

In the original article, there was a mistake in **Figure 6** as published.

The second image of bottom panel of **Figure 6D** was misused. The corrected **Figure 6** appears below.

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Copyright © 2021 Gong, Sun, Gao, Han, Liu, Zhao, Jing, Yao and Sun. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.




FIGURE 6 | Chloroquine suppress TBM1-induced vacuolation and cell death in CRC cells. **(A)** SW480 were treated with 8 μ M TBM1 for different time or starvation or 1 μ M rapamycin for 24 h. Protein expression of LC3-I and LC3-II was detected by Western blotting. Beta-actin were served as loading control. **(B)** SW480 cells were pre-cultured with or without chloroquine (CQ) (10 μ M) for 1 h before exposure to TBM1 for 24 h. LC3 processing was assessed using Western blotting analysis. **(C)** After treatment with SW480 cells for different time as **(B)**, morphological changes were observed by phase-contrast microscopy. **(D)** SW480 cells were pre-cultured with or without 3-MA (3 mM) for 1 h before exposure to TBM1 for different time. Morphological changes were observed using phase-contrast microscopy. **(E)** The number of vacuoles of the same cultures as **(B)** for 48 h was counted from three different imagines. **(F)** Cell viability was measured by MTT assay ($n = 5$). $\#p < 0.01$.