



# Corrigendum: *Moringa oleifera* Alkaloids Inhibited PC3 Cells Growth and Migration Through the COX-2 Mediated Wnt/ $\beta$ -Catenin Signaling Pathway

Jing Xie<sup>1,2†</sup>, Feng-xian Luo<sup>1,3†</sup>, Chong-ying Shi<sup>1,2†</sup>, Wei-wei Jiang<sup>4</sup>, Ying-yan Qian<sup>1,5</sup>, Ming-rong Yang<sup>1,5</sup>, Shuang Song<sup>1,2</sup>, Tian-yi Dai<sup>2</sup>, Lei Peng<sup>3</sup>, Xiao-yu Gao<sup>3</sup>, Liang Tao<sup>1,3</sup>, Yang Tian<sup>1,2,3\*</sup> and Jun Sheng<sup>6\*</sup>

## OPEN ACCESS

### Edited and reviewed by:

Rajasekaran Subbiah,  
ICMR-National Institute for Research  
in Environmental Health, India

### \*Correspondence:

Jun Sheng  
shengjun\_ynau@163.com  
Yang Tian  
tianyanyang1208@163.com

<sup>†</sup>These authors have contributed  
equally to this work

### Specialty section:

This article was submitted to  
Ethnopharmacology,  
a section of the journal  
Frontiers in Pharmacology

**Received:** 19 August 2021

**Accepted:** 03 September 2021

**Published:** 16 September 2021

### Citation:

Xie J, Luo F-x, Shi C-y, Jiang W-w,  
Qian Y-y, Yang M-r, Song S, Dai T-y,  
Peng L, Gao X-y, Tao L, Tian Y and  
Sheng J (2021) Corrigendum: *Moringa*  
*oleifera* Alkaloids Inhibited PC3 Cells  
Growth and Migration Through the  
COX-2 Mediated Wnt/ $\beta$ -Catenin  
Signaling Pathway.  
*Front. Pharmacol.* 12:760933.  
doi: 10.3389/fphar.2021.760933

<sup>1</sup>College of Food Science and Technology, Yunnan Agricultural University, Kunming, China, <sup>2</sup>Engineering Research Center of Development and Utilization of Food and Drug Homologous Resources, Ministry of Education, Yunnan Agricultural University, Kunming, China, <sup>3</sup>National Research and Development Professional Center for Moringa Processing Technology, Yunnan Agricultural University, Kunming, China, <sup>4</sup>College of Science, Yunnan Agricultural University, Kunming, China, <sup>5</sup>Yunnan Province Engineering Research Center of Functional Food of Homologous of Drug and Food, Yunnan Agricultural University, Kunming, China, <sup>6</sup>Key Laboratory of Pu-er Tea Science, Ministry of Education, Yunnan Agricultural University, Kunming, China

**Keywords:** *Moringa oleifera* alkaloids, prostate cancer, PC3 cells, cell growth and migration, COX-2-wnt/ $\beta$ -catenin signaling pathway

## A Corrigendum on

### *Moringa oleifera* Alkaloids Inhibited PC3 Cells Growth and Migration Through the COX-2 Mediated Wnt/ $\beta$ -Catenin Signaling Pathway


by Xie, J., Luo, F-x., Shi, C-y., Jiang, W-w., Qian, Y-y., Yang, M-r., Song, S., Dai, T-y., Peng, L., Gao, X-y., Tao, L., Tian, Y., and Sheng, J. (2020). *Front. Pharmacol.* 11:523962. doi: 10.3389/fphar.2020.523962

In the original article, there was a mistake in **Figure 1C** as published. Some images in **Figure 1C** were mistakenly duplicated. The corrected **Figure 1** appears below.

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

**Publisher's Note:** All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2021 Xie, Luo, Shi, Jiang, Qian, Yang, Song, Dai, Peng, Gao, Tao, Tian and Sheng. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.



**FIGURE 1 |** The effect of MOA on the proliferation of PC3 and RWPE-1 cells. **(A)** The half maximal inhibitory concentration (IC<sub>50</sub>) of ten tumor cell lines following treatment with MOA (0–320 μg/ml) for 48 h. **(B)** Cell viability of PC3 and RWPE-1 cells following MOA (0–320 μg/ml) treatment for 24, 48, and 72 h. **(C,D)** Analysis of the colony formation ability of PC3 cells following treatment with MOA (0, 20, 40, and 80 μg/ml) for 24, 48, and 72 h. The cellular colony formation rates are expressed as fold changes. Results are expressed as the mean ± SEM of three independent experiments. \*\*\*p < 0.001 vs. 0 μg/ml.