



# RETRACTED: Corrigendum: ErHuang Formula Improves Renal Fibrosis in Diabetic Nephropathy Rats by Inhibiting CXCL6/JAK/STAT3 Signaling Pathway

Yu-Li Shen<sup>1,2†</sup>, Yi-ping Jiang<sup>3†</sup>, Xiao-Qin Li<sup>1,4†</sup>, Su-Juan Wang<sup>5</sup>, Ming-Hua Ma<sup>6</sup>, Chun-Yan Zhang<sup>7</sup>, Jian-Yong Zhu<sup>8</sup>, Khalid Rahman<sup>8</sup>, Li-Jun Zhang<sup>1\*</sup>, Xin Luan<sup>1,2\*</sup> and Hong Zhang<sup>1\*</sup>

## OPEN ACCESS

### Edited and reviewed by:

Adolfo Andrade-Cetto,  
National Autonomous University of  
Mexico, Mexico

### \*Correspondence:

Li-Jun Zhang  
zhanglijun0407@163.com,  
Xin Luan  
luanxin@shutcm.edu.cn,

Hong Zhang  
hqzhang51@126.com

†These authors have contributed  
equally to this work

### Specialty section:

This article was submitted to  
Ethnopharmacology,  
a section of the journal  
Frontiers in Pharmacology

Received: 22 October 2021

Accepted: 29 October 2021

Published: 18 November 2021

### Citation:

Shen Y-L, Jiang Y-p, Li X-Q, Wang S-J,  
Ma M-H, Zhang C-Y, Zhu J-Y,  
Rahman K, Zhang L-J, Luan X and  
Zhang H (2021) Corrigendum:  
ErHuang Formula Improves Renal  
Fibrosis in Diabetic Nephropathy Rats  
by Inhibiting CXCL6/JAK/STAT3  
Signaling Pathway.  
Front. Pharmacol. 12:789192.  
doi: 10.3389/fphar.2021.789192

<sup>1</sup>Institute of Interdisciplinary Integrative Medicine Research, Shanghai University of Traditional Chinese Medicine, Shanghai, China, <sup>2</sup>Shuguang Hospital, Shanghai University of Traditional Chinese Medicine, Shanghai, China, <sup>3</sup>Department of Pharmacognosy, School of Pharmacy, Second Military Medical University, Shanghai, China, <sup>4</sup>School of Pharmacy, Chengdu University of Traditional Chinese Medicine, Chengdu, China, <sup>5</sup>Department of Drug Preparation, Hospital of TCM and Hui Nationality Medicine, Ningxia Medical University, Wuzhong, China, <sup>6</sup>Department of Pharmacy, Yangpu Hospital, Tongji University School of Medicine, Shanghai, China, <sup>7</sup>Central Laboratory, Seventh People's Hospital, Shanghai University of Traditional Chinese Medicine, Shanghai, China, <sup>8</sup>School of Pharmacy and Biomolecular Sciences, Faculty of Science, Liverpool John Moores University, Liverpool, United Kingdom

**Keywords:** erhuang formula, renal fibrosis, diabetic nephropathy, mechanism, JAK/STAT3

## A Corrigendum on

### ErHuang Formula Improves Renal Fibrosis in Diabetic Nephropathy Rats by Inhibiting CXCL6/JAK/STAT3 Signaling Pathway


by Shen, Y-L., Jiang, Y-P., Li X-Q., Wang, S-J., Ma, M-H., Zhang, C-Y., Zhu, J-Y., Rahman, K., Zhang, L-J., Luan, X., and Zhang, H. (2020). *Front. Pharmacol.* 10:1596. doi: 10.3389/fphar.2019.01596

In the original article, there was a mistake during production in **Figure 4** as published. **Figure 4** is identical to **Figure 3**. The corrected **Figure 4** appears below.

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

**Publisher's Note:** All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2021 Shen, Jiang, Li, Wang, Ma, Zhang, Zhu, Rahman, Zhang, Luan and Zhang. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.



**FIGURE 4** | EHF inhibited the proliferation of NRK-49F cells and decreased the expression of CXCL6 and CXCR1. **(A)** CCK8 analysis of NRK-49F cell proliferation. **(B), (D)** The mRNA and protein expressions of CXCL6 and CXCR1 in different EHF dose groups were examined by qRT-PCR and Western blotting. GAPDH was served as a loading control. **(C), (E)** The effect of 0.5 mg/ml EHF on the expression of CXCL6 and CXCR1 in CXCL6 overexpression NRK-49F cells was evaluated by qRT-PCR and Western blotting. \* $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$  versus control group.