



OPEN ACCESS

EDITED AND REVIEWED BY
Chang-Hee Suh,
Ajou University, Republic of Korea

*CORRESPONDENCE

Bin Wang,
✉ gztcmwangbin1973@163.com
Liangliang Xu,
✉ xull-2016@gzucm.edu.cn

[†]These authors have contributed equally to this work and share first authorship

SPECIALTY SECTION

This article was submitted to Integrative and Regenerative Pharmacology, a section of the journal Frontiers in Pharmacology

RECEIVED 20 January 2023

ACCEPTED 17 February 2023

PUBLISHED 27 February 2023

CITATION

Liu Y, Fu B, Li X, Chen C, Li X, Xu L and Wang B (2023), Corrigendum: Bushen huoxue decoction inhibits RANKL-stimulated osteoclastogenesis and glucocorticoid-induced bone loss by modulating the NF- κ B, ERK, and JNK signaling pathways.
Front. Pharmacol. 14:1148908.
doi: 10.3389/fphar.2023.1148908

COPYRIGHT

© 2023 Liu, Fu, Li, Chen, Li, Xu and Wang. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Corrigendum: Bushen huoxue decoction inhibits RANKL-stimulated osteoclastogenesis and glucocorticoid-induced bone loss by modulating the NF- κ B, ERK, and JNK signaling pathways

Yamei Liu^{1†}, Binlan Fu^{1,2†}, Xiaoman Li^{1,2}, Chen Chen³, Xican Li³, Liangliang Xu^{4*} and Bin Wang^{5*}

¹School of Basic Medical Science, Guangzhou University of Chinese Medicine, Guangzhou, China, ²Laboratory of Orthopedics and Traumatology, Lingnan Medical Research Center, Guangzhou University of Chinese Medicine, Guangzhou, China, ³School of Chinese Herbal Medicine, Guangzhou University of Chinese Medicine, Guangzhou, China, ⁴The First Affiliated Hospital of Guangzhou University of Chinese Medicine, Guangzhou, China, ⁵Department of Traumatology, The Third Affiliated Hospital of Guangzhou University of Chinese Medicine, Guangzhou, China

KEYWORDS

bushen huoxue decoction, osteoporosis, NF- κ B pathway, ERK pathway, JNK pathway

A Corrigendum on

Bushen huoxue decoction inhibits RANKL-stimulated osteoclastogenesis and glucocorticoid-induced bone loss by modulating the NF- κ B, ERK, and JNK signaling pathways

by Liu Y, Fu B, Li X, Chen C, Li X, Xu L and Wang B (2022). *Front. Pharmacol.* 13:1007839. doi: 10.3389/fphar.2022.1007839

In the published article, there was an error in [Figure 5](#) as published. In [Figure 5E](#), the position of the experimental images of DEX group and DEX + BHD group is inverted, that is, the images of DEX group should appear in DEX + BHD group, while the images of DEX + BHD group should appear in DEX group. The corrected [Figure 5](#) and its caption appear below.

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.




FIGURE 5
 BHD affects osteoclast formation and differentiation by suppressing the expression of the NF- κ B signaling pathway in vivo. (A–E) Representative images of Immunohistochemical staining of decalcified bone sections. Scale bar = 20 μ m.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated

organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.