



OPEN ACCESS

EDITED BY

Mohamed Chahine,
Laval University, Canada

REVIEWED BY

Jon Silva,
Washington University in St. Louis,
United States

Jean-sebastien Rougier,
University of Bern, Switzerland

Robert S. Kass,
Columbia University, United States

*CORRESPONDENCE

Michael Toft Overgaard,
✉ mto@bio.aau.dk

RECEIVED 24 April 2023

ACCEPTED 25 July 2023

PUBLISHED 16 August 2023

CITATION

Brohus M, Busuioc A-O, Wimmer R,
Nyegaard M and Overgaard MT (2023),
Calmodulin mutations affecting
Gly114 impair binding to the Na_v1.5 IQ-
domain.

Front. Pharmacol. 14:1210140.

doi: 10.3389/fphar.2023.1210140

COPYRIGHT

© 2023 Brohus, Busuioc, Wimmer,
Nyegaard and Overgaard. This is an
open-access article distributed under the
terms of the [Creative Commons
Attribution License \(CC BY\)](#). The use,
distribution or reproduction in other
forums is permitted, provided the original
author(s) and the copyright owner(s) are
credited and that the original publication
in this journal is cited, in accordance with
accepted academic practice. No use,
distribution or reproduction is permitted
which does not comply with these terms.

Calmodulin mutations affecting Gly114 impair binding to the Na_v1.5 IQ-domain

Malene Brohus ¹, Ana-Octavia Busuioc¹, Reinhard Wimmer ¹,
Mette Nyegaard ² and Michael Toft Overgaard ^{1*}

¹Department of Chemistry and Bioscience, Aalborg University, Aalborg, Denmark, ²Department of Health Science and Technology, Aalborg University, Gistrup, Denmark

Missense variants in *CALM* genes encoding the Ca²⁺-binding protein calmodulin (CaM) cause severe cardiac arrhythmias. The disease mechanisms have been attributed to dysregulation of RyR2, for Catecholaminergic Polymorphic Ventricular Tachycardia (CPVT) and/or Ca_v1.2, for Long-QT Syndrome (LQTS). Recently, a novel *CALM2* variant, G114R, was identified in a mother and two of her four children, all of whom died suddenly while asleep at a young age. The G114R variant impairs closure of Ca_v1.2 and RyR2, consistent with a CPVT and/or mild LQTS phenotype. However, the children carrying the *CALM2* G114R variant displayed a phenotype commonly observed with variants in Na_v1.5, i.e., Brugada Syndrome (BrS) or LQT3, where death while asleep is a common feature. We therefore hypothesized that the G114R variant specifically would interfere with Na_v1.5 binding. Here, we demonstrate that CaM binding to the Na_v1.5 IQ-domain is severely impaired for two CaM variants G114R and G114W. The impact was most severe at low and intermediate Ca²⁺ concentrations (up to 4 μM) resulting in more than a 50-fold reduction in Na_v1.5 binding affinity, and a smaller 1.5 to 11-fold reduction at high Ca²⁺ concentrations (25–400 μM). In contrast, the arrhythmogenic CaM-N98S variant only induced a 1.5-fold reduction in Na_v1.5 binding and only at 4 μM Ca²⁺. A non-arrhythmogenic I10T variant in CaM did not impair Na_v1.5 IQ binding. These data suggest that the interaction between Na_v1.5 and CaM is decreased with certain CaM variants, which may alter the cardiac sodium current, I_{Na}. Overall, these results suggest that the phenotypic spectrum of calmodulinopathies may likely expand to include BrS- and/or LQT3-like traits.

KEYWORDS

calmodulin, calmodulinopathy, arrhythmogenic, cardiac ion-channel regulation, calmodulin target binding, experimental variant interpretation, SCN5A, NaV1.5

Introduction

The cytosolic calcium (Ca²⁺) binding protein calmodulin (CaM) serves as a critical mediator of intra-cellular Ca²⁺ signals in a multitude of physiological processes (Chin and Anthony, 2000; Xia and Storm, 2005; Clapham, 2007; Sorensen et al., 2013; Berchtold and Villalobo, 2014). The multifaceted nature of CaM comes from its ubiquitous expression and its ability to interact with, and relay information to, more than 350 cellular target proteins (Yap et al., 2000; Tidow and Nissen, 2013).

This extraordinary versatility is due to the two lobes of CaM, the N-lobe and C-lobe, each containing two Ca²⁺-binding EF hands (Figure 1A). The lobes differ in Ca²⁺-affinity and

-kinetics, allowing CaM to respond to changes in Ca²⁺ over a broad range of concentration and time. This range is even further expanded by target-specific changes in Ca²⁺ binding-affinities and -kinetics upon CaM binding to protein targets (Villaruel et al., 2014; Søndergaard et al., 2015a; Søndergaard et al., 2015b).

The cellular importance of CaM Ca²⁺-sensing and integrity is highlighted by the protein's unique genetic architecture and evolutionary conservation. Mammals have three independent genes (CALM1-3) that all encode an identical CaM protein.

Moreover, the protein sequence is invariant in all vertebrates, underpinning the extreme selection pressure against any amino acid variation in this central Ca²⁺-sensor protein (Figure 1B) (Berchtold et al., 1993; Toutenhoofd and Strehler, 2000; Friedberg and Rhoads, 2001).

As a result, genetic variation in all three CALM genes is ultra-rare. Until the first human missense variant was discovered in 2012, and linked to a severe cardiac arrhythmia (Catecholaminergic Polymorphic Ventricular Tachycardia (CPVT)) and sudden




FIGURE 1

(A) Crystal structure of Ca²⁺-bound calmodulin (CaM) (PDB ID: 1CLL) with EF hands I (red), II (orange), III (blue), and IV (cyan) indicated along with their corresponding alpha-helices (α1-8). Beta-sheets are shown in green and flexible loops/unstructured regions in yellow. Amino acid residues known to harbor arrhythmogenic substitutions (Crotti et al., 2019) are shown in purple stick representation, with residue G114 highlighted with a yellow circle. Ca²⁺-ions are shown as black spheres. (B) CaM protein sequence alignment. Red highlights indicate amino acid differences or gaps in the alignment. Residue G114 is indicated by a blue frame. Numbering is according to immature human CaM (including the initial M).

cardiac death (SCD) (Nyegaard et al., 2012), mutations in the *CALM* genes were considered incompatible with life (Jensen et al., 2018). Since the initial discovery of human missense variants linked to CPVT, mutations have also been identified in individuals affected by long QT syndrome (LQTS) (Crotti et al., 2013) and Idiopathic Ventricular Fibrillation (IVF) (Marsman et al., 2014). In 2019, Crotti and co-workers presented an extensive collection of 28 unique CaM variants, identified in 74 carriers, in the International Calmodulinopathy Registry (Crotti et al., 2019). *CALM* variant carriers in this registry present with cardiac arrhythmia phenotypes including LQTS (49%), CPVT (28%), overlap LQTS/CPVT (4%), and a few cases of IVF, sudden unexplained death (SUD), or atypical phenotypes (Crotti et al., 2019).

Clinical characteristics of the calmodulinopathies include an early age of onset (a mean of 1.5 years for LQTS and 6 years for CPVT) and a high risk of a major arrhythmic event (68%), such as cardiac arrest or SCD/SUD (Crotti et al., 2019). The molecular mechanisms underlying the two main phenotypes has largely been ascribed to specific dysregulation of the two primary cardiac Ca^{2+} channels, $\text{Ca}_v1.2$ for LQTS, and RyR2 for CPVT-like phenotypes (Limpitikul et al., 2014; Yin et al., 2014; Søndergaard et al., 2019; 2020; 2017; Nyegaard and Overgaard, 2019; Holt et al., 2020). The broad phenotypic spectrum caused by CaM variants, including mechanistically different cardiac arrhythmias, is likely a consequence of CaM serving as a key regulator of multiple cardiac ion-channels, besides $\text{Ca}_v1.2$ and RyR2, that control cardiac excitation-contraction coupling. Indeed, given the vast number of CaM-regulated proteins, the phenotypic spectrum of calmodulinopathies is likely to expand even further as more carriers are discovered (Jensen et al., 2018; Urrutia et al., 2019).

In 2019, a novel *CALM2* variant, G114R (immature protein numbering including initial Met), was identified in an Australian mother and two of her four children. Over a 10-year period, all four children died suddenly and unexpectedly while asleep, at ages ranging from 19 days to 18 months (Brohus et al., 2021). Further, the two children carrying the G114R variant had infections at the time of death, implying a potential presence of fever. In 2021, we showed that the G114R variant impairs CaM's ability to bind Ca^{2+} -ions and to interact with and regulate $\text{Ca}_v1.2$ and RyR2, with an impact suggesting an arrhythmogenic potential consistent with CPVT, IVF, or mild LQTS (Brohus et al., 2021).

Death while asleep or at rest have only been observed in a small subset of CaM variant carriers, and mainly for CaM variants with a severe impairment of Ca^{2+} binding and/or Ca^{2+} -dependent inactivation of $\text{Ca}_v1.2$, larger than the effect imposed by G114R (Brohus et al., 2021). Therefore, the phenotype of the children carrying the *CALM2* G114R variant to some degree represents an expansion of the known clinical manifestations of CaM variant carriers. The phenotype more closely resembles that of carriers of missense variants in the cardiac sodium channel, $\text{Na}_v1.5$, for whom major arrhythmic events or death while asleep is a common feature (Schwartz et al., 2001; Postema and Wilde, 2008; Takigawa et al., 2008). Given that CaM is critical for $\text{Na}_v1.5$ function, we hypothesized that the G114R variant would specifically interfere with $\text{Na}_v1.5$ binding.

The $\text{Na}_v1.5$ channel is implicated in both Brugada Syndrome (BrS) and LQT3, arrhythmic diseases that result from divergent molecular mechanisms. Intriguingly, both phenotypes can be caused by $\text{Na}_v1.5$ channel mutations that perturb the interaction with and

modulation by CaM (Shah et al., 2006; Yan et al., 2017; Urrutia et al., 2019; Kang et al., 2021; Wu and Liang, 2021). In some cases, the same $\text{Na}_v1.5$ variant causes both phenotypes, which alludes to the difficulty in variant genotype-phenotype interpretation (Supplementary Figure S1).

The gating of $\text{Na}_v1.5$ is modulated by CaM in a bi-directional manner: Both channel activation (peak current), fast inactivation, and persistent current depend on CaM. Several CaM binding domains (CaMBDs) have been identified, but their individual roles in the bi-directional modulation by CaM is still unclear (Kang et al., 2021). While the primary CaM binding site is an IQ-motif located in the C-terminal domain (CTD) of $\text{Na}_v1.5$ (Chagot and Chazin, 2011; C; Wang et al., 2014; Gabelli et al., 2014; C; Wang et al., 2012; Gabelli et al., 2016), CaM has also been shown to interact with a preIQ-domain in the CTD (Yoder et al., 2019), the “inactivation gate” in the DIII-DIV linker (Potet et al., 2009; Sarhan, Van Petegem, and Ahern, 2009; Sarhan et al., 2012; Johnson et al., 2018), and an N-terminal domain (NTD) (Wang et al., 2020) (Figure 2A).

In this study, we illustrate that CaM-G114 is located exactly in the binding interface between CaM and the IQ-domain of $\text{Na}_v1.5$ and demonstrate that CaM variants G114R and G114W impair the interaction with the IQ-domain in a Ca^{2+} -dependent manner with the largest impact occurring at a free Ca^{2+} -concentration range of 3 nM–4 μM . Thus, the apoCaM interaction with the $\text{Na}_v1.5$ -IQ domain is impaired for these CaM variants and their Ca^{2+} -sensing ability in the CaM/ $\text{Na}_v1.5$ -IQ complex has markedly changed.

Materials and methods


Materials and methods can be found in the [Supplementary Material](#).

Results

The CaM-G114 residue is extremely evolutionarily conserved and located at the CaM/ $\text{Na}_v1.5$ IQ-domain interface

CaM-G114 is the terminating residue of the second helix of EF hand III and it so far constitutes the only amino acid residue known to harbor a mutation in the loop between EF hands III and IV (L113-T118) (Figure 1A) (Bycroft et al., 2018; Crotti et al., 2019; Chen et al., 2022). A protein sequence alignment of CaM from different species shows that residue 114 is extremely evolutionarily conserved (Figure 1B, blue square). It is a glycine in all species investigated, including yeast, emphasizing the universal importance of its integrity.

High resolution structures of CaM (green) in complex with the $\text{Na}_v1.5$ -CTD (blue) reveal that CaM-G114 (purple) is located at the interface between CaM and the channel (Figure 2). In the apo-form, only the C-lobe of CaM binds to the $\text{Na}_v1.5$ IQ-domain (Figure 2B), whereas in the Mg^{2+} - and Ca^{2+} -bound forms, CaM wraps around the $\text{Na}_v1.5$ CTD with both its lobes (Figure 2C, D). In all cases, the interaction brings CaM-G114 and the IQ-domain into close



proximity (Figure 2, insets). We therefore hypothesized that substitution of residue G114 would affect the binding between CaM and the Nav_v1.5 channel.


The G114R and G114W variants impair the interaction between CaM and the Nav_v1.5 IQ-domain

To test the hypothesis that the integrity of CaM-G114 is essential for the interaction with the Nav_v1.5 IQ-domain, we monitored the fluorescence anisotropy (FA) signal of the TAMRA-labeled IQ-domain during titration with CaM at eight different Ca²⁺-concentrations, resulting in eight binding curves for each CaM variant (Figure 3A). The CaM variant N98S was included as an arrhythmic control, known to cause both LQTS and CPVT

(Nyegaard et al., 2012; Makita et al., 2014; Jiménez-Jáimez et al., 2016). Another CaM variant, CaM-I10T, identified in the UK Biobank resource (Bycroft et al., 2018), was included as a non-arrhythmic control.

The interaction between CaM-WT and the Nav_v1.5 IQ-domain depends on the level of Ca²⁺, apparent as a change in FA values for the CaM-saturated IQ-domain from low (Figure 3A, blue) to high (Figure 3A, red) Ca²⁺ concentrations. The assay thus allowed us to explore the Ca²⁺-dependency of the interaction, by determining the maximum FA plateau (FA_{max}) (Figure 3B, Supplementary Table S1) and the binding affinity (Figure 3C, Supplementary Table S2) at each of the eight Ca²⁺-concentrations.

The FA signal is a measure of the tumbling rate of the TAMRA-labeled Nav_v1.5 IQ-domain (Rossi and Taylor, 2011). As more CaM is added, more CaM/Nav_v1.5 IQ-domain complex forms, and the FA signal will increase (due to a reduced tumbling rate of the IQ-



domain) until reaching FA_{max} which represents the tumbling rate of the saturated complex (Figure 3A). The tumbling rate depends on the conformation of the complex, and a plot of the FA_{max} value as a function of $[Ca^{2+}]_{free}$ reveals a Ca²⁺-dependent increase in FA_{max} of the CaM-WT/Nav_v1.5 IQ-domain complex, in turn reflecting a Ca²⁺-induced conformational change (Figure 3B, black, Supplementary Table S1). The Ca²⁺-dependent conformational change is accompanied by a 10-fold increase in the CaM-WT binding affinity of the Nav_v1.5 IQ-domain (Figure 3C, black, Supplementary Table S2).

Curiously, the Ca²⁺-dependent development in the FA signal for the CaM-G114 variants is very different from CaM-WT. First, no FA_{max} plateau is reached at free Ca²⁺ concentrations < 4 μM (Figure 3A, B).

Second, the FA_{max} values ≥ 4 μM free Ca²⁺ are significantly different from those of the CaM-WT/Nav_v1.5-IQ complex (Figure 3B, red and blue, Supplementary Table S1). Interestingly, the change in FA_{max} imposed by the two G114 substitutions occurs in opposite directions relative to CaM-WT at saturating Ca²⁺ (FA_{max} increases for CaM-G114R and decreases for CaM-G114W, Supplementary Table S3).

Moreover, the CaM-G114R and -G114W mutations cause a dramatic reduction in IQ-domain binding affinity compared to CaM-WT (Figure 3C, red and blue, Supplementary Table S2). At Ca²⁺-concentrations below 4 μM, the affinity is reduced to an extent where the dissociation constant could not be accurately determined ($K_D < 5$ μM) (Figure 3C, stapled line). However, the data demonstrates that the affinity is reduced at least 47-fold compared to CaM-WT at

these Ca^{2+} -concentrations and estimates of K_D -values could be determined by assuming an identical FA_{max} value at all Ca^{2+} concentrations (Figure 3C, see Methods section for details, Supplementary Table S2, S4). At free Ca^{2+} -concentrations at and above 4 μM , the IQ-domain affinity of CaM-G114R and -G114W is still significantly reduced compared to CaM-WT, but to a smaller extent (1.4 to 11-fold) (Figure 3C; Supplementary Table S4).

In contrast to the CaM-G114 variants, the Ca^{2+} -dependent increase in FA_{max} observed for the CaM-WT/IQ-domain complex is also apparent for the CaM-I10T/IQ-domain complex (Figure 3B, orange). Like for CaM-WT, the FA_{max} value for the CaM-N98S/IQ-domain complex increases with Ca^{2+} , but the transition is shifted to higher Ca^{2+} -concentrations (Figure 3B, green, Supplementary Tables S1, S3). Further, the arrhythmogenic CaM-N98S variant only reduced the $\text{Na}_V1.5$ affinity significantly (1.5-fold compared to CaM-WT) at intermediate 4 μM free Ca^{2+} (Figure 3C; Supplementary Tables S2, S4). This effect is consistent with the observed 4-fold reduction in CaM C-lobe Ca^{2+} affinity imposed by the N98S substitution, but different from the G114R and G114W substitutions, although they have a similar impact on C-lobe Ca^{2+} binding (3- and 7-fold reduction compared to CaM-WT) (Brohus et al., 2021). The non-arrhythmogenic CaM-I10T variant displayed no difference in IQ-domain affinity compared to CaM-WT across any of the Ca^{2+} concentrations tested (Figure 3C, orange). These results are consistent with the observation that both residue I10 and N98 are located away from the CaM/ $\text{Na}_V1.5$ IQ-domain binding interface (Supplementary Figure S2).

The dramatic effect of the G114 variants on $\text{Na}_V1.5$ affinity at low Ca^{2+} concentrations (≤ 200 nM) appear specific to the IQ-domain, as the effect of CaM-G114R and G114W on the $\text{Na}_V1.5$ NTD were much smaller at the corresponding Ca^{2+} concentrations and comparable in magnitude to the effects of arrhythmogenic N98S (within a 4-fold difference from CaM-WT) (Supplementary Figure S3, S4; Supplementary Table S5, S6). As observed for the IQ-domain, the interaction between the non-arrhythmogenic CaM-I10T and the $\text{Na}_V1.5$ NTD did not differ from the CaM-WT/NTD interaction.

Discussion

In this study, we use an FA-based assay to investigate the Ca^{2+} -dependent interactions between CaM-WT and the $\text{Na}_V1.5$ IQ-domain and the recently identified CaM binding domain in the $\text{Na}_V1.5$ NTD, and how these are affected by mutations in CaM.

Intriguingly, the CaM/ $\text{Na}_V1.5$ IQ-domain interaction displays a very different Ca^{2+} sensitivity profile compared to the interactions between CaM and the $\text{Na}_V1.5$ NTD, $\text{Ca}_V1.2$ IQ-domain, and RyR2-CaMBD2 (Wang et al., 2018; Brohus et al., 2019; Wang et al., 2020; Brohus et al., 2021). While the binding affinity increases 10-fold for the CaM-WT/ $\text{Na}_V1.5$ IQ-domain complex from low nM to high μM Ca^{2+} -concentrations, the affinities of the CaM-WT/ $\text{Na}_V1.5$ NTD, CaM-WT/ $\text{Ca}_V1.2$ IQ-domain, and CaM-WT/RyR2-CaMBD2 complexes increase more than 1000-fold across the same Ca^{2+} -range (Supplementary Figure S5) (Brohus et al., 2021). Moreover, apoCaM binds to the $\text{Na}_V1.5$ IQ-domain with high affinity (117 nM—in good agreement with affinities determined by others (Shah et al., 2006; Yan et al., 2017)), much higher than to the $\text{Na}_V1.5$ NTD (825-fold), the $\text{Ca}_V1.2$ IQ-domain (26-fold), and RyR2 CaMBD2 (7-fold). For Ca^{2+} -saturated

CaM, the interaction with the $\text{Na}_V1.5$ IQ-domain is more than 100-fold weaker than the interaction with the CaM binding domains from RyR2 and $\text{Ca}_V1.2$ (Brohus et al., 2021). These results corroborate an essential role of apoCaM in modulating the $\text{Na}_V1.5$ channel via the IQ-domain (Kang et al., 2021).

A potential dysregulation of $\text{Na}_V1.5$, caused by human CaM mutations, has previously been investigated for a handful of LQTS-causing CaM variants (D96V, D130G, F142L, E141G) (Yin et al., 2014; Boczek et al., 2016; Rocchetti et al., 2017; Tarasov et al., 2023). However, the results for these variants have been largely unremarkable. Co-expression of CaM and human $\text{Na}_V1.5$ in tsA102 cells, and subsequent whole-cell patch clamp recordings, showed no effect on channel function for any of the CaM variants investigated. Only when expressing a fetal $\text{Na}_V1.5$ splice variant, CaM-D130G caused a 7.5-fold increase in persistent Na^+ current, and only at 1 μM free Ca^{2+} . Moreover, native Na^+ currents from fetal mouse cardiomyocytes were not affected by CaM-D130G (Yin et al., 2014). Along the same lines, co-expression of CaM-E141G and $\text{Na}_V1.5$ in tsA102 cells, and subsequent whole-cell patch clamp recordings, caused a 1.7-fold increase in persistent Na^+ current, but the effect was no longer apparent when co-expressed with CaM-WT (Boczek et al., 2016). Since these studies, the $\text{Na}_V1.5$ channel has been under the radar in terms of studying its implication in calmodulinopathies. However, and interestingly, Tarasov and co-workers recently demonstrated that CaM-D96V specifically increased the late current of the $\text{Na}_V1.6$ isoform, but not of $\text{Na}_V1.5$, speculating that this was due to a reduced CaM affinity for the $\text{Na}_V1.6$ IQ-domain (Tarasov et al., 2023).

We have previously shown that CaM-G114R and -G114W reduce the affinity of CaM for both the $\text{Ca}_V1.2$ IQ-domain and for RyR2-CaMBD2 at low to medium Ca^{2+} -concentrations (Brohus et al., 2021). Here we show that both mutations also reduce CaM's affinity for the $\text{Na}_V1.5$ IQ-domain, but the effect is dramatically larger than for the CaMBDs of $\text{Ca}_V1.2$ and RyR2, particularly at free Ca^{2+} -concentrations ≤ 200 nM, the physiological Ca^{2+} concentration in the cardiomyocytes at rest, where the CaM/ $\text{Na}_V1.5$ IQ-domain interaction is essentially abolished.

The interaction between apoCaM and $\text{Na}_V1.5$ is critical for channel function, by tuning channel activity. ApoCaM binding to the $\text{Na}_V1.5$ CTD causes an increase in peak channel open probability as well as a decrease in persistent channel open probability, effects that have divergent implications in disease (Kang et al., 2021). Disruption or weakening of apoCaM binding reduces peak open probability of the channel, corresponding to a loss-of-function effect, such as that observed with the BrS phenotype. However, impaired apoCaM binding can also lead to an increase in persistent $\text{Na}_V1.5$ late current, corresponding to a gain-of-function effect, such as that observed with the LQT3 phenotype (Yan et al., 2017; Kang et al., 2021). The dramatic reduction in the affinity of apoCaM for the $\text{Na}_V1.5$ IQ-domain, caused by substitution of CaM-G114 could thus result in similar divergent effects, and may provide a mechanistic explanation for the mixed phenotypic pattern observed for carriers of CaM-G114 mutations, and potentially other residues affecting $\text{Na}_V1.5$ binding.

But how can a mutation in one of six CaM-encoding alleles display a dominant effect through an ion-channel if the affinity for this channel is dramatically reduced? Several points provide hints towards a possible explanation. The intracellular pool of CaM is limited in cardiomyocytes, suggesting a dynamic competition among CaM target binding sites (Persechini and Stemmer, 2002;




FIGURE 4

Schematic overview of the phenotypic spectrum of the main calmodulin (CaM)-regulated cardiac ion-channels responsible for the cardiac action potential and excitation-contraction coupling. Ion-channels are shown and colored according to the permeating ion: the ryanodine receptor (RyR2, green) and the voltage-gated calcium ($Ca_v1.2$, green), sodium ($Na_v1.5$, red), and potassium ($K_v7.1$, blue) channels. CaM is shown as yellow dumbbells. The involvement of multiple potassium channels in shaping the cardiac action potential is represented by multiple copies of this channel. Clinical phenotypes associated with intrinsic channel mutations are given for the CaM-regulated channels.

Wu et al., 2007). In the GTEx transcript database (GTEx Consortium, 2013), *CALM2* represents ~50% of the total *CALM* transcript pool in ventricular tissue, and a heterozygous *CALM2* missense mutation may thus be present in 25% of the CaM protein pool. Since the impact of a specific CaM mutation differs between targets, the available CaM protein will be redistributed accordingly. Targets affected by a large reduction in CaM binding affinity may experience a larger deficit in CaM saturation than expected from the CaM-mutant/CaM-WT protein ratio, thereby increasing the risk of experiencing a “haploinsufficiency-like” phenotype. This notion is corroborated by a study investigating LQT3 mutations within the $Na_v1.5$ IQ-domain (Yan et al., 2017). The study demonstrated that these variants increase the persistent Na^+ current amplitude of $Na_v1.5$ in whole-cell patch clamp recordings in HEK-cells, and that IQ-domains containing these mutations reduce the CaM binding affinity. Overexpression of CaM-WT rescued the increased current for these LQT3- $Na_v1.5$ channels (Yan et al., 2017).

Another possible explanation for a dominant effect of the CaM-G114 variants is that CaM binds to other parts of the $Na_v1.5$ channel than the IQ-domain. One example is binding of apoCaM to the preIQ-domain with high affinity (~40 nM) (Yoder et al., 2019). Such binding may anchor the CaM-G114 variant to the channel and mediate a pathogenic effect through a compromised IQ-domain binding, induced by the C-lobe mutation.

Intrinsic mutations in $Na_v1.5$ are responsible for BrS and LQTS3, arrhythmogenic conditions both known for cardiac events to frequently occur during rest/sleep (Schwartz et al., 2001; Postema and Wilde, 2008; Takigawa et al., 2008). Additionally, for BrS, fever has been established as a trigger of these events (Adler et al., 2013; Michowitz et al., 2018). Some

of these arrhythmogenic channel mutations occur in the CaM-binding IQ-domain of the $Na_v1.5$ CTD and perturb the CaM/ $Na_v1.5$ interaction (Supplementary Figure S1) (Shah et al., 2006; Yan et al., 2017; Kang et al., 2021; Wu and Liang, 2021). This, together with the impaired apoCaM/ $Na_v1.5$ IQ-domain interaction presented in this work, opens a possible mechanistic explanation for the clinical presentation observed for the CaM-G114R carriers, who died suddenly at a very young age while asleep, potentially triggered by a fever from the infections they each had at the time of death (Brohus et al., 2021). Other intrinsic $Na_v1.5$ BrS/LQTS mutations occur in the NTD of the channel, a domain for which the role of CaM has only recently been explored (Wang et al., 2020). Wang and others demonstrated the ability of CaM to interact with the NTD of $Na_v1.5$ and discussed the potential implication of altered CaM binding in the presence of intrinsic channel disease mutations. We find that the binding of CaM to the $Na_v1.5$ -NTD depends dramatically on Ca^{2+} -concentration (Supplementary Figures S3, S4). When the Ca^{2+} -concentration approaches μM range, the CaM/ $Na_v1.5$ -NTD affinity increases and is comparable to that of the IQ-domain, supporting a potential Ca^{2+} -triggered role of the $Na_v1.5$ NTD in CaM regulation of channel activity.

It is not surprising that the phenotypic range of calmodulinopathies may not yet be fully mapped, as CaM interacts with a myriad of cardiac target proteins, that may or may not be affected by specific CaM missense mutations. In addition to CaM mutation effects on $Na_v1.5$, evidence of $K_v7.1$ effects are accumulating, further expanding the phenotypic spectrum of calmodulinopathies. Kato and others described a family of 14 CaM-N138K carriers who displayed a variably expressed LQTS phenotype from asymptomatic carriers to carriers experiencing sudden death as children (Kato et al., 2022). In support of the LQTS phenotype,

the CaM-N138K variant caused an impairment of Ca_v1.2 inactivation by whole-cell patch clamp recordings of HEK-cells. However, the variant also caused an unexpected potentiation of the K_v7.1 current by the same technique in CHO cells, providing a possible explanation for the variably expressed LQTS phenotype, by countering the Ca_v1.2 effects (Kato et al., 2022). Another comprehensive study, involving 13 arrhythmogenic CaM variants, revealed differential effects of the CaM variants on K_v7.1 binding affinity, channel trafficking, and channel gating (activation) (Kang et al., 2023). Interestingly, as the only one of the 13 variants, CaM-G114W diminished the interaction with the K_v7.1 channel, both at resting and elevated Ca²⁺ concentrations, and induced an increase in K_v7.1 trafficking to the cell membrane (Kang et al., 2023).

In conclusion, the data presented in this study warrants a potential expansion of the phenotypic spectrum of calmodulinopathies. Moreover, these results emphasize our incomplete understanding of the molecular mechanisms possible for calmodulinopathy-related diseases and point to the complexity in variant interpreting due to the mixed phenotypes caused by individual CaM mutations. Molecularly, the multifaceted effects of CaM mutations may act additively or synergistically, thereby contributing to compound and mixed phenotypic expressions. Also, given the high number of CaM-binding targets in cardiomyocytes, the likelihood of observing variably expressed phenotypes in CaM mutation carriers increases, compared to carriers of single ion-channel (or single pathway effecting protein) mutations with pure ‘classical’ phenotypes (Figure 4). This brings the calmodulinopathies to the forefront of scientific research into personalized medicine. Careful interrogation of the *CALM* genes in large cohorts of sequenced individuals with unexplained BrS-like or atypical heart arrhythmia phenotypes should be performed to confirm an expansion of the phenotypic spectrum of calmodulinopathies to include *CALM*-BrS. Such knowledge will allow for earlier and more accurate diagnosis and treatment of individuals with calmodulinopathies.

Data availability statement

The original contributions presented in the study are included in the article/Supplementary Files, further inquiries can be directed to the corresponding author.

Ethics statement

Written informed consent was obtained from the individuals, and minors’ legal guardian/next of kin, for the

publication of any potentially identifiable images or data included in this article.

Author contributions

MN, RW, and MO contributed to conception and design of the study. MB and MO organized the laboratory work, which was carried out by A-OB. MB wrote the first draft of the manuscript. All authors contributed to the article and approved the submitted version.

Funding

This work was supported by a research grant from the Lundbeck Foundation (no. R324-2019-1933), and a research grant from the Independent Danish Research Foundation, Council for Natural Sciences (no. 2032-00333B).

Conflict of interest

MN and MO acted as expert witnesses in the 2022 New South Wales Inquiry into the convictions of the Australian woman carrying the CaM-G114R variant.

The remaining authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher’s note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Supplementary material

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fphar.2023.1210140/full#supplementary-material>

References

- Adler, A., Guy, T., Heller, K., Zeltser, D., Ohayon, T., Rozovski, U., et al. (2013). Fever-induced Brugada pattern: how common is it and what does it mean? *Heart rhythm*. 10 (9), 1375–1382. doi:10.1016/j.hrthm.2013.07.030
- Berchtold, M. W., and Villalobo, A. (2014). The many faces of calmodulin in cell proliferation, programmed cell death, autophagy, and cancer. *Biochimica Biophysica Acta* 1843 (2), 398–435. doi:10.1016/j.bbamcr.2013.10.021
- Berchtold, M. W., Egli, R., Rhyner, J. A., Hameister, H., and Strehler, E. E. (1993). Localization of the human bona fide calmodulin genes CALM1, CALM2, and CALM3 to chromosomes 14q24-q31, 2p21.1-p21.3, and 19q13.2-q13.3, 4611-P21.3, and 19q13.2-Q13.3. *Genomics* 16 (2), 461–465. doi:10.1006/geno.1993.1211
- Boczek, N. J., Gomez-Hurtado, N., Ye, D., Calvert, M. L., Tester, D. J., Kryshtal, D., et al. (2016). Spectrum and prevalence of CALM1-CALM2-and CALM3-encoded calmodulin variants in long QT syndrome and functional characterization of a novel long QT syndrome-associated calmodulin missense variant, E141G. *Circ. Cardiovasc. Genet.* 9 (2), 136–146. doi:10.1161/CIRCGENETICS.115.001323
- Brohus, M., Arsov, T., Wallace, D. A., Jensen, H. H., Nyegaard, M., Crotti, L., et al. (2021). Infanticide vs. Inherited cardiac arrhythmias. *Europace* 23 (3), 441–450. doi:10.1093/europace/euaa272
- Brohus, M., Søndergaard, M. T., Michael, T., van Petegem, F., and Overgaard, M. T. (2019). Ca²⁺-Dependent calmodulin binding to cardiac ryanodine receptor (RyR2) calmodulin-binding domains. *Biochem. J.* 476 (2), 193–209. doi:10.1042/BCJ20180545

- Bycroft, C., Freeman, C., Petkova, D., Gavin, B., Elliott, L. T., Sharp, K., et al. (2018). The UK Biobank resource with deep phenotyping and genomic data. *Nature* 562 (7726), 203–209. doi:10.1038/s41586-018-0579-z
- Chagot, B., and Chazin, W. J. (2011). Solution NMR structure of apo-calmodulin in complex with the IQ motif of human cardiac sodium channel NaV1.5. *J. Mol. Biol.* 406 (1), 106–119. doi:10.1016/j.jmb.2010.11.046
- Chen, S., Francioli, L. C., Goodrich, J. K., Collins, R. L., Wang, Q., Alfoldi, J., et al. (2022). A genome-wide mutational constraint map quantified from variation in 76,156 human genomes. *BioRxiv*. doi:10.1101/2022.03.20.485034
- Chin, D., and Anthony, R. (2000). Calmodulin: A prototypical calcium sensor. *Trends Cell. Biol.* 10 (8), 322–328. doi:10.1016/s0962-8924(00)01800-6
- Clapham, D. E. (2007). Calcium signaling. *Cell*. 131 (6), 1047–1058. doi:10.1016/j.cell.2007.11.028
- Crotti, L., Johnson, C. N., Graf, E., De Ferrari, G. M., Cuneo, B. F., Ovadia, M., et al. (2013). Calmodulin mutations associated with recurrent cardiac arrest in infants. *Circulation* 127 (9), 1009–1017. doi:10.1161/CIRCULATIONAHA.112.001216
- Crotti, L., Spazzolini, C., Tester, D. J., Ghidoni, A., Beckmann, B. M., Behr, E. R., et al. (2019). Calmodulin mutations and life-threatening cardiac arrhythmias: insights from the international calmodulinopathy registry. *Eur. Heart J.* 40 (35), 2964–2975. doi:10.1093/eurheartj/ehz311
- Friedberg, F., and Rhoads, A. R. (2001). Evolutionary aspects of calmodulin. *IUBMB Life* 51 (4), 215–221. doi:10.1080/152165401753311753
- Gabelli, S. B., Jesse, B., Tomaselli, G. F., and Amzel, L. M. (2016). Calmodulin and Ca(2+) control of voltage gated Na(+) channels. *Channels (Austin, Tex.)* 10 (1), 45–54. doi:10.1080/19336950.2015.1075677
- Gabelli, S. B., Bianchet, M. A., Farinelli, F., Srinivas, A., Yoder, J., Jean, J., et al. (2014). Regulation of the NaV1.5 cytoplasmic domain by calmodulin. *Nat. Commun.* 5, 5126. doi:10.1038/ncomms6126
- GTEx Consortium (2013). The genotype-tissue expression (GTEx) project. *Nat. Genet.* 45 (6), 580–585. doi:10.1038/ng.2653
- Holt, C., Hamborg, L., Lau, K., Brohus, M., Larsen, K. T., Sommer, C., et al. (2020). The arrhythmogenic N53I variant subtly changes the structure and dynamics in the calmodulin N-terminal domain, altering its interaction with the cardiac ryanodine receptor. *J. Biol. Chem.* 295 (22), 7620–7634. doi:10.1074/jbc.RA120.013430
- Jensen, H. H., Brohus, M., Nyegaard, M., and Overgaard, M. T. (2018). Human calmodulin mutations. *Front. Mol. Neurosci.* 11, 396. doi:10.3389/fnmol.2018.00396
- Jiménez-Jáimez, J., Ortega, Á., Macías-Ruiz, R., Francesca Perin, M., Perin, F., Rodríguez-Vázquez del Rey, M. M., et al. (2016). Calmodulin 2 mutation N98S is associated with unexplained cardiac arrest in infants due to low clinical penetrance electrical disorders. *PLoS One* 11 (4), e0153851. doi:10.1371/journal.pone.0153851
- Johnson, C. N., Potet, F., Thompson, B., Chazin, W. J., Glazer, A. M., Voehler, M. W., et al. (2018). A mechanism of calmodulin modulation of the human cardiac sodium channel. *Struct. Lond. Engl.* 26 (5), 683–694. doi:10.1016/j.str.2018.03.005
- Kang, P. W., Chakouri, N., Diaz, J., Tomaselli, G. F., Yue, D. T., and Ben-Johny, M. (2021). Elementary mechanisms of calmodulin regulation of NaV1.5 producing divergent arrhythmic phenotypes. *Proc. Natl. Acad. Sci. U. S. A.* 118 (21), e2025085118. doi:10.1073/pnas.2025085118
- Kang, P. W., Woodbury, L., Angsutararux, P., Sambare, N., Shi, J., Marras, M., et al. (2023). Arrhythmia-associated calmodulin variants interact with KCNQ1 to confer aberrant membrane trafficking and function. *Prepr. Serv. Biol.* 2023, 526031. doi:10.1101/2023.01.28.526031
- Kato, K., Isbell, H. M., Fressart, V., Denjoy, I., Amal, D., Itoh, H., et al. (2022). Novel CALM3 variant causing calmodulinopathy with variable expressivity in a 4-generation family. *Circulation. Arrhythmia Electrophysiol.* 15 (3), e010572. doi:10.1161/CIRCEP.121.010572
- Limpitikul, W. B., Dick, I. E., Yue, D. T., Overgaard, M. T., and George, A. L. (2014). Calmodulin mutations associated with long QT syndrome prevent inactivation of cardiac L-type Ca(2+) currents and promote proarrhythmic behavior in ventricular myocytes. *J. Mol. Cell. Cardiol.* 74, 115–124. doi:10.1016/j.yjmcc.2014.04.022
- Makita, N., Yagihara, N., Crotti, L., Johnson, C. N., Beckmann, B.-M., Roh, M. S., et al. (2014). Novel calmodulin mutations associated with congenital arrhythmia susceptibility. *Circ. Cardiovasc. Genet.* 7 (4), 466–474. doi:10.1161/CIRCGENETICS.113.000459
- Marsman, R. F., Barc, J., Beekman, L., Alders, M., Dooijes, D., Arthur van den Wijngaard, et al. (2014). A mutation in CALM1 encoding calmodulin in familial idiopathic ventricular fibrillation in childhood and adolescence. *J. Am. Coll. Cardiol.* 63 (3), 259–266. doi:10.1016/j.jacc.2013.07.091
- Michowitz, Y., Milman, A., Sarquella-Brugada, G., Andorin, A., Champagne, J., Postema, P. G., et al. (2018). Fever-related arrhythmic events in the multicenter survey on arrhythmic events in Brugada syndrome. *Heart rhythm.* 15 (9), 1394–1401. doi:10.1016/j.hrthm.2018.04.007
- Nyegaard, M., Overgaard, M. T., Søndergaard, M. T., Vranas, M., Behr, E. R., Hildebrandt, L. L., et al. (2012). Mutations in calmodulin cause ventricular Tachycardia and sudden cardiac death. *Am. J. Hum. Genet.* 91 (4), 703–712. doi:10.1016/j.ajhg.2012.08.015
- Nyegaard, M., and Overgaard, M. T. (2019). The international calmodulinopathy registry: recording the diverse phenotypic spectrum of un-CALM hearts. *Eur. Heart J.* 40 (35), 2976–2978. doi:10.1093/eurheartj/ehz463
- Persechini, A., and Stemmer, P. M. (2002). Calmodulin is a limiting factor in the cell. *Trends Cardiovasc. Med.* 12 (1), 32–37. doi:10.1016/s1050-1738(01)00144-x
- Postema, P. G., and Wilde, A. A. M. (2008). Arrhythmias in Brugada syndrome: changing throughout day and season? *Heart rhythm.* 5 (11), 1528–1529. doi:10.1016/j.hrthm.2008.08.033
- Potet, F., Chagot, B., Anghelescu, M., Viswanathan, P. C., Stepanovic, S. Z., Kupersmidt, S., et al. (2009). Functional interactions between distinct sodium channel cytoplasmic domains through the action of calmodulin. *J. Biol. Chem.* 284 (13), 8846–8854. doi:10.1074/jbc.M806871200
- Rocchetti, M., Sala, L., Dreizehnter, L., Crotti, L., Sinnecker, D., Mura, M., et al. (2017). Elucidating arrhythmogenic mechanisms of long-QT syndrome CALM1-F142L mutation in patient-specific induced pluripotent stem cell-derived cardiomyocytes. *Cardiovasc. Res.* 113 (5), 531–541. doi:10.1093/cvr/cvx006
- Rossi, A. M., and Taylor, C. W. (2011). Analysis of protein-ligand interactions by fluorescence polarization. *Nat. Protoc.* 6 (3), 365–387. doi:10.1038/nprot.2011.305
- Sarhan, M. F., and Ahern, C. A. (2009). A double tyrosine motif in the cardiac sodium channel domain III-IV linker couples calcium-dependent calmodulin binding to inactivation gating. *J. Biol. Chem.* 284 (48), 33265–33274. doi:10.1074/jbc.M109.052910
- Sarhan, M. F., Tung, C., and Ahern, C. A. (2012). Crystallographic basis for calcium regulation of sodium channels. *Proc. Natl. Acad. Sci. U. S. A.* 109 (9), 3558–3563. doi:10.1073/pnas.1114748109
- Schwartz, P. J., Priori, S. G., Spazzolini, C., Moss, A. J., Vincent, G. M., Napolitano, C., et al. (2001). Genotype-phenotype correlation in the long-QT syndrome: gene-specific triggers for life-threatening arrhythmias. *Circulation* 103 (1), 89–95. doi:10.1161/01.cir.103.1.89
- Shah, V. N., Tammy, L., Chazin, W. J., Williams, C. K., and Balsler, J. R. (2006). Calcium-dependent regulation of the voltage-gated sodium channel HH1: intrinsic and extrinsic sensors use a common molecular switch. *Proc. Natl. Acad. Sci. U. S. A.* 103 (10), 3592–3597. doi:10.1073/pnas.0507397103
- Søndergaard, M. T., Liu, Y., Brohus, M., Guo, W., Nani, A., Carvajal, C., et al. (2019). Diminished inhibition and facilitated activation of RyR2-mediated Ca2+ release is a common defect of arrhythmogenic calmodulin mutations. *FEBS J.* 286 (22), 4554–4578. doi:10.1111/febs.14969
- Søndergaard, M. T., Liu, Y., Guo, W., Wei, J., Wang, R., Brohus, M., et al. (2020). Role of cardiac ryanodine receptor calmodulin-binding domains in mediating the action of arrhythmogenic calmodulin N-domain mutation N54I. *FEBS J.* 287 (11), 2256–2280. doi:10.1111/febs.15147
- Søndergaard, M. T., Liu, Y., Larsen, K. T., Nani, A., Tian, X., Holt, C., et al. (2017). The arrhythmogenic calmodulin p.Phe142Leu mutation impairs C-domain Ca2+ binding but not calmodulin-dependent inhibition of the cardiac ryanodine receptor. *J. Biol. Chem.* 292 (4), 1385–1395. doi:10.1074/jbc.M116.766253
- Søndergaard, M. T., Sorensen, A. B., Skov, L. L., Kjaer-Sorensen, K., Bauer, M. C., Nyegaard, M., et al. (2015a). Calmodulin mutations causing catecholaminergic polymorphic ventricular Tachycardia confer opposing functional and biophysical molecular changes. *FEBS J.* 282 (4), 803–816. doi:10.1111/febs.13184
- Søndergaard, M. T., Tian, X., Liu, Y., Wang, R., Chazin, W. J., Chen, S. R. W., et al. (2015b). Arrhythmogenic calmodulin mutations affect the activation and termination of cardiac ryanodine receptor-mediated Ca2+ release. *J. Biol. Chem.* 290 (43), 26151–26162. doi:10.1074/jbc.M115.676627
- Sorensen, A. B., Søndergaard, M. T., and Overgaard, M. T. (2013). Calmodulin in a heartbeat. *FEBS J.* 280 (21), 5511–5532. doi:10.1111/febs.12337
- Takigawa, M., Noda, T., Shimizu, W., Miyamoto, K., Okamura, H., Satomi, K., et al. (2008). Seasonal and circadian distributions of ventricular fibrillation in patients with Brugada syndrome. *Heart rhythm.* 5 (11), 1523–1527. doi:10.1016/j.hrthm.2008.08.022
- Tarasov, M., Struckman, H. L., Olgar, Y., Miller, A., Demirtas, M., Bogdanov, V., et al. (2023). NaV1.6 dysregulation within myocardial T-tubules by D96V calmodulin enhances proarrhythmic sodium and calcium mishandling. *J. Clin. Investigation* 133 (7), e152071. doi:10.1172/JCI152071
- Tidow, H., and Nissen, P. (2013). Structural diversity of calmodulin binding to its target sites. *FEBS J.* 280 (21), 5551–5565. doi:10.1111/febs.12296
- Toutenhoofd, S. L., and Strehler, E. E. (2000). The calmodulin multigene family as a unique case of genetic redundancy: multiple levels of regulation to provide spatial and temporal control of calmodulin pools? *Cell. Calcium* 28 (2), 83–96. doi:10.1054/ceca.2000.0136
- Urrutia, J., Aguado, A., Muguza-Montero, A., Núñez, E., Malo, C., Casis, O., et al. (2019). The crossroad of ion channels and calmodulin in disease. *Int. J. Mol. Sci.* 20 (2), 400. doi:10.3390/ijms20020400
- Villarreal, A., Taghialatela, M., Bernardo-Seisdedos, G., Alaimo, A., Agirre, J., Alberdi, A., et al. (2014). The ever changing moods of calmodulin: how structural plasticity entails transductional adaptability. *J. Mol. Biol.* 426 (15), 2717–2735. doi:10.1016/j.jmb.2014.05.016
- Wang, C., Chung, B. C., Yan, H., Wang, H. G., Lee, S. Y., and Pitt, G. S. (2014). Structural analyses of Ca2+/CaM interaction with NaV channel C-termini reveal mechanisms of calcium-dependent regulation. *Nat. Commun.* 5 (1), 4896. doi:10.1038/ncomms5896

- Wang, C., Chung, B. C., Yan, H., Lee, S. Y., and Pitt, G. S. (2012). Crystal structure of the ternary complex of a NaV C-terminal domain, a fibroblast growth factor homologous factor, and calmodulin. *Struct. Lond. Engl.* 20 (7), 1167–1176. doi:10.1016/j.str.2012.05.001
- Wang, K., Brohus, M., Holt, C., Overgaard, M. T., Wimmer, R., and Filip Van Petegem (2020a). Arrhythmia mutations in calmodulin can disrupt cooperativity of Ca²⁺ binding and cause misfolding. *J. Physiology* 598 (6), 1169–1186. doi:10.1113/JP279307
- Wang, K., Holt, C., Lu, J., Brohus, M., Larsen, K. T., Overgaard, M. T., et al. (2018). Arrhythmia mutations in calmodulin cause conformational changes that affect interactions with the cardiac voltage-gated calcium channel. *Proc. Natl. Acad. Sci. U. S. A.* 115 (45), E10556–E10565. doi:10.1073/pnas.1808733115
- Wang, Z., Vermij, S. H., Sottas, V., Anna, S., Ross-Kaschitza, D., Zaklyazminskaya, E. V., et al. (2020b). Calmodulin binds to the N-terminal domain of the cardiac sodium channel Nav1.5. *Channels (Austin, Tex)* 14 (1), 268–286. doi:10.1080/19336950.2020.1805999
- Wu, X., and Donald, M. (2007). Free and bound intracellular calmodulin measurements in cardiac myocytes. *Cell Calcium* 41 (4), 353–364. doi:10.1016/j.ceca.2006.07.011
- Wu, X., and Liang, H. (2021). Calmodulin interactions with voltage-gated sodium channels. *Int. J. Mol. Sci.* 22 (18), 9798. doi:10.3390/ijms22189798
- Xia, Z., and Storm, D. R. (2005). The role of calmodulin as a signal integrator for synaptic plasticity. *Nat. Rev. Neurosci.* 6 (4), 267–276. doi:10.1038/nrn1647
- Yan, H., Wang, C., Marx, S. O., and Pitt, G. S. (2017). Calmodulin limits pathogenic Na⁺ channel persistent current. *J. General Physiology* 149 (2), 277–293. doi:10.1085/jgp.201611721
- Yap, K. L., Kim, J., Truong, K., Sherman, M., Yuan, T., and Ikura, M. (2000). Calmodulin target database. *J. Struct. Funct. Genomics* 1 (1), 8–14. doi:10.1023/a:1011320027914
- Yin, G., Hassan, F., Ayman, R. H., Murphy, L. L., Crotti, L., Peter, J., et al. (2014). Arrhythmogenic calmodulin mutations disrupt intracellular cardiomyocyte Ca²⁺ regulation by distinct mechanisms. *J. Am. Heart Assoc.* 3 (3), e000996. doi:10.1161/JAHA.114.000996
- Yoder, J. B., Ben-Johny, M., Farinelli, F., Srinivasan, L., Shoemaker, S. R., Tomaselli, G. F., et al. (2019). Ca²⁺-Dependent regulation of sodium channels Nav1.4 and Nav1.5 is controlled by the post-IQ motif. *Nat. Commun.* 10 (1), 1514. doi:10.1038/s41467-019-09570-7