



TDHF Theory and Its Extensions for the Multinucleon Transfer Reaction: A Mini Review

Kazuyuki Sekizawa*

Center for Transdisciplinary Research, Institute for Research Promotion, Niigata University, Niigata, Japan

Time-dependent Hartree-Fock (TDHF) theory has been a powerful tool in describing a variety of complex nuclear dynamics microscopically without empirical parameters. In this contribution, recent advances in nuclear dynamics studies employing TDHF and its extensions are briefly reviewed, in line with the study of multinucleon transfer (MNT) reactions. The latter lies at the core of this Research Topic, whose application for the production of extremely neutron-rich nuclei has been extensively discussed in recent years. Having in mind the ongoing theoretical developments, it is envisaged how microscopic theories may contribute to the future MNT study.

Keywords: low-energy heavy-ion reactions, multinucleon transfer, quasifission, fusion, neutron-rich nuclei, TDHF, TDDFT

OPEN ACCESS

Edited by:

Angela Bonaccorso,
Istituto Nazionale di Fisica Nucleare,
Sezione di Pisa, Italy

Reviewed by:

Stefan Typel,
Technische Universität Darmstadt,
Germany
Kazuhiro Yabana,
University of Tsukuba, Japan

*Correspondence:

Kazuyuki Sekizawa
sekizawa@phys.sc.niigata-u.ac.jp

Specialty section:

This article was submitted to
Nuclear Physics,
a section of the journal
Frontiers in Physics

Received: 14 December 2018

Accepted: 04 February 2019

Published: 12 March 2019

Citation:

Sekizawa K (2019) TDHF Theory and
Its Extensions for the Multinucleon
Transfer Reaction: A Mini Review.
Front. Phys. 7:20.
doi: 10.3389/fphy.2019.00020

1. INTRODUCTION

Multinucleon transfer (MNT) reactions may be a promising means to produce yet-unknown, neutron-rich unstable nuclei, whose production is difficult by other methods. This possibility makes the study of MNT both fascinating and important. For instance, besides the fundamental interest in nuclear structures such as shell evolution [1] and shape transitions [2, 3], properties of extremely neutron-rich nuclei are crucial to fully understanding the detailed scenario of r -process nucleosynthesis [4]. Needless to say, the historic detection of gravitational waves from a merger of two neutron stars, GW170817 [5], together with radiation from a short γ -ray burst, GRB 170817A [6], followed by a kilonova [7] makes it a timely and imperative task to unveil inherent properties of the nuclei far away from stability. Furthermore, the superheavy nuclei produced so far are neutron-deficient ones, and the predicted island of stability has not yet been reached [8, 9]. The production of neutron-rich superheavy nuclei in the island is highly desired, as it would provide a new stringent constraint for microscopic theories. Therefore, the study of MNT reactions is listed as one of the key subjects at the current and future RI beam facilities, such as RIBF (RIKEN, Japan) [10], HIRFL-CSR and HIAF (IMP, China)¹, RAON (RISP, Korea) [11], DRIB (FLNR, Russia)², SPIRAL2 (GANIL, France) [12], FAIR (GSI, Germany)³, SPES (INFN, Italy)⁴, FRIB (MSU, USA) [13], and so on.

This review briefly digests recent advances of microscopic reaction theories to stimulate new ideas for the future study of MNT reactions. In section 2, the theoretical framework of TDHF is succinctly recalled. In section 3, the current status of the MNT study with TDHF is given. In section 4, recent advances of theoretical approaches are outlined, along with a discussion on possible future applications. The article is concluded in section 5. The readers are intended to refer to other articles in this Research Topic for discussions on experimental as well as other theoretical studies.

¹<http://english.imp.cas.cn/Home2017/>

²<http://flerovlab.jinr.ru/flnr/dribs.html>

³<https://fair-center.eu/>

⁴<https://web.infn.it/spes/>

2. THE TDHF THEORY

The TDHF theory was first proposed by Dirac [14] and its application to nuclear physics started in the 1970s [15–22]. The TDHF theory can be formally derived based on the time-dependent variational principle (various derivations can be found in a recent review [23]). In TDHF, the wave function is a Slater determinant for all times, and the Pauli principle is thus automatically ensured. The TDHF equations read as follows:

$$i\hbar \frac{\partial \phi_i(\mathbf{r}\sigma q, t)}{\partial t} = \hat{h}[\rho(t)]\phi_i(\mathbf{r}\sigma q, t), \quad (1)$$

where $\phi_i(\mathbf{r}\sigma q, t)$ is the single-particle wave function of i th nucleon at position \mathbf{r} with spin σ and isospin q . $\hat{h}[\rho(t)] = \frac{\delta E}{\delta \rho}$ is the single-particle Hamiltonian with ρ being the one-body density. $E[\rho]$ is the total energy, which may be regarded as an energy density functional (EDF) in the context of time-dependent density functional theory (TDDFT)⁵ [27–31]. The EDF is constructed so as to reproduce static properties of finite nuclei in a wide mass region and also the basic nuclear matter properties (see e.g., [32–35] and references therein). There is no adjustable parameter in TDHF, once an EDF is given. The non-linearity arises because $\hat{h}[\rho(t)]$ contains the mean-field potential which depends on densities generated by all the nucleons, e.g., $\rho(\mathbf{r}, t) = \sum_{i,\sigma,q} |\phi_i(\mathbf{r}\sigma q, t)|^2$. Such non-linear couplings between single-particle and collective degrees of freedom give rise to the so-called one-body dissipation mechanism (known as wall-and-window formulas [36]). Note that as is evident from a derivation of TDHF from the lowest order truncation of the Bogoliubov-Born-Green-Kirkwood-Yvon (BBGKY) hierarchy [23], where two-body correlations are neglected, two-body dissipations associated with nucleon-nucleon collisions are absent in TDHF. They are, however, expected to play a minor role at low energy due to the Pauli exclusion principle.

TDHF is notably versatile—by changing initial condition, external potential, boundary condition, etc., one can study a wide variety of phenomena: not only collective excitations in the linear response regime [37–46], where TDHF is formally equivalent to the random phase approximation (RPA), but also induced fission [47–52], dynamics of exotic configurations like α -chain structure [53], toroidal nucleus [54, 55], nuclear pasta formation [56–59], as well as heavy-ion reactions, such as nucleon transfer [60–74], quasifission (QF) [75–87], and fusion [88–104], and so on (see [22, 23, 26, 31, 105, 106], for review papers).

In the case of heavy-ion reactions, the initial wave function is composed of projectile and target nuclei in their ground state, which is obtained by self-consistently solving the static Hartree-Fock equations [$i\hbar\partial/\partial t \rightarrow \varepsilon_i$ in Equation 1]. Those wave

functions are placed in a computational box, with a sufficiently large relative distance, boosted with a proper relative momentum. The time evolution according to Equation (1) then allows us to follow reaction dynamics in real space and real time. As the theory deals with the single-particle wave functions of nucleons, not only dynamic effects such as nucleon transfer and internal excitations, but also structural effects such as static/dynamic shell effects and shape evolution are naturally incorporated into the description. Spin-orbit coupling is known to play an important role in energy dissipation processes in heavy-ion reactions [107–110]. The effects of the tensor term on nuclear dynamics were also investigated recently [101, 111–114]. TDHF enables us to study rich and complex physics of low-energy heavy-ion reactions from nucleonic degrees of freedom.


3. TDHF THEORY FOR MULTINUCLEON TRANSFER REACTIONS

3.1. Multinucleon Transfer Reactions

The MNT reaction may be regarded as a non-equilibrium quantum transport process of nucleons during a collision. With the help of the particle-number projection technique [64], transfer probabilities can be deduced from a TDHF wave function after collision. In Sekizawa and Yabana [65, 66], a range of reactions at energies around the Coulomb barrier were studied within TDHF, for which precise experimental data are available [115–118]. Those reactions are characterized by different neutron-to-proton ratios, N/Z , and charge product, $Z_P Z_T$. TDHF identified two distinct transfer mechanisms: (1) a fast isospin equilibration process and (2) transfer of many nucleons associated with dynamics of neck formation and its breaking. The neck breaking dynamics emerge at small impact parameters, especially in reactions with a large $Z_P Z_T$ ($\gtrsim 1,600$). The latter may be regarded as a precursor to QF, where the system starts evolving toward the mass equilibrium. Comparisons with measured cross sections showed that TDHF works fairly well in accuracy comparable to other model predictions. However, effects of secondary neutron evaporation become substantial, when many protons are transferred. Combining TDHF with a statistical compound-nucleus deexcitation model, GEMINI++ [119], it became possible to compute production cross sections after secondary deexcitation processes. With the method, dubbed TDHF+GEMINI, it was shown that the inclusion of deexcitation effects substantially improves agreement with the experimental data [72] (see **Figures 1A,B**). (For more details, see, e.g., [69, 72]).

The description is still, however, not perfect, especially for channels where both neutrons and protons are removed. It would be improved when one includes missing one-body fluctuations into the description (cf. section 4.2). Nevertheless, TDHF may be used, taking advantage of its non-empirical nature, to explore the optimal or novel reaction mechanism for producing new neutron-rich unstable nuclei, at least qualitatively. Recently, other groups have also implemented TDHF+GEMINI: e.g.

⁵ Here a local EDF (like Skyrme) has been assumed, which makes the TDHF Equation (1) local in space, as it is currently used in most practical applications. In general, $E = \int \mathcal{E}(\mathbf{r})d\mathbf{r}$ is composed of various local densities, $\xi \equiv \{\rho, \tau, \dots\}$, and the TDHF Equation (1) should read: $i\hbar \frac{\partial \phi_i(\mathbf{r}\sigma q, t)}{\partial t} = \int \sum_k \frac{\delta \mathcal{E}[\xi(\mathbf{r}, t)]}{\delta \xi_k(\mathbf{r}, t)} \frac{\delta \xi_k(\mathbf{r}, t)}{\delta \phi_i^*(\mathbf{r}\sigma q, t)} d\mathbf{r}' = \sum_{\sigma'} \hat{h}_{\sigma\sigma'}^{(q)}(\mathbf{r}, t)\phi_i(\mathbf{r}\sigma' q, t)$. In such a case, the single-particle Hamiltonian can have spin dependence as well as differential operators (for the explicit form, see, e.g., [24–26]).



production of $N = 126$ neutron-rich nuclei was investigated for the $^{132}\text{Sn}+^{208}\text{Pb}$ reaction [122].

3.2. Quasifission Processes

To synthesize the heaviest elements and also to explore the predicted island of stability, it is crucial to establish deep understanding of the main competitive process, the *quasifission*. The QF process is characterized by a large amount of mass transfer, full energy dissipation, and a long contact time (typically several to tens of zeptoseconds). The characteristic observables of QF are correlations between fragment masses with scattering angles, known as mass-angle distribution (MAD), or with the total kinetic energy (TKE) of outgoing fragments, sometimes called mass-energy distribution (MED) (see e.g., [121, 123–126]). In recent applications of TDHF, it has been shown that TDHF can quantitatively describe main QF dynamics, consistent with experimental observations [70, 78, 80, 81] (see **Figures 1C,D**). An intriguing indication is that in collisions involving an actinide nucleus, TKE becomes larger (smaller energy dissipation) for

tip collisions as compared to side collisions [70, 84], perhaps due to shell effects of ^{208}Pb [127]. This may be in contrast to the naive consideration, where one expects a compact shape at scission for side collisions, which would result in larger TKE for side collisions.

The revival of interest in the MNT reaction was initiated by the seminal work by Zagrebaev and Greiner [128–137]. In their work, the importance of shell effects in MNT processes was emphasized. For example, in the $^{238}\text{U}+^{248}\text{Cm}$ reaction, ^{238}U tends to evolve toward doubly-magic ^{208}Pb , giving the rest of the 30 nucleons to the heavier partner, resulting in the production of transcurium nuclei as primary reaction products. The latter process is called *inverse* (anti-symmetrizing) QF (IQF), since the ordinary QF drives the system toward the mass equilibrium. Clearly, it is essential to correctly include static and dynamic effects of nuclear shells to describe IQF. The IQF was also observed in TDHF. In Kedziora and Simenel [76], a typical “tip-on-side” orientation in collisions of two actinide nuclei, $^{232}\text{Th}+^{250}\text{Cf}$, was found to result in IQF, where about 15 nucleons in the tip of ^{232}Th were transferred to the side of ^{250}Cf . Another type of IQF was reported in Sekizawa [73], where complex surface vibration modes are induced in tip collisions of $^{238}\text{U}+^{124}\text{Sn}$ at energies well above the

⁶ Website of Nuclear Reactions Video (NRV) Project. Available online at: <http://nrv.jinr.ru/nrv/webnrv/grazing/>

Coulomb barrier, which resulted in an abrupt development of a neck, forming a smaller subsystem, leading to transfer of around 27 nucleons from ^{124}Sn to ^{238}U .

An important message here is that those IQF processes emerge as a main reaction outcome in TDHF. Since larger effects of fluctuations and correlations, that give rise to a wider fragment mass distribution, are expected in such damped collisions [105, 138–140], novel IQF processes may pave new pathways to unexplored territories in the nuclear chart far away from stability.

4. NEW TECHNIQUES AND EXTENSIONS

4.1. Extraction of Macroscopic Physical Ingredients

The potential energy surface in a colliding nuclear system is often a key ingredient in phenomenological approaches. With the constrained Hartree-Fock technique [141], one can compute the adiabatic potential energy in a composite system. However, it differs, in general, from the potential in the entrance channel of a collision, where incident-energy-dependent dynamic effects come into play. Such a dynamic potential should be derived from microscopic theories, e.g., TDHF, where dynamic effects such as shape deformation (or necking), nucleon transfer, and inelastic excitations through the one-body dissipation mechanism are naturally taken into account. A method named dissipative-dynamics TDHF (DD-TDHF) [82, 142, 143] allows us to extract from TDHF not only a dynamic potential, but also a friction coefficient, based on a mapping of TDHF trajectories onto a set of classical equations of motion. Density-constrained TDHF (DC-TDHF) [88] is another way to extract a potential. In the latter approach, the density distribution obtained from TDHF is used as a constraint for Hartree-Fock calculations, leading to an adiabatic potential along the TDHF dynamical trajectory (see e.g., [53, 89–101], for various applications). Energy dependence as well as transition from dynamic to adiabatic potentials may be important as well. Later, the idea of the density constraint was also applied to extract a bare (without density distortion) potential taking full account of the Pauli exclusion principle, which is named density-constrained frozen Hartree-Fock (DC-FHF) [144]. Such a bare potential can be applied to, e.g., coupled-channel calculations for fusion reactions [145, 146]. Indeed, such a connection was established, where excitation energies of low-lying collective excitations were also determined by TDHF [147]. This is a remarkable example of building a bridge between elaborated macroscopic or phenomenological frameworks and those rooted with microscopic degrees of freedom, that may enhance predictability of the former.

4.2. Balian-Vénéroni Variational Principle and TDRPA

While there are, of course, limitations inherent in TDHF (e.g., the absence of many-body correlations, deterministic nature as well as spurious cross-channel couplings due to the single mean-field description [141]), some of them can nowadays be overcome with extended approaches. One such approach can be obtained from the variational principle of Balian and Vénéroni

[148, 149]. The Balian-Vénéroni variational principle enables us to control the variational space according to not only the state in the Schrödinger picture, but also to the observable in the Heisenberg picture. For instance, a variation optimized for a Slater determinant and a one-body observable derives the TDHF equations [105]. By extending the variational space for fluctuations of a one-body observable, one obtains a formula that includes effects of one-body fluctuations on top of the TDHF mean-field trajectory, which may be regarded as time-dependent RPA (TDRPA). It has recently been shown that the description of the width of fragment mass distributions in deep-inelastic collisions, which is severely underestimated in TDHF [17, 18, 150], is substantially improved [151, 152], or even quantitatively agrees with experimental data [153], in TDRPA. It implies that one-body dissipation and fluctuations described by TDHF and TDRPA, respectively, are the predominant mechanisms in deep-inelastic collisions. Here, caution is required: the TDRPA formula in the present form can only be applied to symmetric reactions [153]. To overcome this difficulty, one may extend the derivation based on the Balian-Vénéroni variational principle to include higher-order corrections, or derive a formula generalized for asymmetric reactions from, e.g., the stochastic mean-field (SMF) theory [154, 155], from which one can derive the TDRPA formula in the small fluctuation limit [154]. Note that effects of two-body correlations were also indicated by the time-dependent density-matrix (TDDM) approach [156–160].

4.3. Stochastic Extensions

Another limitation of TDHF lies in its deterministic nature. To correctly describe abundant reaction outcomes as observed experimentally, one may need to introduce quantal, collisional, or thermal fluctuations that induce bifurcations of dynamics, leading to, in the end, minor processes apart from the TDHF mean-field trajectory. In the SMF approach, extensively developed by Ayik and his coworkers [138–140, 154, 155, 161–169], initial fluctuations are introduced in a stochastic manner, which gives rise to an ensemble of final states after dynamic evolutions. For the MNT study, the SMF description can be cast into the Fokker-Planck equations, where transport (drift and diffusion) coefficients are determined by the single-particle orbitals in TDHF. In Yilmaz et al. [169], the SMF approach was also applied to the Ni+Ni reaction, showing quantitative agreement with experimental data [153], comparable to TDRPA. Recently, Bulgac et al. [170] proposed another way to introduce stochasticity by means of external potentials which are random in both space and time. The method was applied to describe induced fission processes, with a simplified orbital-free approach, getting a sufficient width of the fission fragment mass distribution [170]. Further applications of those approaches may be promising for exploring optimal reactions, especially for producing the most exotic isotopes, where correct description of rare processes may be essential.

4.4. Inclusion of Pairing

Despite the known importance of pairing in nuclear structure studies, the effect of pairing on nuclear reaction dynamics has rarely been investigated thus far. While the BCS approximation

has been employed to date [171–174], inclusion of dynamic effects of pairing became possible only quite recently with top-tier supercomputers. One may naively expect that pairing in the nucleus is so fragile and it would only affect tunneling phenomena below the Coulomb barrier, where clear effects of nucleon-nucleon correlations were indeed observed experimentally [62, 175–177]. To the contrary, Magierski et al. [178–180] found unexpectedly large effects associated with the relative phase (or gauge angles) of colliding superfluid nuclei on the reaction outcomes, such as TKE and fusion cross section. In this context, experimental fusion cross sections were analyzed, which indicates fusion barrier width may be increased by the superfluid effect [181]. In the work by Magierski et al., a local formulation of superfluid TDDFT, known as time-dependent superfluid local density approximation (TDSLDA) (see e.g., [182–185]), was employed, which is formally similar to the time-dependent Hartree-Fock-Bogoliubov (TDHFB) approach [141]. The TDSLDA has been extensively developed by Bulgac and his coworkers, which was successfully applied to the unitary Fermi gas [186–191], finite nuclei [178, 192–196], as well as interiors of neutron stars [197]. Hashimoto has developed a Gogny TDHFB code that works with a 2D harmonic oscillator and 1D-Lagrange-mesh hybrid basis [198], which could be applied for head-on collisions [199, 200]. Qualitatively, the same effects of the relative phase as reported in Magierski et al. [178] were observed with their code. We have just entered a new era in which dynamic effects of pairing can be fully incorporated with microscopic nuclear dynamics studies. With the aforementioned sophisticated approaches with the usage of top-tier supercomputers, the dynamic effects of pairing in MNT and QF processes will be unveiled in the near future.

4.5. Remarks on Other Theoretical Approaches

There are many models that have been developed so far for the study of MNT, such as the semi-classical model, like GRAZING [201–205]^{4,7} or CWKB [116–118, 206], the dinuclear system (DNS) model [207–221], the improved quantum molecular dynamics (ImQMD) model [222–225], and the Langevin model [128–137, 226, 227]. There are pros and cons in each of those approaches. For instance, on the one hand, the semi-classical model can describe successive transfer processes at peripheral collisions, while on the other hand it misses deep-inelastic components at small impact parameters. In contrast, the DNS model is capable of describing the dynamic evolution of a composite system, according to a master equation with a potential energy (including shell effects) for mass asymmetry and deformation of the subsystem, that gives rise to a probability distribution for massive nucleon transfer as well as fusion. By construction, however, the latter model assumes a formation of a “di-nucleus” in a potential pocket, and thus it misses (quasi)elastic components at large impact parameters. A simple addition of those model predictions, DNS+GRAZING, was considered in the literature [228, 229]. The ImQMD model

is applicable for both peripheral and damped collisions. By taking into account stochastic nucleon-nucleon collisions, it provides distributions of observables accumulated in a number of simulations. To the author’s knowledge, however, the spin-orbit interaction has been neglected in ImQMD, which prevents a proper description of shell effects in MNT and QF processes. The elaborated Langevin model [226] may be promising in describing complex processes of MNT, QF and fusion, in a unified way. The model contains, however, various parameters that have to be tuned carefully to reproduce available experimental data [226].

A possible future task is to non-empirically determine model ingredients based on microscopic theories: potential energy surface with respect to an arbitrary set of variables, drift, diffusion, and friction coefficients, etc., can be derived from the microscopic approaches, which may lead to, e.g., kind of a TDHF+Langevin approach.

5. SUMMARY AND PROSPECT

Production of neutron-rich heavy nuclei is listed as one of the high-priority subjects of nuclear science today. In this contribution, recent advances in the MNT study, especially those based on microscopic dynamic theories, have been briefly reviewed. The traditional TDHF approach has shown remarkable successes in describing a variety of phenomena in nuclear systems. It has been shown that TDHF is capable of describing the main reaction channels, not only MNT processes in peripheral collisions but also deep-inelastic and QF processes in damped collisions, positioning it as a good starting point for building a fully microscopic theory for low-energy heavy-ion reactions. Moreover, the inclusion of one-body fluctuations significantly improves quantitative agreements of the width of fragment mass distributions with experimental data. With the use of top-tier supercomputers, studies of dynamic effects of pairing have just been started. One of the great advantages of microscopic approaches is their non-empirical nature—one may explore novel pathways toward the dreamed-of production of yet-unknown neutron-rich nuclei. Another possible future direction is to use microscopic approaches to determine phenomenological parameters in other theoretical models. It may be the time to combine knowledge obtained with different approaches to construct the most reliable framework to lead future experiments to the successful production of extremely neutron-rich nuclei, the farthest from the continent of stability.

AUTHOR CONTRIBUTIONS

The author confirms being the sole contributor of this work and has approved it for publication.

ACKNOWLEDGMENTS

The author acknowledges P. D. Stevenson, J. A. Maruhn, C. Simenel, Lu Guo, and A. S. Umar for careful reading of the manuscript and for providing valuable comments on this article.

⁷<http://personalpages.to.infn.it/~nanni/grazing/>

REFERENCES

1. Otsuka T, Gade A, Sorlin O, Suzuki T, Utsuno Y. Evolution of nuclear structure in exotic nuclei and nuclear forces. *arXiv:1805.06501 [nucl-th]*. (2018).
2. Cejnar P, Jolie J, Casten RF. Quantum phase transitions in the shapes of atomic nuclei. *Rev Mod Phys.* (2010) **82**:2155. doi: 10.1103/RevModPhys.82.2155
3. Heyde K, Wood JL. Shape coexistence in atomic nuclei. *Rev Mod Phys.* (2011) **83**:1467. doi: 10.1103/RevModPhys.83.1467
4. Cowan JJ, Sneden C, Lawler JE, Aprahamian A, Wiescher M, Langanke K, et al. Making the heaviest elements in the Universe: a review of the rapid neutron capture process. *arXiv:1901.01410 [astro-ph.HE]*. (2019).
5. Abbott BP, Abbott R, Abbott TD, Acernese F, Ackley K, Adams C. GW170817: observation of gravitational waves from a binary neutron star inspiral. *Phys Rev Lett.* (2017) **119**:161101. doi: 10.1103/PhysRevLett.119.161101
6. Abbott BP, Abbott R, Abbott TD, Acernese F, Ackley K, Adams C. Gravitational waves and gamma-rays from a binary neutron star merger: GW170817 and GRB 170817A. *Astrophys J Lett.* (2017) **848**:L13. doi: 10.3847/2041-8213/aa920c
7. Abbott BP, Abbott R, Abbott TD, Acernese F, Ackley K, Adams C. *Estimating contribution of dynamical ejecta in the Kilonova associated with GW170817.* *Astrophys J Lett.* (2017) **850**:L39. doi: 10.3847/2041-8213/aa9478
8. Hofmann S, Münzenberg G. The discovery of the heaviest elements. *Rev Mod Phys.* (2000) **72**:733. doi: 10.1103/RevModPhys.72.733
9. Oganessian YT, Sobiczewski A, Ter-Akopian GM. Superheavy nuclei: from predictions to discovery. *Phys Scr.* (2017) **92**:023003. doi: 10.1088/1402-4896/aa53c1
10. Sakurai H. Nuclear physics with RI Beam Factory. *Front Phys.* (2018) **13**:132111. doi: 10.1007/s11467-018-0849-0
11. Shin T, Kang BH, Kim GD, Kim YJ, Kwon YK, Park YH, et al. Rare isotope production and experimental systems of RAON. *New Phys.* (2016) **66**:1500. doi: 10.3938/NPSM.66.1500
12. Gales S. SPIRAL2 at GANIL: a world leading ISOL facility at the dawn of the next decade. *Prog Part Nucl Phys.* (2007) **59**:22. doi: 10.1016/j.pnpnp.2006.12.021
13. Abel EP, Avilov M, Ayres V, Birnbaum E, Bollen G, Bonito G. Isotope harvesting at FRIB: additional opportunities for scientific discovery. *arXiv:1812.03984 [nucl-th]*. (2018).
14. Dirac PAM. Note on exchange phenomena in the thomas atom. *Proc Camb Philos Soc.* (1930) **26**:376. doi: 10.1017/S0305004100016108
15. Bonche P, Koonin S, Negele JW. One-dimensional nuclear dynamics in the time-dependent Hartree-Fock approximation. *Phys Rev C.* (1976) **13**:1226. doi: 10.1103/PhysRevC.13.1226
16. Cusson RY, Smith RK, Maruhn JA. Time-dependent Hartree-Fock calculation of the reaction $^{16}\text{O}+^{16}\text{O}$ in three dimensions. *Phys Rev Lett.* (1976) **36**:1166. doi: 10.1103/PhysRevLett.36.1166
17. Koonin SE, Davies KTR, Maruhn-Rezwani V, Feldmeier H, Krieger SJ, Negele JW. Time-dependent Hartree-Fock calculations for $^{16}\text{O}+^{16}\text{O}$ and $^{40}\text{Ca}+^{40}\text{Ca}$ reactions. *Phys Rev C.* (1977) **15**:1359. doi: 10.1103/PhysRevC.15.1359
18. Davies KTR, Maruhn-Rezwani V, Koonin SE, Negele JW. Test of the time-dependent mean-field theory in Kr-induced strongly damped collisions. *Phys Rev Lett.* (1978) **41**:632. doi: 10.1103/PhysRevLett.41.632
19. Flocard H, Koonin SE, Weiss MS. Three-dimensional time-dependent Hartree-Fock calculations: application to $^{16}\text{O}+^{16}\text{O}$ collisions. *Phys Rev C.* (1978) **17**:1682. doi: 10.1103/PhysRevC.17.1682
20. Krieger SJ, Davies KTR. Time-dependent Hartree-Fock calculations of fusion cross sections for the reactions $^{16}\text{O}+^{16}\text{O}$ and $^{40}\text{Ca}+^{40}\text{Ca}$. *Phys Rev C.* (1978) **18**:2567. doi: 10.1103/PhysRevC.18.2567
21. Davies KTR, Sandhya Devi KR, Strayer MR. Time-dependent Hartree-Fock calculations of $^{86}\text{Kr}+^{139}\text{La}$ at $E_{\text{lab}}=505, 610,$ and 710 MeV. *Phys Rev C.* (1979) **20**:1372. doi: 10.1103/PhysRevC.20.1372
22. Negele JW. The mean-field theory of nuclear structure and dynamics. *Rev Mod Phys.* (1982) **54**:913. doi: 10.1103/RevModPhys.54.913
23. Simenel C, Umar AS. Heavy-ion collisions and fission dynamics with the time-dependent Hartree-Fock theory and its extensions. *Prog Part Nucl Phys.* (2018) **103**:19. doi: 10.1016/j.pnpnp.2018.07.002
24. Engel YM, Brink DM, Goeke K, Krieger SJ, Vautherin D. Time-dependent hartree-fock theory with Skyrme's interaction. *Nucl Phys.* (1975) **A249**:215. doi: 10.1016/0375-9474(75)90184-0
25. Dobaczewski J, Dudek J. Time-odd components in the mean field of rotating superdeformed nuclei. *Phys Rev C.* (1995) **52**:1827. doi: 10.1103/PhysRevC.52.1827
26. Stevenson PD, Barton MC. Low-energy heavy-ion reactions and the Skyrme effective interaction. *Prog Part Nucl Phys.* (2019) **104**:142. doi: 10.1016/j.pnpnp.2018.09.002
27. Hohenberg P, Kohn W. Inhomogeneous electron gas. *Phys Rev.* (1964) **136**:B864. doi: 10.1103/PhysRev.136.B864
28. Kohn W, Sham LJ. Self-consistent equations including exchange and correlation effects. *Phys Rev.* (1965) **140**:A1133. doi: 10.1103/PhysRev.140.A1133
29. Runge E, Gross EKH. Density-functional theory for time-dependent systems. *Phys Rev Lett.* (1984) **52**:997. doi: 10.1103/PhysRevLett.52.997
30. van Leeuwen R. Mapping from densities to potentials in time-dependent density-functional theory. *Phys Rev Lett.* (1999) **82**:3863. doi: 10.1103/PhysRevLett.82.3863
31. Nakatsukasa T, Matsuyanagi K, Matsuo M, Yabana K. Time-dependent density-functional description of nuclear dynamics. *Rev Mod Phys.* (2016) **88**:045004. doi: 10.1103/RevModPhys.88.045004
32. Bender M, Heenen PH, Reinhard PG. Self-consistent mean-field models for nuclear structure. *Rev Mod Phys.* (2003) **75**:121. doi: 10.1103/RevModPhys.75.121
33. Kortelainen M, McDonnell J, Nazarewicz W, Reinhard PG, Sarich J, Schunck N, et al. Nuclear energy density optimization: large deformations. *Phys Rev C.* (2012) **85**:024304. doi: 10.1103/PhysRevC.85.024304
34. Kortelainen M, McDonnell J, Nazarewicz W, Olsen E, Reinhard PG, Sarich J, et al. Nuclear energy density optimization: shell structure. *Phys Rev C.* (2014) **89**:054314. doi: 10.1103/PhysRevC.89.054314
35. Bulgac A, Forbes MM, Jin S, Perez RN, Schunck N. Minimal nuclear energy density functional. *Phys Rev C.* (2018) **97**:044313. doi: 10.1103/PhysRevC.97.044313
36. Blocki J, Boneh Y, Nix JR, Randrup J, Robel M, Sierk AJ, et al. One-body dissipation and the super-viscosity of nuclei. *Ann Phys.* (1978) **113**:330. doi: 10.1016/0003-4916(78)90208-7
37. Nakatsukasa T, Yabana K. Linear response theory in the continuum for deformed nuclei: Green's function vs time-dependent Hartree-Fock with the absorbing boundary condition. *Phys Rev C.* (2005) **71**:024301. doi: 10.1103/PhysRevC.71.024301
38. Umar AS, Oberacker VE. Time-dependent response calculations of nuclear resonances. *Phys Rev C.* (2005) **71**:034314. doi: 10.1103/PhysRevC.71.034314
39. Maruhn JA, Reinhard PG, Stevenson PD, Rikovsky Stone J, Strayer MR. Dipole giant resonances in deformed heavy nuclei. *Phys Rev C.* (2005) **71**:064328. doi: 10.1103/PhysRevC.71.064328
40. Ebata S, Nakatsukasa T, Inakura T, Yoshida K, Hashimoto Y, Yabana K. Canonical-basis time-dependent Hartree-Fock-Bogoliubov theory and linear-response calculations. *Phys Rev C.* (2010) **82**:034306. doi: 10.1103/PhysRevC.82.034306
41. Fracasso S, Suckling EB, Stevenson PD. Unrestricted Skyrme-tensor time-dependent Hartree-Fock model and its application to the nuclear response from spherical to triaxial nuclei. *Phys Rev C.* (2012) **86**:044303. doi: 10.1103/PhysRevC.86.044303
42. Pardi CI, Stevenson PD. Continuum time-dependent Hartree-Fock method for giant resonances in spherical nuclei. *Phys Rev C.* (2013) **87**:014330. doi: 10.1103/PhysRevC.87.014330
43. Scamps G, Lacroix D. Systematics of isovector and isoscalar giant quadrupole resonances in normal and superfluid spherical nuclei. *Phys Rev C.* (2013) **88**:044310. doi: 10.1103/PhysRevC.88.044310
44. Scamps G, Lacroix D. Systematic study of isovector and isoscalar giant quadrupole resonances in normal and superfluid deformed nuclei. *Phys Rev C.* (2014) **89**:034314. doi: 10.1103/PhysRevC.89.034314

45. Ebata S, Nakatsukasa T, Inakura T. Systematic investigation of low-lying dipole modes using the canonical-basis time-dependent Hartree-Fock-Bogoliubov theory. *Phys Rev C*. (2014) **90**:024303. doi: 10.1103/PhysRevC.90.024303
46. Ebata S, Nakatsukasa T, Inakura T. Systematic investigation of low-lying dipole modes using the canonical-basis time-dependent Hartree-Fock-Bogoliubov theory. *Phys Rev C*. (2015) **92**:049902. doi: 10.1103/PhysRevC.92.049902
47. Simenel C, Umar AS. Formation and dynamics of fission fragments. *Phys Rev C*. (2014) **89**:031601(R). doi: 10.1103/PhysRevC.89.031601
48. Scamps G, Simenel C, Lacroix D. Superfluid dynamics of ^{258}Fm fission. *Phys Rev C*. (2015) **92**:011602(R). doi: 10.1103/PhysRevC.92.011602
49. Tanimura Y, Lacroix D, Scamps G. Collective aspects deduced from time-dependent microscopic mean-field with pairing: application to the fission process. *Phys Rev C*. (2015) **92**:034601. doi: 10.1103/PhysRevC.92.034601
50. Goddard P, Stevenson P, Rios A. Fission dynamics within time-dependent Hartree-Fock: deformation-induced fission. *Phys Rev C*. (2015) **92**:054610. doi: 10.1103/PhysRevC.92.054610
51. Goddard P, Stevenson P, Rios A. Fission dynamics within time-dependent Hartree-Fock. II. Boost-induced fission. *Phys Rev C*. (2016) **93**:014620. doi: 10.1103/PhysRevC.93.014620
52. Scamps G, Simenel C. Impact of pear-shaped fission fragments on mass-asymmetric fission in actinides. *Nature*. (2018) **564**:382. doi: 10.1038/s41586-018-0780-0
53. Umar AS, Maruhn JA, Itagaki N, Oberacker VE. Microscopic study of the triple- α reaction. *Phys Rev Lett*. (2010) **104**:212503. doi: 10.1103/PhysRevLett.104.212503
54. Ichikawa T, Matsuyanagi K, Maruhn JA, Itagaki N. Pure collective precession motion of a high-spin torus isomer. *Phys Rev C*. (2014) **89**:011305(R). doi: 10.1103/PhysRevC.89.011305
55. Ichikawa T, Matsuyanagi K, Maruhn JA, Itagaki N. High-spin torus isomers and their precession motions. *Phys Rev C*. (2014) **90**:034314. doi: 10.1103/PhysRevC.90.034314
56. Schuetrumpf B, Klatt MA, Iida K, Maruhn JA, Mecke K, Reinhard PG. Time-dependent Hartree-Fock approach to nuclear "pasta" at finite temperature. *Phys Rev C*. (2013) **87**:055805. doi: 10.1103/PhysRevC.87.055805
57. Schuetrumpf B, Iida K, Maruhn JA, Reinhard PG. Nuclear "pasta matter" for different proton fractions. *Phys Rev C*. (2014) **90**:055802. doi: 10.1103/PhysRevC.90.055802
58. Schuetrumpf B, Klatt MA, Iida K, Schröder-Turk GE, Maruhn JA, Mecke K, et al. Appearance of the single gyroid network phase in "nuclear pasta" matter. *Phys Rev C*. (2015) **91**:025801. doi: 10.1103/PhysRevC.91.025801
59. Schuetrumpf B, Nazarewicz W. Twist-averaged boundary conditions for nuclear pasta Hartree-Fock calculations. *Phys Rev C*. (2015) **92**:045806. doi: 10.1103/PhysRevC.92.045806
60. Umar AS, Oberacker VE, Maruhn JA. Neutron transfer dynamics and doorway to fusion in time-dependent Hartree-Fock theory. *Eur Phys J A*. (2008) **37**:245. doi: 10.1140/epja/i2008-10614-6
61. Iwata Y, Otsuka T, Maruhn JA, Itagaki N. Suppression of charge equilibration leading to the synthesis of exotic nuclei. *Phys Rev Lett*. (2010) **104**:252501. doi: 10.1103/PhysRevLett.104.252501
62. Evers M, Dasgupta M, Hinde DJ, Luong DH, Rafiei R, du Rietz R, et al. Cluster transfer in the reaction $^{16}\text{O}+^{208}\text{Pb}$ at energies well below the fusion barrier: a possible doorway to energy dissipation. *Phys Rev C*. (2011) **84**:054614. doi: 10.1103/PhysRevC.84.054614
63. Umar AS, Simenel C, Ye W. Transport properties of isospin asymmetric nuclear matter using the time-dependent Hartree-Fock method. *Phys Rev C*. (2017) **96**:024625. doi: 10.1103/PhysRevC.96.024625
64. Simenel C. Particle transfer reactions with the time-dependent Hartree-Fock theory using a particle number projection technique. *Phys Rev Lett*. (2010) **105**:192701. doi: 10.1103/PhysRevLett.105.192701
65. Sekizawa K, Yabana K. Time-dependent Hartree-Fock calculations for multinucleon transfer processes in $^{40,48}\text{Ca}+^{124}\text{Sn}$, $^{40}\text{Ca}+^{208}\text{Pb}$, and $^{58}\text{Ni}+^{208}\text{Pb}$ reactions. *Phys Rev C*. (2013) **88**:014614. doi: 10.1103/PhysRevC.88.014614
66. Sekizawa K, Yabana K. Time-dependent Hartree-Fock calculations for multinucleon transfer processes in $^{40,48}\text{Ca}+^{124}\text{Sn}$, $^{40}\text{Ca}+^{208}\text{Pb}$, and $^{58}\text{Ni}+^{208}\text{Pb}$ reactions. *Phys Rev C*. (2016) **93**:029902(E). doi: 10.1103/PhysRevC.93.029902
67. Sekizawa K, Yabana K. Particle-number projection method in time-dependent Hartree-Fock theory. *Phys Rev C*. (2014) **90**:064614. doi: 10.1103/PhysRevC.90.064614
68. Sonika, Roy BJ, Parmar A, Pal UK, Kumawat H, Jha V, et al. Multinucleon transfer study in $^{206}\text{Pb}(^{18}\text{O}, x)$ at energies above the Coulomb barrier. *Phys Rev C*. (2015) **92**:024603. doi: 10.1103/PhysRevC.92.024603
69. Sekizawa K, Yabana K. Multinucleon transfer reaction in time-dependent Hartree-Fock theory. *arXiv:1511.08322 [nucl-th]*. (2015).
70. Sekizawa K, Yabana K. Time-dependent Hartree-Fock calculations for multinucleon transfer and quasifission processes in the $^{64}\text{Ni}+^{238}\text{U}$ reaction. *Phys Rev C*. (2016) **93**:054616. doi: 10.1103/PhysRevC.93.054616
71. Sekizawa K, Heinz S. Quasifission dynamics and stability of superheavy systems. *Acta Phys Pol B Proc Suppl*. (2017) **10**:225. doi: 10.5506/APhysPolBSupp.10.225
72. Sekizawa K. Microscopic description of production cross sections including deexcitation effects. *Phys Rev C*. (2017) **96**:014615. doi: 10.1103/PhysRevC.96.014615
73. Sekizawa K. Enhanced nucleon transfer in tip collisions of $^{124}\text{Sn}+^{238}\text{U}$. *Phys Rev C*. (2017) **96**:041601(R). doi: 10.1103/PhysRevC.96.041601
74. Roy BJ, Sawant Y, Patwari P, Santra S, Pal A, Kundu A, et al. Deep-inelastic multinucleon transfer processes in the $^{16}\text{O}+^{27}\text{Al}$ reaction. *Phys Rev C*. (2018) **97**:034603. doi: 10.1103/PhysRevC.97.034603
75. Golabek C, Simenel C. Collision dynamics of two ^{238}U atomic nuclei. *Phys Rev Lett*. (2009) **103**:042701. doi: 10.1103/PhysRevLett.103.042701
76. Kedziora DJ, Simenel C. New inverse quasifission mechanism to produce neutron-rich transfermium nuclei. *Phys Rev C*. (2010) **81**:044613. doi: 10.1103/PhysRevC.81.044613
77. Simenel C, Hinde DJ, du Rietz R, Dasgupta M, Evers M, Lin CJ, et al. Influence of entrance-channel magicity and isospin on quasi-fission. *Phys Lett*. (2012) **B710**:607. doi: 10.1016/j.physletb.2012.03.063
78. Wakhle A, Simenel C, Hinde DJ, Dasgupta M, Evers M, Luong DH, et al. Interplay between quantum shells and orientation in quasifission. *Phys Rev Lett*. (2014) **113**:182502. doi: 10.1103/PhysRevLett.113.182502
79. Oberacker VE, Umar AS, Simenel C. Dissipative dynamics in quasifission. *Phys Rev C*. (2014) **90**:054605. doi: 10.1103/PhysRevC.90.054605
80. Umar AS, Oberacker VE. Time-dependent HF approach to SHE dynamics. *Nucl Phys*. (2015) **A944**:238. doi: 10.1016/j.nuclphysa.2015.02.011
81. Hammerton K, Kohley Z, Hinde DJ, Dasgupta M, Wakhle A, Williams E, et al. Reduced quasifission competition in fusion reactions forming neutron-rich heavy elements. *Phys Rev C*. (2015) **91**:041602(R). doi: 10.1103/PhysRevC.91.041602
82. Washiyama K. Microscopic analysis of fusion hindrance in heavy nuclear systems. *Phys Rev C*. (2015) **91**:064607. doi: 10.1103/PhysRevC.91.064607
83. Umar AS, Oberacker VE, Simenel C. Shape evolution and collective dynamics of quasifission in the time-dependent Hartree-Fock approach. *Phys Rev C*. (2015) **92**:024621. doi: 10.1103/PhysRevC.92.024621
84. Umar AS, Oberacker VE, Simenel C. Fusion and quasifission dynamics in the reactions $^{48}\text{Ca}+^{249}\text{Bk}$ and $^{50}\text{Ti}+^{249}\text{Bk}$ using a time-dependent Hartree-Fock approach. *Phys Rev C*. (2016) **94**:024605. doi: 10.1103/PhysRevC.94.024605
85. Yu C, Guo L. Angular momentum dependence of quasifission dynamics in the reaction $^{48}\text{Ca}+^{244}\text{Pu}$. *Sci China Phys Mech Astron*. (2017) **60**:092011. doi: 10.1007/s11433-017-9063-3
86. Guo L, Shen C, Yu C, Wu Z. Isotopic trends of quasifission and fusion-fission in the reactions $^{48}\text{Ca}+^{239,244}\text{Pu}$. *Phys Rev C*. (2018) **98**:064609. doi: 10.1103/PhysRevC.98.064609
87. Zheng H, Burrello S, Colonna M, Lacroix D, Scamps G. Connecting the nuclear equation of state to the interplay between fusion and quasifission processes in low-energy nuclear reactions. *Phys Rev C*. (2018) **98**:024622. doi: 10.1103/PhysRevC.98.024622
88. Umar AS, Oberacker VE. Heavy-ion interaction potential deduced from density-constrained time-dependent Hartree-Fock calculation. *Phys Rev C*. (2006) **74**:021601(R). doi: 10.1103/PhysRevC.74.021601

89. Umar AS, Oberacker VE, Maruhn JA, Reinhard PG. Microscopic calculation of precompound excitation energies for heavy-ion collisions. *Phys Rev C*. **80**:041601(R) (2009).
90. Umar AS, Oberacker VE, Maruhn JA, Reinhard PG. Entrance channel dynamics of hot and cold fusion reactions leading to superheavy elements. *Phys Rev C*. (2010) **81**:064607. doi: 10.1103/PhysRevC.81.064607
91. Oberacker VE, Umar AS, Maruhn JA, Reinhard PG. Microscopic study of the $^{132,124}\text{Sn}+^{96}\text{Zr}$ reactions: dynamic excitation energy, energy-dependent heavy-ion potential, and capture cross section. *Phys Rev C*. (2010) **82**:034603. doi: 10.1103/PhysRevC.82.034603
92. Umar AS, Oberacker VE, Maruhn JA, Reinhard PG, Microscopic composition of ion-ion interaction potentials. *Phys Rev C*. (2012) **85**:017602. doi: 10.1103/PhysRevC.85.017602
93. Oberacker VE, Umar AS, Maruhn JA, Reinhard PG. Dynamic microscopic study of pre-equilibrium giant resonance excitation and fusion in the reactions $^{132}\text{Sn}+^{48}\text{Ca}$ and $^{124}\text{Sn}+^{40}\text{Ca}$. *Phys Rev C*. (2012) **85**:034609. doi: 10.1103/PhysRevC.85.034609
94. Keser R, Umar AS, Oberacker VE. Microscopic study of Ca + Ca fusion. *Phys Rev C*. (2012) **85**:044606. doi: 10.1103/PhysRevC.85.044606
95. Oberacker VE, Umar AS. Microscopic analysis of sub-barrier fusion enhancement in $^{132}\text{Sn}+^{40}\text{Ca}$ versus $^{132}\text{Sn}+^{48}\text{Ca}$. *Phys Rev C*. (2013) **87**:034611. doi: 10.1103/PhysRevC.87.034611
96. Simenel C, Keser R, Umar AS, Oberacker VE. Microscopic study of $^{16}\text{O}+^{16}\text{O}$ fusion. *Phys Rev C*. (2013) **88**:024617. doi: 10.1103/PhysRevC.88.024617
97. Umar AS, Simenel C, Oberacker VE. Energy dependence of potential barriers and its effect on fusion cross sections. *Phys Rev C*. **89**:034611 (2014). doi: 10.1103/PhysRevC.89.034611
98. Steinbach TK, Vadas J, Schmidt J, Haycraft C, Hudan S, deSouza RT, et al. Sub-barrier enhancement of fusion as compared to a microscopic method in $^{18}\text{O}+^{12}\text{C}$. *Phys Rev C*. (2014) **90**:041603(R). doi: 10.1103/PhysRevC.90.041603
99. Reinhard PG, Umar AS, Stevenson PD, Piekarewicz J, Oberacker VE, Maruhn JA. Sensitivity of the fusion cross section to the density dependence of the symmetry energy. *Phys Rev C*. (2016) **93**:044618. doi: 10.1103/PhysRevC.93.044618
100. Godbey K, Umar AS, Simenel C. Dependence of fusion on isospin dynamics. *Phys Rev C*. (2017) **95**:011601(R). doi: 10.1103/PhysRevC.95.011601
101. Guo L, Godbey K, Umar AS. Influence of the tensor force on the microscopic heavy-ion interaction potential. *Phys Rev C*. (2018) **98**:064607. doi: 10.1103/PhysRevC.98.064607
102. Simenel C, Chomaz Ph, de France G. Fusion process studied with a preequilibrium giant dipole resonance in time-dependent Hartree-Fock theory. *Phys Rev C*. (2007) **76**:024609. doi: 10.1103/PhysRevC.76.024609
103. Bourgin D, Simenel C, Courtin S, Haas F. Microscopic study of $^{40}\text{Ca}+^{58,64}\text{Ni}$ fusion reactions. *Phys Rev C*. (2016) **93**:034604. doi: 10.1103/PhysRevC.93.034604
104. Vo-Phuoc K, Simenel C, Simpson EC. Dynamical effects in fusion with exotic nuclei. *Phys Rev C*. (2016) **94**:024612. doi: 10.1103/PhysRevC.94.024612
105. Simenel C. Nuclear quantum many-body dynamics. *Eur Phys J A*. (2012) **48**:152. doi: 10.1140/epja/i2012-12152-0
106. Nakatsukasa T. Density functional approaches to collective phenomena in nuclei: time-dependent density functional theory for perturbative and non-perturbative nuclear dynamics. *Prog Theor Exp Phys*. (2012) **2012**:01A207. doi: 10.1093/ptep/pts016
107. Umar AS, Strayer MR, Reinhard PG. Resolution of the fusion window anomaly in heavy-ion collisions. *Phys Rev Lett*. (1986) **56**:2793. doi: 10.1103/PhysRevLett.56.2793
108. Reinhard PG, Umar AS, Davies KTR, Strayer MR, Lee SJ. Dissipation and forces in time-dependent Hartree-Fock calculations. *Phys Rev C*. (1988) **37**:1026. doi: 10.1103/PhysRevC.37.1026
109. Umar AS, Strayer MR, Reinhard PG, Davies KTR, Lee SJ. Spin-orbit force in time-dependent Hartree-Fock calculations of heavy-ion collisions. *Phys Rev C*. (1989) **40**:706. doi: 10.1103/PhysRevC.40.706
110. Dai GE, Guo L, Zhao EG, Zhou SG. Dissipation dynamics and spin-orbit force in time-dependent Hartree-Fock theory. *Phys Rev C*. (2014) **90**:044609. doi: 10.1103/PhysRevC.90.044609
111. Iwata Y, Maruhn JA. Enhanced spin-current tensor contribution in collision dynamics. *Phys Rev C*. (2011) **84**:014616. doi: 10.1103/PhysRevC.84.014616
112. Dai DF, Guo L, Zhao EG, Zhou SG. Effect of tensor force on dissipation dynamics in time-dependent Hartree-Fock theory. *Sci China Phys Mech Astron*. (2014) **57**:1618. doi: 10.1007/s11433-014-5536-8
113. Stevenson PD, Suckling EB, Fracasso S, Barton MC, Umar AS. Skyrme tensor force in heavy ion collisions. *Phys Rev C*. (2016) **93**:054617. doi: 10.1103/PhysRevC.93.054617
114. Guo L, Simenel C, Shi L, Yu C. The role of tensor force in heavy-ion fusion dynamics. *Phys Lett*. (2018) **B782**:401. doi: 10.1016/j.physletb.2018.05.066
115. Corradi L, He JH, Ackermann D, Stefanini AM, Pisent A, Beghini S et al. Multinucleon transfer reactions in $^{40}\text{Ca}+^{124}\text{Sn}$. *Phys Rev C*. (1996) **54**:201. doi: 10.1103/PhysRevC.54.201
116. Corradi L, Stefanini AM, He JH, Beghini S, Montagnoli G, Scarlassara F, et al. Evidence of complex degrees of freedom in multinucleon transfer reactions of $^{48}\text{Ca}+^{124}\text{Sn}$. *Phys Rev C*. (1997) **56**:938. doi: 10.1103/PhysRevC.56.938
117. Corradi L, Vinodkumar AM, Stefanini AM, Fioretto E, Prete G, Beghini S, et al. Light and heavy transfer products in $^{58}\text{Ni}+^{208}\text{Pb}$ at the Coulomb barrier. *Phys Rev C*. (2002) **66**:024606. doi: 10.1103/PhysRevC.66.024606
118. Szilner S, Corradi L, Pollarolo G, Beghini S, Behera BR, Fioretto E, et al. Multinucleon transfer processes in $^{40}\text{Ca}+^{208}\text{Pb}$. *Phys Rev C*. (2005) **71**:044610. doi: 10.1103/PhysRevC.71.044610
119. Charity RJ. GEMINI: a code to simulate the decay of a compound nucleus by a series of binary decays. In: *Joint ICTP-AIEA Advanced Workshop on Model Codes for Spallation Reactions*. Vienna: IAEA (2008). p. 139.
120. Corradi L, Stefanini AM, Lin CJ, Beghini S, Montagnoli G, Scarlassara F, et al. Multinucleon transfer processes in $^{64}\text{Ni}+^{238}\text{U}$. *Phys Rev C*. (1999) **59**:261. doi: 10.1103/PhysRevC.59.261
121. Kozulin EM, Knyazheva GN, Itkis IM, Itkis MG, Bogachev AA, Krupa L, et al. Investigation of the reaction $^{64}\text{Ni}+^{238}\text{U}$ being an option of synthesizing element 120. *Phys Lett B*. (2010) **686**:227. doi: 10.1016/j.physletb.2010.02.041
122. Jiang X, Wang N. Production mechanism of neutron-rich nuclei around $N = 126$ in the multi-nucleon transfer reaction $^{132}\text{Sn}+^{208}\text{Pb}$. *Chin Phys C*. (2018) **42**:104105. doi: 10.1088/1674-1137/42/10/104105
123. Töke JT, Bock R, Dai GX, Gobbi A, Gralla S, Hildenbrand KD, et al. Quasi-fission - The mass-drift mode in heavy-ion reactions. *Nucl Phys*. (1985) **A440**:327.
124. Bock R, Chu YT, Dakowski M, Gobbi A, Grosse E, Olmi A, et al. Dynamics of the fusion process. *Nucl Phys*. (1982) **A388**:334.
125. du Rietz R, Hinde DJ, Dasgupta M, Thomas RG, Gasques LR, Evers M, et al. Predominant time scales in fission processes in reactions of S, Ti, and Ni with W: zeptosecond versus attosecond. *Phys Rev Lett*. (2011) **106**:052701. doi: 10.1103/PhysRevLett.106.052701
126. du Rietz R, Williams E, Hinde DJ, Dasgupta M, Evers M, Lin CJ, et al. Mapping quasifission characteristics and timescales in heavy element formation reactions. *Phys Rev C*. (2013) **88**:054618. doi: 10.1103/PhysRevC.88.054618
127. Morjean M, Hinde DJ, Simenel C, Jeung DY, Airiau M, Cook KJ, et al. Evidence for the role of proton shell closure in quasifission reactions from X-Ray fluorescence of mass-identified fragments. *Phys Rev Lett*. (2017) **119**:222502. doi: 10.1103/PhysRevLett.119.222502
128. Zagrebaev V, Greiner W. Unified consideration of deep inelastic, quasi-fission and fusion-fission phenomena. *J Phys G Nucl Part Phys*. (2005) **31**:825. doi: 10.1088/0954-3899/31/7/024
129. Zagrebaev VI, Oganessian YT, Itkis MG, Greiner W. Superheavy nuclei and quasi-atoms produced in collisions of transuranium ions. *Phys Rev C*. (2006) **73**:031602(R). doi: 10.1103/PhysRevC.73.031602
130. Zagrebaev V, Greiner W. Shell effects in damped collisions: a new way to superheavies. *J Phys G Nucl Part Phys*. (2007) **34**:2265. doi: 10.1088/0954-3899/34/11/004

131. Zagrebaev V, Greiner W. Low-energy collisions of heavy nuclei: dynamics of sticking, mass transfer and fusion. *J Phys G Nucl Part Phys.* (2007) **34**:1. doi: 10.1088/0954-3899/34/1/001
132. Zagrebaev V, Greiner W. Production of new heavy isotopes in low-energy multinucleon transfer reactions. *Phys Rev Lett.* (2008) **101**:122701. doi: 10.1103/PhysRevLett.101.122701
133. Zagrebaev V, Greiner W. Synthesis of superheavy nuclei: a search for new production reactions. *Phys Rev C.* (2008) **78**:034610. doi: 10.1103/PhysRevC.78.034610
134. Zagrebaev VI, Greiner W. Production of heavy and superheavy neutron-rich nuclei in transfer reactions. *Phys Rev C.* (2011) **83**:044618. doi: 10.1103/PhysRevC.83.044618
135. Zagrebaev VI, Greiner W. Production of heavy trans-target nuclei in multinucleon transfer reactions. *Phys Rev C.* (2013) **87**:034608. doi: 10.1103/PhysRevC.87.034608
136. Zagrebaev VI, Fornal B, Leoni S, Greiner W. Formation of light exotic nuclei in low-energy multinucleon transfer reactions. *Phys Rev C.* (2014) **89**:054608. doi: 10.1103/PhysRevC.89.054608
137. Zagrebaev VI, Greiner W. Cross sections for the production of superheavy nuclei. *Nucl Phys.* (2015) **A944**:257. doi: 10.1016/j.nuclphysa.2015.02.010
138. Ayik S, Yilmaz B, Yilmaz O. Multinucleon exchange in quasifission reactions. *Phys Rev C.* (2015) **92**:064615. doi: 10.1103/PhysRevC.92.064615
139. Ayik S, Yilmaz B, Yilmaz O, Umar AS, Turan G. Multi-nucleon transfer in central collisions of $^{238}\text{U}+^{238}\text{U}$. *Phys Rev C.* (2017) **96**:024611. doi: 10.1103/PhysRevC.96.024611
140. Ayik S, Yilmaz B, Yilmaz O, Umar AS. Quantal diffusion description of multinucleon transfers in heavy-ion collisions. *Phys Rev C.* (2018) **97**:054618. doi: 10.1103/PhysRevC.97.054618
141. Ring P, Schuck P. *The Nuclear Many-Body Problem.* Berlin; Heidelberg: Springer-Verlag (1980). doi: 10.1007/978-3-642-61852-9
142. Washiyama K, Lacroix D. Energy dependence of the nucleus-nucleus potential close to the Coulomb barrier. *Phys Rev C.* (2008) **78**:024610. doi: 10.1103/PhysRevC.78.024610
143. Washiyama K, Lacroix D, Ayik S. One-body energy dissipation in fusion reactions from mean-field theory. *Phys Rev C.* (2009) **79**:024609. doi: 10.1103/PhysRevC.79.024609
144. Simenel C, Umar AS, Godbey K, Dasgupta M, Hinde DJ. How the Pauli exclusion principle affects fusion of atomic nuclei. *Phys Rev C.* (2017) **95**:031601(R). doi: 10.1103/PhysRevC.95.031601
145. Hagino K, Rowley N, Kruppa AT. A program for coupled-channels calculations with all order couplings for heavy-ion fusion reactions. *Comput Phys Commun.* (1999) **123**:143. doi: 10.1016/S0010-4655(99)00243-X
146. Hagino K, Takigawa N. Subbarrier fusion reactions and many-particle quantum tunneling. *Prog Theor Phys.* (2012) **128**:1061. doi: 10.1143/PTP.128.1061
147. Simenel C, Dasgupta M, Hinde DJ, Williams E. Microscopic approach to coupled-channels effects on fusion. *Phys Rev C.* (2013) **88**:064604. doi: 10.1103/PhysRevC.88.064604
148. Balian R, Vénéroni M. Time-dependent variational principle for predicting the expectation value of an observable. *Phys Rev Lett.* (1981) **47**:1353. doi: 10.1103/PhysRevLett.47.1353
149. Balian R, Vénéroni M. Time-dependent variational principle for predicting the expectation value of an observable. *Phys Rev Lett.* (1981) **47**:1765. doi: 10.1103/PhysRevLett.47.1765.3
150. Dasso CH, Dossing T, Pauli HC. On the mass distribution in Time-Dependent Hartree-Fock calculations of heavy-ion collisions. *Z Phys A.* (1979) **289**:395. doi: 10.1007/BF01409391
151. Broomfield JMA, Stevenson PD. Mass distributions beyond TDHF. *AIP Conf Proc.* (2009) **1098**:133. doi: 10.1063/1.3108767
152. Simenel C. Particle-number fluctuations and correlations in transfer reactions obtained using the Balian-Vénéroni variational principle. *Phys Rev Lett.* (2011) **106**:112502. doi: 10.1103/PhysRevLett.106.112502
153. Williams E, Sekizawa K, Hinde DJ, Simenel C, Dasgupta M, Carter IP, et al. Exploring zeptosecond quantum equilibration dynamics: from deep-inelastic to fusion-fission outcomes in $^{58}\text{Ni}+^{60}\text{Ni}$ reactions. *Phys Rev Lett.* (2018) **120**:022501. doi: 10.1103/PhysRevLett.120.022501
154. Ayik S. A stochastic mean-field approach for nuclear dynamics. *Phys Rev Lett.* (2008) **B658**:174. doi: 10.1016/j.physletb.2007.09.072
155. Lacroix D, Ayik S. Stochastic quantum dynamics beyond mean field. *Eur Phys J A.* (2014) **50**:95. doi: 10.1140/epja/i2014-14095-8
156. Shun-jin W, Cassing W. Explicit treatment of N-body correlations within a density-matrix formalism. *Ann Phys (NY).* (1985) **159**:328. doi: 10.1016/0003-4916(85)90116-2
157. Gong M, Tohyama M. Application of a time-dependent density-matrix formalism. *Z Phys A.* (1990) **335**:153. doi: 10.1007/BF01294470
158. Tohyama M, Umar AS. Fusion window problem in time-dependent Hartree-Fock theory revisited. *Phys Rev C.* (2002) **65**:037601. doi: 10.1103/PhysRevC.65.037601
159. Tohyama M, Umar AS. Two-body dissipation effects on the synthesis of superheavy elements. *Phys Rev C.* (2016) **93**:034607. doi: 10.1103/PhysRevC.93.034607
160. Wen K, Barton MC, Rios A, Stevenson PD. Two-body dissipation effect in nuclear fusion reactions. *Phys Rev C.* (2018) **98**:014603. doi: 10.1103/PhysRevC.98.014603
161. Ayik S, Washiyama K, Lacroix D. Fluctuation and dissipation dynamics in fusion reactions from a stochastic mean-field approach. *Phys Rev C.* (2009) **79**:054606. doi: 10.1103/PhysRevC.79.054606
162. Washiyama K, Ayik S, Lacroix D. Mass dispersion in transfer reactions with a stochastic mean-field theory. *Phys Rev C.* (2009) **80**:031602(R). doi: 10.1103/PhysRevC.80.031602
163. Yilmaz B, Ayik S, Lacroix D, Washiyama K. Nucleon exchange mechanism in heavy-ion collisions at near-barrier energies. *Phys Rev C.* (2011) **83**:064615. doi: 10.1103/PhysRevC.83.064615
164. Yilmaz B, Ayik S, Lacroix D, Yilmaz O. Nucleon exchange in heavy-ion collisions within a stochastic mean-field approach. *Phys Rev C.* (2014) **90**:024613. doi: 10.1103/PhysRevC.90.024613
165. Ayik S, Yilmaz O, Yilmaz B, Umar AS, Gokalp A, Turan G, Lacroix D. Quantal description of nucleon exchange in a stochastic mean-field approach. *Phys Rev C.* (2015) **91**:054601. doi: 10.1103/PhysRevC.91.054601
166. Ayik S, Yilmaz O, Yilmaz B, Umar AS. Quantal nucleon diffusion: central collisions of symmetric nuclei. *Phys Rev C.* (2016) **94**:044624. doi: 10.1103/PhysRevC.94.044624
167. Tanimura Y, Lacroix D, Ayik S. Microscopic phase-space exploration modeling of 258Fm spontaneous fission. *Phys Rev Lett.* (2017) **118**:152501. doi: 10.1103/PhysRevLett.118.152501
168. Tanimura Y, Lacroix D, Ayik S. Microscopic phase-space exploration modeling of 258Fm spontaneous fission. *Phys Rev Lett.* (2018) **121**:059902(E). doi: 10.1103/PhysRevLett.121.059902
169. Yilmaz B, Ayik S, Yilmaz O, Umar AS. Multinucleon transfer in $^{58}\text{Ni}+^{60}\text{Ni}$ and $^{60}\text{Ni}+^{60}\text{Ni}$ in a stochastic mean-field approach. *Phys Rev C.* (2018) **98**:034604. doi: 10.1103/PhysRevC.98.034604
170. Bulgac A, Jin S, Stetcu I. Unitary evolution with fluctuations and dissipation. *arXiv:1805.08908 [nucl-th]*. (2018).
171. Scamps G, Lacroix D, Bertsch GF, Washiyama K. Pairing dynamics in particle transport. *Phys Rev C.* (2012) **85**:034328. doi: 10.1103/PhysRevC.85.034328
172. Scamps G, Lacroix D. Effect of pairing on one- and two-nucleon transfer below the Coulomb barrier: a time-dependent microscopic description. *Phys Rev C.* (2013) **87**:014605. doi: 10.1103/PhysRevC.87.014605
173. Ebata S, Nakatsukasa T. Pairing effects in nuclear fusion reaction. *JPS Conf Proc.* (2014) **1**:013038. doi: 10.7566/JPSCP.1.013038
174. Ebata S, Nakatsukasa T. Repulsive aspects of pairing correlation in nuclear fusion reaction. *JPS Conf Proc.* (2015) **6**:020056. doi: 10.7566/JPSCP.6.020056
175. Corradi L, Szilner S, Pollarolo G, Colo G, Mason P, Farnea E. Single and pair neutron transfers at sub-barrier energies. *Phys Rev C.* (2011) **84**:034603. doi: 10.1103/PhysRevC.84.034603
176. Montanari D, Corradi L, Szilner S, Pollarolo G, Fioletto E, Montagnoli G. Neutron pair transfer in $^{60}\text{Ni}+^{116}\text{Sn}$ far below the coulomb barrier. *Phys Rev Lett.* (2014) **113**:052501. doi: 10.1103/PhysRevLett.113.052501

177. Montanari D, Corradi L, Szilner S, Pollarolo G, Goasduff A, Mijatović T. Pair neutron transfer in $^{60}\text{Ni}+^{116}\text{Sn}$ probed via γ -particle coincidences. *Phys Rev C*. (2016) **93**:054623. doi: 10.1103/PhysRevC.93.054623
178. Magierski P, Sekizawa K, Wlazłowski G. Novel role of superfluidity in low-energy nuclear reactions. *Phys Rev Lett*. (2017) **119**:042501. doi: 10.1103/PhysRevLett.119.042501
179. Sekizawa K, Magierski P, Wlazłowski G. Solitonic excitations in collisions of superfluid nuclei. *Proc Sci*. (2017) 214. doi: 10.22323/1.281.0214
180. Sekizawa K, Wlazłowski G, Magierski P. Solitonic excitations in collisions of superfluid nuclei: a qualitatively new phenomenon distinct from the Josephson effect. *EPJ Web Conf*. (2017) **163**:00051. doi: 10.1051/epjconf/201716300051
181. Scamps G. Examining empirical evidence of the effect of superfluidity on the fusion barrier. *Phys Rev C*. (2018) **97**:044611. doi: 10.1103/PhysRevC.97.044611
182. Bulgac A, Magierski P, Forbes MM. The unitary fermi gas: from Monte Carlo to density functionals. In: Zwirger W, editor. *BCS-BEC Crossover and the Unitary Fermi Gas*. Lecture Notes in Physics. Heidelberg: Springer (2012). p. 305–73.
183. Bulgac A. Time-dependent density functional theory and real-time dynamics of fermi superfluids. *Ann Rev Nucl Part Sci*. (2013) **63**:97. doi: 10.1146/annurev-nucl-102212-170631
184. Magierski P. Nuclear reactions and superfluid time dependent density functional theory. *arXiv:1606.02225 [nucl-th]*. (2016).
185. Magierski P, Grinevičute J, Sekizawa K. Pairing dynamics and time-dependent density functional theory. *Acta Phys Pol B*. (2018) **49**:281. doi: 10.5506/APhysPolB.49.281
186. Bulgac A, Yoon S. Large amplitude dynamics of the pairing correlations in a unitary fermi gas. *Phys Rev Lett*. (2009) **102**:085302. doi: 10.1103/PhysRevLett.102.085302
187. Bulgac A, Luo YL, Magierski P, Roche KJ, Yu Y. Real-time dynamics of quantized vortices in a unitary fermi superfluid. *Science*. (2011) **332**:1288. doi: 10.1126/science.1201968
188. Bulgac A, Luo YL, Roche KJ. Quantum shock waves and domain walls in the real-time dynamics of a superfluid unitary fermi gas. *Phys Rev Lett*. (2012) **108**:150401. doi: 10.1103/PhysRevLett.108.150401
189. Wlazłowski G, Sekizawa K, Marchwiany M, Magierski P. Suppressed solitonic cascade in spin-imbalanced superfluid fermi gas. *Phys Rev Lett*. (2018) **120**:253002. doi: 10.1103/PhysRevLett.120.253002
190. Bulgac A, Forbes MM, Kelley MM, Roche KJ, Wlazłowski G. Quantized superfluid vortex rings in the unitary fermi gas. *Phys Rev Lett*. (2014) **112**:025301. doi: 10.1103/PhysRevLett.112.025301
191. Wlazłowski G, Bulgac A, Forbes MM, Roche KJ. Life cycle of superfluid vortices and quantum turbulence in the unitary Fermi gas. *Phys Rev A*. (2015) **91**:031602(R). doi: 10.1103/PhysRevA.91.031602
192. Stetcu I, Bulgac A, Magierski P, Roche KJ. Isovector giant dipole resonance from the 3D time-dependent density functional theory for superfluid nuclei. *Phys Rev C*. (2011) **84**:051309(R). doi: 10.1103/PhysRevC.84.051309
193. Stetcu I, Bertulani C, Bulgac A, Magierski P, Roche KJ. Relativistic coulomb excitation within time dependent superfluid local density approximation. *Phys Rev Lett*. (2015) **114**:012701. doi: 10.1103/PhysRevLett.114.012701
194. Bulgac A, Magierski P, Roche KJ, Stetcu I. Induced fission of ^{240}Pu within a real-time microscopic framework. *Phys Rev Lett*. (2016) **116**:122504. doi: 10.1103/PhysRevLett.116.122504
195. Bulgac A, Jin S, Roche KJ, Schunck N, Stetcu I. *Fission Dynamics*. arXiv:1806.00694 [nucl-th] (2018).
196. Stetcu I, Bulgac A, Jin S, Roche KJ, Schunck N. *Real Time Description of Fission*. arXiv:1810.04024 [nucl-th]. (2018).
197. Wlazłowski G, Sekizawa K, Magierski P, Bulgac A, Forbes MM. Vortex pinning and dynamics in the neutron star crust. *Phys Rev Lett*. (2016) **117**:232701. doi: 10.1103/PhysRevLett.117.232701
198. Hashimoto Y. Time-dependent Hartree-Fock-Bogoliubov calculations using a Lagrange mesh with the Gogny interaction. *Phys Rev C*. (2013) **88**:034307. doi: 10.1103/PhysRevC.88.034307
199. Hashimoto Y, Scamps G. Gauge angle dependence in time-dependent Hartree-Fock-Bogoliubov calculations of $^{20}\text{O}+^{20}\text{O}$ head-on collisions with the Gogny interaction. *Phys Rev C*. (2016) **94**:014610. doi: 10.1103/PhysRevC.94.014610
200. Scamps G, Hashimoto Y. Transfer probabilities for the reactions $^{14,20}\text{O}+^{20}\text{O}$ in terms of multiple time-dependent Hartree-Fock-Bogoliubov trajectories. *Phys Rev C*. (2017) **96**:031602(R).
201. Winther A. Grazing reactions in collisions between heavy nuclei. *Nucl Phys*. (1994) **A572**:191. doi: 10.1016/0375-9474(94)90430-8
202. Winther A. Dissipation, polarization and fluctuation in grazing heavy-ion collisions and the boundary to the chaotic regime. *Nucl Phys*. (1995) **A594**:203. doi: 10.1016/0375-9474(95)00374-A
203. Dasso CH, Pollarolo G, Winther A. Systematics of isotope production with radioactive beams. *Phys Rev Lett*. (1994) **73**:1907. doi: 10.1103/PhysRevLett.73.1907
204. Dasso CH, Pollarolo G, Winther A. Particle evaporation following multinucleon transfer processes with radioactive beams. *Phys Rev C*. (1995) **52**:2264. doi: 10.1103/PhysRevC.52.2264
205. Yanez R, Loveland W. Predicting the production of neutron-rich heavy nuclei in multinucleon transfer reactions using a semi-classical model including evaporation and fission competition, GRAZING-F. *Phys Rev C*. (2015) **91**:044608.
206. Vigezzi E, Winther A. On the application of complex trajectories to direct heavy-ion reactions. *Ann Phys*. (1989) **192**:432. doi: 10.1016/0003-4916(89)90145-0
207. Adamian GG, Antonenko NV, Zubov AS. Production of unknown transactinides in asymmetry-exit-channel quasifission reactions. *Phys Rev C*. (2005) **71**:034603. doi: 10.1103/PhysRevC.71.034603
208. Penionzhkevich YE, Adamian GG, Antonenko NV. Towards neutron drip line via transfer-type reactions. *Phys Lett*. (2005) **B621**:119. doi: 10.1016/j.physletb.2005.05.085
209. Penionzhkevich YE, Adamian GG, Antonenko NV. Production of neutron-rich Ca isotopes in transfer-type reactions. *Eur Phys J A*. (2006) **27**:187. doi: 10.1140/epja/i2005-10252-6
210. Feng ZQ, Jin GM, Li JQ. Production of heavy isotopes in transfer reactions by collisions of $^{238}\text{U}+^{238}\text{U}$. *Phys Rev C*. (2009) **80**:067601. doi: 10.1103/PhysRevC.80.067601
211. Adamian GG, Antonenko NV, Sargsyan VV, Scheid W. Possibility of production of neutron-rich Zn and Ge isotopes in multinucleon transfer reactions at low energies. *Phys Rev C*. (2010) **81**:024604. doi: 10.1103/PhysRevC.81.024604
212. Adamian GG, Antonenko NV, Sargsyan VV, Scheid W. Predicted yields of new neutron-rich isotopes of nuclei with $Z = 64-80$ in the multinucleon transfer reaction $^{48}\text{Ca}+^{238}\text{U}$. *Phys Rev C*. (2010) **81**:057602. doi: 10.1103/PhysRevC.81.057602
213. Adamian GG, Antonenko NV, Lacroix D. Production of neutron-rich Ca, Sn, and Xe isotopes in transfer-type reactions with radioactive beams. *Phys Rev C*. (2010) **82**:064611. doi: 10.1103/PhysRevC.82.064611
214. Mun MH, Adamian GG, Antonenko NV, Oh Y, Kim Y. Production cross section of neutron-rich isotopes with radioactive and stable beams. *Phys Rev C*. (2014) **89**:034622. doi: 10.1103/PhysRevC.89.034622
215. Mun MH, Adamian GG, Antonenko NV, Oh Y, Kim Y. Toward neutron-rich nuclei via transfer reactions with stable and radioactive beams. *Phys Rev C*. (2015) **91**:054610. doi: 10.1103/PhysRevC.91.054610
216. Zhu L, Feng ZQ, Zhang FS. Production of heavy neutron-rich nuclei in transfer reactions within the dinuclear system model. *J Phys G Nucl Part Phys*. (2015) **42**:085102. doi: 10.1088/0954-3899/42/8/085102
217. Zhu L, Su J, Xie WJ, Zhang FS. Production of neutron-rich transcalifornium nuclei in ^{238}U -induced transfer reactions. *Phys Rev C*. (2016) **94**:054606. doi: 10.1103/PhysRevC.94.054606
218. Feng ZQ. Production of neutron-rich isotopes around $N = 126$ in multinucleon transfer reactions. *Phys Rev C*. (2017) **95**:024615. doi: 10.1103/PhysRevC.95.024615
219. Zhu L, Su J, Xie WJ, Zhang FS. Theoretical study on production of heavy neutron-rich isotopes around the $N = 126$ shell closure in radioactive beam induced transfer reactions. *Phys Lett*. (2017) **B767**:437. doi: 10.1016/j.physletb.2017.01.082
220. Li C, Wen P, Li J, Zhang G, Li B, Zhang FS. Production of heavy neutron-rich nuclei with radioactive beams in multinucleon transfer reactions. *Nucl Sci Tech*. (2017) **28**:110. doi: 10.1007/s41365-017-0266-z
221. Zhu L, Zhang FS, Wen PW, Su J, Xie WJ. Production of neutron-rich nuclei with $Z = 60-73$ in reactions induced by Xe isotopes. *Phys Rev C*. (2017) **96**:024606. doi: 10.1103/PhysRevC.96.024606

222. Zhao K, Li Z, Wang N, Zhang Y, Li Q, Wang Y, Wu X. Production mechanism of neutron-rich transuranium nuclei in $^{238}\text{U}+^{238}\text{U}$ collisions at near-barrier energies. *Phys Rev C*. (2015) **92**:024613. doi: 10.1103/PhysRevC.92.024613
223. Li C, Zhang F, Li J, Zhu L, Tian J, Wang N, et al. Multinucleon transfer in the $^{136}\text{Xe}+^{208}\text{Pb}$ reaction. *Phys Rev C*. (2016) **93**:014618. doi: 10.1103/PhysRevC.93.014618
224. Wang N, Guo L. New neutron-rich isotope production in $^{154}\text{Sm}+^{160}\text{Gd}$. *Phys Lett*. (2016) **B760**:236. doi: 10.1016/j.physletb.2016.06.073
225. Zhao K, Li Z, Zhang Y, Wang N, Li Q, Shen C et al. Production of unknown neutron-rich isotopes in $^{238}\text{U}+^{238}\text{U}$ collisions at near-barrier energy. *Phys Rev C*. (2016) **94**:024601. doi: 10.1103/PhysRevC.94.024601
226. Karpov AV, Saiko VV. Modeling near-barrier collisions of heavy ions based on a Langevin-type approach. *Phys Rev C*. (2017) **96**:024618. doi: 10.1103/PhysRevC.96.024618
227. Saiko V, Karpov A. Analysis of multinucleon transfer reactions involving spherical and statically deformed nuclei using a Langevin-type approach. *Phys Rev C*. (2019) **99**:014613. doi: 10.1103/PhysRevC.99.014613
228. Welsh T, Loveland W, Yanez R, Barrett JS, McCutchan EA, Sonzogni AA, et al. Modeling multi-nucleon transfer in symmetric collisions of massive nuclei. *Phys Lett*. (2017) **B771**:119. doi: 10.1016/j.physletb.2017.05.044
229. Wen PW, Li C, Zhu L, Lin CJ, Zhang FS. Mechanism of multinucleon transfer reaction based on the GRAZING model and DNS model. *J Phys G Nucl Part Phys*. (2017) **44**:115101. doi: 10.1088/1361-6471/aa8b07

Conflict of Interest Statement: The author declares that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2019 Sekizawa. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.