



Preliminary Study on the *In vitro* and *In vivo* Effects of *Asparagopsis taxiformis* Bioactive Phycoderivates on Teleosts

Fabio Marino^{1*}, Gianfranco Di Caro², Concetta Gugliandolo¹, Antonio Spanò¹, Caterina Faggio¹, Giuseppa Genovese¹, Marina Morabito¹, Annamaria Russo¹, Davide Barreca¹, Francesco Fazio² and Andrea Santulli³

¹ Department of Chemical, Biological, Pharmaceutical and Environmental Sciences, University of Messina, Messina, Italy, ² Department of Veterinary Sciences, University of Messina, Messina, Italy, ³ Laboratory of Marine Biochemistry and Ecotoxicology, Department of Earth and Sea Sciences, University of Palermo, Trapani, Italy

OPEN ACCESS

Edited by:

Francesca Carella,
University of Naples Federico II, Italy

Reviewed by:

Ivona Mladineo,
Istituto Ortopedico Rizzoli (IRCCS),
Croatia

Donatella Volpatti,
University of Udine, Italy

*Correspondence:

Fabio Marino
marinof@unime.it

Specialty section:

This article was submitted to
Aquatic Physiology,
a section of the journal
Frontiers in Physiology

Received: 15 April 2016

Accepted: 23 September 2016

Published: 25 October 2016

Citation:

Marino F, Di Caro G, Gugliandolo C, Spanò A, Faggio C, Genovese G, Morabito M, Russo A, Barreca D, Fazio F and Santulli A (2016) Preliminary Study on the *In vitro* and *In vivo* Effects of *Asparagopsis taxiformis* Bioactive Phycoderivates on Teleosts. *Front. Physiol.* 7:459. doi: 10.3389/fphys.2016.00459

Several compounds from marine organisms have been studied for their potential use in aquaculture. Among the red algae, *Asparagopsis taxiformis* is considered one of the most promising species for the production of bioactive metabolites with numerous proposed applications. Here, the *in vitro* antibacterial activity, the easy handling and the absence of adverse effects on marine fish species are reported. Depending on the seasonal period of sampling, ethanol extracts of *A. taxiformis* exhibited significantly different inhibitory activity against fish pathogenic bacteria. The extract obtained in late spring showed strong antibacterial activity against *Aeromonas salmonicida* subsp. *salmonicida*, *Vibrio alginolyticus*, and *V. vulnificus*, and moderate activity against *Photobacterium damsela* subsp. *damsela*, *P. damsela* subsp. *piscicida*, *V. harveyi* and *V. parahaemolyticus*. Sea bass and gilthead sea bream were fed with pellets supplied with the alga and algal extracts. The absence of undesired effects on fish was demonstrated. Hematological and biochemical investigations allowed to confirm that the whole alga and its extracts could be proposed for a future application in aquaculture.

Keywords: *Asparagopsis taxiformis*, *Dicentrarchus labrax*, *Sparus aurata*, antibacterial activity, hematological parameters, oxidative stress, toxicity

INTRODUCTION

In the last decades marine organisms have been considered as potential and promising sources of pharmaceuticals (Newman et al., 2003; Blunt et al., 2008; Mayer et al., 2011). Particularly, seaweeds are considered as a source of bioactive metabolites characterized by a wide spectrum of effects (Fouladvand et al., 2011). Compounds with immunostimulant, cytostatic, antiviral, anti-helminthic, antifungal, antibacterial activities have been detected in all phyla of macroalgae (Ballesteros et al., 1992; Val et al., 2001; Smit, 2004; Talarico et al., 2004; Cumashi et al., 2007; Shanmughapriya et al., 2008; Dhargalkar and Verlecar, 2009; Kamenarska et al., 2009; Vallinayagam et al., 2009; Allmendinger et al., 2010; Bouhlal et al., 2010; Chanda et al., 2010; Vonthron-Sénécheau et al., 2011; Barman et al., 2013; de Jesus Raposo et al., 2015).

Among red algae, *Asparagopsis* species (Bonnemaisoniales, Rhodophyta) showed potent antifungal, antibacterial, and antiprotozoal attributes (Burreson et al., 1975; McConnell and Fenical, 1977; Bansemir et al., 2006; Salvador et al., 2007; Genovese et al., 2009, 2012; Jiao et al., 2011; Manilal et al., 2012). The genus *Asparagopsis* produces several halogenated compounds, such as haloforms, methanes, ketones, acetates, and acrylates, which are assumed as bio-active compounds (McConnell and Fenical, 1977; Woolard et al., 1979; Bansemir et al., 2006). In general, the production of biologically-active metabolites is linked to the ability to store them into specialized storage structures in order to avoid autotoxicity (McKey, 1979). Members in the family Bonnemaisoniaceae, to which *Asparagopsis* species belong, form “vesicle” or “gland cells” (Wolk, 1968; Young, 1977; Womersley, 1998; Paul et al., 2006). The pungent aroma of these algae is due to an essential oil composed mainly of bromoform with smaller amounts of other bromine, chlorine, and iodine-containing methane, ethane, ethanol, acetaldehydes, acetones, 2-acetoxypropanes, propenes, epoxypropanes, acroleins, and butenones, stored in vacuoles within gland cells (Burreson et al., 1976; El-Baroty et al., 2007). *Asparagopsis taxiformis* (Delile) Trevisan de Saint-Léon occurs along tropical and warm temperate coasts, and shows disjoint Atlantic, Mediterranean, and Indo-Pacific populations. It exhibits a strong invasive behavior and therefore it was included in the list of the “Worst Invasives in the Mediterranean Sea” (Zenetos et al., 2010). Recent phylogeographic approaches have shown that this species consists of a number of cryptic species, as well as many other marine algae (Andreakis et al., 2015). At present, *Asparagopsis* species are cultivated for dermo-cosmetical and parapharmaceutical purposes in Atlantic Europe, especially in France, Portugal, Ireland, as well as in Hawaii, Indonesia, Philippines, and New Zealand (Kraan and Barrington, 2005; Mata et al., 2012). Biomass yield proved to be high in nutrients rich water, such as the effluents of fish farms, where the alga acts as a biofilter in integrated multi-trophic aquaculture (IMTA, Barrington et al., 2009).

Natural marine products, including those of algal origin, have enormous pharmacological potentialities, especially when addressing the increasing phenomenon of bacterial antibiotic resistance and the collateral effects that synthetic drugs have both on humans and environment. Although several studies focused on the antimicrobial properties of macroalgae, few studies reported on bacterial pathogens relevant in aquaculture (Val et al., 2001; Liao et al., 2003; Bansemir et al., 2006).

Since several studies underlined the variability in the production of antimicrobials among species within an algal genus, we proposed to use DNA labels to identify unambiguously the algal populations used in applied research.

To verify positive or adverse effects of natural substances, both *in vitro* and *in vivo* trials are needed prior to approach large scale applications in aquaculture. As concerns the *in vivo* studies, hematological profile is an important parameter to be considered as pathophysiological indicator of the whole body status.

The maintenance of a constant volume is a homeostatic requirement in animal cells, when exposed to osmotic gradients inducing a temporary haemodilution or haemoconcentration.

Often the cell volume is compromised by the variation in the concentration of intracellular osmolytes, as a consequence of different processes (e.g., secretion, ion gradients, accumulation of nutrients, cell growth, and proliferation), and a variety of pathologic conditions. Cell volume regulation following cell-swelling, due to osmotic influx of water, involves the release of organic solutes, such as amino acids and ions, through the activation of K^+ channels and/or anion channels, KCl cotransport, or parallel activation of K^+/H^+ exchange and Cl^-/HCO_3^- exchange, and the osmotically obliged water efflux (Faggio et al., 2011). This regulatory response is called “regulatory volume decrease” (RVD). Volume-regulatory behavior has been observed in: cells isolated from digestive glands of *Mytilus galloprovincialis* (Torre et al., 2013), *Gobius niger* enterocytes (Trischitta et al., 2004), *Anguilla anguilla* enterocytes (Trischitta et al., 2005), turbot hepatocytes (Ollivier et al., 2006), Ehrlich ascites tumour cells (Hoffmann, 1984), and mammalian red blood cells (Ellory et al., 1985) exposed to osmotic stress.

The work was finalized to investigate: (i) the *in vitro* antibacterial activity of extracts obtained from a DNA barcoded population of *A. taxiformis* collected from the Straits of Messina against fish pathogens, and (ii) the possible effects of dietary administration of the whole alga and its crude extracts on sea bass and gilthead sea bream, by evaluating the RVD of fish hepatocytes, the cytotoxic effects of algal extracts on both red blood cells and isolated hepatocytes of fish, the hematological profiles and the erythrocytes response to oxidative stress.

MATERIALS AND METHODS

Algae Sampling, Identification and Processing

Plants of *A. taxiformis* were collected from a marine site ($38^{\circ}12'3.28''N$, $15^{\circ}33'35.63''E$), along northeastern Sicilian coast of the Straits of Messina (Italy) in April (ASP001), May (ASP002), June (ASP003), and July (ASP004) 2012-2013, throughout the period in which the alga is present in the field. Fresh plants were washed in sterile seawater, manually cleaned of epiphytes and frozen at $-20^{\circ}C$ (for the use within few days) or lyophilized (for longer use).

A subsample of each collection was used to obtain COI-5' sequences according to standard DNA barcoding protocols for red algae (Manghisi et al., 2010). In details, DNA extraction was performed by a Proteinase K protocol and the COI-5' region was PCR amplified using the primers GazF1 and GazR1 (Saunders, 2005).

Sequencing reactions were performed by an external company (Macrogen Europe, The Netherlands). Forward and reverse sequence reads were assembled with the software ChromasPro (v. 1.41, Technelysium Pty Ltd).

A multiple sequence alignment was constructed in MacClade 4.0 (Maddison and Maddison, 2000), including sequences of *Asparagopsis* species downloaded from GenBank, and subjected to distance analysis in PAUP* 4b10 for the Macintosh (Swofford, 2002) to define species attributions.

For crude extract preparation, frozen or lyophilized plants of *A. taxiformis* were soaked in absolute ethanol at room temperature for 48 h. Extracts were dried with a Rotavapor® at low temperature (35°C) to preserve volatile compounds from evaporation.

Ethanol extracts obtained in April (ASP001), May (ASP002), June (ASP003), and July (ASP004) were used for the *in vitro* antibacterial and cytotoxicity assays, while both ethanol extracts and lyophilized plants were used for the *in vivo* trials.

In vitro Experiments

Antibacterial Activity

Ethanol extracts of *A. taxiformis* (ASP001, ASP002, ASP003, and ASP004) were tested against the following lab strains of fish and shellfish pathogenic bacteria: *Aeromonas salmonicida* subsp. *salmonicida* (13M), *Photobacterium damsela* subsp. *damsela* (Pdd), *P. damsela* subsp. *piscicida* (Pdp), *Vibrio alginolyticus* (Ag37), *V. harveyi* (G5), *V. parahaemolyticus* (L12G), and *V. vulnificus* (OPV10) which are part of the Culture Collection housed at the Dept of Chemistry, Biology, Pharmaceutical and Environmental Sciences of the University of Messina (Italy). The strains were previously isolated and identified as reported in Gugliandolo et al. (2008, 2011). They were maintained at -20°C in Tryptone Soy Broth (Difco) supplemented with 1% (w/v) NaCl (TSB1) and 50% (v/v) glycerol.

Antibacterial activity was evaluated by using the standard disk diffusion method (Kirby Bauer test), as accepted by the National Committee for Clinical Laboratory Standards (NCCLS 2000) and commercially available disks (6 mm in diameter, Oxoid). Strains were overnight grown onto plates of Tryptone Soya Agar (TSA) (Oxoid) amended with 1% of NaCl (TSA1), for 24 h at 25°C. Each strain was suspended in 3 ml of 0.9% NaCl solution with a turbidity optically comparable to that of the 0.5 McFarland standard (1.5×10^8 bacteria/ml) and suspensions (100 µl) were inoculated onto triplicate plates of TSA1.

Each extract (100 mg, equivalent to 3 g of lyophilized algal sample) was dissolved in 1000 µl of absolute ethanol, and 20 µl were applied to sterile filter paper disks. After solvent evaporation, the disks (containing 2 mg of the extract) were placed onto the inoculated plates. Disks soaked with ethanol and submitted to evaporation were used as negative control, and those containing chloramphenicol (30 µg) and tetracycline (30 µg) (Difco, Becton Dickinson and Company, USA) were used as positive control. Plates were incubated overnight at 25°C. The diameter of complete inhibition zone was measured and means and standard deviations ($n = 3$) were calculated. Inhibition zones ≥ 15 mm in diameter were stated as strong, from 9 to 14 mm as moderate and ≤ 8 mm as weak in terms of activities (Bansemir et al., 2006).

Minimum inhibitory concentration (MIC) values were determined for the most active extracts, by using the serial dilution assay, as accepted by the European Committee for Antimicrobial Susceptibility Testing (EUCAST) of the European Society of Clinical Microbiology and Infectious Diseases (ESCMID) (2003). Serial dilutions of each extract (16, 8, 4, 2, 1, 0.5, and 0.25 mg/ml) were prepared in tubes of TSB1 and then inoculated with suitable aliquots of overnight cultured bacterial

strain in TSB1. To verify the inhibition activity of the extract, 100 µl from all tubes without visible bacterial growth were plated onto TSA1, and incubated overnight at 25°C.

Isolation of *Dicentrarchus labrax* Hepatocytes and Regulation Volume Decrease (RVD) Experiments

Ten *Dicentrarchus labrax* specimens (total length 23 ± 2.6 cm; weight 143 ± 10.2 g): control basal diet and ten specimens fed diets enriched with algal ethanolic extracts (total length 20 ± 1.6 cm; weight 100 ± 4.3 g) for 2 weeks were sacrificed with an overdose of MS222 on 0.7 g/l ratio. The liver was removed and subsequently hepatocytes were isolated by collagenase (type IV-activity P125 CDU/mg; CDU = collagenase digestion units, Sigma Aldrich, St. Louis, MO, USA) digestion methods, which required perfusion of the portal vein as described previously (Ollivier et al., 2006). Isolated hepatocytes were maintained in a 370 mOsm/kg physiological saline solution: (millimoles/l): 172 NaCl, 3.4 KCl, 0.8 MgSO₄, 1.5 CaCl₂, 5 NaHCO₃, 0.33 Na₂HPO₄, 0.44 KH₂PO₄, 5 glucose and 10 Hepes (pH 7.4 and 18°C). Afterwards, the suspension was filtered by 200 µm and 75 µm nylon filters. The cells were suspended again in physiological saline solution and washed twice by centrifugation (1700 g/10 min/4°C). Prior to the experiments the cells were maintained under slight agitation for at least 1 h at 18°C.

In order to immobilize cells, they were seeded onto the bottom of a thermostated Plexiglas chamber (18°C) coated with poly-L-lysine (0.01%) and left to adhere for 2 min on slide. For RVD experiment the isolated cells were visualized and measured by the method described in a previous paper (Faggio et al., 2011). One drop of cell suspension (5×10^6 cells ml⁻¹), was placed on a glass slide pretreated with poly-lysine to facilitate cell adhesion (to fix the cells). Two thin strips of double-sided adhesive were placed at the upper and lower edges of the glass slide to support the cover slip and to create an interspace in which the hypotonic experimental solution was then added (millimoles/l: 124 NaCl, 3.4 KCl, 0.8 MgSO₄, 1.5 CaCl₂, 5 NaHCO₃, 0.33 Na₂HPO₄, 0.44 KH₂PO₄, 5 glucose, and 10 Hepes; pH 7.4). Hepatocyte images, obtained using a CCD camera (DVC Canada color, 1300, 12 bit cooled, slow scan) connected to the microscope were captured at 60 s intervals during 3 min before and 1 min after initiating the hypotonic shock, thereafter they were captured every 5 min for 60 min. Cell images were digitized using a color video camera (Sony). For each slide six/seven cells were counted. The cell areas for each experimental condition (A_{exp}) were compared to the areas measured in isotonic solution (A_i) at the beginning of the experiment. Consequently, the data were reported as relative area (A_{exp}/A_i).

Cytotoxicity Assays

In order to verify if algal extracts administered to fish exert cytotoxic effects, 100 isolated hepatocytes of *D. labrax* (fed diet enriched with algal extract) were measured to verify the viability which was measured by the Trypan blue exclusion method and compared to control group, constituted of cells isolated from control fish (Matozzo et al., 2016). The quality of their lysosomes was evaluated by neutral red retention assay (NR) (which measures the stability of lysosomal membrane). The dye

retention was evaluated after an incubation period of 30 and 60 min, according to Repetto et al. (2008). The lysosomes of vital hepatocytes were stained red. All experiments were carried in triplicates.

In vivo Trials

The *in vivo* study was carried out at Centre for Experimental Fish Pathology of Sicily (C.I.S.S.), Establishment for Users recognized by the Italian Ministry of Health for experimental activity on aquatic organisms (according to D.L. 26/2014), at the Department of Veterinary Sciences, University of Messina. Fish species used in this study were *D. labrax* and *S. aurata* and all experimental procedures were performed in compliance with EU Directive 63/2010 and D.L. 26/2014 after project evaluation by the competent authorities (authorization n. 587/2016).

Pellets Preparation for Fish Feeding

Lyophilized *A. taxiformis* samples, collected from April to July, or its ethanolic extracts were daily administered to fish by feeding them with commercial feed (Skretting) supplemented in lab. Control pellets and pellets amended with alga or algal extracts, were prepared by adding each compound to the ingredients of the basal diet. The uniform distribution of the ingredients in the feed was obtained by mixing pellets with water through magnetic stirrers Heating (VELPA-Scientifica) at 70°C and 4 Stirrer, until obtaining a damp mixture with soft consistency, and subsequently adding algae or algal extracts to the enriched pellet and continuing to mix the mixture at 7 Stirrer for 30 min. The obtained compound was dried at room temperature for 24 h. The feed was then minced and sieved to produce a crumble size suitable to sea bass and gilthead sea bream feeding. The dosage used was 1 g algae/10 g feed and 0.1 g extract/1 g feed.

Different experimental cycles, summarized in **Tables 1, 2**, were carried out on both adult fish and juveniles to verify possible adverse effects on fish growth obtain enough data to achieve statistically significant results.

Each group of adult Mediterranean sea bass and juveniles were reared in 500 l tanks, whereas gilthead sea bream, coming from a local commercial fish farm, were reared in 100 l tanks. Each tank was supplied by an independent biological and mechanical filtration system, artificial aeration and illumination with neon to assure perfect conditions of animal welfare. The water was prepared using a synthetic salt (Blue treasure, Sea Salt) obtained by mixing pure elements to obtain marine water chemically and biologically excellent. Water parameters in all tanks were checked daily by a multiparametric probe (HI9829 Aquaprobe, Hanna instruments, Padova, Italy) and the following values were maintained for all the 2 months of experiments: temperature 20–22°C; salinity 30‰; pH 8~. Moreover, a photoperiod of 12 h light/12 h dark was provided. In the first trial 80 sea bass subdivided in 4 fish groups were used.

G0. 20 juvenile Mediterranean sea bass, of about 6 months in age, with a median weight of 15 g and a median length of 10 cm, were randomly selected by a pool of healthy subjects coming from a local commercial fish farm; after quarantine, fish were successively transferred to the experimental tank for the trial and were fed with pellet supplied with algae.

As a control group, further 20 age-matched (G1) specimens were used and fed with commercial pellet. G5. 20 adult Mediterranean sea bass (*D. labrax*), 12 months aged, median weight 100 g, 20 cm in length, were fed with pellets supplied with algal extract. Water parameters as previously referred.

G6. 20 sea bass fed with control pellet.

All the experimental groups were treated respectively for a 8 weeks (G0) and 2 weeks (G5 and G6) long period. The shorter time of the trial with algal extract was related to initial appearance of adverse effects on behavior which suggested to anticipate the sacrifice according to general guidelines for animal welfare during toxicological studies.

Water parameters and health status of fish, as well as their vitality and appetite, were daily checked and all possible variations were registered.

In the second experimental cycle, 60 gilthead sea bream, coming from a local commercial fish farm, were subdivided in three groups (G7–G9) each constituted of 20 fish, median weight 7 g and a median length of 5 cm. G7 was supplied with algae, G8 with algal extract, G9 was used as age-matched normal control. The third experimental trial was performed repeating the feed protocols already described using sea bass and sea bream alevins. A total of 120 fish (60 sea bass and 40 sea bream) 50 g in medium weight were used subdivided in groups of 20. Three groups of sea bass (G2 alga, G3 algal extract, G4 normal control) and 3 groups of sea bream (G10 alga, G12 control, G11 algal extract).

Fish were constantly monitored for a period of 10 min after feed administration to evaluate feed palatability and effective food consumption.

Sample Collection

Blood samples were obtained before and at the end of each trial on both treated fish and control fish. For hematological analysis only fish ranging in weight from 50 to 100 g were considered. Fish ($n = 10$ for each group) from G 7, 8, and 9 (for *D. labrax*) and from G 10, 11, and 12 (*S. aurata*) were deeply anesthetized with MS222 (0.4 g/l) and successively underwent venipuncture for blood collection. Blood samples were obtained from caudal vein using a 18 G \times 1 ½ syringe and collected into micro tubes (Miniplast 0.6 mL, LP Italiana Spa, Milano) containing EDTA (1.26 mg/0.6 mL).

All samples were analyzed in triplicate by the same operator. After the last blood sampling, fish were euthanatized with MS222 (0.7 g/l) and tissue samples were collected from skin, gills, heart, liver, gut, spleen, kidney, and brain for routine histopathology. Briefly, 5 μ m thick paraffin sections were obtained and stained with hematoxylin-eosin and Giemsa methods, mounted on slides and observed by light microscope.

Hematological Analyses

The hematological profile was measured using an automated hematology analyzer (HeCo Vet C, SEAC, Florence, Italy) with a method already used both in *S. aurata* and *D. labrax* (Fazio et al., 2012a,b, 2013, 2015). Hemogram included the white blood cell count (WBC), red blood cell count (RBC), hematocrit (Hct), hemoglobin concentration (Hgb), mean corpuscular volume (MCV), mean corpuscular hemoglobin

TABLE 1 | Sea bass experimental groups submitted to the *in vivo* trials.

Group	Species	Number	Weight	Length	Supplementation	Duration
G.0	<i>Dicentrarchus labrax</i>	20	15 g	10 cm	Algae 1 g/10 g feed	2 months
G.1	<i>Dicentrarchus labrax</i>	20	15 g	10 cm	Control	2 months
G.2	<i>Dicentrarchus labrax</i>	20	50 g	14 cm	Algae 1 g/10 g feed	2 months
G.3	<i>Dicentrarchus labrax</i>	20	50 g	15 cm	Algal extract 0.1 g/1 g feed	2 weeks
G.4	<i>Dicentrarchus labrax</i>	20	50 g	16 cm	Control	2 months
G.5	<i>Dicentrarchus labrax</i>	20	100 g	20 cm	Algal extract 0.1 g/1 g feed	2 weeks
G.6	<i>Dicentrarchus labrax</i>	20	100 g	20 cm	Control	2 weeks

TABLE 2 | Sea bream experimental groups submitted to the *in vivo* trials.

Group	Species	Number	Weight	Length	Supplementation	Duration
G.7	<i>Sparus aurata</i>	20	7 g	5 cm	Algae 1 g/10 g feed	2 months
G.8	<i>Sparus aurata</i>	20	7 g	5 cm	Algal extract 0.1 g/1 g feed	2 weeks
G.9	<i>Sparus aurata</i>	20	7 g	5 cm	Control	2 months
G.10	<i>Sparus aurata</i>	20	50 g	14 cm	Algae 1 g/10 g feed	2 months
G.11	<i>Sparus aurata</i>	20	50 g	14 cm	Algal extract 0.1 g/1 g feed	2 weeks
G.12	<i>Sparus aurata</i>	20	50 g	14 cm	Control	2 months

(MCH), mean corpuscular hemoglobin concentration (MCHC), and thrombocyte count (TC).

Erythrocyte Oxidative Stress

At the end of the feeding period, both *D. labrax* and *S. aurata*, 100 and 50 g in weight respectively were anesthetized and divided into 2 groups ($n = 8$) (control and treated with dietary whole *A. taxiformis*). The blood was collected and inserted in distinct tubes containing EDTA (1.26 mg/0.6 ml). The plasma and erythrocytes were separated by centrifugation at 1020 g for 5 min and the red blood cells (RBC) were washed with isosmotic solution.

The evaluation of the levels of erythrocytary hemolysis and met-Hb, membrane peroxidation, determination of the total reducing power (FRAP) and dosage of acetylcholinesterase on erythrocyte membrane were carried out according to Ficarra et al. (2013, 2016), Scala et al. (2015) and Carelli-Alinovi et al. (2016). Nitrites and nitrates (as an index of stability of the phospholipidic bilayer), quantization of the total thiols and glutathione (index of the cells redox state and onset of oxidative injuries) were performed utilizing a commercial determination kit supplied by Sigma Aldrich.

Statistical Analysis

To assess the significant differences in *in vitro* antibacterial activity among the extracts, one-way analysis of variance

(ANOVA) was applied by using SPSS software v.17 (SPSS Italia, Rome, Italy).

In vivo analytical data, represented as mean (M) \pm standard error of the main (SEM), are the averages of three analyses carried out by the same operator. Hematological samples exhibited parallel displacement to the standard curve. The overall intra-assay coefficient of variation was <9%.

Data obtained for different hematological parameters were tested for normality using Kolmogorov-Smirnov test. $P < 0.05$ was considered statistically significant.

One-way analysis of Variance (ANOVA) was used to determinate a statistically significant effect of treatments on all hematological parameters obtained respect to control group for *S. aurata* and *D. labrax*. Bonferroni's multiple comparison test was used for *post hoc* comparison.

Data were analyzed using statistical software prism v. 5.00 (Graphpad Software Ltd., USA, 2003).

RESULTS

Algal Extracts

Algal specimens were identified using DNA barcoding. Eight COI5' sequences were generated, one per each collection. All obtained sequences were identical among them and compared to a sample previously sequenced from the same marine site

(Genbank accession number JN642177). The population under investigation resulted as *A. taxiformis* “lineage 2,” as named in Andreakis et al. (2007).

Ethanol extracts and lyophilized plants were produced and provided for *in vivo* and *in vitro* trials. No evaluation of the efficiency of the extraction procedures was performed due to the nature of algae, being the amount of bioactive molecules variable in time in relation to harvest period, site and other variables. This efficiency will be evaluated when bioactive molecules will be purified.

In vitro Experiments

Antibacterial Activity

No inhibition activity was observed using extracts from algal samples collected in June (ASP003) and July (ASP004), while those collected in April (ASP001) and in May (ASP002) showed antibacterial activities.

Differently from ASP001, ASP002 was active against the whole panel of target pathogens. The inhibition activity of the two extracts was significantly different ($P < 0.01$) as compared to control.

ASP002 (2 mg/disk) showed strong antibacterial activity against *V. alginolyticus* (Ag37) (17.2 ± 1.5 mm), *V. vulnificus* (OPV10) (17.1 mm \pm 1.1), and *Aer. salmonicida* (13M) (15.2 ± 1.0 mm), and moderate activity against *P. damselae* (Pdd) (13.8 ± 1.8), *P. piscicida* (Pdp) (13.8 ± 0.8), *V. harveyi* (G5) (12.4 ± 1.0), and *V. parahaemolyticus* (L12G) (12.2 ± 1.3).

ASP001 showed a moderate activity against *V. alginolyticus* (Ag37) (12.8 ± 1.2), *Aer. salmonicida* (13M) (12.5 ± 0.1), and *V. harveyi* (G5) (12.0 ± 1.5).

The MIC values of the ASP002 extract were the following: 4 mg/ml against *Aer. salmonicida* (13M), *P. damselae* (Pdd), and *P. piscicida* (Pdp), and 2 mg/ml against *V. harveyi* (G5).

RVD Experiments

Isolated hepatocytes of *D. labrax*, fed with control basal diet, exposed to the rapid change of osmolarity, initially increased their size and then tended to return to their initial volume. As showed in **Figure 1** the hepatocytes reached their maximum swelling, corresponding to an increase of 15% of volume, after 5 min exposure to the hypotonic solution. Thereafter, they exhibited an RVD response. No significant difference due to fed diets enriched with algae extract was observed.


Cytotoxicity Assay

The results obtained by Trypan blue exclusion method, supported by the results obtained by Neutral Red (NR), revealed no apparent changes in cellular viability neither in *D. labrax* isolated hepatocytes fed with control basal diet (95% of cellular viability), nor in *D. labrax* isolated hepatocytes fed diets enriched with algal extracts (95% of cellular viability).

In vivo Trials

Fish Feeding

During administering of all the different specific feeds no adverse effects, such as mortality and pathological symptoms, nor significant modifications of the water parameters were



detected. Only when checking the effects of algal extracts, slight behavioral modifications were detected probably due to the higher concentration of the ethanolic extracts in different bioactive molecules, with both positive and negative effects, as compared to the whole algae. Both the remaining specific supplements used, alga and its extracts, did not modify palatability of pellet, although, only at the beginning of the administration, in *S. aurata* a slight delay (less than 2 min) in eating pellet was registered.

Histological results revealed no undesirable changes in all the tissues of the fish studied.

Hematological Analyses

Mean values \pm SEM of hematological parameters obtained in control group and experimental groups for *D. labrax* and *S. aurata* are reported in **Tables 3, 4** respectively.

In the control group all hematological parameters were within the physiological range of *S. aurata* and *D. labrax* reported in previous research (Faggio et al., 2014; Fazio et al., 2015).

For *D. labrax* (**Table 3**) no statistical significant different were observed between experimental groups (G2 and G3) respect to the control group (G4) for all parameters, with the only exception of Hb that showed a significantly reduction in experimental groups (G2 and G3) in the respect to the control group (G4). For *S. aurata* (**Table 4**) ANOVA analysis showed significant changes in all hematological parameters registered in the experimental groups (G10 and G11) respect to the control group (G12), except for MCH and MCHC. In particular, RBC, Hgb, and Hct values were lower in groups G10 and G11 respect to the control group (G12), while MCV showed an increase in G11 respect to the control group (G12).

The mean values of WBC in fish fed with alga (G10) or algal extracts (G11) were $62.17 \times 10^3 \mu\text{l}$ and $54.97 \times 10^3 \mu\text{l}$ respectively, which were significantly lower than that registered in the control group ($74.70 \times 10^3 \mu\text{l}$). Fish subjected to administration of alga (G10) showed significantly lower numbers

TABLE 3 | Mean values \pm SEM of hematological parameters of *D. labrax* ($n = 10$ for each group) obtained in Control Group (G4, gray labeled) and Experimental Groups (G2 and G3).

Parameters	Control group	Experimental groups	
	G4	G2	G3
RBC ($\times 10^6/\mu\text{l}$)	3.50 \pm 0.22 ^a	3.42 \pm 0.90 ^a	3.48 \pm 0.50 ^a
Hgb (g/dL)	9.80 \pm 0.54 ^a	9.27 \pm 0.30 ^b	9.30 \pm 0.54 ^b
Hct (%)	41.40 \pm 1.20 ^a	40.70 \pm 1.30 ^a	41.50 \pm 1.60 ^a
MCV (fL)	118.00 \pm 2.70 ^a	119.00 \pm 1.60 ^a	119.25 \pm 2.00 ^a
MCH (pg/cell)	28.00 \pm 1.00 ^a	26.32 \pm 0.80 ^a	26.72 \pm 1.20 ^a
MCHC (g/dL)	23.67 \pm 0.90 ^a	22.11 \pm 0.70 ^a	22.41 \pm 1.00 ^a
WBC ($\times 10^3/\mu\text{l}$)	70.20 \pm 3.54 ^a	68.40 \pm 1.54 ^a	67.00 \pm 2.10 ^a
TC ($\times 10^3/\mu\text{l}$)	54.00 \pm 0.54 ^a	52.00 \pm 0.32 ^a	53.20 \pm 0.27 ^a

Means without the same alphabetical characters within the same parameters represent statistical differences ($P < 0.05$).

TABLE 4 | Mean values \pm SEM of hematological parameters of *S. aurata* ($n = 10$ for each group) obtained in Control Group (G12, Gray labeled) and experimental Groups (G10 and G11).

Parameters	Control group	Experimental groups	
	G12	G10	G11
RBC ($\times 10^6/\mu\text{l}$)	3.77 \pm 0.13 ^a	3.10 \pm 0.11 ^b	3.42 \pm 0.08 ^b
Hgb (g/dL)	11.75 \pm 0.29 ^a	9.61 \pm 0.19 ^b	10.67 \pm 0.10 ^b
Hct (%)	43.62 \pm 1.37 ^a	35.92 \pm 1.01 ^b	41.73 \pm 0.56 ^b
MCV (fL)	116.10 \pm 2.65 ^a	117.5 \pm 3.78 ^a	129.40 \pm 1.27 ^b
MCH (pg/cell)	31.29 \pm 0.48 ^a	31.20 \pm 1.06 ^a	31.40 \pm 0.68 ^a
MCHC (g/dL)	27.07 \pm 0.48 ^a	26.89 \pm 0.60 ^a	25.62 \pm 0.37 ^a
WBC ($\times 10^3/\mu\text{l}$)	74.70 \pm 2.15 ^a	62.17 \pm 3.74 ^b	54.97 \pm 2.04 ^b
TC ($\times 10^3/\mu\text{l}$)	36.70 \pm 0.56 ^a	33.10 \pm 0.83 ^b	36.60 \pm 1.32 ^a

Means with different alphabetical characters within the same parameters represent statistical differences ($P < 0.05$).


of TC ($33.10 \times 10^3 \mu\text{l}$) than that of the control group ($36.70 \times 10^3 \mu\text{l}$).

Oxidative Erythrocyte Stress

A significant increase of the total reducing capability (FRAP method) of plasma was demonstrated (Figure 2) on sea bass fed for 2 months with the whole *A. taxiformis*. The results obtained with sea bass, that are almost completely superimposable with those obtained with sea bream, are depicted.

This result seems to be confirmed by the experimental evidences obtained by: (i) the lipid peroxidation tests (Figure 3A), (ii) the trials of integrity of membrane, based on determination of acetylcholinesterase activity (Figure 3B), (iii) the quantification of met-Hb formed (Figure 3C) and (iv) the total thiols contents in both serum and cytoplasm of erythrocytes (Figures 3D,E). In our experiment we did not found significant variation between fish fed with the whole *A. taxiformis* and control fish.

When the main endogenous antioxidants were evaluated in fish treated with alga, the amounts of reduced glutathione and

**FIGURE 2 | Determination of FRAP in the plasma performed on control (a) or on specimens of sea bass fed with algae (b) ($n = 8$). **Showed a significant difference $P < 0.05$.**

of nitrous metabolites showed no detectable variations in both plasma and cytoplasm of erythrocytes (Figures 4A–D).



Thus, fish fed with whole *A. taxiformis*, did not show significant alterations in the physiological and biochemical processes analyzed in sea bass and sea bream.

DISCUSSION

Marine algal compounds represent a source of novel bioactive metabolites with unexpected potentialities. The synthesis of antimicrobial compounds by seaweeds may vary with geographic location, ecological factors, and physiological conditions (active growth or sexual maturity) (Genovese et al., 2012 and reference listed therein).

Depending on the period of sampling, probably due to a different content of biomolecules, ethanolic extracts of *A. taxiformis* exhibited significantly different inhibitory activity against fish pathogenic bacteria. The extract obtained in late spring was active against the whole panel of pathogens, including *P. damsela* subsp. *damsela* and *P. damsela* subsp. *piscicida*, considered resistant to most antibiotics, *V. harveyi*, an emerging pathogen of fish and shrimps, and *V. parahaemolyticus* and *V. vulnificus*, recognized as causative agents of human diseases associated with the consumption of raw shellfish. Seasonal differences in the production of bioactive compounds could be associated with the maximum growth period of algae (Hornsey and Hide, 1974), which actually occurs in late spring for *A. taxiformis* at our latitude. The antibacterial activity (evaluated as inhibition zones and MIC values) of crude extracts here reported against *Aer. salmonicida*, *P. damsela* subsp. *damsela*, *P. damsela* subsp. *piscicida* and *V. harveyi* was lower than that of antibiotics used as control. However, future investigations should attempt to purify active compounds from crude extracts and to identify their chemical composition.

The whole alga and its extracts did not show any cytotoxicity effects on fish hepatocytes, confirming our results previously reported (Genovese et al., 2012).



S. aurata fed with whole algae and algal extracts showed significant decreased in RBC, Hgb, and hematocrit (%) (Hct) in comparison with those of the control group. These results are in accordance with those for European Sea bass juveniles fed garlic powder diets at dietary inclusion levels of 4 and 6%, reported by Irkin et al. (2014), the only study available to date for a comparison. The numbers of white blood cells (WBC), which have important implications for the immune response and the ability of the animal to fight infection, significantly decreased in both *D. labrax* and *S. aurata* treated with whole algae and algal extracts, as compared to the control group. In

S. aurata, the results showed a statistically significant lower WBC values in experimental groups (G10 and G11) respect to control group, whereas TC showed statistically significant lower values in G10 respect to G11 and control group. In fish WBC together with TC play an important role for the defense response (Stosik et al., 2001; Kollner et al., 2004; Tavares-Dias and Moraes, 2004; Passantino et al., 2005; Tavares-Dias et al., 2007, 2008; Martins et al., 2008; Prasad and Charles, 2010; Prasad and Priyanka, 2011), therefore these findings suggest a negative effect of algae at this specify dose in body's immune system. Observed negative impacts of algal extracts on the

hematological parameters, would lead to discourage the use of such compounds as food supplements. Nevertheless, such negative effect could be related to a dose/response effect, so that it is possible that further trials could detect an hormetic response after lower dose exposure (Calabrese and Baldwin, 2002; Calabrese, 2008). Moreover, both positive and negative effects on total WBC count have been registered in fish fed with different concentrations of *Fusarium moniliforme* (Lumlertdacha et al., 1995).

The individuation of the minimum effective dose will let start the future precompetitive development of the algal extracts for a field evaluation on a wide scale and the full application in aquaculture.

Differently, fish fed with both alga and algal extracts maintained high levels of endogenous antioxidants, evaluated as total amount of reduced glutathione, in serum and in the cytoplasm of erythrocytes, indicating a lower inclination to suffer oxidative damages of all the compounds tested. The presence of bioactive substances in the algal extracts could allow the fish to maintain a high reducing redox state, eliciting a cellular welfare ready to contrast the oxidative injuries.

The supplementation with alga and algal extracts did not modify growth rate (data not shown) and zootechnical performances, even if the short period of the challenges (2 weeks/2 months) did not allow us to definitely assess the nutritional significance of the algal supplements. Further long time investigations are needed to clarify the nutritional potential of *A. taxiformis* and its extracts on Mediterranean marine fish species.

REFERENCES

- Allmendinger, A., Spavieri, J., Kaiser, M., Casey, R., Hingley-Wilson, S., Lalvani, A., et al. (2010). Antiprotozoal, antimycobacterial and cytotoxic potential of twenty-three British and Irish red algae. *Phytother. Res.* 24, 1099–1103. doi: 10.1002/ptr.3094
- Andreakis, N., Costello, P., Zanolla, M., Saunders, G. W., and Mata, L. (2015). Endemic or introduced? phylogeography of *Asparagopsis* (Florideophyceae) in Australia reveals multiple introductions and a new mitochondrial lineage. *J. Phycol.* 52, 141–147. doi: 10.1111/jpy.12373
- Andreakis, N., Procaccini, G., Maggs, C., and Kooistra, W. H. C. F. (2007). Phylogeography of the invasive seaweed *Asparagopsis* (Bonnemaisoniales, Rhodophyta) reveals cryptic diversity. *Mol. Ecol.* 16, 2285–2299. doi: 10.1111/j.1365-294X.2007.03306.x
- Ballesteros, E., Martin, D., and Uriz, M. J. (1992). Biological activity of extracts from some mediterranean macrophytes. *Bot. Mar.* 35, 481–485. doi: 10.1515/botm.1992.35.6.481
- Bansemir, A., Blume, M., Schröder, S., and Lindequist, U. (2006). Screening of cultivated seaweeds for antibacterial activity against fish pathogenic bacteria. *Aquaculture* 252, 79–84. doi: 10.1016/j.aquaculture.2005.11.051
- Barman, D., Nen, P., Mandal, S. C., and Kumar, V. (2013). Aquaculture health management: a new approach. *J. Mar. Sci. Res. Dev.* 3:139. doi: 10.4172/2155-9910.1000139
- Barrington, K., Chopin, T., and Robinson, S. (2009). Integrated multi-trophic aquaculture (IMTA) in marine temperate waters. Integrated mariculture: a global review. *FAO Fish. Aquacul. Tech. Paper* 529, 7–46.
- Blunt, J. W., Copp, B. R., Hu, W. P., Munro, M. H. G., Northcote, P. T., and Prinsep, M. R. (2008). Marine natural products. *Nat. Prod. Rep.* 25, 35–94. doi: 10.1039/b701534h

Our findings suggest that the whole alga *A. taxiformis* and its extracts, with demonstrated antibacterial activity, could be proposed for future applications in aquaculture, as alternative tool in contrasting most of common fish pathogens. Furthermore, the results obtained could be used to create in the next future a new industrial field for the production of supplements useful for a modern and eco-sustainable fish farming.

AUTHOR CONTRIBUTIONS

FM and GD were involved *in vivo* experiments; CG and ASp were responsible for *in vitro* antimicrobial evaluation; CF performed cytotoxicity experiments; MM and GG selected algal species for *in vitro* and *in vivo* evaluation; AR and DB carried out the Evaluation of oxidative erythrocyte stress; FF accomplished the hematological analyses, whereas ASa reviewed the manuscript.

FUNDING

Project IZS SI 04 2014, Department of Chemical, Biological, Pharmaceutical and Environmental Sciences, University of Messina.

ACKNOWLEDGMENTS

All the research activities have been carried out under the financial support obtained from the project PON INNOVAQUA (PON02_00451_3362185) from Italian Ministry of University and Research (MIUR).

- Bouhlal, R., Riadi, H., Martínez, J., and Bourgougnon, N. (2010). The antibacterial potential of the seaweeds (Rhodophyceae) of the strait of gibraltar and the mediterranean coast of morocco. *Afr. J. Biotechnol.* 9, 6365–6372. doi: 10.5897/AJB09.1911
- Burreson, B. J., Moore, R. E., and Roller, P. (1975). Haloforms in the essential oil of the alga *Asparagopsis taxiformis* (Rhodophyta). *Tetrahedron Lett.* 16, 473–476. doi: 10.1016/S0040-4039(00)71897-1
- Burreson, B. J., Moore, R. E., and Roller, P. P. (1976). Volatile halogen compounds in the alga *Asparagopsis taxiformis* (Rhodophyta). *J. Agric. Food Chem.* 24, 856–861. doi: 10.1021/jf60206a040
- Calabrese, E. J. (2008). Hormesis: why it is important to toxicology and toxicologists. *Environ. Toxicol. Chem.* 27, 1451–1474. doi: 10.1897/07-541.1
- Calabrese, E. J., and Baldwin, L. A. (2002). Defining hormesis. *Hum. Exp. Toxicol.* 21, 91–97. doi: 10.1191/0960327102ht217oa
- Carelli-Alinovi, C., Ficarra, S., Russo, A., Giunta, E., Barreca, D., Galtieri, A., et al. (2016). Involvement of acetylcholinesterase and protein kinase C in the protective effect of caffeine against b-amyloid-induced alterations in red blood cells. *Biochimie* 121, 52–59. doi: 10.1016/j.biochi.2015.11.022
- Chanda, S., Dave, R., Kaneria, M., and Nagani, K. (2010). “Seaweeds: a novel, untapped source of drugs from sea to combat Infectious diseases,” in *Current Research Technology and Education Topics in Applied Microbiology and Microbial Biotechnology*, ed A. Méndez-Vilas (Badajoz: Formatex Research Center), 473–480.
- Cumashi, A., Ushakova, N. A., Preobrazhenskaya, M. E., D’Incecco, A., Piccoli, A., Totani, L., et al. (2007). A comparative study of the anti-inflammatory, anticoagulant, antiangiogenic, and antiadhesive activities of nine different fucoidans from brown seaweeds. *Glycobiology* 17, 541–552. doi: 10.1093/glycob/cwm014

- de Jesus Raposo, M. F., de Morais, A. M. B., and de Morais, R. M. S. C. (2015). Marine polysaccharides from algae with potential biomedical applications. *Mar. Drugs* 13, 2967–3028. doi: 10.3390/md13052967
- Dhargalkar, V. K., and Verlecar, X. N. (2009). Southern Ocean seaweeds: a resource for exploration in food and drugs. *Aquaculture* 287, 229–242. doi: 10.1016/j.aquaculture.2008.11.013
- El-Baroty, G. S., Moussa, M. Y., Shallah, M. A., Ali, M. A., Sabh, A. Z., and Shalaby, E. A. (2007). Contribution to the aroma, biological activities, minerals, protein, pigments and lipid contents of the red alga: *Asparagopsis taxiformis* (Delile) Trevisan. *J. Appl. Sci. Res.* 3, 1825–1834.
- Ellory, J. C., Hall, A. C., and Stewart, G. W. (1985). Volume-sensitive cation fluxes in mammalian red cells. *Mol. Physiol.* 8, 235–246.
- European Committee for Antimicrobial Susceptibility Testing (EUCAST) of the European Society of Clinical Microbiology and Infectious Diseases (ESCMID) (2003). Determination of minimum inhibitory concentrations (MICs) of antibacterial agents by broth dilution. *Clin. Microbiol. Infect.* 9, 1–7. doi: 10.1046/j.1469-0691.2003.00790.x
- Faggio, C., Piccione, G., Marafioti, S., Arfuso, F., Trischitta, F., Fortino, G., et al. (2014). Monthly variations of haematological parameters of *Sparus aurata* and *Dicentrarchus labrax* reared in Mediterranean land off-shore tanks. *Cah. Biol. Mar.* 55, 437–443.
- Faggio, C., Torre, A., Pelle, E., Raffa, F., Villari, V., and Trischitta, F. (2011). Cell volume regulation following hypotonic shock in hepatocytes isolated from *Sparus aurata*. *Comp. Biochem. Physiol.* 158, 143–149. doi: 10.1016/j.cbpa.2010.10.002
- Fazio, F., Faggio, C., Marafioti, S., Torre, A., Sanfilippo, M., and Piccione, G. (2012b). Comparative study of hematological profile on *Gobius niger* in two different habitat sites: Faro Lake and Tyrrhenian Sea. *Cah. Biol. Mar.* 53, 213–219.
- Fazio, F., Filiciotto, F., Marafioti, S., Di Stefano, V., Assenza, A., Placenti, F., et al. (2012a). Automatic analysis to assess hematological parameters in farmed gilthead sea bream (*Sparus aurata* Linnaeus, 1785). *Mar. Freshw. Behav. Physiol.* 45, 63–73. doi: 10.1080/10236244.2012.677559
- Fazio, F., Marafioti, S., Torre, A., Sanfilippo, M., Panzera, M., and Faggio, C. (2013). Haematological and serum protein profiles of *Mugil cephalus*: effect of two different habitats. *Ichthyol. Res.* 60, 36–42. doi: 10.1007/s10228-012-0303-1
- Fazio, F., Saoca, C., Casella, S., Fortino, G., and Piccione, G. (2015). Relationship between blood parameters and biometric indices of *Sparus aurata* and *Dicentrarchus labrax* cultured in onshore tanks. *Mar. Freshw. Behav. Physiol.* 48, 289–296. doi: 10.1080/10236244.2015.1041239
- Ficarra, S., Misiiti, F., Russo, A., Carelli-Alinovi, C., Bellocco, E., Barreca, D., et al. (2013). Antiepileptic carbamazepine drug treatment induces alteration of membranes in red blood cells: possible positive effects on metabolism and oxidative stress. *Biochimie* 95, 833–841. doi: 10.1016/j.biochi.2012.11.018
- Ficarra, S., Tellone, E., Pirolli, D., Russo, A., Barreca, D., Galtieri, A., et al. (2016). Insights into the properties of the two enantiomers of trans- δ -viniferin, a resveratrol derivative: antioxidant activity, biochemical and molecular modeling studies of its interactions with hemoglobin. *Mol. BioSyst.* 12, 1276–1286. doi: 10.1039/C5MB00897B
- Fouladvand, M., Barazesh, A., Farokhzad, F., Malekizadeh, H., and Sartavi, K. (2011). Evaluation of *in vitro* anti-Leishmanial activity of some brown, green and red algae from the Persian Gulf. *Eur. Rev. Med. Pharmacol. Sci.* 15, 597.
- Genovese, G., Faggio, C., Gugliandolo, C., Torre, A., Spanò, A., Morabito, M., et al. (2012). *In vitro* evaluation of antibacterial activity of *Asparagopsis taxiformis* from the Straits of Messina against pathogens relevant in aquaculture. *Mar. Environ. Res.* 73, 1–6. doi: 10.1016/j.marenvres.2011.10.002
- Genovese, G., Tedone, L., Hamann, M. T., and Morabito, M. (2009). The Mediterranean red alga *Asparagopsis*: a source of compounds against leishmaniasis. *Mar. Drugs* 7, 361–366. doi: 10.3390/md7030361
- Gugliandolo, C., Irrera, G. P., Lentini, V., and Maugeri, T. L. (2008). Pathogenic *Vibrio*, *Aeromonas* and *Arcobacter* spp. associated with copepods in the Straits of Messina (Italy). *Mar. Poll. Bull.* 56, 580–606. doi: 10.1016/j.marpolbul.2007.12.001
- Gugliandolo, C., Lentini, V., Spanò, A., and Maugeri, T. L. (2011). Conventional and molecular methods to detect bacterial pathogens in mussels. *Lett. Appl. Microbiol.* 52, 15–21. doi: 10.1111/j.1472-765X.2010.02959.x
- Hoffmann, E. K. (1984). Volume-induced increase of K^+ and Cl^- permeabilities in Ehrlich ascites tumor cells. Role of internal Ca^{++} . *J. Memb. Biol.* 78, 211–222. doi: 10.1007/BF01925969
- Hornsey, I. S., and Hide, D. (1974). The production of antimicrobial compounds by British marine algae. I. Antibiotic-producing marine algae. *Br. Phycol. J.* 9, 353–361. doi: 10.1080/00071617400650421
- Irkin, L. C., Yigit, M., Yilmaz, S., and Maita, M. (2014). Toxicological evaluation of dietary garlic (*Allium sativum*) powder in European sea bass *Dicentrarchus labrax* juveniles. *Food Nutr. Sci.* 5, 989–996. doi: 10.4236/fns.2014.511109
- Jiao, G., Yu, G., Zhang, J., and Ewart, H. S. (2011). Chemical structures and bioactivities of sulfated polysaccharides from marine algae. *Mar. Drugs* 9, 196–223. doi: 10.3390/md9020196
- Kamenarska, Z., Serkedjieva, J., Najdenski, H., Stefanov, K., Tsvetkova, I., Dimitrova-Konaklieva, S., et al. (2009). Antibacterial, antiviral, and cytotoxic activities of some red and brown seaweeds from the Black Sea. *Bot. Mar.* 52, 80–86. doi: 10.1515/BOT.2009.030
- Kollner, B., Fischer, U., Rombout, J. H. W. M., Taverne-Thiele, J. J., and Hansen, J. D. (2004). Potential involvement of rainbow trout thrombocytes in immune functions: a study using a panel of monoclonal antibodies and RT-PCR. *Dev. Comp. Immunol.* 28, 1049–1062. doi: 10.1016/j.dci.2004.03.005
- Kraan, S., and Barrington, K. A. (2005). Commercial farming of *Asparagopsis armata* (Bonnemaisoniaceae, Rhodophyta) in Ireland, maintenance of an introduced species? *J. Appl. Phycol.* 17, 103–110. doi: 10.1007/s10811-005-2799-5
- Liao, W.-R., Lin, J.-Y., Shieh, W.-Y., Jeng, W.-L., and Huang, R. (2003). Antibiotic activity of lectins from marine algae against marine vibrios. *J. Ind. Microbiol. Biotechnol.* 30, 433–439. doi: 10.1007/s10295-003-0068-7
- Lumlerdacha, S., Lovell, R. T., Shelby, R. A., and Kempainen, B. W. (1995). Growth, hematology, and histopathology of channel catfish, *Ictalurus punctatus*, fed toxins from *Fusarium moniliforme*. *Aquaculture* 130, 201–218. doi: 10.1016/0044-8486(94)00219-E
- Maddison, D. R., and Maddison, W. P. (2000). *MacClade 4: Analysis of Phylogeny and Character Evolution, 4.0.*, Sunderland, MA: Sinauer Associates, Inc.
- Manghisi, A., Morabito, M., Bertuccio, C., Le Gall, L., Couloux, A., Cruaud, C., et al. (2010). Is routine DNA barcoding an efficient tool to reveal introductions of alien macroalgae? A case study of *Agardhiella subulata* (Solieriaceae, Rhodophyta) in cape peloro lagoon (Sicily, Italy). *Cryptogam. Algol.* 31, 423–433.
- Manilal, A., Selvin, J., and George, S. (2012). *In vivo* therapeutic potentiality of red seaweed, *Asparagopsis* (Bonnemaisoniales, Rhodophyta) in the treatment of Vibriosis in *Penaeus monodon* Fabricius. *Saudi J. Biol. Sci.* 19, 165–175. doi: 10.1016/j.sjbs.2011.12.003
- Martins, M. L., Mourinho, J. L. P., Amaral, G. V., Vieira, F. N., Dotta, G., Jatobá, A. M. B., et al. (2008). Haematological changes in *Nile tilapia* experimentally infected with *Enterococcus* spp. *Braz. J. Biol.* 68, 631–637. doi: 10.1590/S1519-69842008000300025
- Mata, L., Gaspar, H., and Santos, R. (2012). Carbon/nutrient balance in relation to biomass production and halogenated compound content in the red alga *Asparagopsis taxiformis* (Bonnemaisoniaceae). *J. Phycol.* 48, 248–253. doi: 10.1111/j.1529-8817.2011.01083.x
- Matozzo, V., Pagano, M., Spinelli, A., Caicci, F., and Faggio, C. (2016). *Pinna nobilis*: a big bivalve with big haemocytes? *Fish Shellfish Immunol.* 55, 529–534. doi: 10.1016/j.fsi.2016.06.039
- Mayer, A. M. S., Rodríguez, A. D., Berlink, R. G. S., and Fusetani, N. (2011). Marine pharmacology in, 2007–8: marine compounds with antibacterial, anticoagulant, antifungal, anti-inflammatory, antimalarial, antiprotozoal, antituberculosis, and antiviral activities; affecting the immune and nervous system, and other miscellaneous mechanisms of action. *Comp. Biochem. Physiol.* 153, 191–222. doi: 10.1016/j.cbpc.2010.08.008
- McConnell, O., and Fenical, W. (1977). Halogen chemistry of the red alga *Asparagopsis*. *Phytochemistry* 16, 367–374. doi: 10.1016/0031-9422(77)80067-8
- McKey, D. (1979). “The distribution of secondary compounds within plant,” in *Herbivores: Their Interactonion with Secondary Plant Metabolites*, eds G. A. Rosenthal and D. H. Janzen (San Diego, CA: Academic Press), 56–133.
- Newman, D. J., Cragg, G. M., and Snader, K. M. (2003). Natural products as sources of new drugs over the period, 1981–2002. *J. Nat. Prod.* 66, 1022–1037. doi: 10.1021/np030096l

- Ollivier, H., Pichavant-Rafini, K., Puill-Stephan, E., Calves, P., Nonnote, L., and Nonnote, G. (2006). Effects of hyposmotic stress on exocytosis in isolated turbot, *Scophthalmus maximus*, hepatocytes. *J. Comp. Physiol. B* 176, 643–652. doi: 10.1007/s00360-006-0087-6
- Passantino, L., Cianciotta, A., Patruno, R., Ribaud, M. R., Jirillo, E., and Passantino, G. F. (2005). Do fish thrombocytes play an immunological role? Their cytoenzymatic profiles and function during an accidental piscine candidiasis in aquarium. *Immunopharmacol. Immunotoxicol.* 27, 345–356. doi: 10.1081/IPH-200067959
- Paul, N. A., Cole, L., de Nys, R., and Steinberg, P. D. (2006). Ultrastructure of the gland cells of the red alga *Asparagopsis armata* (Bonnemaisoniaceae). *J. Phycol.* 42, 637–645. doi: 10.1111/j.1529-8817.2006.00226.x
- Prasad, G., and Charles, S. (2010). Haematology and leucocyte enzyme cytochemistry of a threatened yellow catfish *Horabagrus brachysoma* (Gunther, 1864). *Fish. Physiol. Biochem.* 36, 435–443. doi: 10.1007/s10695-009-9313-y
- Prasad, G., and Priyanka, G. L. (2011). Effect of fruit rind extract of *Garcinia gummi-gutta* on haematology and plasma biochemistry of catfish *Pangasianodon hypophthalmus*. *Asian J. Biochem.* 6, 240–251. doi: 10.3923/ajb.2011.240.251
- Repetto, G., Del Peso, A., and Zurita, J. L. (2008). Neutral red uptake assay for the estimation of cell viability/cytotoxicity. *Nat. Protoc.* 3, 1125–1131. doi: 10.1038/nprot.2008.75
- Salvador, N., Gomez Garreta, A., Lavelli, L., and Ribera, M. A. (2007). Antimicrobial activity of Iberian macroalgae. *Sci. Mar.* 71, 101–113. doi: 10.3989/scimar.2007.71n1101
- Saunders, G. W. (2005). Applying DNA barcoding to red macroalgae: a preliminary appraisal holds promise for future applications. *Philos. Trans. R. Soc. Lond. B* 360, 1879–1888. doi: 10.1098/rstb.2005.1719
- Scala, A., Ficarra, S., Russo, A., Barreca, D., Giunta, E., Galtieri, A., et al. (2015). A new erythrocyte-based biochemical approach to predict the antiproliferative effects of heterocyclic scaffolds: the case of indolone. *Biochim. Biophys. Acta* 1850, 73–79. doi: 10.1016/j.bbagen.2014.09.022
- Shanmughapriya, S., Manilal, A., Sujith, S., Selvin, J., Kiran, G., and Natarajaseenivasan, K. (2008). Antimicrobial activity of seaweeds extracts against multiresistant pathogens. *Ann. Microbiol.* 58, 535–541. doi: 10.1007/BF03175554
- Smit, A. J. (2004). Medicinal and pharmaceutical uses of seaweed natural products: a review. *J. Appl. Phycol.* 16, 245–262. doi: 10.1023/B:JAPH.0000047783.36600.ef
- Stosik, H., Deptula, W., and Travnicek, M. (2001). Studies on the number and ingesting ability of thrombocytes in sick carps (*Cyprinus carpio*). *Vet. Med. Czech.* 46, 12–16.
- Swofford, D. L. (2002). *PAUP*. Phylogenetic Analysis Using Parsimony (*and Other Methods)*, 4.0b10. Sunderland, MA: Sinauer Associates Inc.
- Talarico, L. B., Zibetti, R. G. M., Faria, P. C. S., Scolaro, L. A., Duarte, M. E. R., Noseda, M. D., et al. (2004). Anti-herpes simplex virus activity of sulfated galactans from the red seaweeds *Gymnogongrus griffithsiae* and *Cryptonemia crenulata*. *Int. J. Biol. Macromol.* 34, 63–71. doi: 10.1016/j.ijbiomac.2004.03.002
- Tavares-Dias, M., Barcellos, J. F. M., Marcon, J. L., Menezes, G. C., Ono, E. A., and Affonso, E. G. (2007). Hematological and biochemical parameters for the pirarucu *Arapaima gigas* Schinz, 1822 (Osteoglossiformes, Arapaimidae) in net cage culture. *Electron. J. Ichthyol.* 2, 61–68.
- Tavares-Dias, M., and Moraes, F. R. (2004). *Hematology of Teleosts Fish, 1st Edn.*, Ribeirão Preto: Villimpress (In Portuguese).
- Tavares-Dias, M., Moraes, F. R., and Martins, M. L. (2008). Hematological assessment in four Brazilian teleost fish with parasitic infections, collected in fee fishing from Franca, São Paulo, Brazil. *Bol. Inst. Pesca.* 34, 189–196.
- Torre, A., Trischitta, F., Corsaro, C., Mallamace, D., and Faggio, C. (2013). Digestive cells from *Mytilus galloprovincialis* show a partial regulatory volume decrease following acute hypotonic stress through mechanisms involving inorganic ions. *Cell Biochem. Funct.* 31, 489–495. doi: 10.1002/cbf.2925
- Trischitta, F., Denaro, M. G., and Faggio, C. (2004). Ion transport in the intestine of *Gobius niger* in both isotonic and hypotonic conditions. *J. Exp. Zool. A* 301, 49–62. doi: 10.1002/jez.a.20002
- Trischitta, F., Denaro, M. G., and Faggio, C. (2005). Cell volume regulation following hypotonic stress in the intestine of the eel, *Anguilla anguilla*, is Ca²⁺-dependent. *Comp. Biochem. Physiol. B.* 140, 359–367. doi: 10.1016/j.cbpc.2004.10.011
- Val, A., Platas, G., Basilio, A., Cabello, A., Gorrochategui, J., Suay, I., et al. (2001). Screening of antimicrobial activities in red, green and brown macroalgae from Gran Canaria (Canary Islands, Spain). *Int. Microbiol.* 4, 35–40. doi: 10.1007/s101230100006
- Vallinayagam, K., Arumugam, R., Ragupathi Raja Kannan, R., Thirumaran, G., and Anantharaman, P. (2009). Antibacterial Activity of Some Selected Seaweeds from Pudumadam Coastal Regions. *Global Journal of Pharmacology* 3, 50–52.
- Vonhron-Sénécheau, C., Kaiser, M., Devambe, I., Vastel, A., Mussio, I., and Rusig, A. M. (2011). Antiprotozoal activities of organic extracts from french marine seaweeds. *Mar. Drugs* 9, 922–933. doi: 10.3390/md9060922
- Wolk, C. P. (1968). Role of bromine in the formation of the refractile inclusions of the vesicle cells of the Bonnemaisoniaceae (Rhodophyta). *Planta* 78, 371–375. doi: 10.1007/BF00387095
- Womersley, H. B. S. (1998). *The Marine Benthic Flora of Southern Australia*. Part IIIC. Adelaide, SA: State Herbarium of Southern Australia.
- Woolard, F. X., Moore, R. E., and Roller, P. P. (1979). Halogenated acetic and acrylic acids from the red alga *Asparagopsis taxiformis*. *Phytochemistry* 18, 617–620. doi: 10.1016/S0031-9422(00)84271-5
- Young, D. N. (1977). *Comparative Fine Structure and Histochemistry of Vesiculate Cells in Selected Red Algae*, Ph.D., University of California, Berkeley.
- Zenetos, A., Gofas, S., Verlaque, M., Inar, M. E., Raso, J. E. G., Bianchi, C. N., et al. (2010). Alien species in the Mediterranean Sea by 2010. A contribution to the application of European Union's Marine Strategy Framework Directive (MSFD). Part I. Spatial distribution. *Mediterr. Mar. Sci.* 11, 381–493. doi: 10.12681/mms.87

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2016 Marino, Di Caro, Gugliandolo, Spanò, Faggio, Genovese, Morabito, Russo, Barreca, Fazio and Santulli. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.