



Feeding Relationship between *Octopus vulgaris* (Cuvier, 1797) Early Life-Cycle Stages and Their Prey in the Western Iberian Upwelling System: Correlation of Reciprocal Lipid and Fatty Acid Contents

Sílvia Lourenço^{1,2,3*}, Álvaro Roura^{4,5}, María-José Fernández-Reiriz⁴, Luís Narciso⁶ and Ángel F. González⁴

¹ Interdisciplinary Centre of Marine and Environmental Research, Cruise Terminal of the Port of Leixões, Porto, Portugal, ² Divisão de Serviços de Investigação da Direção Regional das Pescas e Aquacultura da RAM, Centro de Maricultura da Calheta, Calheta, Portugal, ³ Oceanic Observatory of Madeira, Agência Regional para o Desenvolvimento da Investigação Tecnologia e Inovação, Funchal, Portugal, ⁴ Instituto de Investigaciones Marinas (CSIC), Vigo, Spain, ⁵ Department of Ecology, Environment and Evolution, La Trobe University, Melbourne, VIC, Australia, ⁶ Mare—Marine and Environmental Sciences Centre, Faculdade de Ciências da Universidade de Lisboa, Lisbon, Portugal

OPEN ACCESS

Edited by:

Giovanna Ponte,
CephRes and Stazione Zoologica
(SZN), Italy

Reviewed by:

Juan A. Estefanell,
Unemployed, Spain
Oscar Monroig,
University of Stirling, United Kingdom

*Correspondence:

Sílvia Lourenço
slourenco@ciimar.up.pt

Specialty section:

This article was submitted to
Invertebrate Physiology,
a section of the journal
Frontiers in Physiology

Received: 22 February 2017

Accepted: 19 June 2017

Published: 19 July 2017

Citation:

Lourenço S, Roura Á,
Fernández-Reiriz M-J, Narciso L and
González ÁF (2017) Feeding
Relationship between *Octopus
vulgaris* (Cuvier, 1797) Early Life-Cycle
Stages and Their Prey in the Western
Iberian Upwelling System: Correlation
of Reciprocal Lipid and Fatty Acid
Contents. *Front. Physiol.* 8:467.
doi: 10.3389/fphys.2017.00467

Under the influence of the Western Iberian upwelling system, the Iberian Atlantic coast holds important hatcheries and recruitment areas for *Octopus vulgaris*. Recently identified as an octopus hatchery, the Ría de Vigo harbors an important mesozooplankton community that supports *O. vulgaris* paralarvae during the first days of their planktonic stage. This study represents a preliminary approach to determine the nutritional link between wild *O. vulgaris* hatchlings, paralarvae and their zooplankton prey in the Ría de Vigo, by analyzing their lipid class content and fatty acid profiles. The results show that octopus hatchlings are richer in structural lipids as phospholipids and cholesterol, while the zooplankton is richer in reserve lipids like triacylglycerol and waxes. Zooplankton samples are also particularly rich in C18:1n9 and 22:6n3 (DHA), that seem to be successfully incorporated by *O. vulgaris* paralarvae thus resulting in a distinct fatty acid profile to that of the hatchlings. On the other hand, content in C20:4n6 (ARA) is maintained high through development, even though the zooplankton is apparently poorer in this essential fatty acid, confirming its importance for the development of *O. vulgaris* paralarvae. The content in monounsaturated fatty acids, particularly C18:1n7, and the DHA: EPA ratio are suggested as trophic markers of the diet of *O. vulgaris* paralarvae.

Keywords: *Octopus vulgaris*, paralarvae, fatty acids, lipid content, zooplankton, prey-predator relationship

INTRODUCTION

The common octopus (*Octopus vulgaris* Cuvier, 1797) is the most important commercially harvested octopus worldwide. With global landing estimates of 42,457 ton/year (FAO, 2016), it is consumed in many countries of Asia, Latin-America and Mediterranean. The high market demand and value, along with the biological characteristics like the short life span, high growth rates and

high food conversion, makes *O. vulgaris* a desirable species for aquaculture production (Vaz-Pires et al., 2004). However, after decades of research the low paralarvae survival remains an important constraint for industrial farming (Iglesias et al., 2007; Garrido et al., 2016b).

The Iberian Atlantic coast is an important hatchery and recruitment area for *O. vulgaris* (Moreno et al., 2014; Guerra et al., 2015). Here, *O. vulgaris* paralarvae find the optimal environmental conditions to grow favored by the strong summer upwelling (Moreno et al., 2009; Roura et al., 2016). In the first days of life, the paralarvae combine endogenous (yolk) with exogenous feeding, preying mainly upon larval stages of crustaceans of the families Crangonidae, Alpheidae, Brachyura, Paguridae, Thalassinidae, Porcellanidae, Cladocera, Copepoda, and Euphausiidae, but also fish larvae and Cnidaria (Roura et al., 2012; Olmos-Pérez et al., 2017). In fact, during summer, all these potential prey are naturally “enriched” with essential fatty acids (EFA) by the seasonal coastal upwelling where the frequent diatom and dinoflagellate blooms are responsible for the production of polyunsaturated fatty acids (PUFA) as 20:5n3 (eicosapentaenoic acid, EPA) and 22:6n3 (docosahexaenoic acid, DHA). The phytoplankton fatty acid (FA) composition and, particularly, the ratios PUFA (n-3)/(n-6) and EPA/DHA will influence the FA composition of the linked trophic levels like meso- and microzooplankton and planktivorous fishes (Dalsgaard et al., 2003) like *Sardina pilchardus* (Garrido et al., 2008) and to certain extent tissues and eggs of higher trophic levels species as *O. vulgaris* (Lourenço et al., 2014). In fact, the FA composition of muscle and eggs in marine organisms reflects to certain level the biochemical and ecological conditions of ecosystems and can be used to identify food web interactions (Bergé and Barnathan, 2005) being used as qualitative markers, or biomarkers, to trace or confirm predator-prey relationships (Dalsgaard et al., 2003; Budge et al., 2006).

From the metabolic perspective, marine lipids have key roles in the physiology and reproductive processes of heterotrophic organisms. The neutral lipids triacylglycerols and wax esters, are energy reserves that produce free fatty acids through oxidation, which will be incorporated into phospholipids and again in fat reserves (Budge et al., 2006). Phospholipids are the building blocks for the membrane lipid bilayer. The lipids facilitate the absorption of fat-soluble vitamins (e.g., Vitamins A, D, E, and K), and play an important role in the production and regulation of eicosanoids (Bergé and Barnathan, 2005). Cholesterol is the predominant sterol in cephalopod's lipid reserves (Sieiro et al., 2006) and it is precursor of steroid hormones including cortisol, corticosterone, and cortisone. From these, cortisol has an important role in stress responses and is involved in the regulation of the carbohydrates and protein metabolism (Tocher and Glencross, 2015). Despite the low content of lipids in cephalopod body composition (6% dw in muscle, 24% in digestive gland of adults, Sieiro et al., 2006) and 12% dw of the paralarvae (Navarro and Villanueva, 2003), lipids have critical roles in cephalopod metabolism and development (Navarro and Villanueva, 2000; Okumura et al., 2005; Miliou et al., 2006; Seixas et al., 2010; Monroig et al., 2013; Reis et al., 2015). The lipid-rich nervous system of hatchlings represents approximately one

quarter of the animal's fresh weight (Navarro et al., 2014) and the long-chain PUFA, namely EPA, DHA, and C20:4n6 (arachidonic acid, ARA) are identified as EFA for cephalopods, particularly in early life-cycle stages (Monroig et al., 2012; Reis et al., 2014). In fact, several studies have suggested that *O. vulgaris* paralarvae require prey of low lipid content, rich in polar lipids, long-chain PUFA, and cholesterol content (Navarro and Villanueva, 2000, 2003; Okumura et al., 2005; Seixas et al., 2008).

Despite the extended knowledge about the environmental physical factors that drive the distribution, abundance, and recruitment success of *O. vulgaris* paralarvae (González et al., 2005; Otero et al., 2008, 2009; Moreno et al., 2009; Roura et al., 2013, 2016), there are few studies regarding the nutritional profile and requirements of wild *O. vulgaris* paralarvae and their natural prey. In recent years, major efforts have been conducted to understand the nutritional needs for paralarvae in captivity (Garrido et al., 2016b) and their fatty acid profile in the wild (Estefanell et al., 2013; Garrido et al., 2016a), however the nutritional link between them and their prey in natural conditions is still largely unknown.

To fulfill this gap, this study aimed to identify the lipid class content of wild *O. vulgaris* hatchlings and paralarvae and that of their potential preys—i.e., the mesozooplankton community—in the Ría de Vigo (NW Spain). The contents in phospholipids, cholesterol, triacylglycerol, free fatty acids, and wax esters were determined in the mesozooplankton samples and *O. vulgaris* hatchlings samples. The FA profile was evaluated in the mesozooplankton, hatchlings and paralarvae samples in terms of individual FA, saturated FA (SFA), monounsaturated (MUFA), polyunsaturated (PUFA), n-6 highly unsaturated FA (n-6), and n-3 highly unsaturated FA (n-3). Based in significant dissimilarities analyses, trophic markers were selected and compared between the zooplankton, hatchlings and paralarvae, aiming to understand which FA were incorporated into planktonic *O. vulgaris* paralarvae through their diet.

MATERIALS AND METHODS

Zooplankton Sampling

A total of 12 mesozooplankton samples were collected at 5 m depth of the Ría de Vigo (NW Spain, **Figure 1**) in three surveys conducted under the LARECO project (CTM2011-25929) in autumn 2012, September 17th (d1); October 1st (d2); and October 5th (d3) in the outer part of the Ría de Vigo. Samples were collected with a multitrawl (MultiNet[®]) sampler (0.71 × 0.71 m opening frame, 200 μm mesh), East (inn samples) and West of Cies Islands (outer samples) and visually examined on board, looking for *Octopus vulgaris* paralarvae, which were manually sorted. Six zooplankton samples ($n = 6$) were washed with sea water and filtered with a 1,000 μm sieve and frozen at -80°C , freeze dried during 48 h and stored again at -80°C for further analytical methods (see below). The zooplankton size selection was supported by the evidence that *O. vulgaris* paralarvae feed preferentially upon prey >1 mm (Passarella and Hopkins, 1991; Villanueva, 1994; Villanueva et al., 1996; Iglesias et al., 2006; Roura et al., 2010). The remaining samples were fixed in 70% ethanol and then used to identify the




FIGURE 1 | Geographical position of the sampling area in the Ría de Vigo, Spain Northwest Atlantic coast. Legend: lines indicate the sampling transects at east (inner) and west (out) of Cies Islands. The symbol * indicates the sampling site for the *Octopus vulgaris* paralarvae collected during diving.

mesozooplankton community cohabiting with the paralarvae. Organisms were identified under a binocular (Nikon SMZ800) or inverted microscope (Nikon Eclipse TS100) to the lowest possible taxonomic level. The community (holoplankton/meroplankton) ratio was determined based in the number of species identified and classified as holoplankton or meroplankton accordingly to Roura et al. (2013).

The *O. vulgaris* paralarvae collected ($n = 44$) were pooled in a single sample and stored at -20°C in a Methanol: Dichloromethane (2:1) solution to avoid long-term degradation of lipids.

To determine the basal biochemical profile of *O. vulgaris* paralarvae before external feeding, newly hatched paralarvae (hereafter called hatchlings) were obtained from ripe eggs from a single female batch collected by scuba diving in October 9th 2012 off the Ría de Vigo (site coordinates: $42^{\circ}14'N$, $8^{\circ}54'W$, **Figure 1**). The hatchlings were pooled and analyzed in duplicate.

Biochemical Methods

Lipids were first extracted from each zooplankton samples and from the single hatchlings with chloroform: methanol (1:2) and after centrifugation, the precipitate was re-extracted with chloroform: methanol (2:1). Both supernatants were subsequently washed with chloroform: methanol: water (8:4:3) as described by Fernández-Reiriz et al. (1989). Total lipids were quantified following the method described by Marsh and Weinstein (1966) with a tripalmitine standard (Sigma Aldrich Inc., Buchs, Switzerland). Wax esters (WAXES), triglycerides (TAG), free fatty acids (FFA), cholesterol (CHL), and phospholipids (PL) content were determined by thin-layer chromatography (TLC)/densitometry. Silica gel 60 W plates (Merck 16486), with a size of 20×20 cm and a layer thickness of 0.25 mm, were used. Samples were applied by automatic

TLC sampler (Camag 27220). The chromatographic staining was conducted accordingly to Freeman and West (1966). The plates were stained with a 10% CuSO_4 solution in 0.85% H_3PO_4 by heating to 180°C (Bitman and Wood, 1982). Standards employed for the quantitative analysis of the WAXES, TAG, FFA, and CHOL were oleyl oleate, triolein, oleic acid, and cod liver oil (CHOL, Sigma), respectively. A standard obtained from *Mytilus galloprovincialis* was used for PL. The plates were scanned with a Shimadzu CS9000 densitometer, using a monochromatic 370 nm beam of 0.4×0.4 mm working in the zigzag mode, reading the whole spot, and with automatic autozero for baseline correction. All solvents, reagents and fatty acid standards used in this work were of analytic grade (Merck, Darmstadt, and Sigma). FA content of total lipids fraction of zooplankton, *O. vulgaris* hatchlings and paralarvae was determined converting total lipids into FA methyl esters (FAME), accordingly to the method described by Lepage and Roy (1984). Fatty acids methyl esters (FAME) were analyzed by gas chromatography. Peaks corresponding to FAME were identified by comparison of their retention times with standard mixtures and the concentration of each fatty acid or fatty acid group was expressed as % FAME.

Statistical Analysis

Zooplankton samples were identified according to the correspondent transect (out or inn) and sampling day (d1, d2, and d3) resulting in the following sampling code: Out_d1, out_d2, out_d3, inn_d1, inn_d2, and inn_d3. The zooplankton sample composition, lipid classes and FA content were analyzed using metric multidimensional techniques aiming to identify dissimilarities between groups. Prior to analysis, zooplankton abundance data was transformed $\log(x+1)$ and screened to select the taxa that appeared at least in 10% of the samples. Zooplankton dissimilarity matrix was calculated using the

Bray-Curtis dissimilarity index and analyzed with principal coordinate analysis (PCO). The species with highest correlation with the first and second coordinate axes were identified as the potential prey group for the lipid analysis (species highlighted in **Table 1**). The lipid class content and FA with mean concentration higher than 1% FAME were normalized, the similarity matrix was determined using Euclidean distance and analyzed with principal component analysis (PCA) (Zuur et al., 2007). The dimension (axes) eigenvalues and FA scores in each dimension obtained were used to select the FA that explained most of the variance (FA in bold in **Table 2**). The zooplankton species, lipid class and FA groups identified were tested for differences related with sampling area and species composition by non-parametric permutational ANOVA (PERMANOVA) considering type I errors. A constrained canonical analysis (CCA) was applied to the set of zooplankton prey using FA as explanatory variables to identify significant correlations between these FA and the zooplankton species.

Following, the lipid class and FA content were compared between zooplankton, *O. vulgaris* hatchlings and paralarvae applying PCA to determine which lipid classes and FA could differentiate between the groups. The groups identified were tested with PERMANOVA for sample type and sample site to test the significance among groups. CCA was applied to the set of most influential FA, using selected trophic markers as explanatory variables to identify significant differences between the FA profile of prey, hatchlings and paralarvae and the zooplankton species selected. The trophic markers selected were Σ SFA (SFA), Σ MUFA (MUFA), Σ PUFA (PUFA), Σ n-6HUFA (n-6), Σ n-3HUFA (n-3), SFA/PUFA; n-3/n-6; DHA/ARA; DHA/EPA). The metric multidimensional analysis was conducted applying the “envfit” function of VEGAN package in R (Oksanen et al., 2013).

RESULTS

The species composition of the zooplankton in the Ría de Vigo (**Table 1**) showed a dominance of holoplankton both in the inner zone (67.65%) and the outer zone (83.61%) with the copepods *Paracalanus parvus*, *Acartia clausii*, and the euphausiid *Nyctiphanes couchii* being the most frequent species. The meroplankton species contributed with 32.35% in the inner zone and 16.39% in the outer zone, with the most frequent larvae being bivalves and gastropod larvae and cirripeds, mainly in the inner zone stations. The holo/meroplankton ratio of the zooplanktonic community ranged between 1.77 in the outer zone and 3.14 in the inner zone indicating that the two sampling groups belonged to the same coastal community. The inner and outer zone zooplankton communities presented similar total lipids and lipid class content, only differing in the concentration of a single FA. The FA C18:1n7 is particularly high in the zooplankton community of the inner zone (blue arrow in the **Figure 2**). The correlation results showed that C18:1n7 was highly correlated with zoaea of different crustaceans and cnidarians.

The lipid class composition of *Octopus vulgaris* hatchlings was significantly different of that of the zooplankton community

(**Figure 3**). The *O. vulgaris* hatchlings were richer in PL, followed by CHOL and low content in FFA and no TAG and WAXES were detected. In general, hatchlings are richer in FA than zooplankton, in detail, the FA profile of both zooplankton and hatchlings (**Table 2**) showed that, while the two groups had similar content of Σ SFA and Σ PUFA, the zooplankton had higher content of Σ MUFA, particularly in C16:1n7, C18:1n7, and C18:1n9. Despite the similarity in the Σ PUFA, zooplankton samples were richer in EPA, while hatchlings and paralarvae had higher content in ARA and DHA.

PCA showed that the lipid class content allowed to separate zooplankton samples from *O. vulgaris* hatchlings, explaining 95% of the model variation (**Figure 4A**) supported by PERMANOVA, $F = 6.67$, p -value = 0.025, 999 perm). The *O. vulgaris* hatchlings were correlated with higher content of CHOL, while the zooplankton samples were correlated with higher content in TAG, FFA, and WAXES (particularly the sample out_d2). Comparing the FA profile of *O. vulgaris* hatchlings with the zooplankton samples, most FA showed different concentrations with exception of C24:0, C22:1n9, C24:1n9, and C18:2n6. Some FA were only identified in *O. vulgaris* hatchlings as the C18:1n9 and C20:2n6, while others were only identified within the zooplankton samples, like C18:3n3 and C20:4n3. The overall FA profile is significantly different when comparing the zooplankton and the *O. vulgaris* hatchlings (PERMANOVA, $F = 139.29$, p -value = 0.01, 999 perm). The biplot in **Figure 4B** shows that the first axis explained about 96% of the variation observed and the zooplankton samples were correlated with higher content of short-chain C14:0, C16:1n7, and the family of C18:0. However C18:0 was positively correlated with *O. vulgaris* hatchlings, as well as the long-chain FA C20:1n9, C22:5n3, and ARA, and the MUFA C17:1.

By comparing the zooplankton samples with *O. vulgaris* hatchlings, some differences arose. The trophic markers selected to compare zooplankton with *O. vulgaris* hatchlings showed significant differences between the two groups (blue arrows in **Figure 4B**), particularly MUFA, n-3/n-6, DHA/ EPA, and DHA/ARA. For instance, n-3/n-6 is two times higher in the zooplankton prey (12.74 ± 0.99) than in *O. vulgaris* hatchlings (7.27 ± 2.33), which influences in the same degree the DHA/ARA (zooplankton 9.88 ± 2.36 ; hatchlings 4.87 ± 2.00).

The FA profile identified in the sample of 40 paralarvae collected in the wild showed that some of the minority FA (<1% FAME) identified in hatchlings were not identified in this older stage (**Table 2**). It is noteworthy that C16:1n7 and C18:1n9 contents were particularly high in the paralarvae in comparison with that of the hatchlings. The ARA content was identical in both, hatchlings and paralarvae, and higher than the zooplankton samples. The DHA content of paralarvae was identical to the zooplankton and lower to that of the hatchlings. EPA content in wild paralarvae was particularly low in comparison with the other groups. Overall, planktonic *O. vulgaris* showed higher concentrations of Σ MUFAs and lower concentrations of Σ PUFAs when compared with hatchlings. The PCA reflected those differences separating the planktonic paralarvae from hatchlings and zooplankton samples, mainly based in the differences in the content of

TABLE 1 | Mesozooplankton community abundance (n/1,000 m³) and % (in parenthesis).

	species code	out_d1	out_d2	out_d3	inn_d1	inn_d2	inn_d3
HOLOPLANKTON							
<i>Acartia clausi</i>	cop_ac	186 (6.93)	90 (6.52)	288 (10.87)	162 (5.25)	549 (24.05)	468 (18.27)
<i>Calanoides carinatus</i>	cop_cca	72 (2.68)	24 (1.74)	6 (0.23)	24 (0.78)	18 (0.79)	3 (0.12)
<i>Calanus helgolandicus</i>	cop_ch	6 (0.22)	12 (0.87)	12 (0.45)	12 (0.39)	3 (0.13)	
<i>Centropages chierchiae</i>	cop_cch	3 (0.11)				3 (0.13)	
<i>Clausocalanus</i> spp.	cop_cla	48 (1.80)	97 (0.65)	3 (0.11)	21 (0.68)	9 (0.39)	
<i>Corycaeus</i> spp.	cop_cor		18 (1.30)	3 (0.11)		24 (1.05)	36 (1.41)
<i>Paracalanus parvus</i>	cop_pp	624 (23.30)	327 (23.70)	390 (14.72)	183 (5.93)	225 (9.86)	372 (14.52)
<i>Paraeuchaeta hebes</i>	cop_peh	54 (2.01)	3 (0.22)				
<i>Paraeuchaeta</i> sp.	cop_pe	3 (0.11)		6 (0.23)			3 (0.12)
<i>Ctenocalanus vanus</i>	cop_cv	30 (1.12)	6 (0.43)	24 (0.91)		24 (1.05)	
<i>Pseudocalanus elongatus</i>	cop_pce	6 (0.22)	6 (0.43)	12 (0.45)	24 (0.78)	30 (1.31)	3 (0.12)
<i>Subeucalanus crassus</i>	cop_sec	3 (0.11)		3 (0.11)		9 (0.39)	3 (0.12)
<i>Temora longicornis</i>	cop_tl	21 (0.78)	3 (0.22)	93 (3.51)	57 (1.85)	9 (0.39)	3 (0.12)
<i>Candacia armata</i>							3 (0.12)
<i>Oithona plumifera</i>		3 (0.11)	3 (0.22)	12 (0.45)	12 (0.39)	12 (0.52)	6 (0.23)
<i>Oncaea media</i>		21 (0.78)		291 (10.99)	99 (3.21)	6 (0.267)	
Harpacticoida					3 (0.10)	6 (0.26)	3 (0.12)
Copepodid stages	cop	72 (2.68)	42 (3.04)		78 (2.53)	45 (1.97)	6 (0.23)
<i>Evadne nordmanni</i>		306 (11.41)	63 (4.57)	9 (0.34)	831 (26.92)	156 (6.83)	54 (2.11)
<i>Podon intermedius</i>			15 (1.09)		21 (0.68)	42 (1.84)	33 (1.29)
<i>Nyctiphanes couchii</i> calyptopa		51 (1.90)	102 (7.39)	600 (22.65)	144 (4.66)	48 (2.10)	24 (0.94)
<i>Nyctiphanes couchii</i> furcilia	nyc_cou	300 (11.18)	129 (9.35)	99 (3.74)	69 (2.24)	69 (3.02)	222 (8.67)
Mysidacea	mys	3 (0.11)		3 (0.11)			
Cnidaria	cnid			27 (1.02)	15 (0.49)		81 (3.16)
Chaetognatha	chaet	108 (4.03)	21 (1.52)	72 (2.72)	66 (2.14)	84 (3.68)	90 (3.51)
Syphonophora	syph	33 (1.23)		30 (1.13)	96 (3.11)		
Tunicata	salp	81 (2.93)	12 (0.86)	30 (1.14)	117 (3.65)	63 (2.68)	156 (5.74)
Platyhelminthes							12 (0.47)
MEROPLANKTON							
Amphioxus		3 (0.11)					
Gammaridea					6 (0.19)	3 (0.13)	3 (0.12)
Cirripida cipris	cirripid	57 (2.12)	3 (0.22)		27 (0.87)	15 (0.66)	24 (0.94)
Polichaeta larvae	polich		9 (0.65)	12 (0.45)	6 (0.19)	12 (0.53)	3 (0.12)
Bivalvia larvae		429 (16.00)	381 (27.61)	312 (11.78)	699 (22.64)	582 (25.49)	642 (25.06)
Gastropoda		57 (2.13)	69 (5.00)	117 (4.41)	84 (2.72)	138 (6.04)	243 (9.48)
Ophiuridea larvae		45 (1.68)	3 (0.22)	3 (0.11)	159 (5.15)	6 (0.26)	
Equinoidea larvae					9 (0.29)		
Cirripida nauplius		18 (0.67)	9 (0.65)	171 (6.46)	51 (1.65)	27 (1.18)	21 (0.82)
Brachyura zoeae	brach_zoea	30 (1.13)	6 (0.43)	15 (0.57)	6 (0.19)	24 (1.05)	18 (0.70)
Crangonidae zoeae	crang_zoea					12 (0.53)	
Paguridae zoeae	pag_zoea				3 (0.10)		6 (0.23)
Palaemonidae zoeae	palam_zoea			3 (0.11)		6 (0.26)	3 (0.12)
Bryozan larvae		6 (0.22)	12 (0.87)			15 (0.66)	9 (0.35)
<i>Pisidia longicornis</i> zoeae	p_long_zoea					3 (0.13)	3 (0.12)
<i>Porcellana platycheles</i> zoeae	p_platy_zoea	3 (0.11)	3 (0.22)				6 (0.23)
<i>Processidae</i> zoeae	process_zoea				3 (0.10)	6 (0.26)	
Fish eggs				3 (0.11)			
Fish larvae						3 (0.13)	
Holoplankton/Meroplankton		1.91	1.77	1.83	2.56	2.11	3.14

Out_d1, out_d2, out_d3, inn_d1, inn_d2, and inn_d3 represent the zooplankton samples. Zooplankton species with individuals bigger than 1 mm were selected as prey for *Octopus vulgaris* paralarvae and analyzed for their nutritional profile (identified with a species code). The species in bold were selected for the constrained canonical analysis (CCA).

TABLE 2 | Fatty acid concentration (mean \pm SD, % FA) of zooplankton community and *Octopus vulgaris* hatchlings in the Ria de Vigo.

	Mesozooplankton	<i>O. vulgaris</i>	
		Hatchlings	Paralarvae
Saturated fatty acids (SFA)			
C14:0	5.46 \pm 0.84 ^a	2.47 \pm 0.17 ^b	2.33
C15:0	0.63 \pm 0.10 ^a	0.33 \pm 0.03 ^b	0.65
C16:0	18.88 \pm 1.21 ^a	19.35 \pm 1.46 ^b	21.97
C17:0	1.35 \pm 1.18 ^a	1.42 \pm 0.11 ^b	1.49
C18:0	4.57 \pm 0.41 ^a	9.96 \pm 0.75 ^b	10.58
C24:0	0.77 \pm 0.11 ^a	0.65 \pm 0.06 ^a	
Σ SFA	31.69 \pm 6.97 ^a	34.19 \pm 2.59 ^a	37.02
Monounsaturated fatty acids (MUFA)			
C15:1		0.37 \pm 0.04 ^b	0.46
C16:1n7	7.18 \pm 1.10 ^a	0.54 \pm 0.05 ^b	1.31
C17:1	1.21 \pm 0.20 ^a	3.29 \pm 0.26 ^b	3.03
C18:1n7	3.15 \pm 0.50 ^a	1.67 \pm 0.19 ^b	1.94
C18:1n9	4.61 \pm 0.51 ^a	2.82 \pm 0.25 ^b	7.65
C20:1n9	0.73 \pm 0.41 ^a	4.09 \pm 0.29 ^b	4.61
C22:1n9	0.62 \pm 0.66 ^a	0.90 \pm 0.14 ^a	
C24:1n9	0.65 \pm 0.09 ^a	0.52 \pm 0.03 ^a	0.63
Σ MUFA	19.40 \pm 2.41 ^a	15.66 \pm 1.38 ^b	21.10
Poli-unsaturated fatty acids (PUFA)			
C18:2n6	1.78 \pm 0.04 ^a	0.37 \pm 0.02 ^a	0.67
C18:4n3	2.61 \pm 0.28 ^a	0.39 \pm 0.03 ^b	
C18:3n3	1.41 \pm 0.12		
C20:2n6		0.76 \pm 0.07	
C20:4n6 ¹ (ARA)	1.79 \pm 0.19 ^a	5.32 \pm 2.40 ^b	5.06
C20:4n3	0.81 \pm 0.12		
C20:5n3 (EPA)	22.23 \pm 1.51 ^a	18.43 \pm 1.44 ^b	15.40
C22:5n3	0.90 \pm 0.05 ^a	1.63 \pm 0.13 ^b	1.78
C22:6n3 (DHA)	17.34 \pm 2.69 ^a	23.25 \pm 1.78 ^b	18.97
Σ PUFA	50.17 \pm 8.17 ^a	50.15 \pm 5.86 ^a	41.88
Σ n-6	3.57 \pm 0.16	6.37 \pm 1.63	5.73
n-3/n-6	12.74 \pm 0.16	7.27 \pm 2.33	3.62
DHA/EPA	0.79 \pm 0.16	1.26 \pm 0.00	1.23
EPA/ARA	12.47 \pm 0.76	3.85 \pm 1.56	3.04
DHA/ARA	9.88 \pm 2.36	4.87 \pm 2.00	3.75

¹The FA C20:4n6 and FA C20:3n3 have the same retention time, and the concentration of FA C20:4n6 is dominant in marine products, the concentration presented here is representative of C20:4n6.

Different superscripts indicate significant statistical differences ($p < 0.05$) between mesozooplankton, *O. vulgaris* hatchlings and paralarvae.

C18:1n9 (Figure 4C). PERMANOVA results showed that the FA profile is different between these three groups in terms of both FA identified and FA content ($F = 85.08$, p -value = 0.004, 999 perm). The trophic markers (blue arrows in the Figure 4C biplot) were highly correlated with axis 1 (95% explained variation), indicating that differences found in these trophic markers ratios were more significant between *O. vulgaris* samples and zooplankton samples than between hatchlings and paralarvae.



DISCUSSION

This study represents the first attempt to analyse the FA contents of *O. vulgaris* paralarvae and that of the zooplankton community where they fed on during the first days of their planktonic life. Moreover, a detailed description of the lipid class composition of wild *O. vulgaris* hatchlings and zooplankton was carried out to understand how they differ. Being aware of the seasonal, regional, and sampling limitations, this study still represents an important snapshot on the nutritional support provided by the zooplankton community to the *O. vulgaris* paralarvae. *Octopus vulgaris* paralarvae are lecithotrophic and in the first days of life, their survival depends of the embryonic yolk which nutritional composition is directly influenced by female's diet (Quintana et al., 2015). After some hours or a few days in the water column the paralarvae start to feed, and with 7 days-old (Garrido et al., 2016a) they are able to feed in a large variety of prey from decapod zoaea, krill, fish larvae, cladocerans, copepods, siphonophores, and jellyfish (Roura et al., 2012; Olmos-Pérez et al., 2017). Here, we observed that the FA profiles of wild *O. vulgaris* hatchlings and paralarvae are different from those of the zooplankton. Given that the zooplankton samples analyzed were constituted by numerous phyla, with different FA and lipid class compositions (Dalsgaard et al., 2003), this difference may be the result of the trophic selection displayed by *O. vulgaris* paralarvae (Roura et al., 2016).

The zooplanktonic samples analyzed during this study included a heterogeneous assemblage of organisms dominated by two copepods *Paracalanus parvus* and *Acartia clausi*, the euphausiid *Nyctiphanes couchii*, chaetognaths and small Tunicata. This assemblage was particularly rich in FFA, TAG, and WAXES. The zooplankton accumulates TAG and WAXES, important energy reserves produced by the microalgae during the frequent upwelling events (Lee et al., 2006). The higher content in TAG in these samples is probably related with the presence of meroplankton species in some samples, particularly cirripeds and brachyuran larvae that are known to storage TAG in large lipid globules (Lee et al., 2006), in opposition to the copepod dominated samples richer in WAXES (Lee et al., 1970, 1971).


This zooplankton community was rich in SFA and PUFA because of the dominance of calanoid copepod species. The higher availability of bacteria, detritus, and green algae during autumn may account for the increase of the content in SFA (~30%) and PUFA (~49%) (Falk-Petersen et al., 2002; Gonçalves et al., 2012). Moreover, MUFA, particularly C18:1n7 had an important role in the nutritional characterization of the zooplankton (see Figures 2, 4B). Despite the relatively low content in comparison with other FA like C18:1n9, increasing concentrations of C18:1n7 might be related with higher abundance of the meroplankton fraction in the zooplankton samples, which is characteristic of coastal communities (Roura et al., 2013). We suggest that this FA can be used as a trophic marker evaluating the contribution of holoplankton and meroplankton to the *O. vulgaris* paralarvae diet.

Several authors have previously shown that newly hatched paralarvae have low lipid content with relatively high PL and CHOL and very low TAG contents (Navarro and Villanueva, 2000; Reis et al., 2015). In comparison with the zooplankton



samples, the hatchlings sample presented higher content of PL and CHOL, and lower content in FFA. The CHOL and PL, important components of cell membranes, have origin in the maternal reserves (Quintana et al., 2015) explaining their relative

high content in the hatchlings. On the other hand, we couldn't detect WAX and TAG in hatchling samples, suggesting a very low content as observed in the work of Navarro and Villanueva (2000). These results show that besides the total lipid contents,



is the proportion and content of some lipids classes that have high relevance for the paralarvae (Navarro et al., 2014; Reis et al., 2015). In this transitional phase, the digestive gland is still developing (Moguel et al., 2010) and is not able to store and digest the neutral lipids as TAG and WAX until 12 days after hatching (Martínez et al., 2011), explaining why, despite being highly energetic nutrients, these lipid classes appear in a very low concentration in the paralarvae. In fact, previous studies on paralarvae nutritional requirements that used *Artemia* as live feed, seem to have produced paralarvae with important shifts from the natural nutritional profile of the paralarvae (including high TAG content), resulting in high paralarvae mortality probably due to the poor essential lipid composition of the *Artemia* (Navarro et al., 2014).

Capturing *O. vulgaris* paralarvae in zooplankton samples is challenging, as it occurs with many other cephalopod paralarvae with pelagic stages (Moreno et al., 2009; Roura et al., 2016). *Octopus vulgaris* paralarvae are among the less abundant meroplanktonic organisms in the zooplanktonic community (Roura et al., 2013; Zaragoza et al., 2015) and it is very difficult to collect high numbers of individuals to conduct biochemical analyses. To overcome this problem, the approach adopted in the present study was to pool all the paralarvae collected in a unique sample, losing individual information. Alternatively, Garrido et al. (2016b) using the same collection method (the multinet sampler) analyzed 10 *O. vulgaris* paralarvae individually, resulting in high variability between individuals. Both approaches are valid, however, some differences arise, particularly in the content of C18:1n9, and DHA with obvious reflection in the content in Σ MUFA and Σ PUFA. To decrease the uncertainty associated to the FA profiles obtained from *O. vulgaris* paralarvae from nature, the sampling approach could be improved by conducting triplicate field samples of pooled paralarvae collected under the same environmental conditions. However, this approach would only be viable by means of increasing the chance of collecting paralarvae. This could be achieved by filtering more water using bongo nets (González et al.,

2005; Roura et al., 2016) or by using light traps, which proved to be quite effective in capturing octopod paralarvae off the NW coast of Australia (Jackson et al., 2008).

DHA and C18:1n9 are essential for *O. vulgaris* paralarvae (Monroig et al., 2013; Reis et al., 2015) and the difference found between this and the study conducted by Garrido et al. (2016b) might be associated with the high variability in mesozooplankton community composition (Roura et al., 2013), together with the variety of prey hunt by the paralarvae (Roura et al., 2012; Olmos-Pérez et al., 2017). High C18:1n9 is common in neutral lipids (e.g. TAG, Viciano et al., 2011) accumulated by decapod zoea (see Figure 3; Letessier et al., 2012), one of the preferential prey of *O. vulgaris*, while DHA is associated with dinoflagellate blooms (Dalsgaard et al., 2003) common during autumn (Crespo et al., 2008) and probably dominated in the plankton community during our sampling season.

In marine larvae, SFA and MUFA are the main substrates to incorporate neutral lipids as TAG to satisfy energy demands, while long chain PUFA are preferentially esterified in structural lipids as the phospholipids in cell membranes (Reis et al., 2015). In this study, the paralarvae had higher content of C16:0, C18:0, C16:1n7, and C18:1n9 than hatchlings. This accumulation in SFA and MUFA was probably related with the diet rich in decapod zoea and other omnivorous and carnivorous holo and meroplankton rich in TAG, consequently in TAG and MUFA (Dalsgaard et al., 2003; Lee et al., 2006). Moreover, ARA content of the paralarvae was similar to that observed in the hatchlings and significantly higher to the prey. The high ARA content was already observed in the mature ovary of females (Rosa et al., 2004; Loureço et al., 2014; Estefanell et al., 2015) and in hatchlings collected off the Gran Canaria Island (Estefanell et al., 2013). Reis et al. (2015) proved that ARA is efficiently incorporated by the paralarvae. In fact, exists a competition mechanism of incorporation of ARA and EPA that are esterified by the same enzymes, and it is this mechanism that is responsible of the high variability in the EPA/ARA obtained for paralarvae in different studies ranging between 0.95 (for paralarvae fed with *Grapsus*

adconsionis in Reis et al., 2015), 2.7 in Garrido et al. (2016b), and 3.04 in the present study.

Trophic markers are used to follow the interactions between prey and predators in the marine trophic marine web. In this study, CCA results showed that ratio of essential DHA/ARA, DHA/EPA, EPA/ARA, n-3/n-6, SFA/PUFA ratios and the content of Σ MUFA and Σ n-6 allowed the discrimination between preys and predators (Budge et al., 2006). In this context, it would be expected to find similar trophic ratios between prey (zooplankton) and predators (paralarvae). In fact, only the paralarvae content in Σ MUFA, Σ n-6 and the DHA:EPA ratio seemed to follow the prey composition. As occurs with their prey, *O. vulgaris* paralarvae seem to have a lower content in DHA and higher content in Σ MUFA, presenting the same tendency presented by feeding experiments where *O. vulgaris* hatchlings were fed with known prey (Iglesias et al., 2013; Reis et al., 2014). It is noteworthy, that the MUFA C18:1n7 and C18:1n9 showed an increase in relation to hatchlings following the pattern of their prey.

Even though the low number of samples analyzed, we believe that the lipidic profile and trophic ratios determined for *O. vulgaris* hatchlings, paralarvae and their potential prey, allowed a first approach to understand the impact of the available prey pool in the nutritional profile (in terms of lipids) of *O. vulgaris* paralarvae. The impact of feeding in the FA content, particularly C18:1n7, C18:1n9, and DHA is notable, showing that Σ MUFA, DHA/EPA, and C18:1n7 can potentially be used as trophic markers of the diet of *O. vulgaris* paralarvae in the wild. Further biochemical and physiological studies targeting the neutral and polar lipid reserves of wild paralarvae and their prey will certainly untangle the nutritional deficiencies obtained under culture conditions for *O. vulgaris* paralarvae.

AUTHOR CONTRIBUTIONS

SL: Contributed with the conception, sampling design of the work, and with acquisition, analysis, and interpretation of data, manuscript drafting and preparation for submission. ÁR: Contributed to the work with acquisition, analysis, and

interpretation of the data, with critical revision of the manuscript, and final approval for the paper submission. MF: Contributed to the present work with the samples biochemical analysis and interpretation, and for the manuscript critical revision in all aspects considering the lipidic analysis. LN: Contributed for the work conception, critical revision and final approval of the version to be published. ÁG: Contributed for the work conception, critical revision, and final approval of the version to be published.

FUNDING

Data collection and analytical analysis were conducted under the LARECO project (CTM2011-25929) and supported partially by FEDER funds and by the Portuguese Foundation for Science and Technology (FCT), through a doctoral fellowship granted to the first author (SFRH/BD/44182/2008). Presently, SL is supported by Agência Regional para o Desenvolvimento da Investigação Tecnologia e Inovação (ARDITI) through a post-doctoral fellowship under the auspices of Project M14-20 - 09-5369-FSE-000001- Bolsa de Pós- Doutorado.

ACKNOWLEDGMENTS

This work is a contribution to the COST (European COoperation on Science and Technology) Action FA1301 “A network for improvement of cephalopod welfare and husbandry in research, aquaculture and fisheries (CephsInAction).” The authors want to acknowledge the crew of RV “Mytilus” and Marcos Regueira during zooplankton surveys, to Manuel Garcia and Jorge Hernández Urcera, the divers that collected the eggs and to Lourdes Nieto for her availability to help with the biochemical analyses. The authors also want to express their sincere acknowledgments to Juan Carlos Navarro (Instituto de Acuicultura “Torre de la Sal”, IATS), Eduardo Almansa (Centro Oceanográfico de Canarias-Instituto Español de Oceanografía), the two reviewers and to the invited editor Giovanna Ponte for the precious comments during the preparation and submission of the manuscript.

REFERENCES

- Bergé, J.-P., and Barnathan, G. (2005). Fatty acids from lipids of marine organisms: molecular biodiversity, roles as biomarkers, biologically active compounds, and economical aspects, *adv. Biochem. Eng. Biotechnol.* 96, 49–125. doi: 10.1007/b135782
- Bitman, J., and Wood, D. L. (1982). An improved copper reagent for quantitative densitometric thin-layer chromatography of lipids. *J. Liq. Chromatogr.* 5, 1155–1162. doi: 10.1080/01483918208067575
- Budge, S. M., Iverson, S. J., and Koopman, H. N. (2006). Studying trophic ecology in marine ecosystems using fatty acid: a primer on analysis and interpretation. *Mar. Mammal Sci.* 22, 759–801. doi: 10.1111/j.1748-7692.2006.00079.x
- Crespo, B., Teixeira, I., Figueiras, F., and Castro, C. (2008). Microplankton composition off NW Iberia at the end of the upwelling season: source areas of harmful dinoflagellate blooms. *Mar. Ecol. Prog. Ser.* 355, 31–43. doi: 10.3354/meps07261
- Dalsgaard, J., St. John, M., Kattner, G., Müller-Navarra, D., and Hagen, W. (2003). Fatty acid trophic markers in the pelagic marine environment. *Adv. Mar. Biol.* 46, 225–340. doi: 10.1016/S0065-2881(03)46005-7
- Estefanell, J., Socorro, J., Izquierdo, M., and Roo, J. (2015). Effect of two fresh diets and sexual maturation on the proximate and fatty acid profile of several tissues in *Octopus vulgaris*: specific retention of arachidonic acid in the gonads. *Aquacult. Nutr.* 21, 274–285. doi: 10.1111/anu.12163
- Estefanell, J., Socorro, J., Ramirez, B., Izquierdo, M., and Roo, J. (2013). Fatty acid profile in eggs and newly hatched paralarvae of *Octopus vulgaris* collected from the wild and after 1-5 days starvation. *Commun. Agric. Applied Biol. Sci.* 78, 119–122. doi: 10.13140/2.1.4164.8644
- FAO (2016). *Fisheries and Aquaculture software. FishStatJ – Software for Fisheries Statistical Time Series*, in *FAO Fisheries and Aquaculture Department* (online). Rome (cited 12 November 2016).
- Falk-Petersen, S., Dahl, T., Scott, C., Sargent, J. R., Gulliksen, B., Kwasiński, S., et al. (2002). Lipid biomarkers and trophic linkages between ctenophores and copepods in Svalbard waters. *Mar. Ecol. Prog. Ser.* 227, 187–194. doi: 10.3354/meps227187
- Fernández-Reiriz, M. J., Perez-Camacho, A., Ferreiro, M. J., Blanco, J., Planas, M., Campos, M. J., et al. (1989). Biomass production and variation in the biochemical profile (Total protein, carbohydrates, RNA, lipids

- and FA) of seven species of marine microalgae. *Aquaculture* 83, 17–27. doi: 10.1016/0044-8486(89)90057-4
- Freeman, C. T., and West, D. (1966). Complete separation of lipid classes on a single thin-layer plate. *J. Lipid Res.* 7, 324–327.
- Garrido, D., Martín, V. M., Rodríguez, C., Iglesias, J., Navarro, J. C., Estévez, A., et al. (2016a). Meta-analysis approach to the effects of live prey on the growth of *Octopus vulgaris* paralarvae under culture conditions. *Rev. Aquacult.* 1–12. doi: 10.1111/raq.12142
- Garrido, D., Navarro, J. C., Perales-Raya, C., Nande, M., Martín, M. V., Iglesias, J., et al. (2016b). Fatty acid composition and age estimation of wild *Octopus vulgaris* paralarvae. *Aquaculture* 464, 564–569. doi: 10.1016/j.aquaculture.2016.07.034
- Garrido, S., Rosa, R., Ben-Hamadou, R., Cunha, M. E., Chicharo, M. A., and van der Lingen, C. D. (2008). Spatio-temporal variability in fatty acid trophic biomarkers in stomach contents and muscle of iberian sardine (*Sardina pilchardus*) and its relationship with spawning. *Mar. Biol.* 154, 1053–1065. doi: 10.1007/s00227-008-9999-7
- Gonçalves, A. M. M., Azeiteiro, U. M., Pardal, M. A., and De Troch, M. (2012). Fatty acid profiling reveals seasonal and spatial shifts in zooplankton diet in a temperate estuary. *Estuar. Coastal Shelf. Sci.* 109, 70–80. doi: 10.1016/j.ecss.2012.05.020
- González, A. F., Otero, J., Guerra, Á., Prego, R., Rocha, F., and Dale, A. W. (2005). Distribution of common octopus and common squid paralarvae in a wind-driven upwelling area (Ria of Vigo, northwestern Spain). *J. Plankton Res.* 27, 271–277. doi: 10.1093/plankt/fbi001
- Guerra, Á., Hernández-Urcera, J., Garcí, M. E., Sestelo, M., Regueira, M., González, A. F., et al. (2015). Spawning habitat selection by *Octopus vulgaris*: new insights for a more effective management of this resource. *Fish. Res.* 167, 313–322. doi: 10.1016/j.fishres.2015.03.011
- Iglesias, J., Fuentes, L., Sánchez, J., Otero, J. J., Moxica, C., and Lago, M. J. (2006). First feeding of *Octopus vulgaris* cuvier 1797 paralarvae using Artemia: effect of prey size, prey density and feeding frequency. *Aquaculture* 261, 817–822. doi: 10.1016/j.aquaculture.2006.08.002
- Iglesias, J., Pazos, G., Fernández, J., Sánchez, F. J., Otero, J. J., Domingues, P., et al. (2013). The effects of using crab zoeae (*Maja brachydactyla*) on growth and biochemical composition of *Octopus vulgaris* (Cuvier 1797) paralarvae. *Aquacult. Int.* 22, 1041–1051. doi: 10.1007/s10499-013-9725-7
- Iglesias, J., Sánchez, F. J., Bersano, J. G. F., Carrasco, J. F., Dhont, J., Fuentes, L., et al. (2007). Rearing of *Octopus vulgaris* paralarvae: present status, bottlenecks and trends. *Aquaculture* 266, 1–15. doi: 10.1016/j.aquaculture.2007.02.019
- Jackson, G. D., Meekan, M. G., Wotherspoon, S., and Jackson, C. H. (2008). Distribution of young cephalopods in the tropical waters of western Australia over two consecutive summers. *ICES J. Mar. Sci.* 65, 140–147. doi: 10.1093/icesjms/fsm186
- Lee, R. E., Hagen, W., and Kattner, G. (2006). Lipid storage in marine zooplankton. *Mar. Ecol. Prog. Ser.* 307, 273–306. doi: 10.3354/meps307273
- Lee, R. F., Nevenzel, J. C., and Paffenhöfer, G.-A. (1971). Importance of wax esters and other lipids in the marine food chain: phytoplankton and copepods. *Mar. Biol.* 9, 99–108. doi: 10.1007/BF00348249
- Lee, R. F., Nevenzel, J. C., Paffenhöfer, G.-A., and Benson, A. A. (1970). The metabolism of wax esters and other lipids by the marine copepod *Calanus helgolandicus*. *J. Lipid Res.* 11, 237–240.
- Lepage, G., and Roy, C. C. (1984). Improved recovery of fatty acid through direct transesterification without prior extraction or purification. *J. Lipid Res.* 25, 1391–1396.
- Letessier, T., Pond, D., McGill, R., Reid, W., and Brierley, A. (2012). Trophic interaction of invertebrate zooplankton on either side of the charlie gibbs fracture zone/subpolar front of the mid-atlantic ridge. *J. Mar. Syst.* 94, 174–184. doi: 10.1016/j.jmarsys.2011.11.020
- Loureño, S., Narciso, L., Gonzalez, Á. F., Pereira, J., Auborg, S., and Xavier, J. C. (2014). Does the trophic habitat influence the biochemical quality of the gonad of *Octopus vulgaris*? Stable isotopes and lipid class contents as bio-indicators of different life-cycle strategies. *Hydrobiologia* 725, 33–46. doi: 10.1007/s10750-013-1717-0
- Marsh, J. B., and Weinstein, D. B. (1966). Simple charring method for determination of lipids. *J. Lipid Res.* 7, 574–576.
- Martínez, R., López-Ripoll, E., Avila-Poveda, O., Santos-Ricalde, R., Mascaró, M., and Rosas, C. (2011). Cytological ontogeny of the digestive gland in post-hatching *Octopus maya*, and cytological background of digestion in juveniles. *Aquat. Biol.* 11, 249–261. doi: 10.3354/ab00305
- Miliou, H., Fintikaki, M., Tzitzinakis, M., Kountouris, T., and Verriopoulos, G. (2006). Fatty acid composition of the common octopus, *Octopus vulgaris*, in relation to rearing temperature and body weight. *Aquaculture* 256, 311–322. doi: 10.1016/j.aquaculture.2006.02.050
- Moguel, C., Mascaró, M., Avila-Poveda, O., Caamal-Monsreal, C., Sanchez, A., Pascual, C., et al. (2010). Morphological, physiological and behavioral changes during post-hatching development of *Octopus maya* (Mollusca: Cephalopoda) with special focus on the digestive system. *Aquat. Biol.* 9, 35–48. doi: 10.3354/ab00234
- Monroig, Ó., Guinot, D., Hontoria, F., Tocher, D. R., and Navarro, J. C. (2012). Biosynthesis of essential fatty acids in *Octopus vulgaris* (Cuvier, 1797): molecular cloning, functional characterisation and tissue distribution of a fatty acyl elongase. *Aquaculture* 360–361, 45–53. doi: 10.1016/j.aquaculture.2012.07.016
- Monroig, Ó., Tocher, D., and Navarro, J. (2013). Biosynthesis of polyunsaturated fatty acids in marine invertebrates: recent advances in molecular mechanisms. *Mar. Drugs* 11, 3998–4018. doi: 10.3390/md11103998
- Moreno, A., dos Santos, A., Piatkowski, U., Santos, A. M. P., and Cabral, H. (2009). Distribution of cephalopod paralarvae in relation to the regional oceanography of the western Iberia. *J. Plankton Res.* 31, 73–91. doi: 10.1093/plankt/fbn103
- Moreno, A., Lourenço, S., Pereira, J., Gaspar, M. B., Cabral, H. N., and Pierce, G. J., et al. (2014). Essential habitats for pre-recruit *Octopus vulgaris* along the portuguese coast. *Fish. Res.* 152, 74–85. doi: 10.1016/j.fishres.2013.08.005
- Navarro, J. C., and Villanueva, R. (2000). Lipid and fatty acid composition of early stages of cephalopods: an approach to their lipid requirements. *Aquaculture* 183, 161–177. doi: 10.1016/S0044-8486(99)00290-2
- Navarro, J. C., and Villanueva, R. (2003). The fatty acid composition of *Octopus vulgaris* paralarvae reared with live and inert food: deviation from their natural fatty acid profile. *Aquaculture* 219, 613–631. doi: 10.1016/S0044-8486(02)00311-3
- Navarro, J. C., Monroig, O., and Sykes, A. V. (2014). “Nutrition as a key factor for cephalopod aqua-culture,” in *Cephalopod Culture*, eds J. Iglesias, L. Fuentes, and R. Villanueva (New York, NY: Springer), 77–96.
- Oksanen, J., Blanchet, F. G., Kindt, R., Legendre, P., Minchin, P. R., O’hara, R. B., et al. (2013). *Vegan: Community Ecology Package. R Package Version 2.0-10*. Available online at: <http://CRAN.R-project.org/package=vegan>
- Okumura, S., Kurihara, A., Iwamoto, A., and Takeuchi, T. (2005). Improved survival and growth in *Octopus vulgaris* paralarvae by feeding large type Artemia and Pacific sandeel, *Ammodytes personatus*. *Aquaculture* 244, 147–157. doi: 10.1016/j.aquaculture.2004.11.044
- Olmos-Pérez, L., Roura, Á., Pierce, G. J., Boyer, S., and González, Á. F. (2017). Diet composition and variability of wild *Octopus vulgaris* and *Alloteuthis media* (Cephalopoda) paralarvae through a metagenomic lens. *Front. Physiol.* 8:321. doi: 10.3389/fphys.2017.00321
- Otero, J., Álvarez-Salgado, X., González, Á. F., Gilcoto, M., and Guerra, Á. (2009). High-frequency coastal upwelling events influence *Octopus vulgaris* larval dynamics on the NW Iberian shelf. *Mar. Ecol. Prog. Ser.* 386, 123–132. doi: 10.3354/meps08041
- Otero, J., Álvarez-Salgado, X., González, Á. F., Miranda, A., Groom, S. B., Cabanas, J. M., et al. (2008). Bottom-up control of common Octopus *Octopus vulgaris* in the Galician upwelling system, northeast Atlantic Ocean. *Mar. Ecol. Prog. Ser.* 362, 181–192. doi: 10.3354/meps07437
- Passarella, K. C., and Hopkins, T. L. (1991). Species composition and food habits of the micronektonic cephalopod assemblage in the Eastern Gulf of Mexico. *B. Mar. Sci.* 49, 638–659.
- Quintana, D., Márquez, L., Arévalo, J. R., Lorenzo, A., and Almansa, E. (2015). Relationships between spawn quality and biochemical composition of eggs and hatchlings of *Octopus vulgaris* under different parental diets. *Aquaculture* 446, 206–216. doi: 10.1016/j.aquaculture.2015.04.023
- Reis, D. B., Acosta, N. G., Almansa, E., Navarro, J. C., Tocher, D. R., Monroig, O., et al. (2014). *In vivo* metabolism of unsaturated fatty acids in *Octopus vulgaris* hatchlings determined by incubation with 14C-labelled fatty acids added directly to seawater as protein complexes. *Aquaculture* 431, 28–33. doi: 10.1016/j.aquaculture.2014.05.016

- Reis, D. B., García-Herrero, I., Riera, R., Felipe, B. C., Rodríguez, C., Sykes, A. V., et al. (2015). An insight on *Octopus vulgaris* paralarvae lipid requirements under rearing conditions. *Aquacult. Nutr.* 21, 797–806. doi: 10.1111/anu.12205
- Rosa, R., Costa, P. R., and Nunes, M. L. (2004). Effect of sexual maturation on the tissue biochemical composition of *Octopus vulgaris* and *O. defilippi* (Mollusca: Cephalopoda). *Mar. Biol.* 145, 563–574. doi: 10.1007/s00227-004-1340-8
- Roura, Á., Álvarez-Salgado, X. A., González, Á. F., Gregori, M., Rosón, G., and Guerra, Á. (2013). Short-term meso-scale variability of mesozooplankton communities in a coastal upwelling system (NW Spain). *Prog. Oceanogr.* 109, 18–32. doi: 10.1016/j.pocean.2012.09.003
- Roura, Á., Álvarez-Salgado, X. A., González, Á. F., Gregori, M., Rosón, G., Otero, J., et al. (2016). Life strategies of cephalopod paralarvae in a coastal upwelling system (NW Iberian Peninsula): insights from zooplankton community and spatio-temporal analyses. *Fish. Oceanogr.* 25, 241–258. doi: 10.1111/fog.12151
- Roura, Á., González, Á. F., Pascual, S., and Guerra, Á. (2010). A molecular approach to identifying the prey of cephalopod paralarvae. *ICES J. Mar. Sci.* 67, 1408–1412. doi: 10.1093/icesjms/fsq051
- Roura, Á., González, Á. F., Redd, K., and Guerra, Á. (2012). Molecular prey identification in wild *Octopus vulgaris* paralarvae. *Mar. Biol.* 159, 1335–1345. doi: 10.1007/s00227-012-1914-9
- Seixas, P., Rey-Méndez, M., Valente, L. M., and Otero, A. (2008). Producing juvenile *Artemia* as prey for *Octopus vulgaris* paralarvae with different microalgal species of controlled biochemical composition. *Aquaculture* 283, 83–91. doi: 10.1016/j.aquaculture.2008.06.019
- Seixas, P., Rey-Méndez, M., Valente, L. M. P., and Otero, A. (2010). High DHA content in *Artemia* is ineffective to improve *Octopus vulgaris* paralarvae rearing. *Aquaculture* 300, 156–162. doi: 10.1016/j.aquaculture.2009.12.021
- Sieiro, M. P., Aubourg, S. P., and Rocha, F. (2006). Seasonal study of the lipid composition in different tissues of the common octopus (*Octopus vulgaris*). *Eur. J. Lipid Sci. Technol.* 108, 479–487. doi: 10.1002/ejlt.200500322
- Tocher, D. R., and Glencross, B. D. (2015). “Lipids and fatty acids,” in *Dietary Nutrients, Additives, and Fish health*, eds C.-S. Lee, C. Lim, D. M. Gatlin, and C. D. Webster (Hoboken, NJ: Wiley Blackwell), 47–94.
- Vaz-Pires, P., Seixas, P., and Barbosa, A. (2004). Aquaculture potential of the common octopus (*Octopus vulgaris* Cuvier, 1797): a review. *Aquaculture* 238, 221–238. doi: 10.1016/j.aquaculture.2004.05.018
- Viciano, E., Iglesias, J., Lago, M. J., Sánchez, F., Otero, J., and Navarro, J. C. (2011). Fatty acid composition of polar and neutral lipid fractions of *Octopus vulgaris* Cuvier, 1797 paralarvae reared with enriched on-grown *Artemia*: polar and neutral lipid fatty acids of octopus. *Aquac. Res.* 42, 704–709. doi: 10.1111/j.1365-2109.2010.02605.x
- Villanueva, R. (1994). Decapod crab zoeae as food for rearing cephalopod paralarvae. *Aquaculture* 128, 143–152. doi: 10.1016/0044-8486(94)90109-0
- Villanueva, R., Nozais, C., and Boletzky, S. V. (1996). Swimming behaviour and food searching in planktonic *Octopus vulgaris* Cuvier from hatching to settlement. *J. Exp. Mar. Biol. Ecol.* 208, 169–184. doi: 10.1016/S0022-0981(96)02670-6
- Zaragoza, N., Quetglas, A., and Moreno, A. (2015). *Identification guide for Cephalopod Paralarvae from the Mediterranean Sea*. ICES Cooperative Research Report.
- Zuur, A. F., Ieno, E. N., and Smith, G. M. (2007). *Analysing Ecological Data*. New York, NY: Springer.

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2017 Loureço, Roura, Fernández-Reiriz, Narciso and González. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.