



# Analysis of Mitochondrial Calcium Retention Capacity in Cultured Cells: Permeabilized Cells Versus Isolated Mitochondria

Sehwan Jang<sup>1</sup>, Xavier R. Chapa-Dubocq<sup>1</sup>, Silvia Fossati<sup>2</sup> and Sabzali Javadov<sup>1\*</sup>

<sup>1</sup>Department of Physiology, University of Puerto Rico School of Medicine, San Juan, PR, United States, <sup>2</sup>Alzheimer's Center at Temple, Lewis Katz School of Medicine, Temple University, Philadelphia, PA, United States

## OPEN ACCESS

### Edited by:

Miguel A. Aon,  
National Institute on Aging, National  
Institutes of Health (NIH),  
United States

### Reviewed by:

Elena N. Dedkova,  
University of California, Davis,  
United States  
An-Chi Wei,  
National Taiwan University, Taiwan

### \*Correspondence:

Sabzali Javadov  
sabzali.javadov@upr.edu

### Specialty section:

This article was submitted to  
Mitochondrial Research,  
a section of the journal  
Frontiers in Physiology

**Received:** 10 September 2021

**Accepted:** 17 November 2021

**Published:** 07 December 2021

### Citation:

Jang S, Chapa-Dubocq XR,  
Fossati S and Javadov S (2021)  
Analysis of Mitochondrial Calcium  
Retention Capacity in Cultured Cells:  
Permeabilized Cells Versus  
Isolated Mitochondria.  
Front. Physiol. 12:773839.  
doi: 10.3389/fphys.2021.773839

In response to various pathological stimuli, such as oxidative and energy stress accompanied by high  $\text{Ca}^{2+}$ , mitochondria undergo permeability transition (PT) leading to the opening of the non-selective PT pores (PTP) in the inner mitochondrial membrane. Opening of the pores at high conductance allows the passage of ions and solutes <1.5 kD across the membrane, that increases colloid osmotic pressure in the matrix leading to excessive mitochondrial swelling. Calcium retention capacity (CRC) reflects maximum  $\text{Ca}^{2+}$  overload of mitochondria that occurs just before PTP opening. Quantification of CRC is important for elucidating the effects of different pathological stimuli and the efficacy of pharmacological agents on the mitochondria. Here, we performed a comparative analysis of CRC in mitochondria isolated from H9c2 cardioblasts, and in permeabilized H9c2 cells *in situ* to highlight the strengths and weaknesses of the CRC technique in isolated cell mitochondria vs. permeabilized cells. The cells were permeabilized by digitonin or saponin, and the  $\text{Ca}^{2+}$ -sensitive fluorescence probe Calcium Green-5N was used in both preparations. Results demonstrated the interference of dye-associated fluorescence signals with saponin and the adverse effects of digitonin on mitochondria at high concentrations. Analysis of the CRC in permeabilized cells revealed a higher CRC in the saponin-permeabilized cells in comparison with the digitonin-permeabilized cells. In addition, the mitochondrial CRC in saponin-permeabilized cells was higher than in isolated mitochondria. Altogether, these data demonstrate that the quantification of the mitochondrial CRC in cultured cells permeabilized by saponin has more advantages compared to the isolated mitochondria.

**Keywords:** permeabilized cells, mitochondria, calcium retention capacity, permeability transition pore, mitochondrial swelling

## INTRODUCTION

Mitochondria are the cell powerplants that provide over 90% of ATP required for cell metabolism. Also, mitochondria play a pivotal role in the maintenance of ion homeostasis, cell growth, redox signaling, and cell death. The metabolism and function of mitochondria are regulated by changes in the matrix volume associated with ion fluxes, particularly  $\text{Ca}^{2+}$ , across the inner

mitochondrial membrane. At low concentrations,  $\text{Ca}^{2+}$  induces negligible matrix swelling (Ichas and Mazat, 1998) and stimulates mitochondrial bioenergetics through fatty acid oxidation, tricarboxylic acid cycle, and oxidative phosphorylation (Halestrap, 1994; Tarasov et al., 2012). However, high  $\text{Ca}^{2+}$  causes excessive mitochondrial swelling leading to mitochondrial dysfunction and cell death. The main mechanism of mitochondrial swelling involves the opening of non-selective channels known as permeability transition pores (PTPs) in low-conductance (physiological) and high-conductance (pathological) modes in the inner mitochondrial membrane (Ichas and Mazat, 1998; Kwong and Molkentin, 2015). Mitochondrial swelling is driven by a high colloid osmotic pressure in the matrix of mitochondria exerted by high  $\text{Ca}^{2+}$  and non-diffusible matrix proteins. Quantification of the extent of mitochondrial swelling is important for the analysis of mitochondrial damage in response to various pathological stimuli.

The  $\text{Ca}^{2+}$  retention capacity (CRC) is broadly used to quantify the extent of PTP opening since CRC indicates the maximum  $\text{Ca}^{2+}$  uptake that mitochondria reach before PTP opening. Therefore, the amount of external  $\text{Ca}^{2+}$  that induces  $\text{Ca}^{2+}$  release through the PTP reflects the CRC of mitochondria that corresponds to the maximum mitochondrial swelling. The CRC can be quantified in isolated mitochondria and intact cells (without isolation of mitochondria). Although the fundamental knowledge on the structural organization, metabolism, and function of mitochondria, as well as their response to a wide range of diseases, was acquired using isolated cell or tissue mitochondria, the use of isolated mitochondria has several disadvantages (Kuznetsov et al., 2008; Dedkova and Blatter, 2012; Salabei et al., 2014). First, the isolation of mitochondria requires a relatively large quantity of cells or tissues since some parts are lost in the isolation process (centrifugation and washing). Second, isolated mitochondria do not represent all populations of mitochondria since mitochondria localized in certain subcellular compartments (e.g., intrafibrillar mitochondria) are not isolated by homogenization. Third, mitochondria, especially dysfunctional (fragile) mitochondria from pathological cells/tissues are partially damaged or lost during the isolation process. Fourth, isolation from the essential intracellular environment has severe effects on the morphology, metabolism, and function of mitochondria and changes their sensitivity to exogenous factors.

In this study, we evaluated the CRC in intact cells *in situ* permeabilized by two different biological detergents and in mitochondria isolated from the cultured cells to clarify the advantages and disadvantages of each technique.

## MATERIALS AND METHODS

### Animals

Adult Sprague Dawley male rats (275–325 g) were purchased from Taconic (Hillside, NJ, United States). All experiments were performed according to protocols approved by the UPR Medical Sciences Campus Institutional Animal Care and Use

Committee and conformed to the National Research Council Guide for the Care and Use of Laboratory Animals published by the US National Institutes of Health (2011, eighth edition).

### Cells

H9c2 embryonic rat cardioblastic cells were cultured according to the manufacturer's recommendations (American Type Culture Collection, Manassas, VA). Briefly, the cells were cultured in DMEM based modified media containing 4 mM L-glutamine, 4.5 g/L glucose, 1 mM sodium pyruvate, and 1.5 g/L sodium bicarbonate supplemented with 10% fetal bovine serum and 1% antibiotic solution (HyClone) and maintained in 95% air and 5%  $\text{CO}_2$  at 37°C. Cells maintained within 80–90% confluence from passages 3–10 were used for experiments.

### Isolation of Mitochondria From Rat Hearts

The isolation of mitochondria was adopted and modified from previous studies (Chapa-Dubocq et al., 2020). Briefly, heart ventricles were cut and homogenized using a Polytron homogenizer in ice-cold sucrose buffer containing (in mM): 300 sucrose, 20 Tris-HCl, and 2 EGTA, pH 7.2, and supplemented with 0.05% BSA. The heart homogenate was centrifuged at 2000×g for 3 min to remove cell debris. The supernatant was centrifuged at 10,000×g for 6 min to precipitate mitochondria and then washed again under the same conditions in sucrose buffer (BSA-free). The final pellet containing mitochondria was resuspended in 300 μl of sucrose buffer.

### Isolation of Mitochondria From Cells

To isolate mitochondria from cultured cells, H9c2 cells were trypsinized and pelleted at 200×g for 7 min (Jang and Javadov, 2018). Pellet was resuspended in ice-cold sucrose buffer containing (in mM): 300 sucrose, 10 Tris-HCl, and 2 EGTA; pH 7.4. Cells were centrifuged at 2,500×g for 5 min at 4°C, the pellet was resuspended in the sucrose buffer. To disrupt the plasma membrane and expose mitochondria, cells were plunged using a 27G needle until all cells were successfully lysed. The cell lysate was then centrifuged at 400×g for 5 min and the supernatant was collected. The mitochondria were concentrated by centrifugation at 10,000×g for 5 min and finally dissolved with sucrose buffer.

### Permeabilization of Cells

The basic principles of the cell permeabilization technique for analysis of mitochondrial function *in situ* have been described in detail elsewhere (Kuznetsov et al., 2008; Dedkova and Blatter, 2012; Salabei et al., 2014). Cells were freshly harvested using trypsin-EDTA then permeabilized in sucrose buffer (300 mM sucrose, 10 mM Tris-HCl, 2 mM EGTA, pH 7.4) containing saponin or digitonin for 10 min on ice. After the permeabilization, cells were washed with equilibration buffer (100 mM sucrose, 10 mM Tris-HCl, 10 μM EGTA, pH 7.4), then resuspended in incubation buffer (200 mM sucrose, 10 mM Tris-MOPS, 5 mM α-ketoglutarate, 2 mM malate, 1 mM Pi, 10 μM EGTA-Tris, pH 7.4) containing 100 nM Calcium Green-5 N.

## Fluorescence Imaging

To evaluate the effects of permeabilization on mitochondrial function, live H9c2 cells grown to 70–80% confluence were treated with saponin or digitonin for 10 min. Then, the cells were further incubated for 30 min with 300 nM DAPI for visualization of the nucleus and 30 nM Mitotracker Red, a membrane potential-dependent dye. As a positive control, 0.1% Triton X-100 was applied to induce irreversible permeabilization of the plasma membrane leading to structural collapse of the cells. Cell images were captured by an Olympus IX73 microscope with LUCPLFN10X objective using Cellsense Dimension (Olympus) software.

## CRC Assay

The CRC was quantified by the  $\text{Ca}^{2+}$ -sensitive fluorescence dyes Oregon Green 488 BAPTA-1, Calcium Green-5N, or Fluo-5N that measure extramitochondrial  $\text{Ca}^{2+}$  fluorescence in the assay buffer. Briefly, freshly isolated mitochondria (0.5 mg/ml) or permeabilized fresh cells were incubated at 37°C in 0.1 ml of incubation buffer (200 mM sucrose, 10 mM Tris-MOPS, 5 mM  $\alpha$ -ketoglutarate, 2 mM malate, 1 mM Pi, 10  $\mu\text{M}$  EGTA-Tris, pH 7.4) containing one of the fluorescence dyes. Calcium was added to increase matrix  $\text{Ca}^{2+}$  load and the fluorescence intensity was recorded by CLARIOStar microplate reader (BMG Labtech).

## Mitochondrial PTP Opening

The swelling of mitochondria as an indicator of PTP opening in the presence or absence of  $\text{Ca}^{2+}$  was determined freshly isolated mitochondria (50  $\mu\text{g}$ ) by monitoring the decrease in light scattering at 525 nm as previously described, with minor modifications (Rodríguez-Graciani et al., 2020). The swelling buffer contained (in mM): 125 KCl, 20 MOPS, 10 Tris-HCl, 0.001 EGTA, and 2  $\text{KH}_2\text{PO}_4$ , pH 7.1. The swelling curves were normalized to control and presented as an absorbance ratio.

## Statistical Analysis

Data values are presented as mean  $\pm$  SE. Student's *t*-test was used to compare differences between two groups.  $p < 0.05$  was considered as statistically significant. The number of biological samples but not technical replicates were used as a sample size.

## RESULTS

### Analysis of Mitochondrial CRC by Different $\text{Ca}^{2+}$ -Sensitive Fluorescent Dyes


In the first set of experiments, we assessed the sensitivity of three different fluorescent dyes to  $\text{Ca}^{2+}$  in isolated heart mitochondria. The CRC was measured in 50  $\mu\text{g}$  of mitochondria by using Oregon Green 488 BAPTA-1, Calcium Green-5N or Fluo-5N that possess a dissociation constant ( $K_d$  for  $\text{Ca}^{2+}$ ) of 0.17  $\mu\text{M}$ , 15  $\mu\text{M}$ , or 90  $\mu\text{M}$ , respectively (Figure 1). Results exhibited a different sensitivity of the dyes to  $\text{Ca}^{2+}$ . The Oregon Green 488 BAPTA-1 dye was the most sensitive and CRC reached the maximum at  $\sim 60 \mu\text{M}$   $\text{Ca}^{2+}$  (Figure 1A) whereas Calcium Green-5N demonstrated the maximum CRC at 200  $\mu\text{M}$ .

(Figure 1B). The CRC of mitochondria was less sensitive to Fluo-5N and could not reach the maximum at 280  $\mu\text{M}$  (Figure 1A). Moreover, inhibition of the mitochondrial swelling by sanglifehrin A, a PTP blocker that inhibits cyclophilin D activity, affected differently the fluorescence intensity of the dyes; inhibition of the CRC was more obvious in the presence of Calcium Green-5N or Fluo-5N (Figures 1B,C) suggesting that the inhibitory effect of sanglifehrin A on mitochondrial swelling depends on the sensitivity of the dyes to  $\text{Ca}^{2+}$  as well as on the concentration of  $\text{Ca}^{2+}$ . It should be noted that decreased CRC was associated with increased PTP opening in the isolated cardiac mitochondria (Figures 1D–F).  $\text{Ca}^{2+}$  induces inner membrane expansion and outer membrane rupture (Strubbe-Rivera et al., 2021), leading to the swelling of mitochondria. The increase in matrix volume is accompanied by a decrease in the intensity of light scattered (Tedeschi and Harris, 1955). Thus, these data demonstrate that due to differences in  $K_d$  for  $\text{Ca}^{2+}$ , the different  $\text{Ca}^{2+}$ -sensitive fluorescent dyes can be applied for measurement of the CRC at low (Oregon Green 488 BAPTA-1:  $K_d = 170 \text{ nM}$ ), medium (Calcium Green-5N:  $K_d = 14 \mu\text{M}$ ), or high (Fluo-5N:  $K_d = 90 \mu\text{M}$ ) concentrations of  $\text{Ca}^{2+}$ . Moreover, we experimentally demonstrated the relationship between mitochondrial swelling and  $\text{Ca}^{2+}$  efflux in response to  $\text{Ca}^{2+}$ -induced PTP opening.

### The Effects of Permeabilization on Mitochondria



Based on the sensitivity to  $\text{Ca}^{2+}$  (Figures 1A–C), we used Calcium Green-5N to assess the permeabilization capabilities, potential effects of digitonin and saponin, and the CRC in H9c2 cardioblasts. DAPI, a nuclear staining dye that does not enter non-permeabilized cells, was used to estimate the effectiveness of the detergents to permeabilize the cells. Analysis of permeabilization capabilities demonstrated that both biological detergents at high concentrations have toxic effects on mitochondria, inducing loss of membrane potential (Figures 2A,C,D). The optimal concentrations of the detergents to induce permeabilization of the cells with no effects on mitochondrial functional activity were different for saponin and digitonin. Quantification of DAPI fluorescence to total cells found that saponin levels of 25  $\mu\text{g}/\text{ml}$  or higher and all digitonin levels used were capable of permeabilizing the cellular membrane (Figures 2A–C). Analysis of Mitotracker Red displayed mitochondria viability ranging from 5 to 100  $\mu\text{g}/\text{ml}$  of saponin as well as digitonin levels between 5 and 10  $\mu\text{g}/\text{ml}$ . Saponin effectively permeabilized the cells at concentrations 25, 50, and 100  $\mu\text{g}/\text{ml}$  (Figure 2A), whereas optimal permeabilization of the cells by digitonin was observed at 5 and 10  $\mu\text{g}/\text{ml}$  (Figure 2C).

To determine whether the detergents have interference with Calcium Green-5N (Figure 3; Supplementary Figures S2A,B), the fluorescence intensity of the dye was measured in the assay buffer (no cells) by adding digitonin or saponin at concentrations of 50, 100, and 200  $\mu\text{g}/\text{ml}$ . Saponin at 200  $\mu\text{g}/\text{ml}$  demonstrated interference with Calcium Green-5N as evidenced by saturated fluorescence signal in the absence of  $\text{Ca}^{2+}$  (Figure 3A; Supplementary Figure S2A) whereas digitonin had no interference with the dye at all concentrations (Figure 3B; Supplementary Figure S2B).



**FIGURE 1 |** CRC analysis in mitochondria isolated from adult Sprague Dawley rat hearts. The CRC was measured using the  $Ca^{2+}$  sensitive fluorescent dyes, 100nM Oregon Green 488 BAPTA-1 (A), 100nM Calcium Green-5N (B), and 500nM Fluo-5N (C). Mitochondria (50  $\mu g$ ) were exposed to repetitive application of 20  $\mu M$  (2nmol) of  $Ca^{2+}$  every 2 min (arrows) to increase matrix  $Ca^{2+}$  load. The PTP dependence of the CRC was determined by adding 0.5  $\mu M$  sangliferhin A (SfA, a cyclophilin D inhibitor) to the incubation media. \* $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$  vs. control (Con).  $n = 3$  per group. (D–F) CRC and mitochondrial swelling analysis in mitochondria isolated from adult Sprague Dawley rat hearts. The CRC was measured using the  $Ca^{2+}$  sensitive fluorescent dyes, 500nM Fluo-5N. Mitochondria (50  $\mu g$ ) were exposed to 100 (D), 200 (E), and 300 (F)  $\mu M$  (10, 20, 30nmol) of  $Ca^{2+}$  at 5 min (arrows) to increase matrix  $Ca^{2+}$  load.  $n = 3$  per group.





## Analysis of the CRC in Permeabilized H9c2 Cardiomyocytes and Isolated Mitochondria

Next, we analyzed the CRC in permeabilized cells *in situ* and in isolated mitochondria *in vitro* to choose the technique which can be used for accurate quantification of mitochondrial CRC in cultured cells. First, we measured the CRC in H9c2 cardiomyocytes permeabilized by saponin or digitonin (Figure 4A; Supplementary Figure S2C). The cells permeabilized by digitonin at all concentrations (10, 50, and 100  $\mu\text{g}/\text{mL}$ ) had lower CRC than saponin-permeabilized cells. The cells permeabilized with 50 and 100  $\mu\text{g}/\text{mL}$  saponin required approximately 9 and 10 nmol  $\text{Ca}^{2+}$ , respectively, to induce a massive release of  $\text{Ca}^{2+}$  whereas nearly 6, 7, and 8 nmol  $\text{Ca}^{2+}$  were required to open the PTP/ $\text{Ca}^{2+}$  release from mitochondria of cells permeabilized by 10, 50, and 100  $\mu\text{g}/\text{mL}$  of digitonin, respectively.


Considering the negative effects of digitonin on the CRC of mitochondria (Figure 4A; Supplementary Figure S2C), in the next set of experiments, we used saponin-permeabilized cells for comparison of mitochondrial CRC in the permeabilized cells versus isolated mitochondria. An equal number of H9c2 cells ( $0.6 \times 10^6$ ) were used for permeabilization of cells and isolation of mitochondria. Notably, mitochondrial mass was not decreased significantly ( $<7\%$ ,  $p < 0.974$ ) in isolated mitochondria in comparison with the permeabilized cells as evidenced by citrate synthase activities in both samples. Results demonstrated that the CRC in permeabilized cells was higher compared to the isolated mitochondria (Figure 4B; Supplementary Figure S2D); 10 nmol  $\text{Ca}^{2+}$  was required to

trigger PTP opening/massive  $\text{Ca}^{2+}$  release in the cells permeabilized with 50  $\mu\text{g}/\text{mL}$  saponin whereas isolated mitochondria started swelling at 6 nmol  $\text{Ca}^{2+}$  indicating at their lower CRC. Thapsigargin, a non-competitive inhibitor of the SERCA, was used to estimate the contribution of sarcoplasmic reticulum to the CRC of mitochondria. The mitochondrial CRC of saponin-permeabilized cells was slightly reduced in the presence of 1  $\mu\text{M}$  thapsigargin. As expected, it had no effect on the CRC of the isolated mitochondria (Figure 4B; Supplementary Figure S2D).

## DISCUSSION

In this study, we attempted to compare two techniques for the measurement of the CRC of mitochondria in cultured cells. Results showed that mitochondrial CRC measured in permeabilized cells is higher than that in isolated mitochondria. Also, we found the optimal concentrations of the biological detergents saponin and digitonin required for effective permeabilization of the H9c2 cells with no toxic effects on mitochondrial function ( $\Delta\Psi_m$ ).

A few techniques have been developed for the quantification of mitochondrial PTP/CRC *in vivo* in cells and tissues. 2-deoxy[ $^3\text{H}$ ] glucose (DOG) entrapment technique can be applied to measure PTP opening in isolated perfused hearts *in vivo* (Javadov et al., 2005; Ciminelli et al., 2006). In cultured cells, mitochondrial PTP/CRC can be measured directly in intact cells by calcein, a cell-permeant fluorescent probe, the intensity of which is quenched strongly by metal ions, such as  $\text{Co}^{2+}$ , in the cytosol. Hence, the fluorescence quenching in response



**FIGURE 4 |** CRC analysis in permeabilized cells *in situ* and isolated mitochondria *in vitro*. **(A)** CRC in H9c2 cells permeabilized by saponin or digitonin. The cells were permeabilized by saponin (50 and 100  $\mu\text{g}/\text{mL}$ ) or digitonin (10, 50, and 100  $\mu\text{g}/\text{mL}$ ) for 10 min in sucrose buffer on ice.  $n = 3-6$  per group. **(B)** Comparative analysis of CRC in permeabilized cells vs. isolated mitochondria in the presence or absence of 1  $\mu\text{M}$  thapsigargin (TG).  $\text{Ca}^{2+}$  was added every 3 min (arrows) by increments of 1 nmol/injection.  $0.6 \times 10^6$  H9c2 cells were used for each well. Permeabilized cells were washed off the detergents before the analysis. Curves represent the averages.  $n = 6$  per group.

to  $\text{Ca}^{2+}$  release from mitochondria through the PTP can allow quantifying the extent of pore opening (Petronilli et al., 1999; Hernandez et al., 2014; Teixeira et al., 2015). However, the technique for analysis of the PTP by calcein has several weaknesses. First,  $\text{Co}^{2+}$  is a heavy metal and has toxic effects on cells. Second, the technique does not allow to specify whether calcein quenching occurs due to its release through mitochondrial PTPs or results from  $\text{Co}^{2+}$  entry into mitochondria. For example, incubation of cells with calcein in combination with a red-fluorescing potentiometric dye demonstrated that cytosolic calcein can be released from normal mitochondria but enters them upon PTP opening (Jones et al., 2002). Third, calcein-AM entering mitochondria is not cleaved in all cell types such as hepatocytes (Lemasters et al., 1998).

Advantages of the technique for quantification of the CRC in permeabilized cells *in situ* compared to isolated mitochondria can be explained with the fact that during permeabilization intracellular structural organization of the cells is preserved, allowing to analyze all subcellular populations of mitochondria (Harrisseh et al., 2019). Analysis of several types of primary cells and cell lines revealed that mitochondria within individual cells are morphologically heterogeneous with varying sizes, are differently distributed, and they may even have distinct functions (Collins et al., 2002). In contrast, analysis of PTP opening, mitochondrial swelling, and the CRC in isolated mitochondria have certain limitations. Homogenization and centrifugation during the isolation procedure damage mitochondria and induce their swelling (Morikawa et al., 2014). In addition, isolated mitochondria do not present all populations of mitochondria localized in different subcellular compartments.

A large number of studies used permeabilized cells, particularly cardiomyocytes, to investigate mitochondrial bioenergetics and respiration rates (Eimre et al., 2008; Xu et al., 2014), CRC (Harrisseh et al., 2019), membrane potential (Keane et al., 2016), ROS (Pham et al., 2000; Keane et al., 2016), and ions ( $\text{Na}^+$ ,  $\text{Ca}^{2+}$ ; Donoso et al., 1992; Dedkova and Blatter, 2009) *in situ*. Permeabilized cardiomyocytes and cardiac muscle fibers have been shown to maintain natural structural organization and mitochondrial bioenergetics (Kay et al., 1997; Saks et al., 1998), which are consistent with our findings (Supplementary Figure S1). Likewise, intracellular morphology and mitochondrial function of H9c2 cardioblasts remained unaffected by permeabilization (Szalai et al., 2000; Medepalli et al., 2013; Namekata et al., 2017). Mostly, two biological detergents digitonin and saponin were used for the permeabilization of cells. Digitonin disrupts the plasma membrane bilayers by targeting lipid rafts while saponins permeabilize plasma membranes by selectively removing cholesterol from the membranes without affecting membrane proteins. The intracellular environment including the structural and functional integrity of subcellular organelles, and interactions between them remain almost intact in permeabilized cells. Interestingly, thapsigargin had no significant effect on the CRC in permeabilized cells (Figure 4B). This can be explained by the low concentration of ATP (due to its dilution in the assay buffer upon permeabilization), which is not sufficient to activate SERCA

and stimulate  $\text{Ca}^{2+}$  uptake by sarcoplasmic reticulum. Apparently, permeabilization efficacy and toxic effects of saponin and digitonin at given concentrations can be varied for different cell types.

In conclusion, this study highlights the advantages of the quantification of mitochondrial CRC in permeabilized cells compared to isolated mitochondria and establishes optimal concentrations of two biological detergents that are widely used for the permeabilization of cells. The results of the study can be taken into consideration during the quantification of mitochondrial CRC in live cells.

## DATA AVAILABILITY STATEMENT

The original contributions presented in the study are included in the article/Supplementary Material, further inquiries can be directed to the corresponding author.

## ETHICS STATEMENT

The animal study was reviewed and approved by the UPR Medical Sciences Campus Institutional Animal Care and Use Committee and conformed by the National Research Council Guide for the Care and Use of Laboratory Animals published by the US National Institutes of Health (2011, eighth edition).

## AUTHOR CONTRIBUTIONS

SJav and SF conceived the study and wrote the manuscript. SJan and XC-D performed the experiments and contributed the data. All authors contributed to the article and approved the submitted version.

## FUNDING

This study was supported by the grants from the National Institutes of Health (SC1GM128210 and R25GM061838 to SJav; R01NS104127 and R01AG062572 to SF), the National Science Foundation (2006477 to SJav), and the Pennsylvania Department of Health Collaborative Research on Alzheimer's Disease (PA Cure, to SF).

## ACKNOWLEDGMENTS

We thank Jorge Garcia-Baez for assisting the experiments and analysis.

## SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fphys.2021.773839/full#supplementary-material>

## REFERENCES

- Chapa-Dubocq, X. R., Rodriguez-Graciani, K. M., Guzman-Hernandez, R. A., Jang, S., Brookes, P. S., and Javadov, S. (2020). Cardiac function is not susceptible to moderate disassembly of mitochondrial respiratory supercomplexes. *Int. J. Mol. Sci.* 21:1555. doi: 10.3390/ijms21051555
- Ciminelli, M., Asch, A., Bourduas, K., and Buelle, Y. (2006). Short term training attenuates opening of the mitochondrial permeability transition pore without affecting myocardial function following ischemia-reperfusion. *Mol. Cell. Biochem.* 291, 39–47. doi: 10.1007/s11010-006-9192-9
- Collins, T. J., Berridge, M. J., Lipp, P., and Bootman, M. D. (2002). Mitochondria are morphologically and functionally heterogeneous within cells. *EMBO J.* 21, 1616–1627. doi: 10.1093/emboj/21.7.1616
- Dedkova, E. N., and Blatter, L. A. (2009). Characteristics and function of cardiac mitochondrial nitric oxide synthase. *J. Physiol.* 587, 851–872. doi: 10.1113/jphysiol.2008.165423
- Dedkova, E. N., and Blatter, L. A. (2012). Measuring mitochondrial function in intact cardiac myocytes. *J. Mol. Cell. Cardiol.* 52, 48–61. doi: 10.1016/j.yjmcc.2011.08.030
- Donoso, P., Mill, J. G., and O'Neill, S.C., and Eisner, D.A., (1992). Fluorescence measurements of cytoplasmic and mitochondrial sodium concentration in rat ventricular myocytes. *J. Physiol.* 448, 493–509. doi: 10.1113/jphysiol.1992.sp019053
- Eimre, M., Paju, K., Pelloux, S., Beraud, N., Roosimaa, M., Kadaja, L., et al. (2008). Distinct organization of energy metabolism in HL-1 cardiac cell line and cardiomyocytes. *Biochim. Biophys. Acta* 1777, 514–524. doi: 10.1016/j.bbabi.2008.03.019
- Halestrap, A. P. (1994). Regulation of mitochondrial metabolism through changes in matrix volume. *Biochem. Soc. Trans.* 22, 522–529. doi: 10.1042/bst0220522
- Harisseh, R., Abrial, M., Chiari, P., Al-Mawla, R., Villedieu, C., Tessier, N., et al. (2019). A modified calcium retention capacity assay clarifies the roles of extra- and intracellular calcium pools in mitochondrial permeability transition pore opening. *J. Biol. Chem.* 294, 15282–15292. doi: 10.1074/jbc.RA119.009477
- Hernandez, J. S., Barreto-Torres, G., Kuznetsov, A. V., Khuchua, Z., and Javadov, S. (2014). Crosstalk between AMPK activation and angiotensin II-induced hypertrophy in cardiomyocytes: the role of mitochondria. *J. Cell. Mol. Med.* 18, 709–720. doi: 10.1111/jcmm.12220
- Ichas, F., and Mazat, J. P. (1998). From calcium signaling to cell death: two conformations for the mitochondrial permeability transition pore. Switching from low- to high-conductance state. *Biochim. Biophys. Acta* 1366, 33–50. doi: 10.1016/S0005-2728(98)00119-4
- Jang, S., and Javadov, S. (2018). Elucidating the contribution of ETC complexes I and II to the respirase formation in cardiac mitochondria. *Sci. Rep.* 8, 17732. doi: 10.1038/s41598-018-36040-9
- Javadov, S., Huang, C., Kirshenbaum, L., and Karmazyn, M. (2005). NHE-1 inhibition improves impaired mitochondrial permeability transition and respiratory function during postinfarction remodeling in the rat. *J. Mol. Cell. Cardiol.* 38, 135–143. doi: 10.1016/j.yjmcc.2004.10.007
- Jones, R. A., Smail, A., and Wilson, M. R. (2002). Detecting mitochondrial permeability transition by confocal imaging of intact cells pinocytically loaded with calcein. *Eur. J. Biochem.* 269, 3990–3997. doi: 10.1046/j.1432-1033.2002.03087.x
- Kay, L., Li, Z., Mericskay, M., Olivares, J., Tranqui, L., Fontaine, E., et al. (1997). Study of regulation of mitochondrial respiration in vivo. An analysis of influence of ADP diffusion and possible role of cytoskeleton. *Biochim. Biophys. Acta* 1322, 41–59. doi: 10.1016/S0005-2728(97)00071-6
- Keane, J., Tajouri, L., and Gray, B. (2016). Recombinant human growth hormone and insulin-like growth factor-1 do not affect mitochondrial derived highly reactive oxygen species production in peripheral blood mononuclear cells under conditions of substrate saturation in-vitro. *Nutr. Metab.* 13:45. doi: 10.1186/s12986-016-0105-y
- Kuznetsov, A. V., Veksler, V., Gellerich, F. N., Saks, V., Margreiter, R., and Kunz, W. S. (2008). Analysis of mitochondrial function in situ in permeabilized muscle fibers, tissues and cells. *Nat. Protoc.* 3, 965–976. doi: 10.1038/nprot.2008.61
- Kwong, J. Q., and Molkenin, J. D. (2015). Physiological and pathological roles of the mitochondrial permeability transition pore in the heart. *Cell Metab.* 21, 206–214. doi: 10.1016/j.cmet.2014.12.001
- Lemasters, J. J., Nieminen, A. L., Qian, T., Trost, L. C., Elmore, S. P., Nishimura, Y., et al. (1998). The mitochondrial permeability transition in cell death: a common mechanism in necrosis, apoptosis and autophagy. *Biochim. Biophys. Acta* 1366, 177–196. doi: 10.1016/S0005-2728(98)00112-1
- Medepalli, K., Alphenaar, B. W., Keynton, R. S., and Sethu, P. (2013). A new technique for reversible permeabilization of live cells for intracellular delivery of quantum dots. *Nanotechnology* 24:205101. doi: 10.1088/0957-4484/24/20/205101
- Morikawa, D., Kanematsu, K., Shibata, T., Haseda, K., Umeda, N., and Ohta, Y. (2014). Detection of swelling of single isolated mitochondrion with optical microscopy. *Biomed. Opt. Express* 5, 848–857. doi: 10.1364/BOE.5.000848
- Namekata, I., Hamaguchi, S., Iida-Tanaka, N., Kusakabe, T., Kato, K., Kawanishi, T., et al. (2017). Fluorescence analysis of the mitochondrial effect of a plasmalemmal Na<sup>+</sup>/Ca<sup>2+</sup> exchanger inhibitor, SEA0400, in permeabilized H9c2 cardiomyocytes. *Biol. Pharm. Bull.* 40, 1551–1555. doi: 10.1248/bpb.b17-00079
- Petronilli, V., Miotto, G., Canton, M., Brini, M., Colonna, R., Bernardi, P., et al. (1999). Transient and long-lasting openings of the mitochondrial permeability transition pore can be monitored directly in intact cells by changes in mitochondrial calcein fluorescence. *Biophys. J.* 76, 725–734. doi: 10.1016/S0006-3495(99)77239-5
- Pham, N. A., Robinson, B. H., and Hedley, D. W. (2000). Simultaneous detection of mitochondrial respiratory chain activity and reactive oxygen in digitonin-permeabilized cells using flow cytometry. *Cytometry* 41, 245–251. doi: 10.1002/1097-0320(20001201)41:4<245::AID-CYTO2>3.0.CO;2-#
- Rodriguez-Graciani, K. M., Chapa-Dubocq, X. R., Macmillan-Crow, L. A., and Javadov, S. (2020). Association between L-OPA1 cleavage and cardiac dysfunction during ischemia-reperfusion injury in rats. *Cell. Physiol. Biochem.* 54, 1101–1114. doi: 10.33594/00000303
- Saks, V. A., Veksler, V. I., Kuznetsov, A. V., Kay, L., Sikk, P., Tiivel, T., et al. (1998). Permeabilized cell and skinned fiber techniques in studies of mitochondrial function in vivo. *Mol. Cell. Biochem.* 184, 81–100. doi: 10.1023/A:1006834912257
- Salabei, J. K., Gibb, A. A., and Hill, B. G. (2014). Comprehensive measurement of respiratory activity in permeabilized cells using extracellular flux analysis. *Nat. Protoc.* 9, 421–438. doi: 10.1038/nprot.2014.018
- Strubbe-Rivera, J. O., Schrad, J. R., Pavlov, E. V., Conway, J. F., Parent, K. N., and Bazil, J. N. (2021). The mitochondrial permeability transition phenomenon elucidated by cryo-EM reveals the genuine impact of calcium overload on mitochondrial structure and function. *Sci. Rep.* 11:1037. doi: 10.1038/s41598-020-80398-8
- Szalai, G., Csordás, G., Hantash, B. M., Thomas, A. P., and Hajnóczky, G. (2000). Calcium signal transmission between ryanodine receptors and mitochondria. *J. Biol. Chem.* 275, 15305–15313. doi: 10.1074/jbc.275.20.15305
- Tarasov, A. I., Griffiths, E. J., and Rutter, G. A. (2012). Regulation of ATP production by mitochondrial Ca(2+). *Cell Calcium* 52, 28–35. doi: 10.1016/j.ceca.2012.03.003
- Tedeschi, H., and Harris, D. L. (1955). The osmotic behavior and permeability to non-electrolytes of mitochondria. *Arch. Biochem. Biophys.* 58, 52–67. doi: 10.1016/0003-9861(55)90092-8
- Teixeira, G., Chiari, P., Fauconnier, J., Abrial, M., Couture-Lepetit, E., Harisseh, R., et al. (2015). Involvement of cyclophilin D and calcium in isoflurane-induced preconditioning. *Anesthesiology* 123, 1374–1384. doi: 10.1097/ALN.0000000000000876
- Xu, A., Szczepanek, K., Maceyka, M. W., Ross, T., Bowler, E., Hu, Y., et al. (2014). Transient complex I inhibition at the onset of reperfusion by extracellular acidification decreases cardiac injury. *Am. J. Physiol. Cell Physiol.* 306, C1142–C1153. doi: 10.1152/ajpcell.00241.2013

**Conflict of Interest:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

**Publisher's Note:** All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2021 Jang, Chapa-Dubocq, Fossati and Javadov. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.