



Severe Acute Respiratory Syndrome Coronavirus 2 Infection in Pregnancy. A Non-systematic Review of Clinical Presentation, Potential Effects of Physiological Adaptations in Pregnancy, and Placental Vascular Alterations

OPEN ACCESS

Edited by:

John D. Imig,
Medical College of Wisconsin,
United States

Reviewed by:

Loren P. Thompson,
University of Maryland, Baltimore,
United States
Anne Prévot,
High School of Health Fribourg,
Switzerland
Eliana Hiromi Akamine,
University of São Paulo, Brazil

*Correspondence:

Paola Ayala-Ramírez
payala@javeriana.edu.co
Marcelo González
mgonzalezo@udec.cl

†These authors have contributed
equally to this work

Specialty section:

This article was submitted to
Vascular Physiology,
a section of the journal
Frontiers in Physiology

Received: 28 September 2021

Accepted: 14 February 2022

Published: 30 March 2022

Citation:

Ayala-Ramírez P, González M, Escudero C, Quintero-Arciniegas L, Giachini FR, Alves de Freitas R, Damiano AE and García-Robles R (2022) Severe Acute Respiratory Syndrome Coronavirus 2 Infection in Pregnancy. A Non-systematic Review of Clinical Presentation, Potential Effects of Physiological Adaptations in Pregnancy, and Placental Vascular Alterations. *Front. Physiol.* 13:785274. doi: 10.3389/fphys.2022.785274

Paola Ayala-Ramírez^{1*†}, Marcelo González^{2,3*†}, Carlos Escudero^{2,4†}, Laura Quintero-Arciniegas^{5,6}, Fernanda R. Giachini^{7,8}, Raiany Alves de Freitas⁸, Alicia E. Damiano^{9,10} and Reggie García-Robles⁶

¹ School of Medicine, Human Genetics Institute, Pontificia Universidad Javeriana, Bogotá, Colombia, ² Group of Research and Innovation in Vascular Health (GRIVAS Health), Chillan, Chile, ³ Laboratorio de Investigación Materno-Fetal (LIMaF), Departamento de Obstetricia y Ginecología, Facultad de Medicina, Universidad de Concepción, Concepción, Chile, ⁴ Laboratory of Vascular Physiology, Department of Basic Sciences, Faculty of Sciences, Universidad del Bio-Bio, Chillan, Chile, ⁵ Perinatal Medicine Seedbed, School of Medicine, Pontificia Universidad Javeriana, Bogotá, Colombia, ⁶ Department of Physiological Sciences, School of Medicine, Pontificia Universidad Javeriana, Bogotá, Colombia, ⁷ Institute of Biological Sciences and Health, Federal University of Mato Grosso, Barra do Garças, Brazil, ⁸ Institute of Biological Sciences, Federal University of Goiás, Goiânia, Brazil, ⁹ Laboratorio de Biología de la Reproducción, Instituto de Fisiología y Biofísica Bernardo Houssay (IFIBIO)- CONICET- Facultad de Medicina, Universidad de Buenos Aires, Buenos Aires, Argentina, ¹⁰ Cátedra de Biología Celular y Molecular, Departamento de Ciencias Biológicas, Facultad de Farmacia y Bioquímica, Universidad de Buenos Aires, Buenos Aires, Argentina

In December 2019, the novel Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) rapidly spread to become a pandemic. To date, increasing evidence has described the potential negative impact of SARS-CoV-2 infection on pregnant women. Although the pathophysiology of coronavirus disease 2019 (COVID-19) is not entirely understood, there is emerging evidence that it causes a severe systemic inflammatory response associated with vascular alterations that could be of special interest considering some physiological changes in pregnancy. Additionally, these alterations may affect the physiology of the placenta and are associated with pregnancy complications and abnormal histologic findings. On the other hand, data about the vaccine against SARS-CoV-2 are limited, but the risks of administering COVID-19 vaccines during pregnancy appear to be minimal. This review summarizes the current literature on SARS-CoV-2 virus infection, the development of COVID-19 and its relationship with physiological changes, and angiotensin-converting enzyme 2 (ACE2) function during pregnancy. We have particularly emphasized evidence coming from Latin American countries.

Keywords: COVID-19, pregnancy, renin-angiotensin-aldosterone system, coronavirus in pregnancy, placenta

INTRODUCTION

In December 2019, an unknown etiology outbreak of pneumonia was described in Wuhan, China. By January 2020, a new type of coronavirus was identified as the primary cause of these pneumonia cases. Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) was identified as a beta coronavirus, the same subtype as its predecessors SARS-CoV and the Middle East Respiratory Syndrome (MERS-CoV). With the rapid spread of cases, the WHO declared a pandemic of coronavirus disease of 2019 (COVID-19) on March 11, 2020.

Severe Acute Respiratory Syndrome Coronavirus 2 causes essential alterations in the cardiovascular system beyond the initial damage to the respiratory system (Zheng et al., 2019). Alterations include the vascular endothelium-mediated recruitment of inflammatory leukocytes that contribute to tissue damage and cytokine release. These alterations are critical drivers of the acute respiratory distress syndrome and disseminated intravascular coagulation and COVID-19-associated cardiovascular complications.

During pregnancy, physiological adaptive changes in the immune, respiratory, cardiovascular, and coagulation systems could modulate COVID-19 presentation. As pregnant women are at high risk of complications and severe disease from infection with other coronaviruses, they were identified as a vulnerable group and were advised to take additional precautions as the COVID-19 pandemic unfolded. Accordingly, later evidence has shown that pregnant women seem to be associated with a greater susceptibility to contagion (Wastnedge et al., 2021), presenting more severe forms of the disease (Liu H. et al., 2020), or a high risk for pregnancy complications (di Mascio et al., 2020). In addition, data have confirmed SARS-CoV-2 vertical transmission, although short- and long-term sequels are still under investigation. Despite that, histological studies of placentas from SARS-CoV-2 positive pregnant women showed poor uterine-placental perfusion, with signs of placental infarcts, atheromas in the decidua vessels chorioangioma, and edema in placental villi (Mulvey et al., 2020; Shanes et al., 2020).

This review summarizes the current literature on SARS-CoV-2 virus infection in pregnancy. In addition, we describe pregnancy-associated adaptations in the angiotensin-converting enzyme 2 (ACE2), a protein identified as SARS-CoV-2 receptor, which could influence the presentation of COVID-19. This review particularly emphasized evidence coming from Latin American countries.


PATHOPHYSIOLOGY OF SEVERE ACUTE RESPIRATORY SYNDROME CORONAVIRUS 2 INFECTION

Severe Acute Respiratory Syndrome Coronavirus 2 is an encapsulated positive single-stranded RNA virus, a coronavirus (Elias et al., 2021). The coronavirus family comprises numerous viruses with the capacity to infect several species (Channappanavar et al., 2014). So far, some of them, including coronaviruses 229E and NL63, display the capacity to affect

humans, generating symptoms of a standard cold (Rabi et al., 2020). The comprising ability of these viruses to bind host cells requires a projection of the membrane structure, called “spike.” This transmembrane structure is composed of a trimetric glycoprotein protuberance, composed of two subunits. The first one, S1, is the binding region to the host cell receptor and, S2 is the region where the virus and the host cellular membranes may fuse (Tang et al., 2020). Person-to-person transmission has been demonstrated through drops, contact, and aerosols spread, but less frequently due to fecal-oral and fomites transmission. The incubation period is around 5 days (5.79–6.97 days) (Elias et al., 2021). For diagnosis, the standard gold technique is the real-time PCR (RT-PCR), which detects the presence of SARS-CoV-2 virus RNA.

At the onset of COVID-19, the most common symptoms are fever, cough, and fatigue, while other symptoms include sputum production, headache, diarrhea, dyspnea, and lymphopenia (Rothan and Byrareddy, 2020). Once inside the host, SARS-CoV-2 activates the innate and adaptive immune responses and elicits a pronounced lymphopenia due to impaired lymphopoiesis and increased lymphocyte apoptosis (Wiersinga et al., 2020). Around 7–8 days after the onset of signs/symptoms, some individuals progress to a more clinically compromised condition and develop pneumonia with respiratory distress requiring hospitalization. In some cases, sepsis appears around days 9–10, and severe acute respiratory syndrome occurs on days 9–12. In some individuals, the start of mechanical ventilation and admission to the Intensive Care Unit (ICU) is required approximately at 10.5 days. Complications, such as heart or kidney injury, usually occur on day 15, secondary infection on day 17, and death or recovery on days 19–22 (Huang et al., 2020; Zhou F. et al., 2020). Some factors that seem to be associated with a higher risk of infection and severe disease are age (>70 years old), male sex, tobacco use, and presence of comorbidities, such as chronic diseases (Cai, 2020; Jin et al., 2020; Jordan et al., 2020; Zheng et al., 2020). Additionally, a high score on the Sequential Organ Failure Assessment (SOFA) and values >1 µg/ml for D-dimer have been associated with higher mortality (Zhou F. et al., 2020).

The mechanisms of SARS-CoV-2 invasion of the host system are illustrated in **Figure 1**. Briefly, the coronavirus virion has structural proteins: nucleocapsid (N), membrane (M), envelope (E), and spike (S) proteins. The entry steps of the viral particles—encompassing attachment to the host cell membrane and fusion—are mediated by the S glycoprotein. S protein is assembled as a homotrimer and is inserted in multiple copies into the virion membrane giving it its crown-like appearance. In addition, the furin-like proteases, transmembrane protease, serine 2 (TMPRSS2), and cathepsin L are involved in the virus invasion process. First, the S protein binds to the ACE2 receptor, after which the virus penetrates the host cells by endocytosis (S1 region) or membrane fusion (S2 region). Cleavage of the S1–S2 boundary is necessary for initiating the membrane-fusion process. After S1–S2 is cleaved, the S2 site activates the fusion process either by TMPRSS2 on the cell surface or by cathepsins in endosomes. Then, the fusion between viral and cellular membranes forms a pore through which viral RNA is released



into the host cell cytoplasm for uncoating and replication (Zhou P. et al., 2020; Jackson et al., 2021).

Interestingly, the main characteristic of COVID-19 is endothelial cell damage (Evans et al., 2020). Thus, infection with SARS-CoV-2 reduces ACE2-mediated regulation of vascular tone and causes endothelial dysfunction at multiple levels, such as inflammatory activation, cytokine storm, leukocyte infiltration, increased permeability, thrombosis, platelet aggregation, vasoconstriction, reactive oxygen species (ROS) production, and apoptosis (Evans et al., 2020).

ANGIOTENSIN-CONVERTING ENZYME 2 AND THE IMPLICATION OF THE RENIN-ANGIOTENSIN SYSTEM ON CORONAVIRUS DISEASE 2019

Angiotensin-converting enzyme 2 is constitutively expressed in several tissues, such as the lung, heart, kidney, and blood vessels (Zou X. et al., 2020). Therefore, SARS-CoV-2 has a particular tropism for the pulmonary system, initiating pronounced respiratory symptoms (Jia et al., 2005). In addition,

the renin-angiotensin system is a critical modulator of vascular function, as a key target of the SARS-CoV-2 infection (Watanabe et al., 2005; Fraga-Silva et al., 2013; Santos, 2014; Sanchis-Gomar et al., 2020). In particular, ACE2 is a pleiotropic peptidase that metabolizes Angiotensin II (Ang II) to Ang 1–7, two peptides with divergent physiological functions. Thus, while Ang II evokes vasoconstrictive, proliferative, and angiogenic effects, Ang 1–7 elicits anti-proliferative, anti-angiogenic, and vasodilator functions (Bharadwaj et al., 2011). Moreover, ACE2 has a protective effect on the endothelium and improves endothelial function, apparently mediated by Ang 1–7 production (Lovren et al., 2008; Fraga-Silva et al., 2013; **Figure 1**). The relevance of ACE2 has been confirmed in animals deficient in this enzyme. Those animals exhibit increased oxidative stress and pro-inflammatory cytokines, compromising cardiovascular function (Wang et al., 2020). Likewise, previous studies on the coronavirus-induced severe acute respiratory syndrome, sepsis, or acid aspiration-induced lung injury have shown that dysfunction of the renin-angiotensin system is involved in these conditions (Zhang et al., 2018; Li H. et al., 2020; Rezaei et al., 2021). Huang et al. (2014) reported that high plasma levels of Ang II were associated with the severity of the disease and also those high levels predicted fatal outcomes due to the H7N9 influenza

virus. In Kuba et al. (2005) described that the function of ACE2 is impaired by the binding of viral protein S, enhancing the Ang II circulating levels, and leading to hemodynamic alterations characterized by vasoconstriction.

The critical role of ACE2 in the COVID-19 pathophysiology raised speculations about ACE2 as a therapeutic target and the implications of the use of renin-angiotensin system inhibitors. Since ACE2 inhibitors and Ang II AT1 receptor antagonists have the potential to increase ACE2 expression, which may aggravate COVID-19, we recommend critical publications regarding the use of renin-angiotensin inhibitors and COVID-19 (Bavishi et al., 2020; Peiró and Moncada, 2020; Sanchis-Gomar et al., 2020; Sommerstein et al., 2020; Vaduganathan et al., 2020). Yet the presence of the soluble and active isoform of ACE2 may be beneficial in COVID-19 patients since soluble ACE2 may compete with the membrane enzyme, limiting its role as a SARS-CoV-2 receptor, and decreasing the Ang II plasma levels (Khodarahmi et al., 2021). As a therapeutic target, the restoration of ACE2 through the administration of recombinant ACE2 may reverse the lung-injury process (Monteil et al., 2020, 2021; Siri wattananon et al., 2021; Zhang et al., 2021).

Therefore, several studies emphasize the critical role of ACE2 in viral infection, the clinical presentation of COVID-19, the potential use of its products (i.e., metabolites) as biomarkers. In addition, the use of drugs that modulate ACE2 has to be taken into account in the treatment and clinical evolution of COVID-19.


PREGNANCY IS A PHYSIOLOGICAL MODULATOR OF ANGIOTENSIN-CONVERTING ENZYME 2 EXPRESSION AND ACTIVITY

Two physiological conditions can modulate ACE2 levels and activity: aging and pregnancy (AlGhatrif, 2020; Huang et al., 2020; Ludvigsson, 2020; Zhou F. et al., 2020). Throughout gestation, a high expression of ACE2 in the human placenta, particularly in the decidua, the syncytiotrophoblast, and the villous stroma, may increase Ang 1–7 blood levels. Specifically, Liu D. et al. (2020) reported that ACE2 expression was upregulated between 6 and 16 weeks and downregulated in term human placentas. Plasmatic levels of ACE2 and Ang 1–7 are significantly augmented during pregnancy (Emanuele et al., 2002; Tamanna et al., 2020; Nobrega Cruz et al., 2021). On the other hand, Ang II was also increased in pregnancy (Emanuele et al., 2002). In this physiological scenario, it is believed that increased expression of ACE2 and blood levels of Ang 1–7 could be counteracting the increased stimulus of Ang II levels. Alternatively, the highest prevalence of a monomeric form of AT1, which is less sensitive to Ang II, may also reduce vascular sensitivity to Ang II. These changes would participate in the adaptive physiological mechanisms of the cardiovascular system during pregnancy, resulting in decreased peripheral vascular resistance and vasodilation of the maternal

vasculature (Gant et al., 1973; AbdAlla et al., 2001; Levy et al., 2008; Marques et al., 2011; Pringle et al., 2011; Stettner et al., 2013), increased aldosterone and Ang II, promoting water and sodium retention, and increased blood volume (Scaife and Mohaupt, 2017). In addition, other critical factors modulated by the renin-angiotensin axis, increased during pregnancy, include the vasodilators prostaglandin E2, nitric oxide (Gant et al., 1980; Corthorn et al., 2006), and bradykinin (Knock and Poston, 1996). Therefore, pregnancy constitutes a physiological condition with major vascular adaptations characterized by reduced systemic vascular resistance that allows the homeostatic control of pregnancy-related hemodynamic changes, including increased cardiac output, expanded blood volume, and reduced blood pressure. Relevance of the renin-angiotensin axis in pregnancy is also remarkable in conditions in which deregulation of this axis impairs endothelial function and leads to pregnancy complications, such as hypertension or pre-eclampsia (Anton et al., 2008; Gilbert et al., 2008). Thus, it is not surprising that pregnant women constitute a potentially vulnerable population in the COVID-19 pandemic, with initial results indicating that the clinical response of pregnant women to COVID-19 could be related to physiological changes in expression levels of ACE2 and reduced sensitivity to Ang II (Anton et al., 2008; Gilbert et al., 2008; Liu D. et al., 2020; Mendoza et al., 2020).

CLINICAL PRESENTATION OF SEVERE ACUTE RESPIRATORY SYNDROME CORONAVIRUS 2 INFECTION IN PREGNANT WOMEN

Some studies have reported that SARS-CoV-2 infection in pregnant women behaves similarly to the general population, contrary to what has been reported with other types of coronavirus infection in the past (Li N. et al., 2020; Yang et al., 2020; Fan et al., 2021). The disease manifests itself with typical symptoms and occurs with different degrees of severity, such as mild disease in 81–86% of cases, severe disease in 9.3–14%, and critical disease in 5%; values close to those reported in the general population (80, 15, and 5%, respectively) (Han et al., 2020). The most common symptoms were fever (65%), cough (60%), and shortness of breath or dyspnea (24%) (Han et al., 2020). Around 5% of mothers were admitted to the ICU; intubation was carried out in 35.87% of patients (Han et al., 2020). The rate of maternal death was <0.01%. Nevertheless, another study estimated the mortality rate in the pregnant population close to 2.7% and an ICU admission rate of 6–8% (Lambelet et al., 2020). In addition, Ellington et al. (2020) reported similar symptoms, with a greater risk of hospitalization, ICU intervention, and mechanical ventilation requirement, yet without an increased risk of death, in pregnant women when compared to non-pregnant women. It is noteworthy that, in this cohort, the group of pregnant women reported a higher frequency of comorbidities (chronic lung disease 22%, diabetes mellitus 15%, and cardiovascular disease 14%) when compared to non-pregnant women (Ellington et al., 2020).



Severe Acute Respiratory Syndrome and MERS have been associated with miscarriage, intrauterine death, fetal growth restriction, and high case fatality rates (Wang et al., 2021). This linkage was also presented in the pandemic of SARS-CoV-2. Thus, infection in pregnant women was associated with a high risk of adverse pregnancy outcomes, such as intrauterine growth restriction, premature rupture of membranes, fetal distress, preterm delivery (Mullins et al., 2020), spontaneous abortion, and stillbirth (Della Gatta et al., 2020). In particular, Han et al. (2020) reported that premature delivery reached 25%. The rate of low birth weight (<2,500 g) was close to 31%, and neonatal intensive care unit (NICU) admission was 24%. Positive nasopharynx swabs or sputum from newborns was <0.01% (Han et al., 2020; **Figure 2**). More recently, a meta-analysis published by Wei et al. (2021) included 42 studies that involve 438,548 pregnant women. They found that COVID-19 was associated with pre-eclampsia [odds ratio (OR) 1.33], preterm birth (OR 1.82), and stillbirth (OR 2.11). In addition, they found that when compared with mild COVID-19, severe COVID-19 was strongly associated with pre-eclampsia (OR 4.16), preterm birth (OR 4.29), gestational diabetes (OR 1.99), and low birth weight

(OR 1.89) (Wei et al., 2021). Another meta-analysis, including twenty-eight studies with 790,954 pregnant women, concluded that SARS-CoV-2 infection during pregnancy was associated with a 58% increased risk of pre-eclampsia. In addition, there was a statistically significant increase in the risk of pre-eclampsia with severe features (OR 1.76, $p < 0.05$), eclampsia (OR 1.97, $p < 0.05$), and Hemolysis, Elevated Liver enzymes, and Low Platelets (HELLP) syndrome (OR 2.10, $p < 0.05$) among pregnant women with SARS-CoV-2 infection, as compared to those without the infection (Papageorghiou et al., 2021). In addition, there is a high prevalence of cesarean delivery, whose main indication seems to be an underlying obstetric condition, such as pre-eclampsia, fetal distress, or premature rupture of membranes, and not the clinical condition of COVID-19 in pregnant women (Della Gatta et al., 2020; Trad et al., 2020).

Conversely, other studies reported that SARS-CoV-2 infection during the first trimester of pregnancy does not seem to predispose to early pregnancy loss (Cosma et al., 2020; Rotshenker-Olshinka et al., 2020; Bortoletto et al., 2021). In addition, Rizzo et al. (2021) reported that SARS-CoV-2 infection did not increase the risk of developing fetal growth restriction. At the same time, Meyer R. et al. (2021) showed a decrease in preterm delivery in pregnant women in Israel, possibly due to the reduction of iatrogenic preterm births, avoidance of infections, or reduced stress levels related to the lockdown policy (Meyer R. et al., 2021). However, the question about the differentiation between spontaneous or iatrogenic preterm delivery remains to be elucidated (Della Gatta et al., 2020). Moreover, there was no association between abnormal umbilical artery Doppler results, defined as a composite of increased S/D ratio, absent end-diastolic velocity, reversed end-diastolic velocity, and COVID-19 infection in growth-restricted pregnancies (Ona et al., 2021).

Differences in these reports depend on several factors. First, we propose that it is relevant to consider that a significant percentage of infected pregnant women could be asymptomatic or become undiagnosed. Remarking this issue, Sutton et al. (2020) reported that in a screening testing of COVID-19 in 215 pregnant women in New York (United States), 211 women were asymptomatic, but 13.7% tested positive to SARS-CoV-2. Thus, the prevalence of COVID-19 in pregnant women may be underestimated and therefore the associated perinatal complications (Lambelet et al., 2020; Sutton et al., 2020).

SEVERE ACUTE RESPIRATORY SYNDROME CORONAVIRUS 2 IN LATIN AMERICA

There are several publications about the SARS-CoV-2 pandemic in Latin America. Most of the articles are reports of a few cases in local hospitals. However, the Iberoamerican Society of Neonatology (SABEN) recruited women with SARS-CoV-2 infection to provide knowledge and experiences on perinatal COVID-19 in Latin America (Sola et al., 2020). Their results described 86 pregnant women with COVID-19 from 11 units of 7 countries: Argentina, Colombia, Ecuador, Equatorial Guinea, Honduras, Peru, and the Dominican Republic. Of these, 68%

women were asymptomatic for COVID-19, and 32% women exhibited symptoms. In total, 89% of symptomatic women had mild symptoms or signs, while 3.5% women had severe respiratory symptoms. Six women were admitted to intensive care, and no woman died. In addition, 94% were term, and 6% were preterm pregnancies. The swab result was positive in 7% of the newborns, with no causalities. Nevertheless, another study that includes 40 obstetric patients diagnosed with COVID-19 from four countries (Peru, Colombia, Bolivia, and Paraguay) reported a maternal mortality rate of 15% (six cases) and a perinatal mortality rate of 2.5% (one case). Associated pathologies included severe pre-eclampsia (25%), HELLP (5%), and gestational hypertension (12.5%), while ten patients received invasive mechanical ventilation since admission to ICU (25%) (Viruez-Soto et al., 2021; **Figure 3**).

In Hispanic women, obesity, advanced maternal age, medical comorbidities, and antepartum admissions related to COVID-19 have been reported as risk factors associated with adverse maternal and neonatal outcomes (Brandt et al., 2020). In Brazil, a report identified 20 COVID-19-related maternal deaths. Symptoms onset was reported during pregnancy for 12 cases, post-partum for three cases, and during the cesarean section for one case (missing data for four). In 16 cases, death occurred in the post-partum period. At least one comorbidity or risk factor was present in 11 cases (missing data for four). Asthma was the most common risk factor (5/11) (Takemoto et al., 2020).

Although Latin America has one of the highest COVID-19 death rates in the world (Editorial The Lancet, 2020), the data worldwide do not show a severe impact on maternal and fetal health (Sánchez-Duque et al., 2020). However, it is worth mentioning that only severe and critically ill patients have been tested for COVID-19, so the number of infected people (including pregnant and post-partum women) is more likely underestimated (Takemoto et al., 2020).

PLACENTAL INFECTION BY SEVERE ACUTE RESPIRATORY SYNDROME CORONAVIRUS 2 AND POTENTIAL PATHOPHYSIOLOGICAL MECHANISM

There is a significant concern regarding eventual transplacental transmission, infection at delivery or through breast milk, and the general care measures for pregnant women and neonates are being systematized. Indeed, convincing pieces of evidence have detected SARS-CoV-2 in the human placenta (Algarroba et al., 2020a; Baud et al., 2020; Best Rocha et al., 2020; Chen et al., 2020; Fenizia et al., 2020; Ferraiolo et al., 2020; Hecht et al., 2020; Hosier et al., 2020; Kirtsman et al., 2020; Kulkarni et al., 2020; Patanè et al., 2020; Richtmann et al., 2020; Smithgall et al., 2020; Vivanti et al., 2020; Hsu et al., 2021; Marinho et al., 2021). Regarding localization of SARS-CoV-2 in the placenta, most of the studies with positive results detected the presence of the virus in the syncytiotrophoblast (Best Rocha et al., 2020; Hecht et al., 2020; Hosier et al., 2020; Patanè et al., 2020; Sisman et al., 2020; Vivanti et al., 2020) and only the study of Hsu et al. (2021) detected the

presence of SARS-CoV-2 in the fetal endothelium. Despite that, other studies did not detect SARS-CoV-2 in the placenta (Edlow et al., 2020; Flores-Pliego et al., 2021; Halici-Ozturk et al., 2021; Levitan et al., 2021; **Table 1**). Therefore, although data are not consistent, there may be vertical transmission in some cases.

Interestingly, there is also evidence for *in utero* transmission of SARS-CoV-2 infection. For instance, detection of SARS-CoV-2 virus in nasopharyngeal samples of newborns and anti-SARS-CoV-2 antibodies detected in the umbilical cord blood in at least one case were reported (Fenizia et al., 2020). In addition, Sisman et al. (2020) presented a preterm infant with placental SARS-CoV-2 infection and positive nasopharyngeal testing at 24 and 48 h of life, who developed fever and mild respiratory disease on the second day of life.

Whether ACE2 and TMPRSS2 are entry mechanisms for SARS-CoV-2 in the placenta remains unclear, although both proteins are expressed in the trophoblast and fetal endothelium. Thus, ACE2 was detected in syncytiotrophoblast (Edlow et al., 2020; Hecht et al., 2020; Taglauer et al., 2020; Lu-Culligan et al., 2021) and endothelium of umbilical arteries (Valdés et al., 2006). Nevertheless, a study conducted in isolated cells from the placenta shows that the co-expression of ACE2 and TMPRSS2 is very low in syncytiotrophoblast, cytotrophoblast, and extravillous trophoblast, especially when it is compared with the expression of cytomegalovirus and Zika virus receptors. These pathogens have been proven to cross the placenta and generate vertical transmission to the fetus (Pique-Regi et al., 2020). In addition, Li M. et al. (2020) reported high expression of ACE2 and TMPRSS2 in syncytiotrophoblast and cytotrophoblast in the single-cell evaluation of trophoblasts in the human placenta. Despite this evidence, it is necessary to investigate whether the expression of both proteins in the placenta may be affected in pregnancy complicated by SARS-CoV-2 infection since a weak expression of TMPRSS2 in the villous endothelium of SARS-CoV-2 positive pregnant women was reported (Edlow et al., 2020; Hecht et al., 2020).

Additionally, fetal sex could affect the placental expression of ACE2. It has been reported that the male sex is associated with a higher ACE2 gene expression (Chlamydas et al., 2021). Moreover, it is known that the ACE2 gene escapes X inactivation (Tukiainen et al., 2017), *in vitro* study shows that 17 β -estradiol, a primarily female sex steroid, can downregulate the ACE2 gene expression in non-pregnant women (Stelzig et al., 2020), providing another hypothesis for the sex-based expression differences of this gene. However, there is no information about sex-dimorphism in the placental expression of ACE2 in pregnant women with COVID-19.

On the other hand, detecting SARS-CoV-2 in placentas from fetal death cases alerted the possible occurrence of placental dysfunction associated with the maternal medical condition (Baud et al., 2020; Hosier et al., 2020; Richtmann et al., 2020). These studies suggest that early infection (first or second trimester of pregnancy) could generate a state of more significant deterioration in the placental function associated with exacerbated inflammation, negatively impacting fetal wellbeing and development. Supporting this pathophysiological explanation, a recent case report from




TABLE 1 | Studies that assess the detection of Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) in placenta around the world indicate the technique used and the state of symptoms.

Method of detection of SARS-CoV-2 in placenta				Presence of COVID-19 symptoms		Total cases	Country	References
qPCR	IHC	ISH	EM	+	-			
Neg	ND	ND	ND	1		1	Turkey	Kalafat et al., 2020
Neg	ND	ND	ND	1		1	China	Peng et al., 2020
Pos(1)	ND	ND	ND	11		11	United States	Penfield et al., 2020
ND	ND	ND	Pos	1		1	United States	Algarroba et al., 2020b
ND	Pos(1)	Pos(2)	ND	ND	ND	19	United States	Hecht et al., 2020
ND	Pos	ND	ND		1	1	United States	Hsu et al., 2021
Pos	ND	ND	ND	1		1	India	Kulkarni et al., 2020
Pos	ND	ND	ND	4	1	5	Brazil	Richtmann et al., 2020
ND	Neg	Neg	ND	50	26	76	United States	Smithgall et al., 2020
Pos	Pos	ND	ND	ND	ND	1	United States	Vivanti et al., 2020
ND	Pos	ND	Pos	1		1	United States	Sisman et al., 2020
ND	Pos	Pos	ND	1	1	2	United States	Best Rocha et al., 2020
Pos	ND	ND	ND		1	1	Italy	Ferraiolo et al., 2020
Pos	Pos	Pos	Pos	1		1	United States	Hosier et al., 2020
ND	Pos	Pos(2)	ND	3		3	Italy	Patanè et al., 2020
Pos	ND	ND	ND	3		3	China	Chen et al., 2020
Pos	ND	ND	ND	1		1	Switzerland	Baud et al., 2020
Pos	ND	ND	ND	1		1	Canada	Kirtsman et al., 2020
Pos (2)	ND	ND	ND	31		31	Italy	Fenizia et al., 2020
ND	ND	Neg	ND	15	32	47	United States	Edlow et al., 2020
Neg	ND	ND	ND	11		11	Mexico	Flores-Pliego et al., 2021
Neg	ND	ND	ND	5	19	24	Turkey	Halici-Ozturk et al., 2021
Pos	Pos	Pos	Pos	1		1	Mexico	Valdespino-Vázquez et al., 2021

qPCR, real-time PCR; IHC, immunohistochemistry; ISH, In situ hybridization; EM, electron microscopy; pos, positive; Neg: negative; ND, not determined.

Brazil showed an association of fetal death at 34 weeks of gestation with histopathological alterations in the placenta. The findings are consistent with intense acute placenta vascular malperfusion, with the detection of SARS-CoV-2 in the placenta, umbilical cord, and fetal tissues (lung, heart, and brain,

among others). Furthermore, the patient had mild symptoms of COVID-19 but, at the time of fetal death, had pro-inflammatory and pro-coagulant syndrome characterized by higher levels of interleukins, ferritin, and D-dimer (Marinho et al., 2021). Therefore, further research is needed to elucidate the

involvement of placental dysfunction in COVID-19 cases and its perinatal consequences.

PLACENTAL VASCULAR DISORDERS ASSOCIATED WITH SEVERE ACUTE RESPIRATORY SYNDROME CORONAVIRUS 2

Regarding the vascular alterations of the placenta, histological studies of placentas from SARS-CoV-2 positive pregnant women showed poor uterine-placental perfusion, with signs of placental infarcts, atheromas in decidua vessels, chorioangioma, and edema in the placental villi (Mulvey et al., 2020; Shanes et al., 2020). Those studies also associated vascular thrombosis in the chorionic plate and decreased capillaries density in the chorionic villi in placentas with SARS-CoV-2 infection. Moreover, Baergen and Heller (2020) described that this viral infection was associated with a high risk of fetal vessel thrombosis and reduced vascularization in the placental microcirculation. Similarly, Patberg et al. (2020) showed that SARS-CoV-2 infection is associated with placental signs of reduced perfusion and villitis of unknown etiology. Another study confirmed these findings, including asymptomatic or mildly symptomatic SARS-CoV-2 positive pregnant women, showing evidence of fetal vascular malperfusion (FVM): chorioangioma, intramural fibrin deposition, and vascular ectasia. Additionally, perivillous fibrin deposition was also significantly higher in placental histopathology (Jaiswal et al., 2021). These results agree with Meyer J. et al.'s (2021) findings showing that 77% of placentas infected with SARS-CoV-2 showed one or more features of maternal vascular malperfusion (MVM). Therefore, growing pieces of evidence, with some exceptions (Zhang et al., 2020; Levitan et al., 2021; Lu-Culligan et al., 2021), have shown placental vascular disorders in pregnant women with COVID-19.

Whether the severity of COVID-19 in pregnancy is related to placental alterations is under investigation. For instance, high trophoblast necrosis was found in pregnant women who required respiratory support or intubation for COVID-19 when compared with non-hypoxic patients (Meyer J. et al., 2021). Moreover, Edlow et al. (2020) found that placental lesions associated with MVM were increased with the severity of COVID-19 in pregnant women. Remarkably, in patients with COVID-19 admitted to the ICU, increased levels of von Willebrand Factor (vWF) antigen and P-selectin in plasma were detected, indicating endothelial dysfunction (Goshua et al., 2020). In addition, in the placenta of pregnant women with severe COVID-19, higher expression of vWF was associated with lower claudin-5 and vascular endothelial (VE-cadherin) in the endothelium from decidua and chorionic villi (Flores-Pliego et al., 2021). These findings suggest that COVID-19 induces endothelial cell injury in the placenta, probably affecting the endothelial barrier, the anti-thrombotic capacity of the endothelium, and the overall function of placental vessels. Compromised placental function, placental hypoxia, and a hypercoagulable state are probably related to the severity of the infection (Ng et al., 2006; Baergen and Heller, 2020;


Shanes et al., 2020; Marinho et al., 2021; Meyer J. et al., 2021; Poisson and Pierone, 2021; **Figure 4**).

In addition, SARS-CoV-2 positive mothers showed higher levels of inflammatory markers. Thus, Fenizia et al. (2020) showed significant increases in inflammatory markers (including interleukin 1 β or interleukin 6) in both placenta and plasma of two women whose placenta was tested positive for SARS-CoV-2. Accordingly, it was previously reported that the presence of SARS-CoV-2 in the placenta generated an increased inflammatory response in the intervillous space (histiocytic intervillitis) and chorionic villi (villitis), with the presence of macrophages (CD68) and T lymphocytes (CD3) in the intervillous space (Hecht et al., 2020; Hosier et al., 2020; Kirtsman et al., 2020; Patanè et al., 2020; Sisman et al., 2020; Smithgall et al., 2020; Vivanti et al., 2020; Hsu et al., 2021). Transcriptome analysis of placentas from pregnant women with COVID-19 showed increased genes associated with immune response. Specifically, Lu-Culligan et al. (2021) found a marked increase of pro-inflammatory genes and chemokines in both immune and non-immune cell types in placentas from COVID-19 cases. In addition, single-cell transcriptome analysis revealed significant enrichment of genes encoding cytotoxic proteins in natural killer (NK) cells, associated with upregulation of the activation marker CD69 in T-cells, increased expression of interferon-induced protein ISG15, and the regulators of nuclear factor- κ B (NF κ B) pathway in endothelial cells (Lu-Culligan et al., 2021).

Despite these findings, the potential implications of vascular alterations of the placenta in COVID-19 cases and fetal development are not entirely understood (Ahlberg et al., 2020; Díaz-Corvillón et al., 2020; Khalil et al., 2020; Shanes et al., 2020). Therefore, there is a demanding necessity to continue gathering information, aiming to elucidate the potential implication of impaired placental environment observed in women with COVID-19 in contributing to adverse perinatal outcomes, particularly in periods of the high spread of the pandemic.

THE IMPACT OF CORONAVIRUS DISEASE 2019 VACCINE ON PREGNANCY

Pregnant women have traditionally been excluded from vaccine trials. Without appropriate evidence about safety and efficacy during pregnancy, they have previously been denied the opportunity to receive vaccines that would have protected them and their offspring. This situation has also been present in the context of the COVID-19 pandemic, and pregnant women, their providers, and health policymakers would have to make unnecessarily tricky decisions because of inadequate evidence about vaccine use in pregnancy. This would lead to less vaccine use and its afforded protections in this population (Beigi et al., 2021). In the V-SAVE pregnancy registry with 3,958 participants who received messenger RNA (mRNA) COVID-19 vaccines, 827 had a completed pregnancy, of which 115 (13.9%) resulted in a pregnancy loss and 712 (86.1%) resulted in a live birth (mostly among pregnant women vaccinated in the third trimester). Adverse neonatal outcomes included preterm



birth (in 9.4%) and small size for gestational age (in 3.2%), and no neonatal deaths were observed. Although not directly comparable, calculated proportions of adverse pregnancy and neonatal outcomes in persons vaccinated against COVID-19 who had a completed pregnancy were similar to incidences reported in studies involving pregnant women that were conducted before the pandemic (Shimabukuro et al., 2021). On the other hand, a recent study in 24,288 singleton pregnancies showed no evident differences, in terms of adverse neonatal and early infant outcomes, between newborns of women who received BNT162b2 mRNA vaccination during pregnancy vs. those of women who were not vaccinated. This study contributes to current evidence in establishing the safety of prenatal vaccine exposure to newborns. However, the interpretation of study findings is limited by the observational design (Goldshtein et al., 2022).

Clinical trials demonstrate that vaccination effectively prevents severity and symptomatic COVID-19 in non-pregnant persons. To highlight, the risks of administering COVID-19 vaccines during pregnancy appear to be minimal. Commonly reported side effects are short-term injection site pain, headache, fever, myalgia, arthralgia, chills, and nausea (National Center for Immunization and Respiratory Diseases, 2021). Moreover, the rate of serious adverse effects has been relatively low (National Center for Immunization and Respiratory Diseases, 2021). In addition, chemical components of the vaccines are not specifically contraindicated in pregnancy (National Center for Immunization and Respiratory Diseases, 2021). About the recommendations, the American College of Obstetrics and Gynecology states that in the absence of data showing that vaccines are contraindicated, then pregnant patients should be immunized (American College of Obstetricians and Gynecologists' Immunization Infectious Disease and Public Health Preparedness Expert Work Group et al.,

2021). The Center for Disease Control has taken a similar position, declaring that the only absolute contraindication to vaccination is an allergy to vaccine components (National Center for Immunization and Respiratory Diseases, 2021). However, the World Health Organization has more reserved recommendations, indicating that vaccination is only indicated in pregnant women who are at high risk for exposure to COVID-19 (healthcare workers, or those with comorbidities that might make disease more severe) (Chavan et al., 2021; World Health Organization, 2021).

CONCLUDING REMARKS

In this manuscript, we have discussed the vulnerability of the pregnant population to COVID-19 infection. This manuscript also described many uncertainties; however, they are not the only ones in this field. For example, the real incidence of COVID-19 in pregnant women is a fundamental question that needs clarification. In addition, whether COVID-19 is associated with a greater risk of severe disease and perinatal complications requires confirmatory population studies. This information would help to generate appropriate public health policies for this particular population.

In addition, we also have remarked that infected placenta with SARS-CoV-2 showed alterations related to inflammatory processes associated with damages to the vascular network. Whether these placental alterations might explain the adverse perinatal outcomes in women with COVID-19 requires confirmatory studies. In addition, although there is vertical transmission in some cases, vertical transmission data are not consistent. We encourage future research to elucidate whether SARS-CoV-2 infection affects fetal programming, as well as the future health of both mother and offspring.

Concerning the vaccine, pregnant women and physicians need to use the limited available data to weigh the benefits and risks of the COVID-19 vaccine during pregnancy, considering the patient's specific risk of SARS-CoV-2 exposure. Currently, there is the absence of evidence that supports pregnancy as a contraindication, and it seems that the benefits of receiving the vaccine far outweigh the unlikely potential harms. However, estimates of global vaccination among pregnant women are yet unknown. Moreover, we remark inequities in the access to the SARS-CoV-2 vaccines worldwide, but, in particular, in Latin America. For example, only 56% of the Latin American people have been vaccinated up to December 2021 (PAHO, 2021), with a significant difference among countries (PAHO, 2022). In addition, it is necessary to delve into how the pandemic has impacted the care of pregnant women in general since substantial and heterogeneous modifications have been reported in maternity services (Jardine et al., 2021).

In conclusion, the available information highlights the greater vulnerability of pregnant women in the context of a pandemic. However, more studies are required to better understand the potential impact of the pandemic on pregnant women, especially in Latin American nations.

REFERENCES

- Abdalla, S., Lother, H., el Massiery, A., and Quitterer, U. (2001). Increased AT(1) receptor heterodimers in preeclampsia mediate enhanced angiotensin II responsiveness. *Nat. Med.* 7, 1003–1009. doi: 10.1038/NM0901-1003
- Ahlberg, M., Neovius, M., Saltvedt, S., Söderling, J., Pettersson, K., Brandkvist, C., et al. (2020). Association of SARS-CoV-2 Test Status and Pregnancy Outcomes. *JAMA* 324:1782. doi: 10.1001/jama.2020.19124
- Algarroba, G. N., Hanna, N. N., Rekawek, P., Vahanian, S. A., Khullar, P., Palaia, T., et al. (2020a). Confirmatory evidence of the visualization of severe acute respiratory syndrome coronavirus 2 invading the human placenta using electron microscopy. *Am. J. Obstet. Gynecol.* 223, 953–954. doi: 10.1016/j.ajog.2020.08.106
- Algarroba, G. N., Rekawek, P., Vahanian, S. A., Khullar, P., Palaia, T., Peltier, M. R., et al. (2020b). Visualization of severe acute respiratory syndrome coronavirus 2 invading the human placenta using electron microscopy. *Am. J. Obstet. Gynecol.* 223, 275–278. doi: 10.1016/j.ajog.2020.05.023
- AlGhatrif, M. (2020). The dilemma of Coronavirus Disease 2019, Aging, and Cardiovascular Disease. *JAMA Cardiol.* 5, 747–748. doi: 10.1001/jamacardio.2020.1329
- American College of Obstetricians and Gynecologists' Immunization Infectious Disease and Public Health Preparedness Expert Work Group, Riley, L., Beigi, R., Jamieson, D., Hughes, B., and Swamy, G. (2021). *COVID-19 Vaccination Considerations for Obstetric-Gynecologic Care | ACOG*. Available online at: https://www.acog.org/clinical/clinical-guidance/practice-advisory/articles/2020/12/covid-19-vaccination-considerations-for-obstetric-gynecologic-care?utm_source=redirect&utm_medium=web&utm_campaign=int (accessed August 24, 2021).
- Anton, L., Merrill, D. C., Neves, L. A. A., Stovall, K., Gallagher, P. E., Diz, D. I., et al. (2008). Activation of local chorionic villi angiotensin II levels but not angiotensin (1-7) in preeclampsia. *Hypertension* 51, 1066–1072. doi: 10.1161/HYPERTENSIONAHA.107.103861
- Baergen, R. N., and Heller, D. S. (2020). Placental Pathology in Covid-19 Positive Mothers: preliminary Findings. *Pediatr. Dev. Pathol.* 23, 177–180. doi: 10.1177/1093526620925569

AUTHOR CONTRIBUTIONS

PA-R, MG, and RG-R conceived, designed, planned, and supervised the manuscript. PA-R, MG, and CE critically review the manuscript and generate the final published version. LQ-A, FG, RA, CE, and AD wrote the manuscript. All authors provided critical feedback and approved the final version.

FUNDING

This work was supported by the Pontificia Universidad Javeriana, Federal University of Mato Grosso, Federal University of Goiás, Universidad de Concepción, and Universidad de Buenos Aires. CE was financially supported by Fondecyt 1200250 (Chile).

ACKNOWLEDGMENTS

Authors thanks all members of RIVATREM (Red Iberoamericana de Alteraciones Vasculares en Transtornos del Embarazo) for all our productive discussion leading to the generation of this manuscript.

- Baud, D., Greub, G., Favre, G., Gengler, C., Jatou, K., Dubruc, E., et al. (2020). Second-Trimester Miscarriage in a Pregnant Woman with SARS-CoV-2 Infection. *JAMA* 323, 2198–2200. doi: 10.1001/jama.2020.7233
- Bavishi, C., Maddox, T., and Messerli, F. (2020). Coronavirus Disease 2019 (COVID-19) Infection and Renin Angiotensin System Blockers. *Am. Med. Assoc.* 5, 745–747. doi: 10.1111/jcmm.12573
- Beigi, R. H., Krubiner, C., Jamieson, D. J., Lyster, A. D., Hughes, B., Riley, L., et al. (2021). The need for inclusion of pregnant women in COVID-19 vaccine trials. *Vaccine* 39:868. doi: 10.1016/J.VACCINE.2020.12.074
- Best Rocha, A., Stroberg, E., Barton, L. M., Duval, E. J., Mukhopadhyay, S., Yarid, N., et al. (2020). Detection of SARS-CoV-2 in formalin-fixed paraffin-embedded tissue sections using commercially available reagents. *Lab. Invest.* 100, 1485–1489. doi: 10.1038/s41374-020-0464-x
- Bharadwaj, M. S., Strawn, W. B., Groban, L., Yamaleyeva, L. M., Chappell, M. C., Horta, C., et al. (2011). Angiotensin-Converting Enzyme 2 Deficiency Is Associated With Impaired Gestational Weight Gain and Fetal Growth Restriction. *Hypertension* 58, 852–858. doi: 10.1161/HYPERTENSIONAHA.111.179358
- Bortoletto, P., Romanski, P. A., Stewart, J., Rosenwaks, Z., and Pfeifer, S. M. (2021). Incidence of first trimester pregnancy loss in the infertile population during the first wave of the coronavirus disease 2019 pandemic in New York City. *FS Rep.* 2, 209–214. doi: 10.1016/J.XFRE.2021.04.005
- Brandt, J. S., Hill, J., Reddy, A., Schuster, M., Patrick, H. S., Rosen, T., et al. (2020). Epidemiology of coronavirus disease 2019 in pregnancy: risk factors and associations with adverse maternal and neonatal outcomes. *Am. J. Obstet. Gynecol.* 224:e389. doi: 10.1016/j.ajog.2020.09.043
- Cai, H. (2020). Correspondence Sex difference and smoking predisposition Smoking or Vaping May Increase the Risk of a Severe Coronavirus Infection. *Elsevier* 2600, 19–20. doi: 10.1111/all.14238.Yang
- Channappanavar, R., Zhao, J., and Perlman, S. (2014). T cell-mediated immune response to respiratory coronaviruses. *Immunol. Res.* 59, 118–128. doi: 10.1007/s12026-014-8534-z
- Chavan, M., Qureshi, H., Karnati, S., and Kollikonda, S. (2021). COVID-19 Vaccination in Pregnancy: the Benefits Outweigh the Risks. *J. Obstet. Gynaecol. Can.* 43:814. doi: 10.1016/J.JOGC.2021.03.010

- Chen, W., Yuan, P., Yang, M., Yan, Z., Kong, S., Yan, J., et al. (2020). SARS-CoV-2 Entry Factors: ACE2 and TMPRSS2 Are Expressed in Peri-Implantation Embryos and the Maternal–Fetal Interface. *Engineering* 6, 1162–1169. doi: 10.1016/j.eng.2020.07.013
- Chlamydas, S., Papavassiliou, A. G., and Piperi, C. (2021). Epigenetic mechanisms regulating COVID-19 infection. *Epigenetics* 16, 263–270. doi: 10.1080/15592294.2020.1796896
- Corthorn, J., Germain, A. A., Chacón, C., Rey, S., Soto, G. X., Figueroa, C. D., et al. (2006). Expression of kallikrein, bradykinin b2 receptor, and endothelial nitric oxide synthase in placenta in normal gestation, preeclampsia, and placenta accreta. *Endocrine* 29, 491–499. doi: 10.1385/ENDO:29:3:491
- Cosma, S., Carosso, A. R., Cusato, J., Borella, F., Carosso, M., Bovetti, M., et al. (2020). Coronavirus disease 2019 and first-trimester spontaneous abortion: a case-control study of 225 pregnant patients. *Am. J. Obstet. Gynecol.* 224:391.e1-391.e7. doi: 10.1016/j.ajog.2020.10.005
- Della Gatta, A. N., Rizzo, R., Pilu, G., and Simonazzi, G. (2020). Coronavirus disease 2019 during pregnancy: a systematic review of reported cases. *Am. J. Obstet. Gynecol.* 223, 36–41. doi: 10.1016/j.ajog.2020.04.013
- di Mascio, D., Khalil, A., Saccone, G., Rizzo, G., Buca, D., Liberati, M., et al. (2020). Outcome of coronavirus spectrum infections (SARS, MERS, COVID-19) during pregnancy: a systematic review and meta-analysis. *Am. J. Obstet. Gynecol. MFM* 2:100107. doi: 10.1016/j.AJOGMF.2020.100107
- Díaz-Corvillón, P., Mönckeberg, M., Barros, A., Illanes, S. E., Soldati, A., Nien, J.-K., et al. (2020). Routine screening for SARS CoV-2 in unselected pregnant women at delivery. *PLoS One* 15:e0239887. doi: 10.1371/journal.pone.0239887
- Editorial The Lancet. (2020). COVID-19 in Latin America: a humanitarian crisis. *Lancet* 396:1463. doi: 10.1016/S0140-6736(20)32328-X
- Eldow, A. G., Li, J. Z., Collier, A. R. Y., Atyeo, C., James, K. E., Boatman, A. A., et al. (2020). Assessment of Maternal and Neonatal SARS-CoV-2 Viral Load, Transplacental Antibody Transfer, and Placental Pathology in Pregnancies During the COVID-19 Pandemic. *JAMA Network Open* 3:e2030455. doi: 10.1001/jamanetworkopen.2020.30455
- Elias, C., Sekri, A., Leblanc, P., Cucherat, M., and Vanhems, P. (2021). The incubation period of COVID-19: a meta-analysis. *Int. J. Infect. Dis.* 104, 708–710. doi: 10.1016/j.IJID.2021.01.069
- Ellington, S., Strid, P., Tong, V. T., Woodworth, K., Galang, R. R., Zambrano, L. D., et al. (2020). Characteristics of Women of Reproductive Age with Laboratory-Confirmed SARS-CoV-2 Infection by Pregnancy Status - United States, January 22–June 7, 2020. *MMWR* 69, 769–775. doi: 10.15585/mmwr.mm6925a1
- Emanuele, N., Ren, J., Lapaglia, N., Steiner, J., and Emanuele, M. A. (2002). Angiotensin-(1-7) in normal and preeclamptic pregnancy. *Endocrine* 18, 239–245. doi: 10.1385/ENDO:18:3:239
- Evans, P. C., Rainger, G. E., Mason, J. C., Guzik, T. J., Osto, E., Stamataki, Z., et al. (2020). Endothelial dysfunction in COVID-19: a position paper of the ESC Working Group for Atherosclerosis and Vascular Biology, and the ESC Council of Basic Cardiovascular Science. *Cardiovasc. Res.* 116:2177–2184. doi: 10.1093/CVR/CVAA230
- Fan, S., Yan, S., Liu, X., Liu, P., Huang, L., and Wang, S. (2021). Human Coronavirus Infections and Pregnancy. *Matern.Fetal Med.* 3, 53–65. doi: 10.1097/FM9.0000000000000071
- Fenizia, C., Biasin, M., Cetin, I., Vergani, P., Mileto, D., Spinillo, A., et al. (2020). Analysis of SARS-CoV-2 vertical transmission during pregnancy. *Nat. Commun.* 11:5128. doi: 10.1038/s41467-020-18933-4
- Ferraiolo, A., Barra, F., Kratochwila, C., Paudice, M., Vellone, V. G., Godano, E., et al. (2020). Report of positive placental swabs for sars-cov-2 in an asymptomatic pregnant woman with covid-19. *Medicina* 56, 1–9. doi: 10.3390/medicina56060306
- Flores-Pliego, A., Miranda, J., Vega-Torreblanca, S., Valdespino-Vázquez, Y., Helguera-Repetto, C., Espejel-Nuñez, A., et al. (2021). Molecular Insights into the Thrombotic and Microvascular Injury in Placental Endothelium of Women with Mild or Severe COVID-19. *Cells* 10:364. doi: 10.3390/cells10020364
- Fraga-Silva, R. A., Costa-Fraga, F. P., Murça, T. M., Moraes, P. L., Martins Lima, A., Lautner, R. Q., et al. (2013). Angiotensin-converting enzyme 2 activation improves endothelial function. *Hypertension* 61, 1233–1238. doi: 10.1161/HYPERTENSIONAHA.111.00627
- Gant, N. F., Daley, G. L., Chand, S., Whalley, P. J., and MacDonald, P. C. (1973). A study of angiotensin II pressor response throughout primigravid pregnancy. *J. Clin. Invest.* 52, 2682–2689. doi: 10.1172/JCI107462
- Gant, N. F., Worley, R. J., Everett, R. B., and MacDonald, P. C. (1980). Control of vascular responsiveness during human pregnancy. *Kidney Int.* 18, 253–258. doi: 10.1038/KI.1980.133
- Gilbert, J. S., LaMarca, B. B., and Granger, J. P. (2008). ACE2 and ANG-(1-7) in the gravid uterus: the new players on the block. *Am. J. Physiol.* 294, 915–916. doi: 10.1152/ajpregu.00018.2008
- Goldstein, I., Steinberg, D. M., Kuint, J., Chodick, G., Segal, Y., Shapiro Ben, et al. (2022). Association of BNT162b2 COVID-19 Vaccination During Pregnancy With Neonatal and Early Infant Outcomes. *JAMA Pediatr.* 10:e220001. doi: 10.1001/JAMAPEDIATRICS.2022.0001
- Goshua, G., Pine, A. B., Meizlish, M. L., Chang, C. H., Zhang, H., Bahel, P., et al. (2020). Endotheliopathy in COVID-19-associated coagulopathy: evidence from a single-centre, cross-sectional study. *Lancet Haematol.* 7, e575–e582. doi: 10.1016/S2352-3026(20)30216-7
- Halici-Ozturk, F., Ocal, F. D., Aydin, S., Tanacan, A., Ayhan, S. G., Altinboga, O., et al. (2021). Investigating the risk of maternal-fetal transmission of SARS-CoV-2 in early pregnancy. *Placenta* 106, 25–29. doi: 10.1016/j.placenta.2021.02.006
- Han, Y., Ma, H., Suo, M., Han, F., Wang, F., Ji, J., et al. (2020). Clinical manifestation, outcomes in pregnant women with COVID-19 and the possibility of vertical transmission: a systematic review of the current data. *J. Perinat. Med.* 48, 912–924. doi: 10.1515/jpm-2020-0431
- Hecht, J. L., Quade, B., Deshpande, V., Mino-Kenudson, M., Ting, D. T., Desai, N., et al. (2020). SARS-CoV-2 can infect the placenta and is not associated with specific placental histopathology: a series of 19 placentas from COVID-19-positive mothers. *Modern Pathol.* 33, 2092–2103. doi: 10.1038/s41379-020-0639-4
- Hosier, H., Farhadian, S. F., Morotti, R. A., Deshmukh, U., Lu-Culligan, A., Campbell, K. H., et al. (2020). SARS-CoV-2 infection of the placenta. *J. Clin. Invest.* 130, 4947–4953. doi: 10.1172/JCI139569
- Hsu, A. L., Guan, M., Johannesen, E., Stephens, A. J., Khaleel, N., Kagan, N., et al. (2021). Placental SARS-CoV-2 in a pregnant woman with mild COVID-19 disease. *J. Med. Virol.* 93, 1038–1044. doi: 10.1002/jmv.26386
- Huang, C., Wang, Y., Li, X., Ren, L., Zhao, J., Hu, Y., et al. (2020). Clinical features of patients infected with 2019 novel coronavirus in Wuhan. *CHIN. Lancet* 395, 497–506. doi: 10.1016/S0140-6736(20)30183-5
- Huang, F., Guo, J., Zou, Z., Liu, J., Cao, B., Zhang, S., et al. (2014). Angiotensin II plasma levels are linked to disease severity and predict fatal outcomes in H7N9-infected patients. *Nat. Commun.* 5:3595. doi: 10.1038/ncomms4595
- Jackson, C. B., Farzan, M., Chen, B., and Choe, H. (2021). Mechanisms of SARS-CoV-2 entry into cells. *Nat. Rev. Mol. Cell Biol.* 2021, 3–20. doi: 10.1038/s41580-021-00418-x
- Jaiswal, N., Puri, M., Agarwal, K., Singh, S., Yadav, R., Tiwary, N., et al. (2021). COVID-19 as an independent risk factor for subclinical placental dysfunction. *Eur. J. Obstet. Gynecol. Reprod. Biol.* 259, 7–11. doi: 10.1016/j.ejogrb.2021.01.049
- Jardine, J., Relph, S., Magee, L., Dadelszen, P., Morris, E., Ross-Davie, M., et al. (2021). Maternity services in the UK during the coronavirus disease 2019 pandemic: a national survey of modifications to standard care. *BJOG* 128, 880–889. doi: 10.1111/1471-0528.16547
- Jia, H. P., Look, D. C., Shi, L., Hickey, M., Pewe, L., Netland, J., et al. (2005). ACE2 Receptor Expression and Severe Acute Respiratory Syndrome Coronavirus Infection Depend on Differentiation of Human Airway Epithelia. *J. Virol.* 79, 14614–14621. doi: 10.1128/jvi.79.23.14614-14621.2005
- Jin, J. M., Bai, P., He, W., Wu, F., Liu, X. F., Han, D. M., et al. (2020). Gender Differences in Patients With COVID-19: focus on Severity and Mortality. *Front. Public Health* 8:152. doi: 10.3389/fpubh.2020.00152
- Jordan, R. E., Adab, P., and Cheng, K. K. (2020). Covid-19: risk factors for severe disease and death. *BMJ* 368:m1198. doi: 10.1136/bmj.m1198
- Kalafat, E., Yaprak, E., Cinar, G., Varli, B., Ozisik, S., Uzun, W., et al. (2020). Lung ultrasound and computed tomographic findings in pregnant woman with COVID-19. *Ultrasound Obstet. Gynecol.* 55, 835–837. doi: 10.1002/UOG.22034
- Khalil, A., von Dadelszen, P., Draycott, T., Ugwumadu, A., O'Brien, P., and Magee, L. (2020). Change in the Incidence of Stillbirth and Preterm Delivery During the COVID-19 Pandemic. *JAMA* 324:705. doi: 10.1001/jama.2020.12746
- Khodarahmi, R., Sayad, B., and Sobhani, M. (2021). The ACE2 as a “rescue protein” or “suspect enzyme” in COVID-19: possible application of the “engineered

- Penfield, C. A., Brubaker, S. G., Limaye, M. A., Lighter, J., Ratner, A. J., Thomas, K. M., et al. (2020). Detection of SARS-CoV-2 in Placental and Fetal Membrane Samples. *Am. J. Obstet. Gynecol. MFM.* 2:100133. doi: 10.1016/j.ajogmf.2020.100133
- Peng, Z., Wang, J., Mo, Y., Duan, W., Xiang, G., Yi, M., et al. (2020). Unlikely SARS-CoV-2 vertical transmission from mother to child: a case report. *J. Infect. Public Health* 13, 818–820. doi: 10.1016/j.jiph.2020.04.004
- Pique-Regi, R., Romero, R., Tarca, A. L., Luca, F., Xu, Y., Alazizi, A., et al. (2020). Does the human placenta express the canonical cell entry mediators for SARS-CoV-2? *eLife* 9:e58716. doi: 10.7554/eLife.58716
- Poisson, T. M., and Pierone, G. (2021). Placental pathology and fetal demise at 35 weeks of gestation in a woman with SARS-CoV-2 infection: a case report. *Case Rep. Womens Health* 30:e00289. doi: 10.1016/j.crwh.2021.e00289
- Pringle, K. G., Tadors, M. A., Callister, R. J., and Lumbers, E. R. (2011). The expression and localization of the human placental prorenin/renin-angiotensin system throughout pregnancy: roles in trophoblast invasion and angiogenesis? *Placenta* 32, 956–962. doi: 10.1016/j.placenta.2011.09.020
- Rabi, F. A., al Zoubi, M. S., Al-Nasser, A. D., Kasasbeh, G. A., and Salameh, D. M. (2020). Sars-cov-2 and coronavirus disease 2019: what we know so far. *Pathogens* 9:231. doi: 10.3390/pathogens9030231
- Rezaei, M., Ziai, S. A., Fakhri, S., and Pouriran, R. (2021). ACE2 receptor expression and severe acute respiratory syndrome and COVID-19. *Journal of Cellular Physiology* 236, 2430–2442. doi: 10.1002/JCP.30041
- Richtmann, R., Torloni, M. R., Oyamada Otani, A. R., Levi, J. E., Crema Tobará, M., de Almeida Silva, C., et al. (2020). Fetal deaths in pregnancies with SARS-CoV-2 infection in Brazil: a case series. *Case Rep. Womens Health* 27:e00243. doi: 10.1016/j.crwh.2020.e00243
- Rizzo, G., Mappa, I., Maqina, P., Bitsadze, V., Khizroeva, J., Makatsarya, A., et al. (2021). Effect of SARS-CoV-2 infection during the second half of pregnancy on fetal growth and hemodynamics: a prospective study. *Acta Obstet. Gynecol. Scand.* 100, 1034–1039. doi: 10.1111/AOGS.14130
- Rothan, H. A., and Byrreddy, S. N. (2020). The epidemiology and pathogenesis of coronavirus disease (COVID-19) outbreak. *J. Autoimmun.* 109:102433. doi: 10.1016/j.jaut.2020.102433
- Rotshenker-Olshinka, K., Volodarsky-Perel, A., Steiner, N., Rubinfeld, E., and Dahan, M. H. (2020). COVID-19 pandemic effect on early pregnancy: are miscarriage rates altered, in asymptomatic women? *Arch. Gynecol. Obstet.* 303, 839–845. doi: 10.1007/s00404-020-05848-0
- Sánchez-Duque, J. A., Arce-Villalobos, L. R., and Rodríguez-Morales, A. J. (2020). Coronavirus disease 2019 (COVID-19) in Latin America: role of primary care in preparedness and response. *Atencion Primaria* 52, 369–372. doi: 10.1016/j.aprim.2020.04.001
- Sanchis-Gomar, F., Lavie, C. J., Perez-Quilis, C., Henry, B. M., and Lippi, G. (2020). Angiotensin-converting enzyme 2 and anti-hypertensives (angiotensin receptor blockers and angiotensin converting enzyme inhibitors) in coronavirus disease 2019 (COVID-19). *Mayo Clin. Proc.* 95, 1222–1230. doi: 10.1016/j.mayocp.2020.03.026
- Santos, R. A. (2014). Angiotensin-(1-7). *Hypertension* 63, 1138–1147. doi: 10.1161/HYPERTENSIONAHA.113.01274
- Scaife, P. J., and Mohaupt, M. G. (2017). Salt, aldosterone and extrarenal Na⁺ - sensitive responses in pregnancy. *Placenta* 56, 53–58. doi: 10.1016/j.placenta.2017.01.100
- Shanes, E. D., Mithal, L. B., Otero, S., Azad, H. A., Miller, E. S., and Goldstein, J. A. (2020). Placental Pathology in COVID-19. *Am. J. Clin. Pathol.* 154, 23–32. doi: 10.1093/ajcp/aqaa089
- Shimabukuro, T. T., Kim, S. Y., Myers, T. R., Moro, P. L., Odubebo, T., Panagiotakopoulos, L., et al. (2021). Preliminary Findings of mRNA Covid-19 Vaccine Safety in Pregnant Persons. *N. Engl. J. Med.* 384, 2273–2282. doi: 10.1056/NEJMOA2104983
- Siriwattananon, K., Manopwisedjaroen, S., Kanjanasirirat, P., Budi Purwono, P., Rattanapisit, K., Shanmugaraj, B., et al. (2021). Development of Plant-Produced Recombinant ACE2-Fc Fusion Protein as a Potential Therapeutic Agent Against SARS-CoV-2. *Front. Plant Sci.* 11:2117. doi: 10.3389/FPLS.2020.604663/BIBTEX
- Sisman, J., Jaleel, M. A., Moreno, W., Rajaram, V., Collins, R. R. J., Savani, R. C., et al. (2020). Intrauterine Transmission of SARS-CoV-2 Infection in a Preterm Infant. *Pediatr. Infect. Dis. J.* 39, e265–e267. doi: 10.1097/INF.0000000000002815
- Smithgall, M. C., Liu-Jarin, X., Hamele-Bena, D., Cemic, A., Mourad, M., Debelenko, L., et al. (2020). Third-trimester placentas of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2)-positive women: histomorphology, including viral immunohistochemistry and in situ hybridization. *Histopathology* 77, 994–999. doi: 10.1111/his.14215
- Sola, A., Rodríguez, S., Cardetti, M., and Dávila, C. (2020). Perinatal COVID-19 in Latin America. *Pan Am. J. Public Health* 44, e47. doi: 10.26633/RPSP.2020.47
- Sommerstein, R., Kochen, M. M., Messerli, F. H., and Gräni, C. (2020). Coronavirus Disease 2019 (COVID-19): do Angiotensin-Converting Enzyme Inhibitors/Angiotensin Receptor Blockers Have a Biphasic Effect? *J. Am. Heart Assoc.* 9:e016509. doi: 10.1161/JAHA.120.016509
- Stelzig, K. E., Canepa-Escaro, F., Schiliro, M., Berdnikovs, S., Prakash, Y. S., and Chiarella, S. E. (2020). Estrogen regulates the expression of SARS-CoV-2 receptor ACE2 in differentiated airway epithelial cells. *Am. J. Physiol* 318, 1280–1281. doi: 10.1152/ajplung.00153.2020
- Stettner, D., Bujak-Gizycka, B., Olszanecki, R., Rytlewski, K., Huras, H., and Korbut, R. (2013). Assessment of angiotensin I metabolism in the human placenta using an LC/MS method. *Folia medica Cracoviensis* 53, 31–39.
- Sutton, D., Fuchs, K., D'Alton, M., and Goffman, D. (2020). Universal screening for SARS-CoV-2 in women admitted for delivery. *N. Engl. J. Med.* 382, 2163–2164. doi: 10.1056/NEJMc2009316
- Taglauer, E., Benarroch, Y., Rop, K., Barnett, E., Sabharwal, V., Yarrington, C., et al. (2020). Consistent localization of SARS-CoV-2 spike glycoprotein and ACE2 over TMPRSS2 predominance in placental villi of 15 COVID-19 positive maternal-fetal dyads. *Placenta* 100, 69–74. doi: 10.1016/j.placenta.2020.08.015
- Takemoto, M. L. S., Menezes, M. O., Andreucci, C. B., Knobel, R., Sousa, L. A. R., Katz, L., et al. (2020). Maternal mortality and COVID-19. *J. Matern. Fetal Neonat. Med.* 16, 1–7. doi: 10.1080/14767058.2020.1786056
- Tamanna, S., Clifton, V. L., Rae, K., van Helden, D. F., Lumbers, E. R., and Pringle, K. G. (2020). Angiotensin Converting Enzyme 2 (ACE2) in Pregnancy: preeclampsia and Small for Gestational Age. *Front. Physiol.* 11:590787. doi: 10.3389/FPHYS.2020.590787
- Tang, T., Bidon, M., Jaimes, J. A., Whittaker, G. R., and Daniel, S. (2020). Coronavirus membrane fusion mechanism offers a potential target for antiviral development. *Antiviral Res.* 178:104792. doi: 10.1016/j.antiviral.2020.104792
- Trad, A. T. A., Ibiroga, E. R., Elrefaei, A., Narang, K., Tonni, G., Picone, O., et al. (2020). SARS-CoV-2 in Pregnancy: where and What Is the Evidence? *SSRN Electron. J.* [preprint]. doi: 10.2139/ssrn.3576828
- Tukiainen, T., Villani, A. C., Yen, A., Rivas, M. A., Marshall, J. L., Satiya, R., et al. (2017). Landscape of X chromosome inactivation across human tissues. *Nature* 550, 244–248. doi: 10.1038/nature24265
- Vaduganathan, M., Vardeny, O., Michel, T., McMurray, J. J. V., Pfeffer, M. A., Ph, D., et al. (2020). Renin – Angiotensin – Aldosterone System Inhibitors in Patients with Covid-19. *N. Engl. J. Med.* 382, 1653–1659.
- Valdés, G., Neves, L. A. A., Anton, L., Corthorn, J., Chacón, C., Germain, A. M., et al. (2006). Distribution of angiotensin-(1-7) and ACE2 in human placentas of normal and pathological pregnancies. *Placenta* 27, 200–207. doi: 10.1016/j.placenta.2005.02.015
- Valdespino-Vázquez, M. Y., Helguera-Repetto, C. A., León-Juárez, M., Villavicencio-Carrisoza, O., Flores-Pliego, A., Moreno-Verduzco, E. R., et al. (2021). Fetal and placental infection with SARS-CoV-2 in early pregnancy. *J. Med. Virol.* 93, 4480–4487. doi: 10.1002/JMV.26965
- Villar, J., Ariff, S., Gunier, R. B., Thiruvengadam, R., Rauch, S., Kholin, A., et al. (2021). Maternal and neonatal morbidity and mortality among pregnant women with and without COVID-19 infection: the INTERCOVID multinational cohort study. *JAMA Pediatr.* 175, 817–826. doi: 10.1001/JAMAPEDIATRICS.2021.1050
- Viruez-Soto, J. A., Olaya-Garay, S. X., Quiñones-Quiñones, E., Murillo-García, D. R., Ibáñez-Rodríguez, C. E., Franco-Hernández, G. A., et al. (2021). COVID-19 y obstetricia crítica en cuatro países de Latinoamérica. *Respirar* 13, 65–71.
- Vivanti, A. J., Vauloup-Fellous, C., Prevot, S., Zupan, V., Suffee, C., do Cao, J., et al. (2020). Transplacental transmission of SARS-CoV-2 infection. *Nat. Commun.* 11:3572. doi: 10.1038/s41467-020-17436-6
- Wang, C. L., Liu, Y. Y., Wu, C. H., Wang, C. Y., Wang, C. H., and Long, C. Y. (2021). Impact of COVID-19 on Pregnancy. *Int. J. Med. Sci.* 18:763. doi: 10.7150/IJMS.49923
- Wang, J., Kaplan, N., Wysocki, J., Yang, W., Lu, K., Peng, H., et al. (2020). The ACE2-deficient mouse: a model for a cytokine storm-driven inflammation. *FASEB J.* 34, 10505–10515. doi: 10.1096/FJ.202001020R

- Wastnedge, E., Reynolds, R., van Boeckel, S., Stock, S., Denison, F., Maybin, J., et al. (2021). PREGNANCY AND COVID-19. *Physiol. Rev.* 101, 303–312. doi: 10.1152/physrev.00024.2020
- Watanabe, T., Barker, T. A., and Berk, B. C. (2005). Angiotensin II and the endothelium: diverse signals and effects. *Hypertension* 45, 163–169. doi: 10.1161/01.HYP.0000153321.13792.b9
- Wei, S. Q., Bilodeau-Bertrand, M., Liu, S., and Auger, N. (2021). The impact of COVID-19 on pregnancy outcomes: a systematic review and meta-analysis. *CMAJ* 193, E540–E548. doi: 10.1503/CMAJ.202604
- Wiersinga, W. J., Rhodes, A., Cheng, A. C., Peacock, S. J., and Prescott, H. C. (2020). Pathophysiology, Transmission, Diagnosis, and Treatment of Coronavirus Disease 2019 (COVID-19): a Review. *JAMA* 324, 782–793. doi: 10.1001/jama.2020.12839
- World Health Organization (2021). *Update on WHO Interim Recommendations on COVID-19 Vaccination of Pregnant and Lactating Women*. Available online at: www.who.int/groups/strategic-advisory-group-of-experts-on-immunization (accessed March 3, 2022).
- Yang, H., Wang, C., and Poon, L. C. (2020). Novel coronavirus infection and pregnancy. *Ultrasound Obstetr. Gynecol.* 55, 435–437. doi: 10.1002/UOG.22006
- Zhang, M., Gao, Y., Zhao, W., Yu, G., and Jin, F. (2018). ACE-2/ANG1-7 ameliorates ER stress-induced apoptosis in seawater aspiration-induced acute lung injury. *Am. J. Physiol.* 315, L1015–L1027. doi: 10.1152/ajplung.00163.2018
- Zhang, P., Salafia, C., Heyman, T., Salafia, C., Lederman, S., and Dygulska, B. (2020). Detection of severe acute respiratory syndrome coronavirus 2 in placentas with pathology and vertical transmission. *Am. J. Obstetr. Gynecol. MFM* 2:100197. doi: 10.1016/j.ajogmf.2020.100197
- Zhang, Z., Zeng, E., Zhang, L., Wang, W., Jin, Y., Sun, J., et al. (2021). Potent prophylactic and therapeutic efficacy of recombinant human ACE2-Fc against SARS-CoV-2 infection in vivo. *Cell Disc.* 7:65. doi: 10.1038/s41421-021-00302-0
- Zheng, Y.-Y., Zhang, J.-Y., and Xie, X. (2019). COVID-19 and the cardiovascular system. *Nat. Rev. Cardiol.* 17, 259–260.
- Zheng, Z., Peng, F., Xua, B., Zhao, J., Liu, H., Peng, J., et al. (2020). Risk factors of critical & mortal COVID-19 cases: a systematic literature review and meta-analysis - ScienceDirect. *J. Infect.* 18, e16–e25.
- Zhou, F., Yu, T., Du, R., Fan, G., Liu, Y., Liu, Z., et al. (2020). Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* 395, 1054–1062. doi: 10.1016/S0140-6736(20)30566-3
- Zhou, P., Yang, X., Lou, W., Wang, X. G., Hu, B., Zhang, L., et al. (2020). A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature* 579, 270–273. doi: 10.1038/s41586-020-2012-7
- Zou, X., Chen, K., Zou, J., Han, P., Hao, J., and Han, Z. (2020). Single-cell RNA-seq data analysis on the receptor ACE2 expression reveals the potential risk of different human organs vulnerable to 2019-nCoV infection. *Front. Med.* 14:185–192. doi: 10.1007/s11684-020-0754-0

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2022 Ayala-Ramírez, González, Escudero, Quintero-Arciniegas, Giachini, Alves de Freitas, Damiano and García-Robles. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.