



Gene regulation by cytokinin in *Arabidopsis*

Wolfram G. Brenner, Eswar Ramireddy, Alexander Heyl* and Thomas Schmülling*

Institute of Biology/Applied Genetics, Dahlem Centre of Plant Sciences, Freie Universität Berlin, Berlin, Germany

Edited by:

Andreas P. M. Weber, University of Düsseldorf, Germany

Reviewed by:

Hitoshi Sakakibara, RIKEN, Japan
Stefan Hoth, Universität Hamburg, Germany

*Correspondence:

Alexander Heyl and Thomas Schmülling, Institute of Biology/Applied Genetics, Dahlem Centre of Plant Sciences, Freie Universität Berlin, Albrecht-Thaer-Weg 6, D-14195 Berlin, Germany.
e-mail: hey@zedat.fu-berlin.de;
thomas.schmuelling@fu-berlin.de

The plant hormone cytokinin realizes at least part of its signaling output through the regulation of gene expression. A great part of the early transcriptional regulation is mediated by type-B response regulators, which are transcription factors of the MYB family. Other transcription factors, such as the cytokinin response factors of the AP2/ERF family, have also been shown to be involved in this process. Additional transcription factors mediate distinct parts of the cytokinin response through tissue- and cell-specific downstream transcriptional cascades. In *Arabidopsis*, only a single cytokinin response element, to which type-B response regulators bind, has been clearly proven so far, which has 5'-GAT(T/C)-3' as a core sequence. This motif has served to construct a synthetic cytokinin-sensitive two-component system response element, which is useful for monitoring the cellular cytokinin status. Insight into the extent of transcriptional regulation has been gained by genome-wide gene expression analyses following cytokinin treatment and from plants having an altered cytokinin content or signaling. This review presents a meta analysis of such microarray data resulting in a core list of cytokinin response genes. Genes encoding type-A response regulators displayed the most stable response to cytokinin, but a number of cytokinin metabolism genes (*CKX4*, *CKX5*, *CYP735A2*, *UGT76C2*) also belong to them, indicating homeostatic mechanisms operating at the transcriptional level. The cytokinin core response genes are also the target of other hormones as well as biotic and abiotic stresses, documenting crosstalk of the cytokinin system with other hormonal and environmental signaling pathways. The multiple links of cytokinin to diverse functions, ranging from control of meristem activity, hormonal crosstalk, nutrient acquisition, and various stress responses, are also corroborated by a compilation of genes that have been repeatedly found by independent gene expression profiling studies. Such functions are, at least in part, supported by genetic studies.

Keywords: cytokinin, gene regulation, transcription factor, cis-element, transcriptomics, meta analysis, signal transduction, regulatory network

FOREWORD

The phytohormone cytokinin regulates numerous developmental and physiological processes, including the activity of shoot and root meristems, reproductive behavior, leaf senescence, and responses to environmental cues, in particular light and nutrients (Argueso et al., 2009; Werner and Schmülling, 2009). Regulation of gene expression is a crucial part of realizing the different activities in a developmental- and tissue-specific context. The results of early efforts to identify cytokinin-regulated genes were summarized more than a decade ago (Schmülling et al., 1997). Since then, several major steps toward a more comprehensive understanding of the early and late events of gene regulation by cytokinin have been made. Some important milestones have been the discovery of type-A response regulator genes as primary cytokinin response genes (Brandstatter and Kieber, 1998; Sakakibara et al., 1998); the description of type-B response regulators as transcription factors mediating the cytokinin response (Sakai et al., 2000, 2001; Hwang and Sheen, 2001); the identification of a cis-acting cytokinin response element (Sakai et al., 2000; Taniguchi et al., 2007); and the genome-wide analysis of the cytokinin response of the *Arabidopsis* transcriptome (Kiba et al., 2004; Brenner et al., 2005). Additional

genome-wide studies addressed more specifically the effects of altered cytokinin signaling, the identification of targets of specific signaling components, or organ-specific responses (Rashotte et al., 2006; Lee et al., 2007; Taniguchi et al., 2007; Yokoyama et al., 2007; Argyros et al., 2008; Heyl et al., 2008; Brenner and Schmülling, submitted). More recent work has linked specific transcriptional responses to the biological activities of cytokinin (Dello Ioio et al., 2008a; Müller and Sheen, 2008; Argueso et al., 2010; Zhao et al., 2010; Bishopp et al., 2011c; Köllmer et al., 2011). Taken together, it has become clear that fine-tuned gene regulation is of utmost importance to realize the many different biological activities of cytokinin in diverse tissues.

In this review, we will firstly briefly describe the state-of-the-art knowledge about the early events involved in cytokinin-dependent gene regulation and the main players, and address the question how this system evolved. Then we will summarize the results of a number of genome-wide transcript profiling studies focusing on cytokinin as a regulator of gene expression, present the results of a meta analysis of microarray data and, finally, compile some of the frequently found cytokinin-regulated genes and discuss their functional context. The vast majority of findings come

from experiments using *Arabidopsis thaliana*, thus we will focus on results from this model plant in this review. The key facts presented in this review are summarized in **Table 1**. For recent reviews addressing various other aspects of the cytokinin signaling system, see To and Kieber (2008), Argueso et al. (2009, 2010), Bishopp et al. (2009, 2011a), Moubayidin et al. (2009), Werner and Schmülling (2009), Hellmann et al. (2010), Kudo et al. (2010), Perilli et al. (2010), Domagalska and Leyser (2011), Heyl et al. (2011), Kiba et al. (2011), and Müller (2011).

THE DISCOVERY OF PRIMARY RESPONSE GENES OF CYTOKININ

The first primary response genes rapidly induced by cytokinin were identified using the method of differential display in *Arabidopsis thaliana* (Brandstatter and Kieber, 1998) and maize (Sakakibara et al., 1998). In *Arabidopsis*, two genes designated *IBC6* and *IBC7* (*INDUCED BY CYTOKININ, IBC*) – now known as type-A response regulator genes *ARR5* and *ARR4*, respectively – were identified as being induced after 2 h of cytokinin treatment (Brandstatter and Kieber, 1998). These genes were upregulated within minutes after cytokinin treatment (D'Agostino et al., 2000) and the response was insensitive to cycloheximide, both hallmarks of primary response genes. At the same time, the maize gene *ZmCip1*, also encoding a type-A response regulator, was found to be induced 30 min after cytokinin treatment (Sakakibara et al., 1998). A number of related *Arabidopsis* type-A response regulator genes – *ARR3*, *ARR6*, *ARR7*, *ARR8*, *ARR9*, *ARR15*, and *ARR16*, which were mostly identified by sequence homology to bacterial response regulators (Imamura et al., 1998; Urao et al., 1998) – were characterized as cytokinin-induced genes with different response kinetics (Imamura et al., 1999; Kiba et al., 1999; D'Agostino et al., 2000). It was shown later that type-A ARR proteins act as negative regulators of cytokinin signaling (Hwang and Sheen, 2001; To et al., 2004). Today, by virtue of genome-wide expression analyses, numerous other response genes of cytokinin have been discovered (see Genome-Wide Analysis of the Cytokinin-Regulated Transcriptome) but most of them lack a functional link to the hormone at present.

TRANSCRIPTION FACTORS REGULATING THE PRIMARY CYTOKININ RESPONSE

The discovery of type-A response regulator genes as primary cytokinin response genes was a strong indication that cytokinin signaling may occur through a two-component signaling system (TCS). In fact, the cytokinin signal is perceived by sensor histidine kinases (Inoue et al., 2001; Suzuki et al., 2001; Heyl et al., 2011), named AHK2, AHK3, and CRE1/AHK4 in *Arabidopsis*, and the following primary signal transduction pathway of cytokinin is a multistep trans-phosphorylation chain similar to the bacterial TCS. Following the transfer of the signal to the nucleus via histidine phosphotransmitter proteins (named AHPs in *Arabidopsis*; Hwang and Sheen, 2001; Punwani et al., 2010), ultimately a family of transcription factors, the type-B response regulators, is activated and these mediate the transcriptional response to cytokinin (Hwang and Sheen, 2001).

The *Arabidopsis* genome encodes 11 type-B ARRs. They belong to the MYB family of transcription factors and all contain an

Table 1 | Key facts about cytokinin-regulated gene expression.

- Type-B response regulators mediate most, if not all, of the immediate-early changes of gene expression induced by cytokinin.
- Other transcription factors, such as cytokinin response factors (CRF), are necessary for co-regulation of the transcriptional cytokinin response.
- During evolution, type-B response regulators appear earlier in plants than cytokinin receptors, which are only found in land plants.
- The core sequence of the only known cytokinin response element of *Arabidopsis* is 5'-GAT(T/C). The extended motif 5'-AAGAT(T/C)TT-3' has been identified as the preferred binding site of type-B response regulator ARR1.
- Genes encoding different type-A response regulators show the most robust response to cytokinin.
- Changes in transcript abundance indicate a feedback of cytokinin signaling on cytokinin metabolism genes.
- Numerous specific, context-dependent transcriptional cascades operate downstream of the two-component signaling system to realize diverse cytokinin activities.
- A large number of genes with diverse functions in regulating developmental and physiological processes respond to cytokinin.

N-terminal response regulator domain. This domain is thought to negatively regulate the activity of the type-B ARRs as a deletion of this domain leads to a constitutively active form of the protein (Sakai et al., 2000; Imamura et al., 2003; Tajima et al., 2004; Taniguchi et al., 2007). This observation led to a model in which the phosphorylation of the response regulator domain causes a conformational change of the type-B ARRs, thereby activating the proteins. Phosphomimics, in which the canonical Asp of the response regulator domain is exchanged with a Glu leading to a constitutive activation, are supposed to have an altered structure and seem to support this model (Hass et al., 2004). However, structural information confirming the proposed mode of action is still missing. This is in contrast to the DNA-binding MYB-domain. NMR spectroscopy has revealed how the DNA-binding domain of ARR10 is in direct contact with the target DNA, 5'-(A/T)GATT-3' (Hosoda et al., 2002). The C-terminal part of the type-B ARRs is often enriched by P/Q-rich acidic domains, which have been shown to be important for their transactivation capacity (Sakai et al., 2000; Imamura et al., 2003). The nuclear localization of the type-B ARRs is consistent with their role as transcription factors (Hwang and Sheen, 2001; Lohrmann et al., 2001; Mason et al., 2004; Dortay et al., 2006).

Phylogenetic analyses based on the receiver domains or DNA-binding domains of type-B ARRs revealed a division into three sub-classes (Tajima et al., 2004; To et al., 2004; Mason et al., 2005; Heyl et al., 2008). The largest sub-class consists of seven members (*ARR1*, *ARR2*, *ARR10*, *ARR12*, *ARR11*, *ARR14*, and *ARR18*) and two other sub-classes containing two members each (*ARR13/ARR21* and *ARR19/ARR20*). The expression patterns of type-B ARR genes reflect this subdivision revealed by phylogenetic analysis. While the members of the largest sub-class are expressed in almost all tissues, members of the other two sub-classes are expressed specifically in reproductive organs (Mason et al., 2004; Tajima et al., 2004; Heyl et al., 2006). The region of expression for

ARR19 and *ARR21* was narrowed down to the chalazal region of the endosperm (Tiwari et al., 2006; Day et al., 2008).

Analyses have shown the functional redundancy of type-B ARR genes (Mason et al., 2005): for example, single knockout mutants of *arr1* and *arr21* displayed no major alteration of their phenotypes (Sakai et al., 2001; Horák et al., 2003). Only the *arr2* knockout mutant showed a slight insensitivity toward cytokinin and ethylene in a hypocotyl elongation assay (Hass et al., 2004). *ARR2* has also been implicated in the regulation of leaf senescence (Kim et al., 2006). In contrast, analysis of multiple knockout mutant lines of type-B ARR genes revealed that *ARR1*, *ARR10*, and *ARR12* mediate the majority of cytokinin action in different processes, such as vascular development, light sensitivity, chlorophyll production, and cell division in the root and shoot (Yokoyama et al., 2007; Argyros et al., 2008; Ishida et al., 2008). Overexpression of full length *ARR1* and *ARR2* genes did not cause notable phenotypic changes (Sakai et al., 2001; Hass et al., 2004). However, overexpression of only the C-terminal part of the protein, including the DNA-binding domain of *ARR1*, *ARR11*, *ARR14*, *ARR20*, and *ARR21*, respectively, caused distinct abnormalities of plant phenotypes during the vegetative stage (*ARR1/ARR11/ARR14*) or the reproductive stage (*ARR20/ARR21*; Sakai et al., 2001; Imamura et al., 2003; Tajima et al., 2004). A compilation of the known phenotypes of single and multiple type-B ARR mutants (and other genes of the TCS) has been published recently (Müller, 2011).

Chimeric repressor technology (Hiratsu et al., 2003) was used in order to investigate the functions of type-B ARRs more generally. The addition of the SRDX peptide (LDLDLELRIGFA) to the C-terminal end of *ARR1* turned this transcriptional activator into a *trans*-dominant transcriptional repressor. Protoplast transactivation assays confirmed that *ARR1-SRDX* represses not only the transactivation activity of *ARR1*, but also of all other type-B ARRs tested, thus demonstrating its *trans*-dominant function (Heyl et al., 2008). Plants expressing the *ARR1-SRDX* construct under the control of the 35S promoter displayed a strong morphological phenotype, reminiscent of the cytokinin deficiency syndrome seen in triple cytokinin receptor mutants or cytokinin oxidase/dehydrogenase (CKX) overexpressors (Werner et al., 2003; Riefler et al., 2006). This phenotype is most likely due to the concerted repression of the activities of the type-B ARR (Heyl et al., 2008). Transcriptional profiling indicated attenuation of the early transcriptional response to cytokinin and suggested that *ARR1* and other type-B ARRs use, at least in part, the same DNA-binding sites. Together, the results showed that most, if not all, of the transcriptional response to cytokinin is mediated, at least partly, by the type-B ARRs (Heyl et al., 2008).

Another group of transcription factors regulating the transcriptional response to cytokinin are the cytokinin response factors (CRF). Rashotte and colleagues showed that members of this subfamily of AP2/ERF transcription factors, which is defined by the presence of a common CRF domain, are involved in the control of gene expression of many primary cytokinin response genes (Rashotte et al., 2006; Rashotte and Goertzen, 2010). *CRF2* and *CRF5* were themselves recognized as primary response genes due to their transcriptional regulation by cytokinin in the presence of cycloheximide (Rashotte et al., 2006). Protoplast experiments showed that all CRFs localize from the cytosol into the nucleus

upon cytokinin treatment. This translocation was dependent on the presence of the cytokinin receptors and the AHPs, but not type-A or type-B ARRs (Rashotte et al., 2006). Yeast two-hybrid and Split-YFP experiments showed homo- and heterodimerization among the CRFs and between the AHPs and the CRFs. Deletion experiments indicated that the CRF domain is both necessary and sufficient to mediate these protein interactions. It has been speculated that homodimerization of the CRFs might be necessary for DNA-binding (Cutcliffe et al., 2011). The phenotypic changes of different multiple *crf* mutants occurred with low penetrance, but point at a role in the development of embryos, cotyledons, and leaves. The severity and the penetrance of the loss-of-function phenotypes increased in higher order mutants indicating functional redundancy (Rashotte et al., 2006).

EVOLUTION OF TRANSCRIPTION FACTORS MEDIATING THE CYTOKININ RESPONSE

In order to get an insight into the evolution of the TCS, a phylogenetic analysis has been carried out for all four participating protein families (Pils and Heyl, 2009). The type-B ARRs are present in all plants species, including algae. Thus, they predate the appearance of the cytokinin receptors and the type-A ARRs, which are only found in land plants (Pils and Heyl, 2009). That raises the questions which functions the type-B ARRs have in those species missing the cytokinin receptors and whether those ancient functions are still performed by type-B ARRs of higher plant species. The CRF protein family has at least eight members in *Arabidopsis* (Rashotte and Goertzen, 2010). A phylogenetic study showed that CRF genes are present in the genomes of all land plants, but are absent from those of algae. The late-diverging land plants separated into two clades, which coincided with the presence or the absence of an additional domain, the so-called TEH (Thr, Glu, His) region. However, the functional relevance of this TEH region is not yet clear (Rashotte and Goertzen, 2010).

Taken together, these data suggest that the type-B ARRs predate the emergence of the cytokinin signaling pathway as we know it from higher plants and were only recruited to it at a later stage of plant evolution. Once the morphology and the developmental program of land plants became more complex, not only did the number of type-B ARRs in the genomes increase (Pils and Heyl, 2009; Schaller et al., 2011), but it was also likely that additional factors, such as the CRFs, were recruited to fine-tune the cytokinin response.

cis-REGULATORY CYTOKININ RESPONSE ELEMENTS REQUIRED FOR THE REGULATION OF PRIMARY RESPONSE GENES

Considering the high conservation of amino acids in the DNA-binding MYB-domain of type-B ARRs (Heyl et al., 2008), it is not surprising that several of them bind to the same or very similar sequence motifs. It has been demonstrated that the DNA-binding domains of both, *ARR1* and *ARR2*, bind *in vitro* to DNA sequences containing the 5'-AGAT(T/C)-3' motif. The actual DNA recognition by the DNA-binding domain appeared to occur mainly at the last four positions of this 5-bp sequence (Sakai et al., 2000). The optimal *in vitro* binding for *ARR10* and *ARR11* was observed for a DNA sequence containing the motifs 5'-AGATT-3' and 5'-GGATT-3', respectively (Hosoda et al., 2002; Imamura et al., 2003).

Later, it was shown that ARR1 requires 2-bp extensions on either side of the core sequence for optimal binding *in vitro* (Taniguchi et al., 2007). Comparing the bases surrounding the 5-bp long core motif, the sequence 5'-AAGAT(T/C)TT-3' was identified as an extended version of the ARR1-binding motif (Taniguchi et al., 2007).

A synthetic cytokinin reporter that was generated to visualize the output of the TCS *in vivo* harbors 24 concatemered copies of a variant of the above-mentioned type-B ARR binding core motif, 5'-(A/G)GATT-3', in sense and antisense direction and a minimal 35S promoter binding motif upstream of a reporter gene (Müller and Sheen, 2008). The presumed advantage of this synthetic cytokinin reporter compared to other marker genes, such as the commonly used type-A response regulator genes, is that its activity is largely independent of other signaling or tissue-specific information, and thus shows solely the TCS output (Müller and Sheen, 2008). Using this synthetic reporter, the local and temporal defined antagonistic interaction between auxin and cytokinin during the establishment of the root stem cell niche was identified (Müller and Sheen, 2008). Similarly, this TCS reporter or an improved version of it has been used to study cytokinin activity in the shoot apical meristem (Gordon et al., 2009; Yoshida et al., 2011).

So far, the functionality of binding motifs identified for ARR1 and other type-B ARRs have been supported mainly by *in vitro* experiments (Sakai et al., 2001; Imamura et al., 2003; Taniguchi et al., 2007) and their use for the TCS reporter (Müller and Sheen, 2008). There are only a very few loss-of-function studies confirming the functionality of these sequences *in planta*. One example is the transcriptional regulation of the *non-symbiotic hemoglobin (NSHB)* genes (Hunt et al., 2001; Ross et al., 2004). It was shown by using transient bombardment assays in tobacco leaf disks that cytokinin activates the *Oryza sativa NSHB2 (OsNSHB2)* promoter and that this regulation can be mediated by ARR1 of *Arabidopsis* (Ross et al., 2004). Deletion and site-directed mutational analyses of the predicted ARR1-binding *cis*-element 5'-AGATT-3' in the *OsNSHB2* promoter confirmed the functionality of this element in the cytokinin response. Deletion led to a dramatic reduction in cytokinin-inducibility of this promoter (Ross et al., 2004). These experiments were not fully conclusive, as the cytokinin-dependent activation of a rice promoter by an *Arabidopsis* transcription factor was tested in tobacco cells. However, it has been shown recently that mutation of this core motif in the *ARR6* promoter strongly reduces its response toward cytokinin in its original host (Ramireddy, 2009).

The afore-mentioned work confirmed the functional relevance of the core motif *in planta*. However, it should be noted that because of its statistically frequent occurrence, this short motif is not diagnostic for the cytokinin responsiveness of a given promoter. It is not significantly enriched in the top 20 cytokinin-responsive promoters within the 1-kb 5' of the translational start site (Figure 1), where it occurs at a frequency of 0.68% in comparison to the promoters of the 20 genes least likely to be regulated by cytokinin, where it has a frequency of 0.57%. The extended motif, 5'-AAGAT(T/C)TT-3', which was identified for ARR1 (Taniguchi et al., 2007), is enriched in the top 20 cytokinin-responsive promoters, where it occurs at a frequency of 0.16% compared to

0.01% in the promoters of the 20 genes least likely to be regulated by cytokinin. However, it is missing in 5 of the top 20 genes (Figure 1). This could indicate that ARR1 is not involved in the transcriptional regulation of all these cytokinin response genes, or that other variants of the extended motif allow for the binding of and transcriptional activation by type-B ARRs.


Another putative cytokinin response element has been identified in cucumber. Expression analysis showed that cytokinin and light activate the transcription of the NADPH-protochlorophyllide reductase (*POR*) gene which has a key role in chlorophyll synthesis (Kuroda et al., 2000). Gel shift experiments demonstrated that the promoter of the *POR* gene contains a *cis*-element (5'-TATTAG-3') which binds nuclear extracts in a cytokinin-dependent manner (Fusada et al., 2005). Mutation of this sequence in the *POR* promoter caused a reduced cytokinin-dependent binding of nuclear proteins and caused a reduced cytokinin responsiveness in transient expression assays (Fusada et al., 2005). However, the protein factor(s) binding to this motif are not known. This new element, mediating part of the cytokinin response, is present in only 6 of the top 20 cytokinin-regulated genes of *Arabidopsis* in the meta analysis, indicating that this motif also is not absolutely required for cytokinin responsiveness or might be species-specific (data not shown).

GENOME-WIDE ANALYSIS OF THE CYTOKININ-REGULATED TRANSCRIPTOME

With the dawn of the transcriptomics age, enabled by the wide availability of microarrays, several attempts have been made to identify more cytokinin-responsive genes. Over time, the themes of the experimental approaches have changed. The first transcriptome studies were purely discovery-based approaches in order to find as many cytokinin-regulated genes as possible for a number of different time points. The focus then shifted toward the identification of genes being targeted directly by single or multiple transcription factors of the type-B response regulator and CRF families. The latest investigations have shown two trends: firstly, the attempt to investigate the transcriptome of smaller expression domains; secondly, an enhanced focus on physiological and developmental downstream processes influenced by cytokinin.

PUBLISHED STUDIES ON THE CYTOKININ-REGULATED TRANSCRIPTOME

A first study investigating cytokinin effects on the transcript level of many genes in 2002 used a microarray containing 8297 probe sets from *Arabidopsis* and assessed gene expression during shoot induction from root callus (Che et al., 2002). As the samples were taken days after transferring the explants to cytokinin-containing shoot induction medium, early changes in gene expression were not identified and only the expression data of known cytokinin and auxin responsive genes were shown. The results confirmed the cytokinin induction of the type-A ARR genes and *CRE1/AHK4*. Another early study focusing on long-term transcriptomic changes used the method of massively parallel signature sequencing and analyzed seedlings harboring an inducible cytokinin synthesizing bacterial *IPT* gene (Hoth et al., 2003). This work also confirmed a number of known cytokinin-regulated genes and it was noted that subsets of genes are regulated independently of tissue type or



developmental stage. Both studies also identified numerous novel genes regulated or potentially regulated by cytokinin.

The first investigation devoted to the immediate-early cytokinin response also used the microarray with 8297 probe sets (Rashotte et al., 2003). Samples were obtained from seedlings treated with different cytokinins (benzyladenine, *trans*-zeatin) in different concentrations (1–5 μM) and for different time periods (15 min to 24 h). The *wol* and *cre1* cytokinin receptor mutants were analyzed, too. More than 70 novel cytokinin-regulated genes were identified. Two cytokinin-responsive “putative AP2 genes” (AT2G46310 and AT4G23750) were discovered, which were classified as CRFs in later research (Rashotte et al., 2006). Homeostatic mechanisms were revealed as two of the cytokinin-induced genes encoded cytokinin metabolic enzymes: the cytokinin oxidase/dehydrogenase CKX4 (Werner et al., 2001) and a cytochrome P450 enzyme (CYP735A2) with cytokinin *trans*-hydroxylase activity (Takei et al., 2004). Another potential pattern

of cytokinin-regulated genes indicated an effect on the redox state (Rashotte et al., 2003).

One year later, a first analysis using a whole-genome microarray featuring 22,810 probe sets (ATH1 array) was published (Kiba et al., 2004). Although the work focused on the type-C (by then still classified as type-A) response regulator ARR22, it also contained information on cytokinin-induced genes in wild-type *Arabidopsis*. There, a large number of additional genes were found to be cytokinin-regulated, including several heat shock-related genes and the histidine kinase gene *AHK1*. This study was extended to plants overexpressing the C-terminal half of the type-B response regulator ARR21 (ARR21-C-ox) causing a constitutive activation of the cytokinin response, in order to contrast ARR22 overexpressors (ARR22-ox) attenuating the cytokinin response (Kiba et al., 2005). A set of 68 robustly cytokinin-responsive genes was generated by comparing transcript levels in wild type, cytokinin-treated wild type, ARR22-ox, and ARR21-C-ox.

Another study using the ATH1 array also appeared in 2005 comparing cytokinin-induced and cytokinin-deficient seedlings (35S:CKX1), with the uninduced wild type as the reference (Brenner et al., 2005). An extensive evaluation of the functional context of the regulated genes identified transcriptional regulation, signal transduction, interaction with other hormones (particularly gibberellin), light signaling, primary metabolism, and transport as cytokinin target processes. It was found that transcription factor genes were largely overrepresented among the cytokinin-regulated genes in comparison to their proportion in the whole genome. Many of these genes were only temporarily induced, while others were upregulated at a late time point, suggesting that transcriptional cascades are triggered during the course of cytokinin action. The MapMan software (Thimm et al., 2004) was used to identify cytokinin-regulated physiological processes. Among all the genes of cytokinin metabolism and signaling, only the type-A response regulators showed a clearly opposite regulation in cytokinin-induced and cytokinin-deficient seedlings. In addition, two processes were identified with an opposite regulation of their genes depending on the cytokinin status, namely metabolism of the regulator of carbon utilization, trehalose-6-phosphate, and valine and leucine metabolism. Finally, the upregulation of several plastidial transcripts was noted. The data generated in this study spawned further research on four cytokinin-regulated transcription factor genes, namely *GATA22* (also known as *CGA1/GNL*), two genes coding for members of the homeodomain zip (HD zip) class II transcription factor family (*HAT4*, *HAT22*), and *bHLH64* (Köllmer et al., 2011).

Transcript profiling of cytokinin-treated *arr1 arr12*, *crf1 crf2 crf5*, and *crf2 crf3 crf6* mutants was conducted in order to identify the target genes of the CRFs. The results led to the conclusion that both, type-B ARR and CRFs, are necessary for cytokinin induction of many of the early cytokinin response genes, because their regulation was attenuated in higher order mutants of either transcription factor family. Interestingly, while the expression of the type-A ARR genes in response to cytokinin treatment was reduced in the *arr1 arr12* mutant, it was unaffected or even slightly upregulated in the *crf* mutants tested (Rashotte et al., 2006).

The reduced induction of cytokinin-regulated genes in ARR7 overexpressing plants showed clearly that ARR7 is a negative regulator of cytokinin signaling (Lee et al., 2007). However, the regulation of only 30% to 50% of all cytokinin-regulated genes was attenuated by ARR7 overexpression, indicating that the protein acts only on part of the cytokinin signaling pathway. The regulation of transcription factor genes, hormone-related and stress-related genes, as well as the finding that the responsiveness of most cytokinin-induced expansin genes and several *LATERAL ORGAN BOUNDARIES DOMAIN (LBD)* genes was attenuated by ARR7, were specifically addressed.

Direct target genes of ARR1 were identified using plants expressing a dexamethasone-inducible variant of ARR1 (Sakai et al., 2001) and an AFLP-based expression profiling method called “high-coverage expression profiling” (HiCEP; Fukumura et al., 2003) as the analytical tool (Taniguchi et al., 2007). An improved sequence motif for ARR1-binding was characterized in the promoters of the 23 ARR1 target genes identified (see cis-Regulatory

Cytokinin Response Elements Required for the Regulation Of Primary Response Genes).

The roots of an *arr10 arr12* mutant were studied using the ATH1 microarray (Yokoyama et al., 2007). This mutant combination was selected because it causes a strong root phenotype, suggesting that ARR10 and ARR12 (together with ARR1) transduce most of the cytokinin signal in the root. It was found that the transcriptional response to cytokinin in this mutant was attenuated for most of the known cytokinin response genes, as well as for several genes newly discovered in this study. The induction of the type-A response regulator genes *ARR15* and *ARR16* was particularly strongly diminished, leading the authors to suggest a functional link between ARR10 and ARR12, on the one hand, and ARR15 and ARR16, on the other hand.

Following a similar approach, a transcription profiling of the shoots of seedlings was part of the characterization of an *arr1 arr10 arr12* triple mutant (Argyros et al., 2008). In these plants, the cytokinin-induced increase in transcript levels of 71 genes, including many of the then well-known primary response genes, was severely attenuated. In addition, the basal transcript levels of almost 90% of these genes were reduced more than twofold, indicating a reduced response to endogenous cytokinin. The cytokinin-induced repression of the transcript levels of a number of genes was also diminished in the mutant plants. These results added convincing evidence that most of the general cytokinin response during the seedling stage is signaled through the three type-B response regulators ARR1, ARR10, and ARR12 (Argyros et al., 2008).

A study looking more broadly into the action of type-B response regulators was carried out using transgenic plants overexpressing a trans-dominant chimeric repressor of type-B response regulator action, ARR1-SRDX (see Transcription Factors Regulating the Primary Cytokinin Response; Heyl et al., 2008). Transcriptome comparison between ARR1-SRDX transgenic seedlings and wild-type seedlings identified transcripts regulated by type-B response regulators. About three-quarters of the genes induced by cytokinin in the wild type were not induced in ARR1-SRDX plants, and a total of 658 genes were found whose transcripts were less abundant in the ARR1-SRDX overexpressing plants. The functional context of these genes indicated that type-B ARRs regulate a number of processes, such as nitrate transport and metabolism, red light signaling, and crosstalk with other hormones (auxin, ethylene, and gibberellin). A possible negative feedback on ARR1 expression, as well as on cytokinin metabolism, was also detected. It was proposed that type-B ARRs are possible nodes for the integration of different signaling pathways (Heyl et al., 2008).

The latest cytokinin-related microarray study so far investigated similarities and differences of the transcriptomic response to cytokinin induction and cytokinin deficiency in roots and shoots using the CATMA microarray (Brenner and Schmölling, submitted). The rationale for this study were the marked differences of the cytokinin effects on roots and shoots which could be reflected by differences in the set of responsive genes. It was found by principal component analysis that the early pattern of the transcriptomic cytokinin response was largely similar in roots and shoots while the later response showed some notable differences: Cytokinin-induced roots had a more shoot-like global transcript

profile and, similarly, cytokinin-deficient shoots had a global transcript profile shifted toward a more root-like fashion. Interestingly, nuclear genes encoding plastid-localized proteins were strongly overrepresented among these differentially regulated genes. These observations suggest that cytokinin promotes shoot identity on the transcript level at least partially independently of morphological organ identity. Several genes with organ-specific responses were identified, including root-specificity for the cytokinin hydroxylase gene *CYP735A2* and shoot specificity for the cell cycle regulator gene *CDKA;1*. However, contrasting with the fundamental difference of the growth response of roots and shoots to the hormone, the vast majority of the cytokinin-regulated transcriptome showed similar response patterns in roots and shoots (Brenner and Schmülling, submitted).

In summary, the microarray studies have shown that the multitude of processes influenced by cytokinin (Werner and Schmülling, 2009) are reflected by changes of the abundance of transcripts encoding proteins of diverse function. There is, in part, little overlap between the genes identified by different studies, indicating their context-dependent regulation. The transcriptomic results can serve as a springboard to the investigation of downstream cytokinin signaling events, as will be discussed further below.

A META ANALYSIS OF PUBLICLY AVAILABLE MICROARRAY DATA SETS

In addition to the published and commented microarray results described in Section “Published Studies on the Cytokinin-Regulated Transcriptome,” a number of unpublished datasets are available in the microarray databases. As of July 2011, there were 28 sets of cytokinin-related microarray-derived data stored in the ArrayExpress database (www.ebi.ac.uk/arrayexpress), and a lot more are probably archived in individual labs inaccessible to the community. *Arabidopsis thaliana* is the experimental system of 25 of the publicly available datasets; one dataset comes from *Medicago truncatula*, one dataset from *Solanum tuberosum* and one from hybrid poplar (*Populus tremula* × *Populus alba*; **Table 2**). Despite great efforts to get the data into a uniform format and high-quality annotation, such as the MIAME standard for the description of a microarray experiment (Brazma et al., 2001), the data are not readily comparable for several reasons. Firstly, different array platforms were used; data from single-color microarrays are not directly comparable with data from two-color arrays. Secondly, the annotation standards are different because data originally submitted to different databases were combined. Thirdly, the experiments submitted had been designed for distinct research purposes and different growth conditions or developmental stages were used. Last but not least, each laboratory has its individual plant growth system resulting in differences in global gene expression (Massonnet et al., 2010) and, presumably, also the cytokinin response. For these reasons, we have chosen eight microarray experiments which have used the same array platform, the same organism (i.e., *Arabidopsis thaliana*), a roughly comparable experimental layout and a sufficient number of replicates, included wild-type data, and had a good overall quality of the data and traceable sample annotations for the meta analysis (**Table 2**).

The meta analysis based on these selected microarray datasets summarizes the genes most robustly regulated by cytokinin

(**Table 3**). At the top of the list, there are 6 of the 10 type-A response regulator genes and 2 more are found among the top 20 cytokinin response genes. *ARR3* ranked in position 66 of the list, while *ARR17* did not have a significant meta-*P*-value and thus does not appear to be regulated by cytokinin. Some other genes of cytokinin metabolism and signaling are also part of this list, including *CKX4* and *CKX5*, which encode cytokinin oxidases/dehydrogenases (Werner et al., 2001), the UDP-glucosyltransferase gene *UGT76C2* encoding a cytokinin-*O*-glycosyltransferase (Hou et al., 2004), the cytochrome P450 gene *CYP735A2* encoding a cytokinin *trans*-hydroxylase involved in *trans*-zeatin synthesis (Takei et al., 2004), and the cytokinin receptor gene *CRE1/AHK4* (Inoue et al., 2001; Suzuki et al., 2001). The other histidine kinase gene in the list, *AHK1*, encodes a sensor of osmotic stress (Urao et al., 1999; Tran et al., 2007; Wohlbach et al., 2008). Other well-characterized genes are also part of this list, such as the cytokinin response factors *CRF2* and *CRF5* (Rashotte et al., 2006). Interestingly, in a co-expression analysis, almost all the genes of the top 25 list that have a known function in cytokinin metabolism and signaling clustered together (**Figure 2A**). Expression of *CYP735A2* clustered separately with *ASL9* (**Figure 2A**). *ASL9* encodes a LOB-domain transcription factor whose overexpression results in a dwarf phenotype and alters the cytokinin response of seedlings, but a loss-of-function phenotype is not known (Kiba et al., 2005; Naito et al., 2007). One other gene that has not yet been functionally linked to cytokinin, *FAF3*, which is a member of the newly established *FANTASTIC FOUR* (*FAF*) gene family (Wahl et al., 2010), is part of the central cluster (**Figure 2A**). *FAF3* is most strongly expressed in the vasculature, where it might have a function in differentiation, which is regulated by cytokinin (Mähönen et al., 2006; Bishopp et al., 2011b). No functional information exists for the genes encoding the ubiquitin-conjugating enzyme 16 (*UBC16*), the metallo-hydroxylase/oxidoreductase superfamily protein (*MHO*), and cytochrome P450 *CYP82F1*. These three genes show, together with *ARR16*, the less related (most distinct) expression profile compared to the known cytokinin-related genes (**Figure 2B**).

We were interested whether the top 20 cytokinin-responsive genes were also targets of other signaling pathways. We, therefore, analyzed their response to different hormones and environmental cues using the data sets available from Genevestigator (Hruz et al., 2008). We selected hormone experiments that employed conditions similar to those of the cytokinin experiments (Goda et al., 2008). The heatmap, shown in **Figure 3**, reveals that several of the top 20 cytokinin response genes are the subject of rapid regulation by other hormones. In most cases, this regulation appears to cause a reduction of the transcript levels rather than an induction, and most genes respond at a later time point to these other hormones than to cytokinin. Treatment with auxin, and partly also with ABA, for example, causes reduced transcript levels of numerous cytokinin-responsive genes after 1–3 h (**Figure 3**). Surprisingly, the strongest and earliest reactions were seen with brassinolide (BL) and methyl jasmonate. Expression of *CKX4* is upregulated and expression of *CRE1/AHK4*, *CRF2*, and *ARR9* is downregulated only 30 min after BL treatment. Stronger and broader effects of the different hormones on the transcript levels of cytokinin-regulated genes are seen at later time points (data not

Table 2 | Cytokinin-related microarray datasets in the ArrayExpress database.

ID	Title	No. of arrays	Species	Array platform	Lab	Used for meta analysis
E-ATMX-3	Transcription profiling of <i>Arabidopsis thaliana</i> wild type and <i>ARR7</i> plants to identify cytokinin response genes at two different time points	16	<i>Arabidopsis thaliana</i>	Affymetrix ATH1	J. Kim, Chonnam National University, South Korea	✓
E-ATMX-11	Transcription profiling of <i>Arabidopsis</i> Col-0 wild-type and <i>35S:ARR1-SRDX</i> transgenic seedlings either induced with 6-Benzyladenine or mock-treated	36	<i>A. thaliana</i>	CATMA v2.4	T. Schmölling, Freie Universität Berlin, Germany	
E-CAGE-49	Transcription profiling of <i>Arabidopsis</i> plants (<i>tup3</i> mutant and Col-0 <i>gl</i>) in cotyledon stage after treatment with cytokinin or auxin for different times	56	<i>A. thaliana</i>	CATMA v2.3	T. Schmölling, Freie Universität Berlin, Germany	
E-CAGE-111	Cytokinin induction of roots and shoots	40	<i>A. thaliana</i>	CATMA v2.3	T. Schmölling, Freie Universität Berlin, Germany	
E-CAGE-169	Transcription profiling of <i>Arabidopsis</i> roots in response to cytokinin treatment and phosphate deprivation	16	<i>A. thaliana</i>	CATMA v2.3	R. Solano, Centro Nacional de Biotechnología, Spain	
E-CAGE-170	Transcription profiling of <i>Arabidopsis</i> roots of mutant in response to cytokinin treatment	12	<i>A. thaliana</i>	CATMA v2.3	R. Solano, Centro Nacional de Biotechnología, Spain	
E-CAGE-191	Transcription profiling of <i>Arabidopsis</i> roots in response to cytokinin treatment and phosphate deprivation	16	<i>A. thaliana</i>	CATMA v2.3	R. Solano, Centro Nacional de Biotechnología, Spain	
E-GEOD-1766	Transcription profiling of <i>Arabidopsis</i> seedlings treated with 6-Benzyl adenine – time series	13	<i>A. thaliana</i>	Affymetrix ATH1	T. Schmölling, Freie Universität Berlin, Germany	✓
E-GEOD-5698	Transcription profiling of <i>Arabidopsis</i> cytokinin treatment of seedlings – AtGenExpress	12	<i>A. thaliana</i>	Affymetrix ATH1	T. Mizuno, Nagoya University, Japan	✓
E-GEOD-5725	Transcription profiling of <i>Arabidopsis</i> <i>Agrobacterium tumefaciens</i> -induced tumor development	4	<i>A. thaliana</i>	Affymetrix ATH1	R. Deeken, Universität Würzburg, Germany	
E-GEOD-6148	Transcription profiling of <i>Arabidopsis</i> cells cultures undergoing transdifferentiation	6	<i>A. thaliana</i>	Affymetrix ATH1	D. Brown, Nottingham <i>Arabidopsis</i> Stock Centre, United Kingdom	
E-GEOD-6832	Transcription profiling of <i>Arabidopsis</i> aerial parts of seedlings after cytokinin	12	<i>A. thaliana</i>	Affymetrix ATH1	H. Sakakibara, RIKEN Plant Science Center, Yokohama	✓
E-GEOD-15049	The cytokinin type-B response regulator <i>PtRR13</i> is a negative regulator of adventitious root development in <i>Populus</i>	39	<i>Populus tremula</i> × <i>Populus alba</i>	Nimblegen <i>Populus trichocarpa</i> array	J. M. Davis, University of Florida, USA	
E-GEOD-20231	Transcript profiling of auxin/cytokinin crosstalk in the <i>Arabidopsis</i> primary root apex	14	<i>A. thaliana</i>	Affymetrix ATH1	K. Ljung, Umeå Plant Science Centre, Sweden	✓
E-GEOD-20232	Cytokinin treatment on roots of seedlings	12	<i>A. thaliana</i>	Affymetrix ATH1	H. Sakakibara, RIKEN Plant Science Center, Yokohama	✓
E-MEXP-125	Transcription profiling of <i>Arabidopsis</i> <i>ARR2</i> mutants vs wild type	5	<i>A. thaliana</i>	Syngenta Virtual Array	K. Harter, Universität zu Köln, Germany	
E-MEXP-254	Transcription profiling of <i>Plasmodiophora brassicae</i> infection of <i>Arabidopsis thaliana</i>	12	<i>A. thaliana</i>	Affymetrix AG	J. Siemens, Technische Universität Dresden, Germany	

(Continued)

Table 2 | Continued

ID	Title	No. of arrays	Species	Array platform	Lab	Used for meta analysis
E-MEXP-344	Transcription profiling of 5 to 7 days old <i>Arabidopsis thaliana</i> seedlings of wild type (Col-0) and the cytokinin-deficient transgenic line <i>35S:CKX1</i>	10	<i>A. thaliana</i>	Affymetrix ATH1	T. Schmülling, Freie Universität Berlin, Germany	
E-MEXP-432	Transcription profiling of inducible overexpression of <i>Arabidopsis</i> meristem regulators by <i>AlcR/AlcA</i> system in continuous light	10	<i>A. thaliana</i>	Affymetrix ATH1	J. U. Lohmann, Max Planck Institute for Developmental Biology, Germany	
E-MEXP-749	Transcription profiling time series of <i>Arabidopsis</i> seedlings treated with the cytokinin benzyladenine or DMSO	13	<i>A. thaliana</i>	Affymetrix AG	J. J. Kieber, University of North Carolina, USA	
E-MEXP-1154	Transcription profiling of wild type and <i>athk1</i> knock-down <i>Arabidopsis</i> plants under dehydration stress conditions	4	<i>A. thaliana</i>	Agilent <i>Arabidopsis 2</i>	K. Yamaguchi-Shinozaki, University of Tokyo, Japan	
E-MEXP-1155	Transcription profiling of wild type, <i>athk1</i> knock-down and <i>athk2/athk3</i> knockout <i>Arabidopsis</i> plants under non-stress conditions	4	<i>A. thaliana</i>	Agilent <i>Arabidopsis 2</i>	K. Yamaguchi-Shinozaki, University of Tokyo, Japan	
E-MEXP-1442	Transcription profiling of <i>Medicago truncatula</i> with benzylaminopurine treatment to the root tips and untreated control plants	4	<i>Medicago truncatula</i>	UNIBI-Mt16kOL1Plus	F. Frugier, Institut des Sciences du Végétal, France	
E-MEXP-1573	Transcription profiling of wild type and <i>arr1, 10, 12</i> mutant <i>Arabidopsis</i> seedlings treated with the cytokinin benzyladenine	8	<i>A. thaliana</i>	Affymetrix ATH1	E. G. Schaller, Dartmouth College, USA	✓
E-MEXP-2270	Transcription profiling of <i>Arabidopsis</i> wild type, <i>35S-ARR7</i> , <i>arr7</i> , and <i>35S-ARR15</i> treated with cytokinin, auxin, or both	32	<i>A. thaliana</i>	Affymetrix ATH1	J. U. Lohmann, Max Planck Institute for Developmental Biology, Germany	✓
E-MEXP-2350	Transcription profiling of <i>Arabidopsis thaliana</i> <i>AP1::AtIPT4</i> transformants vs wild-type plants to identify the candidate genes responsive to cytokinin	6	<i>A. thaliana</i>	Affymetrix ATH1	X. S. Zhang, Shandong Agricultural University, China	
E-MEXP-2921	Transcription profiling by array of potato tuber buds from wild type and <i>AtCKX1</i> expressing tubers treated with gibberellic acid	12	<i>Solanum tuberosum</i>	POCI S <i>tuberosum</i> 44K v2	S. Sonnewald, Friedrich-Alexander-Universität Erlangen-Nürnberg, Germany	
E-NASC-75	Transcription profiling of aerial parts of <i>Arabidopsis</i> seedlings treated with the cytokinin <i>trans-zeatin</i>	12	<i>A. thaliana</i>	Affymetrix ATH1	H. Sakakibara, RIKEN Plant Science Center, Yokohama	

Microarray datasets available in July 2011 are listed here. The datasets used for the meta analysis (Table 3) are indicated in the last column.

shown). Taken together, this supports the idea that the antagonistic and agonistic action of cytokinin and different hormones are, at least partly, realized through transcriptional regulation of the same target genes.

A second analysis for the changes in transcript abundance of the top 20 cytokinin response genes was made in a similar fashion for their responses to different stresses or environmental cues. An arbitrary selection of the resulting heatmaps is shown in Figure 4. Most stress conditions (heat, drought, osmotic and biotic stress, high light irradiation) caused a reduced expression

of cytokinin response genes. An exception is cold stress, which induced an increase of transcript levels in most cases. Short treatments (45 min) with blue or red light did not cause changes of the transcript abundance of the genes analyzed, while UV-B light was repressive, at least after 6 h treatment. As most stress treatments inhibit growth, it could be that the downregulation of cytokinin-responsive genes is an early indication of such growth inhibitory activity. We hypothesize that modulation (i.e., mostly a reduction) of the cytokinin status is a common part of the general stress response, and that this modulation is required to redirect the use

Table 3 | Top 25 genes of a meta analysis across selected transcript profiling experiments of *Arabidopsis thaliana* performed with Affymetrix ATH1 arrays.

AGI	Meta P	Average ratio	Encoded protein
AT1G74890	214.45	6.90	ARR15
AT3G48100	206.35	7.56	ARR5
AT2G40670	189.62	5.37	ARR16
AT1G19050	169.59	4.96	ARR7
AT1G10470	164.68	2.74	ARR4
AT5G62920	163.05	5.34	ARR6
AT1G16530	157.32	3.11	ASL9
AT5G05860	154.99	2.47	UGT76C2
AT3G57040	144.94	2.55	ARR9
AT5G19260	136.46	2.72	Protein of unknown function (DUF3049)
AT4G29740	133.59	3.48	CKX4
AT4G03610	131.02	2.92	Metallo-hydrolase/oxidoreductase superfamily protein
AT4G23750	130.50	2.22	CRF2
AT2G01830	128.24	2.35	CRE1/AHK4
AT2G17820	125.80	2.98	AHK1
AT1G75440	124.56	2.35	UBC16
AT2G46310	124.16	2.64	CRF5
AT1G67110	124.03	6.45	CYP735A2
AT2G41310	122.30	2.72	ARR8
AT2G25160	120.28	5.16	CYP82F1
AT2G40230	119.64	2.26	HXXXD-type acyltransferase family protein
AT1G13420	117.93	8.20	ST4B
AT1G75450	115.82	2.24	CKX5
AT2G30540	114.84	1.99	Thioredoxin superfamily protein
AT3G62930	114.26	2.19	Thioredoxin superfamily protein

Raw data of eight datasets with the accessions E-ATMX-3, E-GEOD-1766, E-GEOD-5698, E-GEOD-6832, E-GEOD-20231, E-GEOD-20232, E-MEXP-1573, and E-MEXP-2270 were downloaded from the ArrayExpress database (www.ebi.ac.uk/arrayexpress). The experiments were re-evaluated using an RMA-based analysis pipeline written in R using the *limma* (Smyth, 2005) and *affy* (Irizarry et al., 2005) packages. P-values for cytokinin-induced wild-type samples vs untreated wild-type samples were calculated. A meta-P-value was calculated for each gene by adding the negative \log_2 of the P-values of the individual experiments. The genes were then ranked by the meta-P-value. A gene having a P-value of 0.01 in all 11 experiments would have a meta-P-value of 53.15. According to this cut off, 1606 genes would be significantly regulated by cytokinin in the selected experiments. The meta-P-value of the last gene in this table corresponds to a P-value of 0.00005 in all eight experiments.

of plant resources for the stress response and no longer to maintain growth. Consistently, there is a tendency that stress responses are accompanied by a reduction of the cytokinin content (Hare et al., 1997).

THE FUNCTIONAL CONTEXT OF CYTOKININ-REGULATED GENES


Regulation of the transcript abundance of a given gene can be an indication for its functional relevance in a cytokinin-regulated process. The gene expression profiling studies described a plethora

of candidate genes. More than 3000 *Arabidopsis* genes in total have been described as being regulated by cytokinin. We were interested to know which genes related to which processes would be found independently by different laboratories. We have screened, therefore, the existing publications and show in **Table 4** an arbitrary and non-exhaustive compilation of genes that were identified at least three times in the 13 existing studies (see Published Studies on the Cytokinin-Regulated Transcriptome; **Table 2**). Some of the genes were identified up to 10 times, indicating that they are highly sensitive to a change in the cytokinin status (**Table 4**). Interestingly, the same cytokinin metabolism genes as in the meta analysis (**Table 3**), namely *CKX4*, *CYP735A2*, and *UGT76C2* (see previous chapter), were found most often by different laboratories. This overlap indicates that similar relevant genes might be found by both approaches supporting the validity of the meta analysis (**Table 3**).

A closer inspection of the functional context of the cytokinin-regulated genes has revealed a number of common themes which provide insights into the great variety of cytokinin actions. The processes that were identified range from crosstalk with other hormones, to different events related to growth (e.g., cell expansion, meristem activity), the redox state, and various biotic and abiotic stresses (**Table 4**). In the following sections, we discuss several of these genes. Detailed discussions of the functional context of some other cytokinin-regulated genes can also be found in several of the transcriptomic studies (Rashotte et al., 2003; Kiba et al., 2004, 2005; Brenner et al., 2005; Brenner and Schmölling, submitted). Importantly, genetic studies have provided evidence in some cases that the indications of a putative functional relevance concluded from cytokinin-sensitive changes of transcript abundance are indeed reflected by a role of the hormone in the respective process.

Crosstalk between cytokinin and other phytohormones

Among the more than 20 auxin-related genes that were found to be regulated by cytokinin are many that are also regulated by auxin, indicating that the auxin–cytokinin crosstalk may often be realized through transcriptional regulation. The list comprises five *AUX/IAA* genes and eight *SAUR*-like genes, including *SAUR-AC1* (**Table 4**). However, only two of these, *SHY2/IAA3* and *AXR3/IAA17*, were found in four independent studies. The *AUX/IAA* genes encode negative regulators of auxin signaling rapidly degraded upon an auxin stimulus through the ubiquitin–proteasome pathway (Dharmasiri et al., 2005; Kepinski and Leyser, 2005). Transcription of *SHY2/IAA3* is directly activated by the type-B response regulator ARR1 in the root meristem (Dello Ioio et al., 2008b). Its activation leads to cell differentiation by auxin redistribution caused by its negative effect on genes encoding PIN auxin efflux carriers. Regulation of *PIN* genes by cytokinin has also been reported by others (Laplaze et al., 2007; Růžicka et al., 2009; Bishopp et al., 2011b,c; Yoshida et al., 2011). The function of *AXR3* in auxin signaling is of a more general type. It was reported to have functions in leaf development, apical dominance, several aspects of root development, and crosstalk with brassinosteroids (Leyser et al., 1996; Pérez-Pérez et al., 2010). The role of the *SAUR*-like genes has remained unclear thus far (Jain et al., 2006), apart from reports about one rice *SAUR* gene, which acts as a negative






FIGURE 3 | Regulation of cytokinin response genes by different hormones. The hormonal meta-profile analysis of the top 20 cytokinin response genes (see **Table 2**) was established using the Genevestigator database (<http://www.genevestigator.com/gv/index.jsp>; Hruz et al., 2008) and modified. Only the hormone experiments described by Goda et al. (2008) were considered as tissues (i.e., seedlings); time points and other experimental conditions were most similar to those used for the cytokinin treatments. The gene expression responses are calculated as ratios between

hormone-treated and mock-treated samples. The resulting heatmap is color coded as indicated and thus reflects the up- (red color) or downregulation (green color) of genes. Each experiment can be retrieved from the NASCArrays database (Craigion et al., 2004) by using their unique ID. Abbreviations used in the figure: d, days; h, hours; min, minutes; μ M, micromolar; CK, cytokinin; IAA, indole-3-acetic acid; ACC, 1-aminocyclopropane-1-carboxylic acid; ABA, abscisic acid; GA, gibberellic acid; BL, brassinolide; MeJa, methyl jasmonate.

regulator of auxin synthesis and transport (Kant and Rothstein, 2009; Kant et al., 2009). No specific role related to cytokinin has been revealed for these genes.

Auxin–cytokinin interaction also plays a role in shoot apical meristem control (Zhao et al., 2010). The absence of meristem control genes, which are known to be regulated by cytokinin, such as *CLV1* (Lindsay et al., 2006; Gordon et al., 2009; Bartrina et al., 2011), from the microarray data is probably due to the fact that either their regulation was not detected, or that the transcripts were too low in abundance, both due to the small size of their respective expression domains in comparison to the tissues which were sampled for the transcriptome studies. Similarly, cytokinin-regulated auxin genes involved in the patterning of the root vasculature, where auxin and cytokinin act in a signaling loop causing the mutual restriction of the active domains of the respective other hormone (Bishopp et al., 2011b,c), were missed, probably for the

same reasons. A possible exception is *FAF3*, a novel gene which was discussed in Section “A Meta Analysis of Publicly Available Microarray Data Sets.”

Cytokinin also interacts with ethylene. It causes the upregulation of four aminocyclopropane-1-carboxylate synthase (*ACS*) genes involved in ethylene biosynthesis, as well as six ethylene-sensitive transcription factor genes of the *ERF/AP2* family. However, none of these genes has been identified in more than two different studies. Whether the upregulation of the transcription factors is directly caused by cytokinin, or whether it is indirectly mediated by increased ethylene levels due to the induction of *ACS* genes, cannot be decided based on the array experiments summarized here. It is known that part of the cytokinin response is mediated through ethylene as a second messenger, such as the regulation of root architecture (Cary et al., 1995; Laplaze et al., 2007; Růžicka et al., 2009).




Table 4 | Selection of developmental and physiological processes in which cytokinin-regulated genes participate.

Annotated function	Examples of regulated genes	AGI	Reference codes*
Abiotic stress: 70 genes, including 10 <i>HSP</i> , 7 peroxidase, 6 thioredoxin, 2 glutaredoxin genes	<i>AHK1</i> <i>ANNAT4</i> <i>GSTU26</i> <i>Thioredoxin</i> <i>Thioredoxin</i>	AT2G17820 AT2G38750 AT1G17190 AT2G30540 AT3G62930	1, 2, 4, 6, 7, 8, 10 3, 4, 6, 7, 8 4, 6, 7, 8, 11, 12 3, 6, 8, 9, 10 4, 6, 8, 9
Auxin action: 30 genes, including 9 <i>SAUR</i> -like and 4 <i>Aux/IAA</i> genes	<i>AXR3</i> <i>IAA3</i>	AT1G04250 AT1G04240	1, 3, 8, 12 3, 5, 7, 9
Biotic stress: 61 genes, including 4 dirigent-like and 6 <i>WRKY</i> genes	<i>Dirigent-like</i> <i>Dirigent-like</i> <i>NLM1</i> <i>YLS9</i>	AT4G11190 AT4G11210 AT4G19030 AT2G35980	3, 4, 6, 8, 9, 11 3, 6, 8, 9 3, 6, 7, 8, 9, 11 3, 7, 8, 11
Cell wall: 42 genes, including 13 expansin, 4 laccase, 6 pectin-modifying, and 5 xyloglucan-modifying genes	<i>EXP1</i>	AT1G69530	1, 2, 3, 5, 6, 8, 9
Cytokinin homeostasis: 10 genes, including 5 cytokinin oxidase/dehydrogenase and 2 purine permease genes	<i>CKX4</i> <i>CYP735A2</i> <i>UGT76C2</i>	AT4G29740 AT1G67110 AT5G05860	2, 3, 4, 6, 8, 9, 10, 11, 12 3, 6, 8, 9, 10, 11 4, 6, 8, 12
Primary metabolism (including regulation): 35 genes, including 10 trehalose-6-phosphate metabolism genes	<i>Cell wall invertase 1</i>	AT3G13790	1, 7, 8, 12
Root apical meristem: 12 genes, including 3 thalianol metabolism genes	<i>THAD</i> <i>THAH</i> <i>THAS</i>	AT5G47990 AT5G48000 AT5G48010	4, 6, 7, 8 6, 7, 8, 12 6, 7, 8
Transcription: 98 genes, including 4 <i>bHLH</i> , 3 <i>CRF</i> , 5 other <i>ERF/AP2</i> , 3 <i>HAT</i> , 9 <i>MYB</i> , 4 <i>WRKY</i> genes	<i>ASL9</i> <i>ATR1</i> <i>CRF2</i> <i>CRF5</i>	AT1G16530 AT5G60890 AT4G23750 AT2G46310	3, 6, 8, 10, 11 1, 2, 6, 8, 12 1, 3, 4, 6, 7, 8, 9, 12 3, 4, 6, 7, 8, 9
Ethylene action: 18 genes, including 4 ACC synthase, 6 ethylene-sensitive <i>ERF/AP2</i> transcription factor genes			
Gibberellin action: 12 genes, including <i>GA4</i> , <i>GNL</i> , <i>GNC</i>			
Light signalling/response: 20 genes, including 4 <i>NPH3</i> family genes			
Nutrient uptake: 17 genes, including 3 <i>AMT</i> , 4 <i>NRT</i> , 2 <i>PHT</i> , 3 <i>SULTR</i> family genes			
Photosynthesis: 12 genes, including 4 <i>LHCB2</i> genes			

All genes listed in 12 medium- and large-scale transcriptomic studies related to cytokinin were collected and functionally annotated to generate this non-exhaustive list of developmental and physiological processes in which cytokinin-regulated genes participate. Functional categories harboring either a particularly large number of genes or genes found in three or more publications were included in this table. Single genes listed in four or more studies, or genes belonging to an operon-like structure were considered. The genes encoding proteins of the cytokinin signaling system have not been listed. Compilations of their roles in various biological processes can be found elsewhere (Heyl et al., 2011; Müller, 2011). *Reference codes: 1, Che et al. (2002); 2, Hoth et al. (2003); 3, Rashotte et al. (2003); 4, Kiba et al. (2004); 5, Brenner et al. (2005); 6, Kiba et al. (2005); 7, Rashotte et al. (2006); 8, Lee et al. (2007); 9, Taniguchi et al. (2007); 10, Yokoyama et al. (2007); 11, Argyros et al. (2008); 12, Heyl et al. (2008).

The interaction of cytokinin with gibberellins (GA) is reflected on the molecular level by the regulation of two GATA transcription factor genes, *GNL/CGA1/GATA22* and *GNC/GATA21*. It was shown in a recent genetic study that both *GNL* and *GNC* are direct targets of the GA signaling pathway, and that both negatively regulate germination, greening, leaf expansion, and flowering time (Richter et al., 2010). Additionally, *GNL* seems to have a role in chloroplast development, as *GNL* overexpressors developed chloroplasts in illuminated roots, while a *gnl* mutant formed fewer chloroplasts in the hypocotyl of seedlings. It has been proposed

that *GNL* integrates signaling pathways of light, GA, and cytokinin (Köllmer et al., 2011). Cytokinin also appears to interfere with GA activity by regulating the GA synthesis gene *GA4* and the GA signaling genes *RGA* and *GAI*, both encoding DELLA repressor proteins. Together, cytokinin action caused a reduction of GA activity (Brenner et al., 2005).

Cytokinin affects different parts of the plant cell

One of the most frequently listed genes in the present microarray studies encodes the expansin *EXP1*. Twelve other expansin

genes were found in at least two microarray studies, along with 18 other cell wall-related genes (Table 4). Expansins are thought to be involved in weakening the structure of the cell wall, which is required for cell expansion, and thus growth. Links between cytokinin and cell wall-related genes have been found previously (Downes and Crowell, 1998). A cytokinin-induced change of wall extensibility has also been directly measured (Thomas et al., 1981), and a negative influence of cytokinin signaling on cell wall thickness has been reported (Jung et al., 2008). However, detailed investigations on the role of *EXPI* and other cell wall-related genes in growth regulation by cytokinin have not yet been published.

Cytokinin has been known to have a positive effect on chloroplast development, performance, and longevity for a long time (Staetler and Laetsch, 1965; Parthier, 1979; Chory et al., 1994; Yaronskaya et al., 2006; Okazaki et al., 2009; Brenner and Schmölling, submitted), and may thus influence photosynthesis. It affects the transcript abundance of genes encoding plastid-localized proteins (Zubo et al., 2008), and a proteomic study identified about 50% of the cytokinin-regulated proteins to be localized in the chloroplast (Cerný et al., 2011). Furthermore, cytokinin appears to exert a protective activity on biochemical processes associated with photosynthesis (Rivero et al., 2009). Four of the six genes found to be induced by cytokinin in at least two publications belong to the *LHCB2* family, encoding components of the light harvesting complexes. This is in accordance with the influence of cytokinin on the relative amount of light harvesting complexes (Siffel et al., 1992; Catský et al., 1993) and on the accumulation of light harvesting complex mRNAs and proteins (Marziani Longo et al., 1990).

Cytokinin as a regulator of metabolism genes

Several genes annotated as linked to root growth but rarely considered in cytokinin-related studies were identified. The *THAD*, *THAH*, and *THAS* genes are required for the synthesis of thalianol, which is a triterpene of unknown function (Field and Osbourn, 2008). The genes are organized in an operon-like fashion (Field and Osbourn, 2008). The operon includes another gene encoding an acyltransferase of unknown function (At5g47980) which has also been found in numerous microarray studies to be regulated by cytokinin (Kiba et al., 2004, 2005; Rashotte et al., 2006; Lee et al., 2007; Argyros et al., 2008; Heyl et al., 2008). Interestingly, plants with a higher thalianol content show a stunted shoot growth but have longer roots (Field and Osbourn, 2008), both phenotypic features reminiscent of cytokinin-deficient plants (Werner et al., 2003). However, the exact role of thalianol in regulating root and shoot growth and its possible interplay with cytokinin is unknown.

A function of cytokinin in regulating uptake and metabolism of different nutrients – nitrogen, sulfate, phosphate, and iron – has been known for some time (Sakakibara et al., 2006; Ribot et al., 2008; Argueso et al., 2009; Igarashi et al., 2009; Rubio et al., 2009). This is supported by the regulation of nitrogen (*AMT*, *NRT*), sulfate (*SULTR*), and phosphate (*PHT*) transporter genes by the hormone (Table 4). In the case of nitrate, genes encoding both high-affinity (*NRT1*) and low-affinity (*NRT2*) transporters (Crawford and Glass, 1998) are regulated by cytokinin. *NRT1.1* – also cytokinin-regulated – functions as a nitrate sensor

(Ho et al., 2009), suggesting an influence of cytokinin on nitrate sensing. The reciprocal signaling relations between nitrogen and cytokinin were reviewed recently (Kiba et al., 2011). One of the three ammonium transporters, *AMT1;3*, has a major role in ammonium-triggered lateral root branching (Lima et al., 2010). The sulfate transporter *SULTR1;2* is responsible for sulfate uptake into the roots, while the exact role of the other two sulfate transporters, *SULTR3;1* and *SULTR3;4*, is not yet clear. *SULTR1;2* and *SULTR1;1* were shown to be negatively regulated by cytokinin in a partially *CRE1/AHK4*-dependent manner, associated by a decrease in sulfate uptake (Maruyama-Nakashita et al., 2004). The functions of the phosphate transporters *PHT1;2* and *PHT1;4* are not restricted to phosphate uptake from the soil into the root, but extend to broader roles in phosphate remobilization and distribution throughout the whole plant (Mudge et al., 2002).

Nineteen genes involved in primary metabolism and its regulation were identified, however, only the one encoding cell wall invertase 1 was found repeatedly (Table 4). Cell wall invertases provide hexose sugars to cells by cleavage of sucrose in the cell wall. The provision of either hexoses or sucrose has a regulatory potential for source–sink relationships and carbon utilization, as well as for development (Sherson et al., 2003). Total cell wall invertase activity was shown to be reduced in the source leaves of cytokinin-deficient plants (Werner et al., 2008), and *cell wall invertase 1* expression was consistently found to be increased by cytokinin in the transcriptomics studies summarized here. Ten of the 19 genes assigned to the category primary metabolism are involved in trehalose-6-phosphate metabolism. Trehalose-6-phosphate is also regulated by source–sink relationships and carbon utilization, and plants with a disturbed trehalose-6-phosphate homeostasis show severe developmental and metabolic defects (Schluepmann et al., 2003, 2004; Dijken et al., 2004). It was shown that trehalose synthesis genes are regulated by cytokinin in an opposite fashion than genes encoding trehalose-degrading enzymes, supporting the idea that trehalose-6-phosphate levels are regulated, at least partly, by cytokinin (Brenner et al., 2005). In summary, the gene regulatory data indicate that cytokinin action on source–sink relationships and carbon utilization could be accomplished through the regulation of cell wall invertase and trehalose-6-phosphate metabolism genes.

Cytokinin and light

Cytokinin regulates the transcript levels of four *NPH3*-like genes, which have an annotated function in light signaling. However, *NPH3* proteins are involved in various signaling pathways. Its founding member interacts with the blue light receptor *NPH1/PHOT1* and with the E3 ubiquitin ligase component *CUL3*; the resulting degradation contributes to photomorphogenesis and leaf positioning (Motchoulski and Liscum, 1999; Pedmale and Liscum, 2007; Inoue et al., 2008). Five other *NPH3*-like genes have been linked with auxin-regulated organogenesis acting through influencing the polar localization of PIN proteins (Trembl et al., 2005; Cheng et al., 2007; Furutani et al., 2007, 2011). The four *NPH3*-like genes found repeatedly to be regulated by cytokinin, two of which are closely related phylogenetically, have not yet been functionally described.

The transcript level of three genes encoding members of the phytochrome signaling pathway is regulated by cytokinin in the combined microarray dataset: *PHYA*, the phytochrome kinase substrate *PKS1*, and the GRAS transcription factor gene *PAT1*. *PHYA* has functions in far-red-dependent de-etiolation, day-length sensing, internode and leaf growth, and shade avoidance (Schepens et al., 2004). *PKS1* is a negative regulator of phytochrome signaling (Fankhauser et al., 1999; Lariguet et al., 2003) and is also necessary for blue light-induced phototropism in a phyA-dependent manner (Lariguet et al., 2006; Boccalandro et al., 2008). *PAT1* is a member of the transcription factor GRAS family and specifically involved in an early stage of phyA signaling (Bolle et al., 2000). The transcriptional regulation of genes involved in phytochrome signaling pathways is consistent with known links between cytokinin and light responses (Argueso et al., 2009; Werner and Schmülling, 2009; Yoshida et al., 2011).

Cytokinin regulation of transcription factor genes

Gene expression studies have revealed very early that cytokinin acts downstream of the TCS through activating transcriptional cascades (Rashotte et al., 2003; Brenner et al., 2005), and several studies have identified functionally relevant cytokinin-regulated transcription factor genes (Argueso et al., 2010). Other cytokinin-regulated transcription factor genes are still looking for a function (*CRF2*, *CRF5*; see above), or have, so far, only been studied by overexpression, such as *ASL9* (Naito et al., 2007), *bHLH64*, *HAT4*, *HAT22*, and *GATA22/CGA1/GNL* (Köllmer et al., 2011). Cytokinin regulates the transcript levels of many other transcription factor genes, and seven of them have been detected in at least four publications describing transcriptomic studies (Table 4). No function is yet known for most of them, such as the zinc finger transcription factor *At1g68360*, which was found to be also regulated by auxin (Pufky et al., 2003) and cold stress (Vergnolle et al., 2005), or the B-box zinc finger transcription factor *STH7* (Datta et al., 2008), which responded to molecules in wildfire smoke that stimulate seed germination (Nelson et al., 2010). The bZIP transcription factor *BZIP11* (formerly known as *ATB2*) is interesting because it has been proposed to be a powerful regulator of carbohydrate metabolism that functions in a growth regulatory network including trehalose-6-phosphate (see Cytokinin as a Regulator of Metabolism Genes; Ma et al., 2011). *BZIP11* controls the sink strength by regulating the expression of genes in young sink tissue (Delatte et al., 2011). In addition, it has been shown to control two genes encoding enzymes in amino acid metabolism, *ASN1* and *ProDH2*. *ATR1* encodes a transcription factor of the MYB family, which is the key regulator of genes involved in the biosynthesis of glucosinolates and auxin from tryptophane (Celenza et al., 2005). Its regulation by cytokinin provides another facet to the cytokinin–auxin crosstalk. *TZF1* encodes a zinc finger protein which has been identified as a regulated gene in studies on the carbon status (Contento et al., 2004; Lin et al., 2011), disease and insect defense (Cartieaux et al., 2003; Ascencio-Ibáñez et al., 2008), stress tolerance (Kasukabe et al., 2004; Lin et al., 2011), development of male and female reproductive organs (Alves-Ferreira et al., 2007; Skinner and Gasser, 2009), and ABA and GA signaling (Lin et al., 2011). Its

mode of action is currently under investigation (Pomeranz et al., 2011)

A role for cytokinin in the stress response

Many cytokinin-responsive genes are also regulated by different abiotic and biotic stresses (Figure 4). One important type of abiotic stress influenced by cytokinin appears to be oxidative stress, as many genes encoding enzymes involved in relieving oxidative stress – peroxidases, thioredoxins, and glutaredoxins – were found to be regulated by the hormone in different microarray studies (Table 4). A beneficial effect of cytokinin in coping with oxidative stress has been described (Zavaleta-Mancera et al., 2007). Two thioredoxin genes, *At2g30540* and *At3g62930*, were particularly often named in the publications listed in Table 4. They were consistently found among the most strongly cytokinin-regulated genes in the meta analysis, namely at position 24 and 25, respectively (Table 3). No precise function has yet been assigned to these genes. Generally, thioredoxins regulate the redox status of disulfide bonds of their target proteins, thereby supplying reducing power for reductases, which detoxify lipid hydroperoxides or repair damaged proteins. In addition to these functions, they may have roles in oxidative stress signaling or as regulators of scavenging mechanisms (Vieira Dos Santos and Rey, 2006). Three other genes related to abiotic stress, *AHK1*, *ANNAT4*, and *GSTU26*, were listed in 6 or 7 of the 13 cytokinin-related transcriptomics studies, suggesting a relevant influence of cytokinin on their related field of action. *AHK1* encodes a histidine kinase receptor structurally related to the cytokinin receptors and functions as a positive regulator of drought and salt stress responses and ABA signaling (Tran et al., 2007). *ANNAT4* encodes an annexin, a calcium-dependent membrane-binding protein that mediates osmotic stress and abscisic acid signal transduction in *Arabidopsis* (Lee et al., 2004). The glutathione *S*-transferase gene *GSTU26* is also induced by low temperature (Nutricati et al., 2006).

A link between cytokinin and biotic stress has been suggested because altered cytokinin levels have been observed after pathogen infection (Clarke et al., 1999; Maksimov et al., 2002) and an altered susceptibility to pathogens due to an altered cytokinin status has been observed (Clarke et al., 1998; Pertry et al., 2009). Therefore, it was not surprising that several defense-related genes were found to be regulated by cytokinin in multiple transcriptomic studies, among them three dirigent-like proteins. They are known to be transcriptionally induced by biotic stress (Fristensky et al., 1985; Görlach et al., 1996; Lee et al., 1996; Ralph et al., 2006) and participate as mediators of stereospecificity in the formation of lignans (Davin et al., 1997; Burlat et al., 2001), which possess antiviral, antimicrobial, and cytotoxic properties (MacRae and Towers, 1984; Mendonça Pauletti et al., 2000; Rahman and Gray, 2002). *NLM1*, also known as *NIP1;1*, encodes an aquaporin homolog that determines antimonite and arsenite sensitivity (Kamiya and Fujiwara, 2009; Kamiya et al., 2009) and also responds to auxin and osmotic stress (Kreps et al., 2002; Goda et al., 2004). *YLS9* is induced in response to various biotic triggers, including flagellin and the oviposition of butterfly eggs (Navarro et al., 2004; Michel et al., 2006; Little et al., 2007; Zhao et al., 2007).

FUTURE PROSPECTS

A wealth of information has been generated by transcriptomic studies creating new hypotheses about the functions of cytokinin and functional mechanisms of its action. Numerous interesting areas deserve further investigation, such as the regulation of ribosomal genes by cytokinin suggesting translational control of cytokinin-induced changes (Brenner et al., 2005; Kiba et al., 2005; Rashotte and Goertzen, 2010; Brenner and Schmölling, submitted). Several developmental processes are affected by mutations in ribosomal genes, including embryogenesis, root growth, and leaf development. This points out the functional relevance of the specific assembly of ribosomal proteins in addition to their fundamental role in translation, presumably because they have an effect on the translation of specific proteins (Weijers et al., 2001; Byrne, 2009; Horiguchi et al., 2011). With the dawn of the era of next-generation sequencing-based transcriptomics, more data can be expected from species whose genomes have not been sequenced thus far. Data from crop species will be of special interest for applied research given the potential of manipulating the cytokinin

status to achieve yield increase (Ashikari et al., 2005; Bartrina et al., 2011) or to ameliorate other plant traits relevant for agricultural performance (Werner et al., 2010; Zalewski et al., 2010). In addition, experimental limitations of the hybridization-based microarray platform, especially regarding annotation dependency, sensitivity, dynamic range, and reproducibility, will be overcome. With the new technologies, the transcriptome of smaller samples can be evaluated, detecting smaller changes in gene expression with higher confidence. Therefore, we are looking forward to an exciting expansion of high-fidelity transcriptomic knowledge across many species. New tools to master the challenge of making proper use of the huge body of data generated will certainly facilitate an understanding of the functions of the networks discovered.

ACKNOWLEDGMENTS

This work was funded by the DFG grant Schm 814/22-1 within the framework of the Arabidopsis Functional Genomics Network (AFGN).

REFERENCES

- Aerts, S., Van Loo, P., Thijs, G., Mayer, H., de Martin, R., Moreau, Y., and De Moor, B. (2005). TOUCAN2: the all-inclusive open source workbench for regulatory sequence analysis. *Nucleic Acids Res.* 33, W393–W396.
- Alves-Ferreira, M., Wellmer, F., Banhara, A., Kumar, V., Riechmann, J. L., and Meyerowitz, E. M. (2007). Global expression profiling applied to the analysis of *Arabidopsis* stamen development. *Plant Physiol.* 145, 747–762.
- Argueso, C. T., Ferreira, F. J., and Kieber, J. J. (2009). Environmental perception avenues: the interaction of cytokinin and environmental response pathways. *Plant Cell Environ.* 32, 1147–1160.
- Argueso, C. T., Raines, T., and Kieber, J. J. (2010). Cytokinin signaling and transcriptional networks. *Curr. Opin. Plant Biol.* 13, 533–539.
- Argyros, R. D., Mathews, D. E., Chiang, Y.-H., Palmer, C. M., Thibault, D. M., Etheridge, N., Argyros, D. A., Mason, M. G., Kieber, J. J., and Schaller, G. E. (2008). Type B response regulators of *Arabidopsis* play key roles in cytokinin signaling and plant development. *Plant Cell* 20, 2102–2116.
- Ascencio-Ibáñez, J. T., Sozzani, R., Lee, T.-J., Chu, T.-M., Wolfinger, R. D., Cella, R., and Hanley-Bowdoin, L. (2008). Global analysis of *Arabidopsis* gene expression uncovers a complex array of changes impacting pathogen response and cell cycle during geminivirus infection. *Plant Physiol.* 148, 436–454.
- Ashikari, M., Sakakibara, H., Lin, S., Yamamoto, T., Takashi, T., Nishimura, A., Angeles, E. R., Qian, Q., Kitano, H., and Matsuoka, M. (2005). Cytokinin oxidase regulates rice grain production. *Science* 309, 741–745.
- Bartrina, I., Otto, E., Strnad, M., Werner, T., and Schmölling, T. (2011). Cytokinin regulates the activity of reproductive meristems, flower organ size, ovule formation, and thus seed yield in *Arabidopsis thaliana*. *Plant Cell* 23, 69–80.
- Bishopp, A., Benková, E., and Helariutta, Y. (2011a). Sending mixed messages: auxin-cytokinin crosstalk in roots. *Curr. Opin. Plant Biol.* 14, 10–16.
- Bishopp, A., Help, H., El-Showk, S., Weijers, D., Scheres, B., Friml, J., Benková, E., Mähönen, A. P., and Helariutta, Y. (2011b). A mutually inhibitory interaction between auxin and cytokinin specifies vascular pattern in roots. *Curr. Biol.* 21, 917–926.
- Bishopp, A., Lehesranta, S., Vatén, A., Help, H., El-Showk, S., Scheres, B., Helariutta, K., Mähönen, A. P., Sakakibara, H., and Helariutta, Y. (2011c). Phloem-transported cytokinin regulates polar auxin transport and maintains vascular pattern in the root meristem. *Curr. Biol.* 21, 927–932.
- Bishopp, A., Help, H., Helariutta, Y., and Kwang, W. J. (2009). “Cytokinin signaling during root development,” in *International Review of Cell and Molecular Biology*, ed. K. Jeon (San Diego: Academic Press), 1–48.
- Boccalandro, H. E., De Simone, S. N., Bergmann-Honsberger, A., Schepens, I., Fankhauser, C., and Casal, J. J. (2008). PHYTOCHROME KINASE SUBSTRATE1 regulates root phototropism and gravitropism. *Plant Physiol.* 146, 108–115.
- Bolle, C., Koncz, C., and Chua, N.-H. (2000). PAT1, a new member of the GRAS family, is involved in phytochrome A signal transduction. *Genes Dev.* 14, 1269–1278.
- Brandstatter, I., and Kieber, J. J. (1998). Two genes with similarity to bacterial response regulators are rapidly and specifically induced by cytokinin in *Arabidopsis*. *Plant Cell* 10, 1009–1020.
- Brazma, A., Hingamp, P., Quackenbush, J., Sherlock, G., Spellman, P., Stoeckert, C., Aach, J., Ansorge, W., Ball, C. A., Causton, H. C., Gaasterland, T., Glenisson, P., Holstege, F. C. P., Kim, I. F., Markowitz, V., Matese, J. C., Parkinson, H., Robinson, A., Sarkans, U., Steffen, S.-K., Stewart, J., Taylor, R., Vilo, J., and Vingron, M. (2001). Minimum information about a microarray experiment (MIAME): toward standards for microarray data. *Nat. Genet.* 29, 365–371.
- Brenner, W. G., Romanov, G. A., Köllmer, I., Bürkle, L., and Schmölling, T. (2005). Immediately and delayed cytokinin response genes of *Arabidopsis thaliana* identified by genome-wide expression profiling reveal novel cytokinin-sensitive processes and suggest cytokinin action through transcriptional cascades. *Plant J.* 44, 314–333.
- Burlat, V., Kwon, M., Davin, L. B., and Lewis, N. G. (2001). Dirigent proteins and dirigent sites in lignifying tissues. *Phytochemistry* 57, 883–897.
- Byrne, M. E. (2009). A role for the ribosome in development. *Trends Plant Sci.* 14, 512–519.
- Carteaux, F., Thibaud, M.-C., Zimmerli, L., Lessard, P., Sarrobert, C., David, P., Gerbaud, A., Robaglia, C., Somerville, S., and Nussaume, L. (2003). Transcriptome analysis of *Arabidopsis* colonized by a plant-growth promoting rhizobacterium reveals a general effect on disease resistance. *Plant J.* 36, 177–188.
- Cary, A. J., Liu, W., and Howell, S. H. (1995). Cytokinin action is coupled to ethylene in its effects on the inhibition of root and hypocotyl elongation in *Arabidopsis thaliana* seedlings. *Plant Physiol.* 107, 1075–1082.
- Celenza, J. L., Quiel, J. A., Smolen, G. A., Merrikh, H., Silvestro, A. R., Normanly, J., and Bender, J. (2005). The *Arabidopsis* ATR1 Myb transcription factor controls indolic glucosinolate homeostasis. *Plant Physiol.* 137, 253–262.
- Che, P., Gingerich, D. J., Lall, S., and Howell, S. H. (2002). Global and hormone-induced gene expression changes during shoot development in *Arabidopsis*. *Plant Cell* 14, 2771–2785.
- Cheng, Y., Qin, G., Dai, X., and Zhao, Y. (2007). NPY1, a BTB-NPH3-like protein, plays a critical role in auxin-regulated organogenesis in *Arabidopsis*. *Proc. Natl. Acad. Sci. U.S.A.* 104, 18825–18829.
- Chory, J., Reinecke, D., Sim, S., Washburn, T., and Brenner, M. (1994). A role for cytokinins in de-etiolation in *Arabidopsis*. *Plant Physiol.* 104, 339–347.

- Clarke, S. F., Burritt, D. J., Jameson, P. E., and Guy, P. L. (1998). Influence of plant hormones on virus replication and pathogenesis-related proteins in *Phaseolus vulgaris* L. infected with white clover mosaic potyvirus. *Physiol. Mol. Plant Pathol.* 53, 195–207.
- Clarke, S. F., McKenzie, M. J., Burritt, D. J., Guy, P. L., and Jameson, P. E. (1999). Influence of white clover mosaic potyvirus infection on the endogenous cytokinin content of bean. *Plant Physiol.* 120, 547–552.
- Contento, A. L., Kim, S.-J., and Bassham, D. C. (2004). Transcriptome profiling of the response of *Arabidopsis* suspension culture cells to suc starvation. *Plant Physiol.* 135, 2330–2347.
- Craigon, D. J., James, N., Okyere, J., Higgins, J., Jotham, J., and May, S. (2004). NASCArrays: a repository for microarray data generated by NASC's transcriptomics service. *Nucleic Acids Res.* 32, D575–D577.
- Crawford, N. M., and Glass, A. D. M. (1998). Molecular and physiological aspects of nitrate uptake in plants. *Trends Plant Sci.* 3, 389–395.
- Cutcliffe, J. W., Hellmann, E., Heyl, A., and Rashotte, A. M. (2011). CRFs form protein–protein interactions with each other and with members of the cytokinin signalling pathway in *Arabidopsis* via the CRF domain. *J. Exp. Bot.* 62, 4995–5002.
- D'Agostino, I. B., Deruere, J., and Kieber, J. J. (2000). Characterization of the response of the *Arabidopsis* response regulator gene family to cytokinin. *Plant Physiol.* 124, 1706–1717.
- Datta, S., Johansson, H., Hettiarachchi, C., Irigoyen, M. L., Desai, M., Rubio, V., and Holm, M. (2008). LZFI/SALT TOLERANCE HOMOLOG3, an *Arabidopsis* B-box protein involved in light-dependent development and gene expression, undergoes cop1-mediated ubiquitination. *Plant Cell* 20, 2324–2338.
- Davin, L. B., Wang, H.-B., Crowell, A. L., Bedgar, D. L., Martin, D. M., Sarkanen, S., and Lewis, N. G. (1997). Stereoselective bimolecular phenoxyl radical coupling by an auxiliary (dirigent) protein without an active center. *Science* 275, 362–367.
- Day, R. C., Herridge, R. P., Ambrose, B. A., and Macknight, R. C. (2008). Transcriptome analysis of proliferating *Arabidopsis* endosperm reveals biological implications for the control of syncytial division, cytokinin signaling, and gene expression regulation. *Plant Physiol.* 148, 1964–1984.
- Delatte, T. L., Sedijani, P., Kondou, Y., Matsui, M., de Jong, G. J., Somssen, G. W., Wiese-Klinkenberg, A., Primavesi, L. F., Paul, M. J., and Schluempmann, H. (2011). Growth arrest by trehalose-6-phosphate: an astonishing case of primary metabolite control over growth by way of the SnRK1 signaling pathway. *Plant Physiol.* 157, 160–174.
- Dello Ioio, R., Linhares, F. S., and Sabatini, S. (2008a). Emerging role of cytokinin as a regulator of cellular differentiation. *Curr. Opin. Plant Biol.* 11, 23–27.
- Dello Ioio, R., Nakamura, K., Moubayidin, L., Perilli, S., Taniguchi, M., Morita, M. T., Aoyama, T., Costantino, P., and Sabatini, S. (2008b). A genetic framework for the control of cell division and differentiation in the root meristem. *Science* 322, 1380–1384.
- Dharmasiri, N., Dharmasiri, S., and Estelle, M. (2005). The F-box protein TIR1 is an auxin receptor. *Nature* 435, 441–445.
- Dijken, A. J. H. V., Schluempmann, H., and Smeeckens, S. C. M. (2004). *Arabidopsis* trehalose-6-phosphate synthase 1 is essential for normal vegetative growth and transition to flowering. *Plant Physiol.* 135, 969–977.
- Domagalska, M. A., and Leyser, O. (2011). Signal integration in the control of shoot branching. *Nat. Rev. Mol. Cell Biol.* 12, 211–221.
- Dortay, H., Mehnert, N., Bürkle, L., Schmülling, T., and Heyl, A. (2006). Analysis of protein interactions within the cytokinin-signaling pathway of *Arabidopsis thaliana*. *FEBS J.* 273, 4631–4644.
- Downes, B. P., and Crowell, D. N. (1998). Cytokinin regulates the expression of a soybean β -expansin gene by a post-transcriptional mechanism. *Plant Mol. Biol.* 37, 437–444.
- Edgar, R., Domrachev, M., and Lash, A. E. (2002). Gene expression omnibus: NCBI gene expression and hybridization array data repository. *Nucleic Acids Res.* 30, 207–210.
- Fankhauser, C., Yeh, K. C., Lagarias, J. C., Zhang, H., Elich, T. D., and Chory, J. (1999). PKS1, a substrate phosphorylated by phytochrome that modulates light signaling in *Arabidopsis*. *Science* 284, 1539–1541.
- Field, B., and Osbourn, A. E. (2008). Metabolic diversification – independent assembly of operon-like gene clusters in different plants. *Science* 320, 543–547.
- Fristensky, B., Riggleman, R. C., Waggoner, W., and Hadwiger, L. A. (1985). Gene expression in susceptible and disease resistant interactions of peas induced with *Fusarium solani* pathogens and chitosan. *Physiol. Mol. Plant Pathol.* 27, 15–28.
- Fukumura, R., Takahashi, H., Saito, T., Tsutsumi, Y., Fujimori, A., Sato, S., Tatsumi, K., Araki, R., and Abe, M. (2003). A sensitive transcriptome analysis method that can detect unknown transcripts. *Nucleic Acids Res.* 31, e94.
- Furutani, M., Kajiwara, T., Kato, T., Trembl, B. S., Stockum, C., Torres-Ruiz, R. N. A., and Tasaka, M. (2007). The gene MACCHI-BOU4/ENHANCER OF PINOID encodes a NPH3-like protein and reveals similarities between organogenesis and phototropism at the molecular level. *Development* 134, 3849–3859.
- Furutani, M., Sakamoto, N., Yoshida, S., Kajiwara, T., Robert, H. S., Friml, J., and Tasaka, M. (2011). Polar-localized NPH3-like proteins regulate polarity and endocytosis of PIN-FORMED auxin efflux carriers. *Development* 138, 2069–2078.
- Fusada, N., Masuda, T., Kuroda, H., Shimada, H., Ohta, H., and Takamiya, K.-I. (2005). Identification of a novel cis-element exhibiting cytokinin-dependent protein binding in vitro; in the 5'-region of NADPH-protochlorophyllide oxidoreductase gene in cucumber. *Plant Mol. Biol.* 59, 631–645.
- Goda, H., Sasaki, E., Akiyama, K., Maruyama-Nakashita, A., Nakabayashi, K., Li, W., Ogawa, M., Yamauchi, Y., Preston, J., Aoki, K., Kiba, T., Takatsuto, S., Fujioka, S., Asami, T., Nakano, T., Kato, H., Mizuno, T., Sakakibara, H., Yamaguchi, S., Nambara, E., Kamiya, Y., Takahashi, H., Hirai, M. Y., Sakurai, T., Shinozaki, K., Saito, K., Yoshida, S., and Shimada, Y. (2008). The AtGenExpress hormone and chemical treatment data set: experimental design, data evaluation, model data analysis and data access. *Plant J.* 55, 526–542.
- Goda, H., Sawa, S., Asami, T., Fujioka, S., Shimada, Y., and Yoshida, S. (2004). Comprehensive comparison of auxin-regulated and brassinosteroid-regulated genes in *Arabidopsis*. *Plant Physiol.* 134, 1555–1573.
- Gordon, S. P., Chickarmane, V. S., Ohno, C., and Meyerowitz, E. M. (2009). Multiple feedback loops through cytokinin signaling control stem cell number within the *Arabidopsis* shoot meristem. *Proc. Natl. Acad. Sci. U.S.A.* 106, 16529–16534.
- Görlach, J., Volrath, S., Knauf-Beiter, G., Hengy, G., Beckhove, U., Kogel, K. H., Oostendorp, M., Staub, T., Ward, E., Kessmann, H., and Ryals, J. (1996). Benzothiadiazole, a novel class of inducers of systemic acquired resistance, activates gene expression and disease resistance in wheat. *Plant Cell* 8, 629–643.
- Hare, P. D., Cress, W. A., and van Staden, J. (1997). The involvement of cytokinins in plant responses to environmental stress. *Plant Growth Regul.* 23, 79–103.
- Hass, C., Lohrmann, J., Albrecht, V., Sweere, U., Hummel, F., Yoo, S. D., Hwang, I., Zhu, T., Schäfer, E., Kudla, J., and Harter, K. (2004). The response regulator 2 mediates ethylene signalling and hormone signal integration in *Arabidopsis*. *EMBO J.* 23, 3290–3302.
- Hellmann, E., Gruhn, N., and Heyl, A. (2010). The more, the merrier: cytokinin signaling beyond *Arabidopsis*. *Plant Signal. Behav.* 5, 1384–1390.
- Heyl, A., Ramireddy, E., Brenner, W. G., Riefler, M., Allemeersch, J., and Schmülling, T. (2008). The transcriptional repressor ARR1-SRDX suppresses pleiotropic cytokinin activities in *Arabidopsis*. *Plant Physiol.* 147, 1380–1395.
- Heyl, A., Riefler, M., Romanov, G. A., and Schmülling, T. (2011). Properties, functions and evolution of cytokinin receptors. *Eur. J. Cell Biol.* (in press). doi:10.1016/j.ejcb.2011.1002.1009.
- Heyl, A., Werner, T., and Schmülling, T. (eds). (2006). "Cytokinin metabolism and signal transduction," in *Plant Hormone Signaling, Annual Plant Reviews*, eds P. Hedden and S. Thomas (Oxford: Blackwell Publishing), 93–123.
- Hiratsu, K., Matsui, K., Koyama, T., and Ohme-Takagi, M. (2003). Dominant repression of target genes by chimeric repressors that include the EAR motif, a repression domain, in *Arabidopsis*. *Plant J.* 34, 733–739.
- Ho, C.-H., Lin, S.-H., Hu, H.-C., and Tsay, Y.-F. (2009). CHL1 functions as a nitrate sensor in plants. *Cell* 138, 1184–1194.
- Horák, J., Brzobohatý, B., and Lexa, M. (2003). Molecular and physiological characterisation of an insertion mutant in the ARR21 putative response regulator gene from *Arabidopsis thaliana*. *Plant Biol.* 5, 245–254.
- Horiguchi, G., Mollá-Morales, A., Pérez-Pérez, J. M., Kojima, K., Robles, P., Ponce, M. R., Micol, J. L., and Tsukaya, H. (2011). Differential

- contributions of ribosomal protein genes to *Arabidopsis thaliana* leaf development. *Plant J.* 65, 724–736.
- Hosoda, K., Imamura, A., Katoh, E., Hatta, T., Tachiki, M., Yamada, H., Mizuno, T., and Yamazaki, T. (2002). Molecular structure of the GARP family of plant Myb-related DNA binding motifs of the *Arabidopsis* response regulators. *Plant Cell* 14, 2015–2029.
- Hoth, S., Ikeda, Y., Morgante, M., Wang, X., Zuo, J., Hanafey, M. K., Gaasterland, T., Tingey, S. V., and Chua, N. H. (2003). Monitoring genome-wide changes in gene expression in response to endogenous cytokinin reveals targets in *Arabidopsis thaliana*. *FEBS Lett.* 554, 373–380.
- Hou, B., Lim, E. K., Higgins, G. S., and Bowles, D. J. (2004). N-glucosylation of cytokinins by glycosyltransferases of *Arabidopsis thaliana*. *J. Biol. Chem.* 279, 47822–47832.
- Hruz, T., Laule, O., Szabo, G., Wessendorp, F., Bleuler, S., Oertle, L., Widmayer, P., Gruissem, W., and Zimmermann, P. (2008). Genevestigator V3: a reference expression database for the meta-analysis of transcriptomes. *Adv. Bioinformatics* 2008, Article ID 420747.
- Hunt, P. W., Watts, R. A., Trevasakis, B., Llewellyn, D. J., Burnell, J., Dennis, E. S., and Peacock, W. J. (2001). Expression and evolution of functionally distinct haemoglobin genes in plants. *Plant Mol. Biol.* 47, 677–692.
- Hwang, I., and Sheen, J. (2001). Two-component circuitry in *Arabidopsis* cytokinin signal transduction. *Nature* 413, 383–389.
- Igarashi, D., Izumi, Y., Dokiya, Y., Totsumi, K., Fukusaki, E., and Ohsumi, C. (2009). Reproductive organs regulate leaf nitrogen metabolism mediated by cytokinin signal. *Planta* 229, 633–644.
- Imamura, A., Hanaki, N., Nakamura, A., Suzuki, T., Taniguchi, M., Kiba, T., Ueguchi, C., Sugiyama, T., and Mizuno, T. (1999). Compilation and characterization of *Arabidopsis thaliana* response regulators implicated in His-to-Asp phosphorelay signal transduction. *Plant Cell Physiol.* 40, 733–742.
- Imamura, A., Hanaki, N., Umeda, H., Nakamura, A., Suzuki, T., Ueguchi, C., and Mizuno, T. (1998). Response regulators implicated in His-to-Asp phosphotransfer signaling in *Arabidopsis*. *Proc. Natl. Acad. Sci. U.S.A.* 95, 2691–2696.
- Imamura, A., Kiba, T., Tajima, Y., Yamashino, T., and Mizuno, T. (2003). In vivo and in vitro characterization of the ARR1 response regulator implicated in the His-to-Asp phosphorelay signal transduction in *Arabidopsis thaliana*. *Plant Cell Physiol.* 44, 122–131.
- Inoue, S.-I., Kinoshita, T., Takemiya, A., Doi, M., and Shimazaki, K.-I. (2008). Leaf positioning of *Arabidopsis* in response to blue light. *Mol. Plant* 1, 15–26.
- Inoue, T., Higuchi, M., Hashimoto, Y., Seki, M., Kobayashi, M., Kato, T., Tabata, S., Shinozaki, K., and Kakimoto, T. (2001). Identification of CRE1 as a cytokinin receptor from *Arabidopsis*. *Nature* 409, 1060–1063.
- Irizarry, R. A., Gautier, L., Bolstad, B. M., Miller, C., Astrand, M., Cope, L. M., Gentleman, R., Gentry, J., Halling, C., Huber, W., MacDonald, J., Rubin-stein, B. I. P., Workman, C., and Zhang, J. (2005). *affy: Methods for Affymetrix Oligonucleotide Arrays*. R Package Version 1.8.1.
- Ishida, K., Yamashino, T., Yokoyama, A., and Mizuno, T. (2008). Three type-B response regulators, ARR1, ARR10 and ARR12, play essential but redundant roles in cytokinin signal transduction throughout the life cycle of *Arabidopsis thaliana*. *Plant Cell Physiol.* 49, 47–57.
- Jain, M., Tyagi, A. K., and Khurana, J. P. (2006). Genome-wide analysis, evolutionary expansion, and expression of early auxin-responsive SAUR gene family in rice (*Oryza sativa*). *Genomics* 88, 360–371.
- Jung, K. W., Oh, S. I., Kim, Y. Y., Yoo, K. S., Cui, M. H., and Shin, J. S. (2008). *Arabidopsis* histidine-containing phosphotransfer factor 4 (AHP4) negatively regulates secondary wall thickening of the anther endothecium during flowering. *Mol. Cells* 25, 294–300.
- Kamiya, T., and Fujiwara, T. (2009). *Arabidopsis* NIP1;1 transports antimone and determines antimone sensitivity. *Plant Cell Physiol.* 50, 1977–1981.
- Kamiya, T., Tanaka, M., Mitani, N., Ma, J. F., Maeshima, M., and Fujiwara, T. (2009). NIP1;1, an aquaporin homolog, determines the arsenite sensitivity of *Arabidopsis thaliana*. *J. Biol. Chem.* 284, 2114–2120.
- Kant, S., Bi, Y.-M., Zhu, T., and Rothstein, S. J. (2009). SAUR39, a small auxin-up RNA gene, acts as a negative regulator of auxin synthesis and transport in rice. *Plant Physiol.* 151, 691–701.
- Kant, S., and Rothstein, S. (2009). Auxin-responsive SAUR39 gene modulates auxin level in rice. *Plant Signal. Behav.* 4, 1174–1175.
- Kasukabe, Y., He, L., Nada, K., Misawa, S., Ihara, I., and Tachibana, S. (2004). Overexpression of spermidine synthase enhances tolerance to multiple environmental stresses and up-regulates the expression of various stress-regulated genes in transgenic *Arabidopsis thaliana*. *Plant Cell Physiol.* 45, 712–722.
- Kepinski, S., and Leyser, O. (2005). The *Arabidopsis* F-box protein TIR1 is an auxin receptor. *Nature* 435, 446–451.
- Kiba, T., Aoki, K., Sakakibara, H., and Mizuno, T. (2004). *Arabidopsis* response regulator, ARR22, ectopic expression of which results in phenotypes similar to the wol cytokinin-receptor mutant. *Plant Cell Physiol.* 45, 1063–1077.
- Kiba, T., Kudo, T., Kojima, M., and Sakakibara, H. (2011). Hormonal control of nitrogen acquisition: roles of auxin, abscisic acid, and cytokinin. *J. Exp. Bot.* 62, 1399–1409.
- Kiba, T., Naitou, T., Koizumi, N., Yamashino, T., Sakakibara, H., and Mizuno, T. (2005). Combinatorial microarray analysis revealing *Arabidopsis* genes implicated in cytokinin responses through the His²Asp phosphorelay circuitry. *Plant Cell Physiol.* 46, 339–355.
- Kiba, T., Taniguchi, M., Imamura, A., Ueguchi, C., Mizuno, T., and Sugiyama, T. (1999). Differential expression of genes for response regulators in response to cytokinins and nitrate in *Arabidopsis thaliana*. *Plant Cell Physiol.* 40, 767–771.
- Kim, H. J., Ryu, H., Hong, S. H., Woo, H. R., Lim, P. O., Lee, I. C., Sheen, J., Nam, H. G., and Hwang, I. (2006). Cytokinin-mediated control of leaf longevity by AHK3 through phosphorylation of ARR2 in *Arabidopsis*. *Proc. Natl. Acad. Sci. U.S.A.* 103, 814–819.
- Köllmer, I., Werner, T., and Schmölling, T. (2011). Ectopic expression of different cytokinin-regulated transcription factor genes of *Arabidopsis thaliana* alters plant growth and development. *J. Plant Physiol.* 168, 1320–1327.
- Kreps, J. A., Wu, Y., Chang, H.-S., Zhu, T., Wang, X., and Harper, J. F. (2002). Transcriptome changes for *Arabidopsis* in response to salt, osmotic, and cold stress. *Plant Physiol.* 130, 2129–2141.
- Kudo, T., Kiba, T., and Sakakibara, H. (2010). Metabolism and long-distance translocation of cytokinins. *J. Integr. Plant Biol.* 52, 53–60.
- Kuroda, H., Masuda, T., Fusada, N., Ohta, H., and Takamiya, K. (2000). Expression of NADPH-protochlorophyllide oxidoreductase gene in fully green leaves of cucumber. *Plant Cell Physiol.* 41, 226–229.
- Laplaze, L., Benkova, E., Casimiro, I., Maes, L., Vanneste, S., Swarup, R., Weijers, D., Calvo, V., Parizot, B., Herrera-Rodriguez, M. B., Offringa, R., Graham, N., Doumas, P., Friml, J., Bogusz, D., Beeckman, T., and Bennett, M. (2007). Cytokinins act directly on lateral root founder cells to inhibit root initiation. *Plant Cell* 19, 3889–3900.
- Lariguet, P., Boccacandro, H. E., Alonso, J. M., Ecker, J. R., Chory, J., Casal, J. J., and Fankhauser, C. (2003). A growth regulatory loop that provides homeostasis to phytochrome A signaling. *Plant Cell* 15, 2966–2978.
- Lariguet, P., Schepens, I., Hodgson, D., Pedmale, U. V., Trevisan, M., Kami, C., de Carbonnel, M., Alonso, J. M., Ecker, J. R., Lisicum, E., and Fankhauser, C. (2006). PHYTOCHROME KINASE SUBSTRATE 1 is a phototropin 1 binding protein required for phototropism. *Proc. Natl. Acad. Sci. U.S.A.* 103, 10134–10139.
- Lee, D. J., Park, J. Y., Ku, S. J., Ha, Y. M., Kim, S., Kim, M. D., Oh, M. H., and Kim, J. (2007). Genome-wide expression profiling of *ARABIDOPSIS* RESPONSE REGULATOR 7 (ARR7) overexpression in cytokinin response. *Mol. Genet. Genomics* 277, 115–137.
- Lee, J., Parthier, B., and Löbler, M. (1996). Jasmonate signalling can be uncoupled from abscisic acid signalling in barley: identification of jasmonate-regulated transcripts which are not induced by abscisic acid. *Planta* 199, 625–632.
- Lee, S., Lee, E. J., Yang, E. J., Lee, J. E., Park, A. R., Song, W. H., and Park, O. K. (2004). Proteomic identification of annexins, calcium-dependent membrane binding proteins that mediate osmotic stress and abscisic acid signal transduction in *Arabidopsis*. *Plant Cell* 16, 1378–1391.
- Leyser, H. M. O., Pickett, F. B., Dharmasiri, S., and Estelle, M. (1996). Mutations in the AXR3 gene of *Arabidopsis* result in altered auxin response including ectopic expression from the SAUR-AC1 promoter. *Plant J.* 10, 403–413.
- Lima, J. E., Kojima, S., Takahashi, H., and von Wirén, N. (2010). Ammonium triggers lateral root branching in *Arabidopsis* in an

- AMMONIUM TRANSPORTER1;3-dependent manner. *Plant Cell* 22, 3621–3633.
- Lin, P.-C., Pomeranz, M. C., Jikumaru, Y., Kang, S. G., Hah, C., Fujioka, S., Kamiya, Y., and Jang, J.-C. (2011). The *Arabidopsis* tandem zinc finger protein AtTZF1 affects ABA- and GA-mediated growth, stress and gene expression responses. *Plant J.* 65, 253–268.
- Lindsay, D. L., Sawhney, V. K., and Bonham-Smith, P. C. (2006). Cytokinin-induced changes in CLAVATA1 and WUSCHEL expression temporally coincide with altered floral development in *Arabidopsis*. *Plant Sci.* 170, 1111–1117.
- Little, D., Gouhier-Darimont, C., Bruesow, F., and Reymond, P. (2007). Oviposition by pierid butterflies triggers defense responses in *Arabidopsis*. *Plant Physiol.* 143, 784–800.
- Lohrmann, J., Sweere, U., Zabaleta, E., Bäurle, I., Keitel, C., Kozma, B. L., Brennicke, A., Schäfer, E., Kudla, J., and Harter, K. (2001). The response regulator ARR2: a pollen-specific transcription factor involved in the expression of nuclear genes for components of mitochondrial complex I in *Arabidopsis*. *Mol. Genet. Genomics* 265, 2–13.
- Ma, J., Hanssen, M., Lundgren, K., Hernández, L., Delatte, T., Ehlert, A., Liu, C.-M., Schluempmann, H., Dröge-Laser, W., Moritz, T., Smeekens, S., and Hanson, J. (2011). The sucrose-regulated *Arabidopsis* transcription factor bZIP11 reprograms metabolism and regulates trehalose metabolism. *New Phytol.* 191, 733–745.
- MacRae, W. D., and Towers, G. H. N. (1984). Biological activities of lignans. *Phytochemistry* 23, 1207–1220.
- Mähönen, A. P., Bishopp, A., Higuchi, M., Nieminen, K. M., Kinoshita, K., Tormakangas, K., Ikeda, Y., Oka, A., Kakimoto, T., and Helariutta, Y. (2006). Cytokinin signaling and its inhibitor AHP6 regulate cell fate during vascular development. *Science* 311, 94–98.
- Maksimov, I. V., Ganiev, R. M., and Khairullin, R. M. (2002). Changes in the levels of IAA, ABA, and cytokinins in wheat seedlings infected with *Tilletia caries*. *Russ. J. Plant Physiol.* 49, 221–224.
- Maruyama-Nakashita, A., Nakamura, Y., Yamaya, T., and Takahashi, H. (2004). A novel regulatory pathway of sulfate uptake in *Arabidopsis* roots: implication of CRE1/WOL/AHK4-mediated cytokinin-dependent regulation. *Plant J.* 38, 779–789.
- Marziani Longo, G. P., Bracale, M., Rossi, G., and Longo, C. P. (1990). Benzyladenine induces the appearance of LHCP-mRNA and of the relevant protein in dark-grown excised watermelon cotyledons. *Plant Mol. Biol.* 14, 569–573.
- Mason, M. G., Li, J., Mathews, D. E., Kieber, J. J., and Schaller, G. E. (2004). Type-B response regulators display overlapping expression patterns in *Arabidopsis*. *Plant Physiol.* 135, 927–937.
- Mason, M. G., Mathews, D. E., Argiros, D. A., Maxwell, B. B., Kieber, J. J., Alonso, J. M., Ecker, J. R., and Schaller, G. E. (2005). Multiple type-B response regulators mediate cytokinin signal transduction in *Arabidopsis*. *Plant Cell* 17, 3007–3018.
- Massonnet, C., Vile, D., Fabre, J., Hannah, M. A., Caldana, C., Lisee, J., Beemster, G. T. S., Meyer, R. C., Messerli, G., Gronlund, J. T., Perkovic, J., Wigmore, E., May, S., Bevan, M. W., Meyer, C., Rubio-Diaz, S., Weigel, D., Micol, J. L., Buchanan-Wollaston, V., Fiorani, F., Walsh, S., Rinn, B., Gruissem, W., Hilsen, P., Hennig, L., Willmitzer, L., and Granier, C. (2010). Probing the reproducibility of leaf growth and molecular phenotypes: a comparison of three *Arabidopsis* accessions cultivated in ten laboratories. *Plant Physiol.* 152, 2142–2157.
- Mendonça Pauletti, P., Araújo, A. R., Young, M. C. M., Giesbrecht, A. M., and da Silva Bolzani, V. (2000). nor-Lignans from the leaves of *Styrax ferrugineus* (Styracaceae) with antibacterial and antifungal activity. *Phytochemistry* 55, 597–601.
- Michel, K., Abderhalden, O., Bruggmann, R. M., and Dudler, R. (2006). Transcriptional changes in powdery mildew infected wheat and *Arabidopsis* leaves undergoing syringolin-triggered hypersensitive cell death at infection sites. *Plant Mol. Biol.* 62, 561–578.
- Motchoulski, A., and Liscum, E. (1999). *Arabidopsis* NPH3: a NPH1 photoreceptor-interacting protein essential for phototropism. *Science* 286, 961–964.
- Moubayidin, L., Di Mambro, R., and Sabatini, S. (2009). Cytokinin-auxin crosstalk. *Trends Plant Sci.* 14, 557–562.
- Mudge, S. R., Rae, A. L., Diatloff, E., and Smith, F. W. (2002). Expression analysis suggests novel roles for members of the Pht1 family of phosphate transporters in *Arabidopsis*. *Plant J.* 31, 341–353.
- Müller, B. (2011). Generic signal-specific responses: cytokinin and context-dependent cellular responses. *J. Exp. Bot.* 62, 3273–3288.
- Müller, B., and Sheen, J. (2008). Cytokinin and auxin interaction in root stem-cell specification during early embryogenesis. *Nature* 453, 1094–1097.
- Naito, T., Yamashino, T., Kiba, T., Koizumi, N., Kojima, M., Sakakibara, H., and Mizuno, T. (2007). A link between cytokinin and ASL9 (ASYMMETRIC LEAVES 2 LIKE 9) that belongs to the AS2/LOB (LATERAL ORGAN BOUNDARIES) family genes in *Arabidopsis thaliana*. *Biosci. Biotechnol. Biochem.* 71, 1269–1278.
- Navarro, L., Zipfel, C., Rowland, O., Keller, I., Robatzek, S., Boller, T., and Jones, J. D. G. (2004). The transcriptional innate immune response to flg22. Interplay and overlap with Avr gene-dependent defense responses and bacterial pathogenesis. *Plant Physiol.* 135, 1113–1128.
- Nelson, D. C., Flematti, G. R., Riseborough, J.-A., Ghisalberti, E. L., Dixon, K. W., and Smith, S. M. (2010). Karrikins enhance light responses during germination and seedling development in *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci. U.S.A.* 107, 7095–7100.
- Nutricati, E., Miceli, A., Blando, F., and De Bellis, L. (2006). Characterization of two *Arabidopsis thaliana* glutathione S-transferases. *Plant Cell Rep.* 25, 997–1005.
- Obayashi, T., Hayashi, S., Saeki, M., Ohta, H., and Kinoshita, K. (2009). ATTED-II provides coexpressed gene networks for *Arabidopsis*. *Nucleic Acids Res.* 37, D987–D991.
- Obayashi, T., Nishida, K., Kasahara, K., and Kinoshita, K. (2011). ATTED-II updates: condition-specific gene coexpression to extend coexpression analyses and applications to a broad range of flowering plants. *Plant Cell Physiol.* 52, 213–219.
- Okazaki, K., Kabeya, Y., Suzuki, K., Mori, T., Ichikawa, T., Matsui, M., Nakanishi, H., and Miyagishima, S.-Y. (2009). The PLASTID DIVISION1 and 2 components of the chloroplast division machinery determine the rate of chloroplast division in land plant cell differentiation. *Plant Cell* 21, 1769–1780.
- Parthier, B. (1979). The role of phytohormones (cytokinin) in chloroplasts development. *Biochem. Physiol. Pflanz.* 174, 173–214.
- Pedmale, U. V., and Liscum, E. (2007). Regulation of phototropic signaling in *Arabidopsis* via phosphorylation state changes in the phototropin 1-interacting protein NPH3. *J. Biol. Chem.* 282, 19992–20001.
- Pérez-Pérez, J. M., Candela, H., Robles, P., López-Torrejón, G., del Pozo, J. C., and Micol, J. L. (2010). A role for AUXIN RESISTANT3 in the coordination of leaf growth. *Plant Cell Physiol.* 51, 1661–1673.
- Perilli, S., Moubayidin, L., and Sabatini, S. (2010). The molecular basis of cytokinin function. *Curr. Opin. Plant Biol.* 13, 21–26.
- Pertry, I., Václavíková, K., Depuydt, S., Galuszka, P., Spíchal, L., Temmerman, W., Stes, E., Schmülling, T., Kakimoto, T., Van Montagu, M. C. E., Strnad, M., Holsters, M., Tarkowski, P., and Vereecke, D. (2009). Identification of *Rhodococcus fascians* cytokinins and their modus operandi to reshape the plant. *Proc. Natl. Acad. Sci. U.S.A.* 106, 929–934.
- Pils, B., and Heyl, A. (2009). Unraveling the evolution of cytokinin signaling. *Plant Physiol.* 151, 782–791.
- Pomeranz, M., Zhang, L., Finer, J., and Jang, J. C. (2011). Can AtTZF1 act as a transcriptional activator or repressor in plants? *Plant Signal. Behav.* 6, 719–722.
- Pufky, J., Qiu, Y., Rao, M., Hurban, P., and Jones, A. (2003). The auxin-induced transcriptome for etiolated *Arabidopsis* seedlings using a structure/function approach. *Funct. Integr. Genomics* 3, 135–143.
- Punwani, J. A., Hutchison, C. E., Schaller, G. E., and Kieber, J. J. (2010). The subcellular distribution of the *Arabidopsis* histidine phosphotransfer proteins is independent of cytokinin signaling. *Plant J.* 62, 473–482.
- Rahman, M. M., and Gray, A. I. (2002). Antimicrobial constituents from the stem bark of *Feronia limonia*. *Phytochemistry* 59, 73–77.
- Ralph, S., Park, J.-Y., Bohlmann, J. R., and Mansfield, S. (2006). Dirigent proteins in conifer defense: gene discovery, phylogeny, and differential wound- and insect-induced expression of a family of DIR and DIR-like genes in spruce (*Picea* spp.). *Plant Mol. Biol.* 60, 21–40.
- Ramirezzy, E. (2009). *Functional Characterization of B-Type Response Regulators of Arabidopsis thaliana*. Dissertation, Freie Universität Berlin, Berlin.
- Rashotte, A., and Goertzen, L. (2010). The CRF domain defines cytokinin response factor proteins in

- plants. *BMC Plant Biol.* 10, 74. doi:10.1186/1471-2229-10-74
- Rashotte, A. M., Carson, S. D., To, J. P., and Kieber, J. J. (2003). Expression profiling of cytokinin action in *Arabidopsis*. *Plant Physiol.* 132, 1998–2011.
- Rashotte, A. M., Mason, M. G., Hutchison, C. E., Ferreira, F. J., Schaller, G. E., and Kieber, J. J. (2006). A subset of *Arabidopsis* AP2 transcription factors mediates cytokinin responses in concert with a two-component pathway. *Proc. Natl. Acad. Sci. U.S.A.* 103, 11081–11085.
- Ribot, C. C., Wang, Y., and Poirier, Y. (2008). Expression analyses of three members of the AtPHO1 family reveal differential interactions between signaling pathways involved in phosphate deficiency and the responses to auxin, cytokinin, and abscisic acid. *Planta* 227, 1025–1036.
- Richter, R., Behringer, C., Müller, I. K., and Schwechheimer, C. (2010). The GATA-type transcription factors GNC and GNL/CGA1 repress gibberellin signaling downstream from DELLA proteins and PHYTOCHROME-INTERACTING FACTORS. *Genes Dev.* 24, 2093–2104.
- Riefler, M., Novak, O., Strnad, M., and Schumlling, T. (2006). *Arabidopsis* cytokinin receptor mutants reveal functions in shoot growth, leaf senescence, seed size, germination, root development, and cytokinin metabolism. *Plant Cell* 18, 40–54.
- Rivero, R. M., Shulaev, V., and Blumwald, E. (2009). Cytokinin-dependent photorespiration and the protection of photosynthesis during water deficit. *Plant Physiol.* 150, 1530–1540.
- Rocca-Serra, P., Brazma, A., Parkinson, H., Sarkans, U., Shojatalab, M., Contrino, S., Vilo, J., Abeygunawardena, N., Mukherjee, G., Holloway, E., Kapushesky, M., Kemmeren, P., Lara, G. G., Oezcimen, A., and Sansone, S. A. (2003). ArrayExpress: a public database of gene expression data at EBI. *C. R. Biol.* 326, 1075–1078.
- Ross, E. J. H., Stone, J. M., Elowsky, C. G., Arredondo-Peter, R., Klucas, R. V., and Sarath, G. (2004). Activation of the *Oryza sativa* non-symbiotic haemoglobin-2 promoter by the cytokinin-regulated transcription factor, ARR1. *J. Exp. Bot.* 55, 1721–1731.
- Rubio, V., Bustos, R., Irigoyen, M. A., Cardona-López, X., Rojas-Triana, M. N., and Paz-Ares, J. (2009). Plant hormones and nutrient signaling. *Plant Mol. Biol.* 69, 361–373.
- Růžicka, K., Šimášková, M., Duclercq, J., Petrášek, J., Zajímalová, E., Simon, S., Friml, J., Van Montagu, M. C. E., and Benková, E. (2009). Cytokinin regulates root meristem activity via modulation of the polar auxin transport. *Proc. Natl. Acad. Sci. U.S.A.* 106, 4284–4289.
- Sakai, H., Aoyama, T., and Oka, A. (2000). *Arabidopsis* ARR1 and ARR2 response regulators operate as transcriptional activators. *Plant J.* 24, 703–711.
- Sakai, H., Honma, T., Aoyama, T., Sato, S., Kato, T., Tabata, S., and Oka, A. (2001). ARR1, a transcription factor for genes immediately responsive to cytokinins. *Science* 294, 1519–1521.
- Sakakibara, H., Suzuki, M., Takei, K., Deji, A., Taniguchi, M., and Sugiyama, T. (1998). A response-regulator homologue possibly involved in nitrogen signal transduction mediated by cytokinin in maize. *Plant J.* 14, 337–344.
- Sakakibara, H., Takei, K., and Hirose, N. (2006). Interactions between nitrogen and cytokinin in the regulation of metabolism and development. *Trends Plant Sci.* 11, 440–448.
- Schaller, G. E., Shiu, S.-H., and Armitage, J. P. (2011). Two-component systems and their co-option for eukaryotic signal transduction. *Curr. Biol.* 21, R320–R330.
- Schepens, I., Duek, P., and Fankhauser, C. (2004). Phytochrome-mediated light signalling in *Arabidopsis*. *Curr. Opin. Plant Biol.* 7, 564–569.
- Schluepmann, H., Pellny, T., Van, D. A., Smeekens, S., and Paul, M. (2003). Trehalose 6-phosphate is indispensable for carbohydrate utilization and growth in *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci. U.S.A.* 100, 6849–6854.
- Schluepmann, H., van Dijken, A., Aghdasi, M., Wobbles, B., Paul, M., and Smeekens, S. (2004). Trehalose mediated growth inhibition of *Arabidopsis* seedlings is due to trehalose-6-phosphate accumulation. *Plant Physiol.* 135, 879–890.
- Schumlling, T., Schäfer, S., and Romanov, G. (1997). Cytokinins as regulators of gene expression. *Physiol. Plant* 100, 505–519.
- Sherson, S. M., Alford, H. L., Forbes, S. M., Wallace, G., and Smith, S. M. (2003). Roles of cell-wall invertases and monosaccharide transporters in the growth and development of *Arabidopsis*. *J. Exp. Bot.* 54, 525–531.
- Šíffel, P., Sindelková, E., Durchan, M., and Zajícová, M. (1992). Photosynthetic characteristics of *Solanum tuberosum* L. plants transformed by genes for synthesis of phytohormones. *Photosynthetica* 27, 441–447.
- Skinner, D., and Gasser, C. (2009). Expression-based discovery of candidate ovule development regulators through transcriptional profiling of ovule mutants. *BMC Plant Biol.* 9, 29. doi:10.1186/1471-2229-9-29
- Smyth, G. K. (2005). “Limma: linear models for microarray data,” in *Bioinformatics and Computational Biology Solutions Using R and Bioconductor*, eds R. Gentleman, V. Carey, S. Dudoit, R. Irizarry, and W. Huber (New York: Springer), 397–420.
- Staetler, D. A., and Laetsch, W. M. (1965). Kinetin-induced chloroplast maturation in cultures of tobacco tissue. *Science* 149, 1387–1388.
- Suzuki, T., Miwa, K., Ishikawa, K., Yamada, H., Aiba, H., and Mizuno, T. (2001). The *Arabidopsis* sensor His-kinase, AHK4, can respond to cytokinins. *Plant Cell Physiol.* 42, 107–113.
- Swarbreck, D., Wilks, C., Lamesch, P., Berardini, T. Z., Garcia-Hernandez, M., Foerster, H., Li, D., Meyer, T., Muller, R., Ploetz, L., Radenbaugh, A., Singh, S., Swing, V., Tissier, C., Zhang, P., and Huala, E. (2008). The *Arabidopsis* Information Resource (TAIR): gene structure and function annotation. *Nucleic Acids Res.* 36, D1009–D1014.
- Tajima, Y., Imamura, A., Kiba, T., Amano, Y., Yamashino, T., and Mizuno, T. (2004). Comparative studies on the type-B response regulators revealing their distinctive properties in the His-to-Asp phosphorelay signal transduction of *Arabidopsis thaliana*. *Plant Cell Physiol.* 45, 28–39.
- Takei, K., Yamaya, T., and Sakakibara, H. (2004). *Arabidopsis* CYP735A1 and CYP735A2 encode cytokinin hydroxylases that catalyze the biosynthesis of trans-zeatin. *J. Biol. Chem.* 279, 41866–41872.
- Taniguchi, M., Sasaki, N., Tsuge, T., Aoyama, T., and Oka, A. (2007). ARR1 directly activates cytokinin response genes that encode proteins with diverse regulatory functions. *Plant Cell Physiol.* 48, 263–277.
- Thimm, O., Blaessing, O., Gibon, Y., Nagel, A., Meyer, S., Krueger, P., Selbig, J., Mueller, L. A., Rhee, S. Y., and Stitt, M. (2004). MAPMAN: A user-driven tool to display genomics data sets onto diagrams of metabolic pathways and other biological processes. *Plant J.* 37, 914–939.
- Thomas, J., Ross, C. W., Chastain, C. J., Koomanoff, N., Hendrix, J. E., and Van Volkenburgh, E. (1981). Cytokinin-induced wall extensibility in excised cotyledons of radish and cucumber. *Plant Physiol.* 68, 107–110.
- Tiwari, S., Spielman, M., Day, R. C., and Scott, R. J. (2006). Proliferative phase endosperm promoters from *Arabidopsis thaliana*. *Plant Biotechnol. J.* 4, 393–407.
- To, J. P., Haberer, G., Ferreira, F. J., Deruere, J., Mason, M. G., Schaller, G. E., Alonso, J. M., Ecker, J. R., and Kieber, J. J. (2004). Type-A *Arabidopsis* response regulators are partially redundant negative regulators of cytokinin signaling. *Plant Cell* 16, 658–671.
- To, J. P. C., and Kieber, J. J. (2008). Cytokinin signaling: two-components and more. *Trends Plant Sci.* 13, 85–92.
- Tran, L.-S. P., Urao, T., Qin, F., Maruyama, K., Kakimoto, T., Shinozaki, K., and Yamaguchi-Shinozaki, K. (2007). Functional analysis of AHK1/ATHK1 and cytokinin receptor histidine kinases in response to abscisic acid, drought, and salt stress in *Arabidopsis*. *Proc. Natl. Acad. Sci. U.S.A.* 104, 20623–20628.
- Treml, B. S., Winderl, S., Radykewicz, R., Herz, M., Schweizer, G. N., Hutzler, P., Glawischnig, E., and Ruiz, R. N. A. T. (2005). The gene ENHANCER OF PINOID controls cotyledon development in the *Arabidopsis* embryo. *Development* 132, 4063–4074.
- Urao, T., Yakubov, B., Satoh, R., Yamaguchi-Shinozaki, K., Seki, M., Hirayama, T., and Shinozaki, K. (1999). A transmembrane hybrid-type histidine kinase in *Arabidopsis* functions as an osmosensor. *Plant Cell* 11, 1743–1754.
- Urao, T., Yakubov, B., Yamaguchi-Shinozaki, K., and Shinozaki, K. (1998). Stress-responsive expression of genes for two-component response regulator-like proteins in *Arabidopsis thaliana*. *FEBS Lett.* 427, 175–178.
- Vergnolle, C., Vaultier, M.-N. I., Tacconat, L., Renou, J.-P., Kader, J.-C., Zachowski, A., and Ruelland, E. (2005). The cold-induced early activation of phospholipase C and D pathways determines the response of two distinct clusters of genes in *Arabidopsis* cell suspensions. *Plant Physiol.* 139, 1217–1233.
- Vieira Dos Santos, C., and Rey, P. (2006). Plant thioredoxins are key actors in the oxidative stress response. *Trends Plant Sci.* 11, 329–334.
- Wahl, V., Brand, L. H., Guo, Y. L., and Schmid, M. (2010). The FAN-TASTIC FOUR proteins influence

- shoot meristem size in *Arabidopsis thaliana*. *BMC Plant Biol.* 10, 285. doi:10.1186/1471-2229-10-285
- Weijers, D., Franke-van Dijk, M., Vencken, R.-J., Quint, A., Hooykaas, P., and Offringa, R. (2001). An *Arabidopsis* minute-like phenotype caused by a semi-dominant mutation in a RIBOSOMAL PROTEIN S5 gene. *Development* 128, 4289–4299.
- Werner, T., Holst, K., Pors, Y., Guivarc'h, A., Mustroph, A., Chriqui, D., Grimm, B., and Schumlling, T. (2008). Cytokinin deficiency causes distinct changes of sink and source parameters in tobacco shoots and roots. *J. Exp. Bot.* 59, 2659–2672.
- Werner, T., Motyka, V., Laucou, V., Smets, R., van Onckelen, H., and Schumlling, T. (2003). Cytokinin-deficient transgenic *Arabidopsis* plants show multiple developmental alterations indicating opposite functions of cytokinins in the regulation of shoot and root meristem activity. *Plant Cell* 15, 2532–2550.
- Werner, T., Motyka, V., Strnad, M., and Schumlling, T. (2001). Regulation of plant growth by cytokinin. *Proc. Natl. Acad. Sci. U.S.A.* 98, 10487–10492.
- Werner, T., Nehnevajova, E., Köllmer, I., Novak, O., Strnad, M., Krämer, U., and Schumlling, T. (2010). Root-specific reduction of cytokinin causes enhanced root growth, drought tolerance, and leaf mineral enrichment in *Arabidopsis* and tobacco. *Plant Cell* 22, 3905–3920.
- Werner, T., and Schumlling, T. (2009). Cytokinin action in plant development. *Curr. Opin. Plant Biol.* 12, 527–538.
- Wohlbach, D. J., Quirino, B. F., and Sussman, M. R. (2008). Analysis of the *Arabidopsis* histidine kinase ATHK1 reveals a connection between vegetative osmotic stress sensing and seed maturation. *Plant Cell* 20, 1101–1117.
- Yaronskaya, E., Vershilovskaya, I., Poers, Y., Alawady, A., Averina, N., and Grimm, B. (2006). Cytokinin effects on tetrapyrrole biosynthesis and photosynthetic activity in barley seedlings. *Planta* 224, 700–709.
- Yokoyama, A., Yamashino, T., Amano, Y., Tajima, Y., Imamura, A., Sakakibara, H., and Mizuno, T. (2007). Type-B ARR transcription factors, ARR10 and ARR12, are implicated in cytokinin-mediated regulation of protoxylem differentiation in roots of *Arabidopsis thaliana*. *Plant Cell Physiol.* 48, 84–96.
- Yoshida, S., Mandel, T., and Kuhlemeier, C. (2011). Stem cell activation by light guides plant organogenesis. *Genes Dev.* 25, 1439–1450.
- Zalewski, W., Galuszka, P., Gasparis, S., Orczyk, W., and Nadolska-Orczyk, A. (2010). Silencing of the HvCKX1 gene decreases the cytokinin oxidase/dehydrogenase level in barley and leads to higher plant productivity. *J. Exp. Bot.* 61, 1839–1851.
- Zavaleta-Mancera, H. A., López-Delgado, H., Loza-Tavera, H., Mora-Herrera, M., Trevilla-García, C., Vargas-Suárez, M., and Ougham, H. (2007). Cytokinin promotes catalase and ascorbate peroxidase activities and preserves the chloroplast integrity during dark-senescence. *J. Plant Physiol.* 164, 1572–1582.
- Zhao, J., Wang, J., An, L., Doerge, R., Chen, Z., Grau, C., Meng, J., and Osborn, T. (2007). Analysis of gene expression profiles in response to *Sclerotinia sclerotiorum* in *Brassica napus*. *Planta* 227, 13–24.
- Zhao, Z., Andersen, S. U., Ljung, K., Doležal, K., Miotk, A., Schultheiss, S. J., and Lohmann, J. U. (2010). Hormonal control of the shoot stem-cell niche. *Nature* 465, 1089–1092.
- Zubo, Y. O., Yamburenko, M. V., Selivankina, S. Y., Shakirova, F. M., Avalbaev, A. M., Kudryakova, N. V., Zubkova, N. K., Liere, K., Kulaeva, O. N., Kusnetsov, V. V., and Börner, T. (2008). Cytokinin stimulates chloroplast transcription in detached barley leaves. *Plant Physiol.* 148, 1082–1093.
- Catský, J., Pospíšilová, J., Machácková, I., Wilhelmová, N., and Šesták, Z. (1993). Photosynthesis and water relations in transgenic tobacco plants with T-DNA carrying gene 4 for cytokinin synthesis. *Biol. Plant.* 35, 393–399.
- Cerný, M., Dycka, F., Bobál'ová, J., and Brzobohatý, B. (2011). Early cytokinin response proteins and phosphoproteins of *Arabidopsis thaliana* identified by proteome and phosphoproteome profiling. *J. Exp. Bot.* 62, 921–937.

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 30 September 2011; paper pending published: 21 October 2011; accepted: 06 January 2012; published online: 31 January 2012.

Citation: Brenner WG, Ramireddy E, Heyl A and Schumlling T (2012) Gene regulation by cytokinin in *Arabidopsis*. *Front. Plant Sci.* 3:8. doi: 10.3389/fpls.2012.00008

This article was submitted to *Frontiers in Plant Physiology*, a specialty of *Frontiers in Plant Science*.

Copyright © 2012 Brenner, Ramireddy, Heyl and Schumlling. This is an open-access article distributed under the terms of the Creative Commons Attribution Non Commercial License, which permits non-commercial use, distribution, and reproduction in other forums, provided the original authors and source are credited.