



# The effects of inorganic nitrogen form and CO<sub>2</sub> concentration on wheat yield and nutrient accumulation and distribution

Eli Carlisle<sup>1\*</sup>, Samuel Myers<sup>2</sup>, Victor Raboy<sup>3</sup> and Arnold Bloom<sup>1</sup>

<sup>1</sup> Department of Plant Sciences, University of California, Davis, CA, USA

<sup>2</sup> Harvard School of Public Health, Harvard University, Boston, MA, USA

<sup>3</sup> Agricultural Research Service, United States Department of Agriculture, Aberdeen, ID, USA

## Edited by:

Jan K. Schjoerring, University of Copenhagen, Denmark

## Reviewed by:

Roberto A. Gaxiola, Arizona State University, USA

Victoria Fernandez, Technical University of Madrid, Spain

## \*Correspondence:

Eli Carlisle, Department of Plant Sciences, University of California, One Shields Avenue, Davis, CA 95616, USA.  
e-mail: ecarlisle@ucdavis.edu

Inorganic N is available to plants from the soil as ammonium (NH<sub>4</sub><sup>+</sup>) and nitrate (NO<sub>3</sub><sup>-</sup>). We studied how wheat grown hydroponically to senescence in controlled environmental chambers is affected by N form (NH<sub>4</sub><sup>+</sup> vs. NO<sub>3</sub><sup>-</sup>) and CO<sub>2</sub> concentration (“subambient,” “ambient,” and “elevated”) in terms of biomass, yield, and nutrient accumulation and partitioning. Wheat supplied with NH<sub>4</sub><sup>+</sup> as a sole N source had the strongest response to CO<sub>2</sub> concentration. Plants exposed to subambient and ambient CO<sub>2</sub> concentrations typically had the greatest biomass and nutrient accumulation under both N forms. In general NH<sub>4</sub><sup>+</sup>-supplied plants had higher concentrations of total N, P, K, S, Ca, Zn, Fe, and Cu, while NO<sub>3</sub><sup>-</sup>-supplied plants had higher concentrations of Mg, B, Mn, and NO<sub>3</sub><sup>-</sup>-N. NH<sub>4</sub><sup>+</sup>-supplied plants contained amounts of phytate similar to NO<sub>3</sub><sup>-</sup>-supplied plants but had higher bioavailable Zn, which could have consequences for human health. NH<sub>4</sub><sup>+</sup>-supplied plants allocated more nutrients and biomass to aboveground tissues whereas NO<sub>3</sub><sup>-</sup>-supplied plants allocated more nutrients to the roots. The two inorganic nitrogen forms influenced plant growth and nutrient status so distinctly that they should be treated as separate nutrients. Moreover, plant growth and nutrient status varied in a non-linear manner with atmospheric CO<sub>2</sub> concentration.

**Keywords:** climate change, wheat, ammonium, nitrate, nutrients, grain, phytate, CO<sub>2</sub>

## INTRODUCTION

Nitrogen (N) is the mineral element that most often limits plant growth and primary productivity in natural and agricultural systems. Plants usually acquire N from the soil in the forms of ammonium (NH<sub>4</sub><sup>+</sup>) and nitrate (NO<sub>3</sub><sup>-</sup>), and management of these forms is vital to agriculture. Wheat can utilize either form alone (Wang and Below, 1992), but mixed N nutrition (e.g., NH<sub>4</sub>NO<sub>3</sub>) typically produces the best grain yields and quality in hydroponically grown (Gentry et al., 1989; Heberer and Below, 1989; Wang and Below, 1995) and field-grown plants (Bock, 1987; Camberato and Bock, 1990).

Ammonium and nitrate affect crops differently when either is supplied as the sole N source (Bloom, 1997). Ammonium requires less energy to assimilate into organic compounds (Bloom, 1997), but can prove toxic if it accumulates to high concentrations within plant tissues (Cox and Reisenauer, 1973; Goyal and Huffaker, 1984). Nitrate is generally the predominant form available in aerated, temperate agricultural soils (Haynes, 1986; Bloom, 1997), and may accumulate within plant tissues to high concentrations without toxicity (Goyal and Huffaker, 1984). In wheat, the N form supplied has been found to influence many physiological parameters profoundly including biomass (Wang and Below, 1995, 1996, 1998; Bloom et al., 2002), leaf area (Bloom et al., 2002), tillering (Chen et al., 1998), seed mass (Wang and Below, 1995), protein content (Wang and Below, 1995), and mineral nutrient

acquisition and distribution (Gashaw and Mugwira, 1981; Wang and Below, 1998), although such differences can vary among cultivars (Gashaw and Mugwira, 1981; Wang and Below, 1995).

The presence of NH<sub>4</sub><sup>+</sup>, as either a sole N source or in mixed N nutrition, increased organic N concentration in shoots, roots, and grain and decreased partitioning of dry matter to the roots in wheat (Wang and Below, 1995). Decreased cation uptake has been found in wheat under NH<sub>4</sub><sup>+</sup> nutrition (e.g., Gashaw and Mugwira, 1981; Wang and Below, 1998), although results varied among cultivars (Gashaw and Mugwira, 1981). For example, NH<sub>4</sub><sup>+</sup> nutrition decreased whole plant and shoot accumulations of K, Cu, Ca, Mg, Fe, Mn, and Zn (Wang and Below, 1998). Nutrient allocation to plant tissues also varied between N forms. NH<sub>4</sub><sup>+</sup>-fed plants distributed a smaller percentage of total P, K, Cu, and B to roots relative to NO<sub>3</sub><sup>-</sup>-fed plants (Wang and Below, 1995, 1998). Also, a greater percentage of reduced N was allocated to the shoots in NH<sub>4</sub><sup>+</sup>-fed plants (Wang and Below, 1995).

Elevated atmospheric concentrations of CO<sub>2</sub> alter growth and N dynamics of wheat and other C<sub>3</sub> plants. Under elevated CO<sub>2</sub>, wheat has lower protein and N concentrations (e.g., Thompson and Woodward, 1994; Bloom et al., 2002; Wu et al., 2004), and lower macro- and micronutrients concentrations (Manderscheid et al., 1995; Fangmeier et al., 1997, 1999; Wu et al., 2004; Högy and Fangmeier, 2008). Grain phytate concentrations are also thought to increase or remain the same under elevated CO<sub>2</sub>,

and in conjunction with decreased concentrations of micronutrients, bioavailable Zn and Fe are expected to decrease even further under elevated CO<sub>2</sub> (Loladze, 2002; Manoj-Kumar, 2011), as these micronutrients form indigestible complexes with phytate. By contrast, wheat yields (Fangmeier et al., 1996; Amthor, 2001; Högy and Fangmeier, 2008), harvest index (HI; Wu et al., 2004), whole plant biomass (Fangmeier et al., 1996; Högy and Fangmeier, 2008), shoot biomass (Fangmeier et al., 1996; Högy et al., 2009), and root biomass (Chaudhuri et al., 1990; Wechsung et al., 1995) typically increase under CO<sub>2</sub> enrichment. In addition, elevated CO<sub>2</sub> concentration can increase tillering (Weigel et al., 1994), nitrogen use efficiency (NUE, Fangmeier et al., 1997), and micro/macronutrient use efficiencies (Manderscheid et al., 1995). The influence of elevated CO<sub>2</sub> on many of these characteristics may vary among cultivars and research protocols (e.g., FACE vs. controlled environment chamber, greenhouse vs. field; Amthor, 2001; Högy and Fangmeier, 2008).

Wheat grown under CO<sub>2</sub> enrichment behaves differently under NO<sub>3</sub><sup>-</sup> and NH<sub>4</sub><sup>+</sup> nutrition. Exposure to elevated CO<sub>2</sub> inhibits NO<sub>3</sub><sup>-</sup> photoassimilation in wheat (Bloom et al., 1989, 2002, 2010; Cousins and Bloom, 2004) as well as in all other C<sub>3</sub> and C<sub>3</sub>-C<sub>4</sub> intermediate plants tested (Bloom et al., 2012). At elevated CO<sub>2</sub>, NH<sub>4</sub><sup>+</sup>-fed plants showed greater increases in leaf area and smaller decreases in shoot protein concentration than NO<sub>3</sub><sup>-</sup>-fed plants (Bloom et al., 2002), which could have consequences for human nutrition. Vegetative plants receiving NH<sub>4</sub><sup>+</sup> had greater shoot, stem, and root biomass at elevated CO<sub>2</sub> (Bloom et al., 2002). Wheat receiving NO<sub>3</sub><sup>-</sup> grew slower at elevated CO<sub>2</sub> than at ambient CO<sub>2</sub> (Bloom et al., 2002). Shoot NO<sub>3</sub><sup>-</sup> concentrations in NH<sub>4</sub><sup>+</sup>-fed plants were undetectable while those in NO<sub>3</sub><sup>-</sup>-fed plants increased by 62% with CO<sub>2</sub> enrichment (Bloom et al., 2002). This increase was associated with an inhibition in NO<sub>3</sub><sup>-</sup> and NO<sub>2</sub><sup>-</sup> reductase activities under elevated CO<sub>2</sub> (Bloom et al., 2002).

The interaction between atmospheric CO<sub>2</sub> concentration and inorganic N form and how it influences plant growth and nutrient concentrations has not been examined in wheat or any other crop species grown to senescence. Here, we grew wheat hydroponically in controlled environment chambers and measured mineral nutrition, biomass, and nutrient allocation in response to three concentrations of atmospheric CO<sub>2</sub> (subambient, ambient, and elevated) and two forms of N nutrition (NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup>). We tested the following hypotheses: (1) plant nutrient concentrations and allocation patterns will respond differently to CO<sub>2</sub> enrichment under the two N forms, and (2) NO<sub>3</sub><sup>-</sup>-fed plants will show a smaller biomass and yield enhancement in response to CO<sub>2</sub> enrichment than NH<sub>4</sub><sup>+</sup>-fed plants as a result of CO<sub>2</sub> inhibition of shoot NO<sub>3</sub><sup>-</sup> assimilation. Also, we observed both differences in the Zn concentration between plants grown on NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup> and a strong dependence of Zn absorption on Zn and phytate concentration, indicating that phytate and bioavailable Zn are affected by N form and CO<sub>2</sub>. Therefore, we used the well supported Miller equation (Miller et al., 2007) to estimate how N and CO<sub>2</sub> might impact a hypothetical human population. Iron, another important micronutrient that forms complexes with phytate, was not analyzed because we observed no significant differences in iron concentrations between the N forms and because how best to

estimate Fe absorption in humans is still uncertain (Welch and Graham, 2004).

## MATERIALS AND METHODS

### EXPERIMENTAL

Wheat seeds (*Triticum aestivum* cv. Veery 10) were surface sterilized for one minute in 2.6% sodium hypochlorite solution and thoroughly rinsed with DDI water. The seeds were then rolled up in germination paper saturated with 10 mM CaSO<sub>4</sub>. The germination paper was placed in a 400 mL beaker with approximately 75 mL of 10 mM CaSO<sub>4</sub> solution, covered with a plastic bag and placed in an incubator (23°C) for four days. Seedlings were transplanted into 20 L tubs filled with an aerated nutrient solution that contained 1 mM CaSO<sub>4</sub>, 1 mM K<sub>2</sub>HPO<sub>4</sub>, 1 mM KH<sub>2</sub>PO<sub>4</sub>, 2 mM MgSO<sub>4</sub>, and 0.2 g L<sup>-1</sup> Fe-NaEDTA and micronutrients (20% of a modified Hoagland's solution with either 0.2 mM KNO<sub>3</sub> or 0.1 mM (NH<sub>4</sub>)<sub>2</sub>HPO<sub>4</sub> as the N source, Epstein and Bloom, 2005). The nutrient solution was replaced weekly and an additional 0.2 mM of NO<sub>3</sub><sup>-</sup> or NH<sub>4</sub><sup>+</sup> - N was added midweek until harvest. The solution volume was maintained by daily addition of deionized water. Solution pH varied between 6.8 and 7.0 for both of the N forms, and the NH<sub>4</sub><sup>+</sup> and the NO<sub>3</sub><sup>-</sup> solutions did not differ by more than 0.1 pH units.

The plants were grown in controlled environment chambers (Conviron, Winnipeg, Canada) set at 23/20°C day/night at 60–70% relative humidity with a photoperiod of 15 h. The photosynthetic flux density was 375 μmol m<sup>-2</sup> s<sup>-1</sup> at plant height. Plants were subjected to one of three CO<sub>2</sub> concentrations: “subambient” (310 ± 30 ppm), “ambient” (410 ± 30 ppm), and “elevated” (720 ± 5 ppm). Subambient CO<sub>2</sub> concentrations were maintained by passing air that entered the growth chamber through wet soda lime, a mixture of KOH, NaOH, and Ca(OH)<sub>2</sub> that was replaced as needed. The elevated CO<sub>2</sub> conditions were maintained in an environmental chamber equipped with non-dispersive infrared analyzers for CO<sub>2</sub> (Horiba model APBA-250E) and valves that added pure CO<sub>2</sub> to the incoming air stream to hold the chamber concentration at 720 ppm.

The wheat was grown until all aboveground parts turned completely yellow. Plant matter was sorted into grain, chaff, shoots, and roots and dried for 48 h at 55°C. Data on kernel number (KN), kernel mass, number of heads, kernels head<sup>-1</sup>, and HI were collected prior to sample preparation for nutrient analysis. A portion of the grain was analyzed for phytate using a modification of the method as described by Haug and Lantzsch (1983). The remainder of the grain as well as the shoots and chaff was bulked into five repetitions per treatment and sent to the UC Davis Analytical Laboratory for nutrient analysis. The roots of plants for each CO<sub>2</sub> × N treatment became entangled within the same tub; therefore, we were unable to separate the roots of the individual plants for analysis. Root data are thus presented as means for each treatment with no standard errors or confidence intervals.

Data were analyzed using PROC MIXED (SAS 9.0 Cary, NC, USA). Nitrogen form and CO<sub>2</sub> factors were treated as fixed independent variables. We used the Tukey–Kramer Honestly Significant Difference test for mean separation. Probabilities less than 0.05 were considered significant. Because some of the transformed variables did not meet the assumption of homogeneity

of variances, but one-way ANOVAs met the ANOVA assumptions, we analyzed the results via one-way ANOVAs to gain some information on the interactions between CO<sub>2</sub> and N form.

### MODELING THE INFLUENCE OF N FORM ON Zn NUTRITION IN THE HUMAN DIET

We used a database derived from the United Nation's Food and Agriculture Organization (FAO)'s national food balance sheets (FBS) to estimate the average daily per capita dietary intake of zinc and phytate from 95 different food commodities in each of 176 countries. This database combines FAO data on per capita intake of food commodities with USDA data on the nutrient or phytate content of each of these commodities. More detailed discussion of the creation of this database for the International Zinc Collaborative Group may be found in Wuehler et al. (2005). Using this database, we produced two datasheets: one containing per capita daily dietary intake of zinc from each food commodity for each country and another containing per capita phytate intake from each food commodity for each country. To calculate total dietary zinc (TDZ) and total dietary phytate (TDP) per country, we summed across the rows of all food commodities for each respective country.

To determine the proportion of a population at risk for zinc deficiency from a hypothetical least developed country (LDC), we first calculated TDP and TDZ values for a set of 44 countries defined by the United Nations as being least developed. We took the mean TDP and TDZ values for these countries to represent a hypothetical "less developed country." To calculate the bioavailable zinc portion (TAZ; usually a small fraction of TDZ) we used the Miller equation (Equation 1: Miller et al., 2007).

$$\text{TAZ} = 0.5 \cdot \left( A_{\text{max}} + \text{TDZ} + K_{\text{R}} \cdot \left( 1 + \frac{\text{TDP}}{K_{\text{P}}} \right) - \sqrt{\left( A_{\text{max}} + \text{TDZ} + K_{\text{R}} \cdot \left( 1 + \frac{\text{TDP}}{K_{\text{P}}} \right) \right)^2 - 4 \cdot A_{\text{max}} \cdot \text{TDZ}} \right)$$

Equation 1 : Miller equation

Mean TDZ and TDP values were converted to mg mmol<sup>-1</sup> and put into the Miller equation to compute the average per capita TAZ in our hypothetical LDC. The variables TDZ, TDP, and TAZ are described above, and  $A_{\text{max}}$ ,  $K_{\text{P}}$ , and  $K_{\text{R}}$  are constants as described in Miller et al. (2007).

We made an assumption that our hypothetical LDC receives half of its phytate and half of its zinc from wheat, which is roughly consistent with many of the LDCs in the FAO database. We analyzed the effect of elevated carbon dioxide levels on TDP, TDZ, and TAZ concentrations in a hypothetical LDC population for both NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup>-supplied wheat. To calculate a new TAZ for wheat grown under elevated CO<sub>2</sub> conditions, we first calculated the percent change in TAZ from ambient to elevated levels for wheat receiving NH<sub>4</sub><sup>+</sup> or NO<sub>3</sub><sup>-</sup>. This computed percent change was then applied to half of the hypothetical TDZ and TDP; meanwhile, the other half of the hypothetical TDZ and TDP remained unmodified. Thus, the total new TDP and TDZ is the sum of the unmodified and modified portions. These new TDP and TDZ

values for both NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup>-supplied wheat were then put into the Miller equation to compute new hypothetical TAZ values for an LDC. Differences and corresponding percent changes between the new TAZ values and the original TAZ value for a LDC were computed to determine the overall affect of elevated CO<sub>2</sub> on TAZ in NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup>-supplied wheat for an average developing world population. TAZ, TDP, and TDZ concentrations can only be compared within a single N form across the CO<sub>2</sub> concentrations due to methodological constraints of the model.

## RESULTS

We divide the results here into three categories: first, biomass and yield data for the shoots, grain, and roots; second, tissue concentrations and whole plant micro- and macronutrient contents; and third, nutrient distribution among the different tissues. Values of the statistical significance of the results were placed into a table (Table 1) in order to improve the readability of the text.

### BIOMASS AND YIELD

Plants supplied NH<sub>4</sub><sup>+</sup> vs. NO<sub>3</sub><sup>-</sup> nutrition reacted differently to CO<sub>2</sub> enrichment (Figure 1; Table 1). Plants supplied NH<sub>4</sub><sup>+</sup> differed across CO<sub>2</sub> treatments for most of the yield and biomass measurements. The greatest values typically were found at ambient CO<sub>2</sub> concentrations. Shoot, chaff, grain yield, number of heads, and KN were greatest at ambient CO<sub>2</sub> levels. Individual kernel mass was greatest under both ambient and elevated CO<sub>2</sub> treatments. HI and kernels head<sup>-1</sup> showed no change across CO<sub>2</sub> treatments. In contrast, biomass and yield measures of NO<sub>3</sub><sup>-</sup>-supplied plants did not differ among the three CO<sub>2</sub> concentrations.

At subambient CO<sub>2</sub>, differences between the NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup> treatments occurred in shoot biomass and three of the yield components: kernel mass, head number, and kernels head<sup>-1</sup>. Ammonium-supplied plants had a larger number of heads while NO<sub>3</sub><sup>-</sup>-supplied plants had greater shoot biomass, kernel mass, and kernels head<sup>-1</sup>. At ambient CO<sub>2</sub>, NH<sub>4</sub><sup>+</sup>-supplied plants had a greater number of heads and greater chaff biomass. Plants supplied NO<sub>3</sub><sup>-</sup> had a larger number of kernels head<sup>-1</sup>. At elevated CO<sub>2</sub>, biomass and yield measures did not differ with N treatment.

### ROOT

Roots had a smaller mean biomass when supplied NH<sub>4</sub><sup>+</sup> than when supplied NO<sub>3</sub><sup>-</sup> at all CO<sub>2</sub> concentrations (Figure 1). Both N treatments had the greatest biomass at ambient CO<sub>2</sub>, and the smallest at subambient CO<sub>2</sub>. The highest root:shoot ratios for both NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup>-supplied plants were observed at ambient and elevated CO<sub>2</sub>. Ammonium-supplied plants always had lower root:shoot ratios and biomasses than NO<sub>3</sub><sup>-</sup>-supplied plants at the same CO<sub>2</sub> concentration.

## NUTRIENTS

### Total plant nutrients

Total plant nutrients generally followed the same trend within N form, although NH<sub>4</sub><sup>+</sup>-supplied plants exhibited a greater diversity of responses to increasing CO<sub>2</sub> concentrations (Table 2). Total plant P, K, B, Ca, Mg, and Zn decreased with increasing CO<sub>2</sub> under NH<sub>4</sub><sup>+</sup>, while S and Mn were highest under ambient CO<sub>2</sub>.

**Table 1 | Results of a series of one-way ANOVAs run on the data.**

Among CO <sub>2</sub> cnc. within an N form	Grain		Shoot			
	NH <sub>4</sub> <sup>+</sup>	NO <sub>3</sub> <sup>-</sup>	NH <sub>4</sub> <sup>+</sup>	NO <sub>3</sub> <sup>-</sup>		
Total N	**	NS	***	NS		
P	*	NS	**	NS		
K	***	NS	NS	***		
S	NS	NS	**	NS		
Ca	***	NS	*	**		
Mg	NS	NS	NS	***		
Zn	NS	NS	***	***		
B	**	*	*	*		
Mn	**	NS	***	NS		
Fe	*	NS	*	*		
Cu	***	NS	NS	***		
NO <sub>3</sub> <sup>-</sup> - N	*	NS	***	***		
Phytate	NS	NS	N/A	N/A		
Between N forms within a CO <sub>2</sub> cnc.	Grain			Shoot		
	Sub	Amb	Elev	Sub	Amb	Elev
Total N	**	NS	NS	NS	NS	**
P	NS	NS	NS	*	NS	**
K	**	NS	**	NS	**	NS
S	*	NS	*	**	NS	***
Ca	***	***	***	NS	NS	NS
Mg	NS	NS	NS	***	***	*
Zn	*	***	*	**	***	***
B	NS	NS	*	***	NS	***
Mn	NS	NS	NS	***	***	*
Fe	NS	NS	NS	NS	NS	NS
Cu	**	**	**	NS	***	*
NO <sub>3</sub> <sup>-</sup> - N	NS	NS	NS	***	***	***
Phytate	**	NS	NS	N/A	N/A	N/A
Among CO <sub>2</sub> cnc. or between N forms	Sub	Amb	Elev	NH <sub>4</sub> <sup>+</sup>	NO <sub>3</sub> <sup>-</sup>	
Yield	NS	*	NS	**	NS	
Shoot	**	NS	NS	**	NS	
Chaff	NS	*	NS	**	NS	
Grain number	NS	NS	NS	*	NS	
Grain mass	***	NS	NS	***	NS	
Grains head <sup>-1</sup>	***	NS	NS	NS	NS	
Heads	**	**	NS	*	NS	
Harvest index	NS	NS	NS	NS	NS	

Differences among CO<sub>2</sub> concentrations within an N form and between N forms at each CO<sub>2</sub> concentration for shoot and grain nutrient concentrations. Differences among CO<sub>2</sub> concentrations within an N form or between N forms at each CO<sub>2</sub> concentration for biomass and yield components. The symbols indicate statistical significance (\*0.05, \*\*0.01, \*\*\*0.001, NS, not significant).

Ammonium-supplied plants had the greatest amounts of Fe and total N at subambient CO<sub>2</sub>. Nitrate-supplied plants accumulated the greatest amounts of total N, P, K, S, B, Ca, Zn, Mn, and Mg at ambient CO<sub>2</sub>. Only three nutrients – K, S, and Fe – had the lowest contents at elevated CO<sub>2</sub>.

### Shoot

Under NH<sub>4</sub><sup>+</sup> supply, plants varied with CO<sub>2</sub> concentration for total N, P, S, Ca, Cu, B, Mn, Zn, and NO<sub>3</sub><sup>-</sup> - N (Table 1; Figure 2). Calcium and Cu were highest under subambient CO<sub>2</sub>. Total N and S were greatest at subambient and elevated CO<sub>2</sub>. Nitrate-N was greatest at ambient CO<sub>2</sub>. Phosphorus was highest at elevated CO<sub>2</sub> concentrations. Boron, Zn, and Mn increased with CO<sub>2</sub> concentration.

Plants supplied NO<sub>3</sub><sup>-</sup> showed significant variation across CO<sub>2</sub> treatments for K, Ca, Mg, B, Fe, Cu, Zn, and NO<sub>3</sub><sup>-</sup> - N (Table 1; Figure 2). Calcium and Cu had the greatest concentrations at subambient CO<sub>2</sub>. The highest concentrations of B, Fe, and Zn occurred at subambient and elevated CO<sub>2</sub>. Potassium concentrations were highest at elevated CO<sub>2</sub>. Nitrate-N increased with CO<sub>2</sub>. Magnesium showed the opposite trend, decreasing with CO<sub>2</sub> concentration.

Differences between N forms were also evident. At subambient CO<sub>2</sub>, NH<sub>4</sub><sup>+</sup>-supplied plants had increased concentrations of P, S, and Zn, while NO<sub>3</sub><sup>-</sup>-supplied plants had greater concentrations of B, Mg, Mn, and NO<sub>3</sub><sup>-</sup> - N (Table 1; Figure 2). Concentrations of K, Zn, and Cu were higher in plant supplied NH<sub>4</sub><sup>+</sup> at ambient CO<sub>2</sub>, while Mg, Mn, and NO<sub>3</sub><sup>-</sup> - N were greater in plants supplied NO<sub>3</sub><sup>-</sup>. At elevated CO<sub>2</sub>, concentrations of N, P, S, and Zn were higher in plants supplied NH<sub>4</sub><sup>+</sup>, while concentrations of B, Mg, Mn, and NO<sub>3</sub><sup>-</sup> - N were greater in plants supplied NO<sub>3</sub><sup>-</sup>.


### Grain

**Grain nutrient concentrations.** Plants supplied NH<sub>4</sub><sup>+</sup> showed significant variation across the CO<sub>2</sub> treatments in the concentrations of total N, P, K, Ca, B, Cu, Fe, Mn, and NO<sub>3</sub><sup>-</sup> - N (Table 1; Figure 3). The greatest concentrations of total N, P, K, Ca, and Cu were found at subambient CO<sub>2</sub>. Iron concentrations were high at both subambient and ambient CO<sub>2</sub>. Boron was equally high at subambient and elevated CO<sub>2</sub>. Manganese was greatest at elevated CO<sub>2</sub>. Nitrate-N decreased with increasing CO<sub>2</sub>.

Significant differences among the NO<sub>3</sub><sup>-</sup>-supplied plants across CO<sub>2</sub> treatments were only observed in S and B. The greatest concentrations of B were found at subambient CO<sub>2</sub>. Sulfur was highest at ambient CO<sub>2</sub>.

Nitrogen form significantly affected grain nutrient concentrations (Table 1; Figure 3). At subambient CO<sub>2</sub>, NH<sub>4</sub><sup>+</sup>-supplied plants had higher concentrations of total N, K, S, Ca, Zn, and Cu than NO<sub>3</sub><sup>-</sup> plants. At ambient CO<sub>2</sub>, Ca, Zn, and Cu were greatest under NH<sub>4</sub><sup>+</sup>. Ammonium-supplied plants also had the highest concentrations of K, S, Ca, Zn, and Cu at elevated CO<sub>2</sub>. At no CO<sub>2</sub> concentration did plants supplied NH<sub>4</sub><sup>+</sup> have significantly lower concentrations of any micro- or macronutrient than those supplied NO<sub>3</sub><sup>-</sup>.

**Phytate and bioavailable Zn.** Phytate was relatively insensitive to CO<sub>2</sub> concentration. Phytate concentrations were highest



**FIGURE 1 | The effect of N form and CO<sub>2</sub> concentration on biomass and yield components of wheat hydroponically grown to senescence.** Closed (NH<sub>4</sub><sup>+</sup>) and open (NO<sub>3</sub><sup>-</sup>) symbols and error bars represent the

back-transformed means and 95% confidence intervals (*n* = 10). Means within N form across CO<sub>2</sub> treatment are significantly different if labeled with different letters. Root-related data do not have error bars.

at subambient CO<sub>2</sub> for NH<sub>4</sub><sup>+</sup>-supplied plants (Figure 4). Sub-ambient CO<sub>2</sub> also produced the lowest phytate concentrations in NO<sub>3</sub><sup>-</sup>-supplied plants. NH<sub>4</sub><sup>+</sup>-supplied plants had greater phytate concentrations than NO<sub>3</sub><sup>-</sup>-supplied plants at subambient CO<sub>2</sub>, but not at the other CO<sub>2</sub> concentrations. Grain from plants grown

under NH<sub>4</sub><sup>+</sup> nutrition had roughly 7, 18, and 8% higher bioavailable Zn than NO<sub>3</sub><sup>-</sup>-supplied plants at subambient, ambient, and elevated CO<sub>2</sub>, respectively (Figure 4).

Based on this phytate and bioavailable Zn data, we modeled how a human population from a LDC would be affected by

**Table 2 | Total plant nutrients (mg plant<sup>-1</sup>) as affected by N form and CO<sub>2</sub> concentration.**

	Sub	Amb	Elev	Sub	Amb	Elev	Sub	Amb	Elev	Sub	Amb	Elev
	<b>Total N</b>			<b>P</b>			<b>K</b>			<b>S</b>		
NH <sub>4</sub> <sup>+</sup>	215.66	191.62	208.56	80.64	73.96	68.69	228.91	202.92	196.33	49.98	50.72	46.82
NO <sub>3</sub> <sup>-</sup>	159.39	210.26	164.88	63.21	85.02	67.75	208.32	259.07	198.79	42.21	50.84	38.25
	<b>B</b>			<b>Ca</b>			<b>Mg</b>			<b>Zn</b>		
NH <sub>4</sub> <sup>+</sup>	0.28	0.25	0.18	23.18	19.55	19.24	42.41	38.62	35.34	0.62	0.54	0.45
NO <sub>3</sub> <sup>-</sup>	0.29	0.41	0.31	21.10	25.54	22.48	45.26	52.45	52.45	0.27	0.48	0.36
	<b>Mn</b>			<b>Fe</b>			<b>Cu</b>					
NH <sub>4</sub> <sup>+</sup>	2.66	2.93	2.24	1.93	1.26	1.47	0.06	0.05	0.05			
NO <sub>3</sub> <sup>-</sup>	2.16	3.54	2.52	2.16	2.71	1.75	0.05	0.06	0.06			

changes in atmospheric CO<sub>2</sub> concentrations (Table 3). The calculations were based on differences among CO<sub>2</sub> concentrations; therefore, modeled TDZ, TDP, and TAZ values cannot be compared between NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup>-supplied grain. Grain from plants supplied the different N forms behaved differently as CO<sub>2</sub> concentration increased. We found that under NH<sub>4</sub><sup>+</sup> supply, TAZ would increase 3.6% with the rise in CO<sub>2</sub> from subambient to ambient, and decrease 1.6% with the rise from ambient to elevated CO<sub>2</sub> (Figure 4). Humans provided NO<sub>3</sub><sup>-</sup>-supplied wheat would experience a decrease in TAZ of 3.5% going from subambient to ambient, and an increase 5.6% from ambient to elevated CO<sub>2</sub> (Figure 4).

### Roots

Ammonium-supplied plants generally showed a trend toward decreasing nutrient concentrations with increasing CO<sub>2</sub> concentration while NO<sub>3</sub><sup>-</sup>-supplied plants varied widely across CO<sub>2</sub> treatments (Figure 5). The decrease in nutrient concentrations under NH<sub>4</sub><sup>+</sup> supply corresponded to an increase in root mass. Nitrate-supplied plants tended to have their highest nutrient concentrations in the ambient and elevated CO<sub>2</sub> treatments. Ammonium-supplied plants had higher concentrations of Zn and Mn across all of the CO<sub>2</sub> treatments, as well as higher total N and Fe at subambient CO<sub>2</sub>. Nitrate-supplied plants typically had higher concentrations of the other nutrients at all CO<sub>2</sub> concentrations.

### Distribution of nutrients

The distribution of nutrients and micronutrients among plant parts followed similar patterns in both the NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup>-supplied plants, although the NH<sub>4</sub><sup>+</sup>-supplied plant distributions were slightly more variable (Table 4). Allocations to root and grain usually were greatest at ambient CO<sub>2</sub>, and those to chaff and shoots at either subambient or elevated CO<sub>2</sub>. Grain typically contained the largest proportion of total N, P, Zn, and Cu, although the organ with the largest percentage of Cu varied with CO<sub>2</sub> treatment among NO<sub>3</sub><sup>-</sup>-supplied plants. Plants at subambient and elevated CO<sub>2</sub> allocated more Cu to the grain, while those at ambient CO<sub>2</sub> allocated more to the roots. In general shoots received the majority of K, S, B, Ca, and Mg for all N and CO<sub>2</sub> treatments. Ammonium-supplied plants allocated slightly more Mn to the roots at subambient CO<sub>2</sub>, but allocated increasing amounts to the shoots at the expense of the roots as CO<sub>2</sub> concentration increased. In contrast, NO<sub>3</sub><sup>-</sup>-supplied plants allocated most of the

Mn to the shoots. Ammonium-supplied plants typically allocated more resources to the chaff while NO<sub>3</sub><sup>-</sup>-supplied plants allocated a greater percentage of elements to the roots.


### DISCUSSION

No other study to our knowledge has examined the influence of N form (NH<sub>4</sub><sup>+</sup> vs. NO<sub>3</sub><sup>-</sup>) on plant nutrient relations at three different atmospheric CO<sub>2</sub> concentrations. Overall, N form affected growth, total plant nutrient contents, and nutrient distribution in senescing wheat shoots, grain, and roots. The influence of NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup> on growth and nutrient status were so distinct that they should be treated as separate nutrients and not bundled into a general category of N nutrition. Wheat size and nutrition at senescence responded to CO<sub>2</sub> concentration in a non-linear manner. As was previously shown (Bloom et al., 2012), we found that plants supplied with NH<sub>4</sub><sup>+</sup> were more responsive to CO<sub>2</sub> concentration than those supplied with NO<sub>3</sub><sup>-</sup>.

Although not explicitly addressed here because of the heterogeneity of variances, interactions between CO<sub>2</sub> and N treatments likely existed for a number of the biomass and nutrient measures. Most nutrient concentrations were generally higher in NH<sub>4</sub><sup>+</sup>-supplied plants, with the exceptions of NO<sub>3</sub><sup>-</sup> - N, Mg, B, and Mn, which were generally higher in NO<sub>3</sub><sup>-</sup>-supplied plants. Phytate, which hinders human absorption of Zn and Fe (Raboy, 2009), showed little variation at ambient and elevated CO<sub>2</sub> between NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup>-supplied plants, which, in conjunction with the observed greater bioavailable of Zn in NH<sub>4</sub><sup>+</sup>-supplied plants, may have consequences for human nutrition. Distribution of nutrients to the shoots, roots, chaff, and grain in response to CO<sub>2</sub> concentration and N form was also non-linear and varied by nutrient.


### BIOMASS AND YIELD

The data support our hypothesis that NO<sub>3</sub><sup>-</sup>-supplied plants would show a more limited biomass and yield enhancement with CO<sub>2</sub> enrichment than NH<sub>4</sub><sup>+</sup>-supplied plants. Nevertheless, mean biomass and yield decreased from ambient to elevated CO<sub>2</sub> in both NO<sub>3</sub><sup>-</sup>- and NH<sub>4</sub><sup>+</sup>-supplied plants in contrast to biomass increases in prior work on wheat seedlings (Bloom et al., 2002). NO<sub>3</sub><sup>-</sup>-supplied plants allocated more biomass to roots and had larger root:shoot ratios than NH<sub>4</sub><sup>+</sup>-supplied plants regardless of CO<sub>2</sub> concentrations as has been reported previously (Wang and Below,



**FIGURE 2 | The effect of N form and CO<sub>2</sub> concentration on shoot nutrient concentrations of wheat grown hydroponically to senescence.** Closed (NH<sub>4</sub><sup>+</sup>) and open (NO<sub>3</sub><sup>-</sup>) symbols represent back-transformed means and 95% confidence intervals (n = 5). Macro- and micronutrients are listed in the upper


left of each frame. Differences are significant within N form if letters are different. Differences between N forms at each CO<sub>2</sub> concentration are generally significantly different if error bars do not overlap (see **Table 1** for statistical significance).



**FIGURE 3 | The effect of N form and CO<sub>2</sub> concentration on grain nutrient concentrations of wheat grown hydroponically to senescence.** Closed (NH<sub>4</sub><sup>+</sup>) and open (NO<sub>3</sub><sup>-</sup>) symbols represent back-transformed means and 95% confidence intervals (*n* = 5). Macro- and micronutrients are listed in the upper

left of each frame. Differences are significant within N form if letters are different. Differences between N forms at each CO<sub>2</sub> concentration are generally significantly different if error bars do not overlap (see **Table 1** for statistical significance).





1995; Bloom et al., 2002), but increased root mass at elevated CO<sub>2</sub> concentration for NO<sub>3</sub><sup>-</sup>-supplied plants reported previously (Bloom et al., 2002) were not observed here. The shoot biomass data suggest that growth differences measured early in the lifespan of wheat supplied with NH<sub>4</sub><sup>+</sup> or NO<sub>3</sub><sup>-</sup> or NH<sub>4</sub><sup>+</sup> (i.e., greater shoot biomass in plants supplied NH<sub>4</sub><sup>+</sup> relative to those supplied NO<sub>3</sub><sup>-</sup> at elevated CO<sub>2</sub> concentrations; Bloom et al., 2002) do not necessarily carry through to senescence. This may be due in part to a shift in NO<sub>3</sub><sup>-</sup> assimilation to the root (Kruse et al., 2003), allowing NO<sub>3</sub><sup>-</sup>-supplied plants to compensate for the decrease in shoot NO<sub>3</sub><sup>-</sup> assimilation that occurs at elevated atmospheric CO<sub>2</sub> concentrations (Bloom et al., 2002, 2010, 2012).

The decrease in yield and biomass measures at elevated CO<sub>2</sub> concentrations does not agree with field observations where wheat yields as well as overall biomass increased with elevated CO<sub>2</sub> (Högy and Fangmeier, 2008; Taub et al., 2008). Similarly, our results that the greatest values for other yield measures (e.g., heads, kernel mass, KN) occurred at ambient CO<sub>2</sub> concentrations varies from the literature. High CO<sub>2</sub> has been found to increase flowering tillers (Havelka et al., 1984; Fangmeier et al., 1996), KN (McKee et al., 1997), and kernel mass (i.e., thousand grain weight; McKee

**Table 3 | Total dietary Zn (TDZ), total dietary phytate (TDP), and total bioavailable Zn (TAZ) of a human population from a hypothetical less developed nation reliant on wheat for 50% of their dietary phytate and Zn as modeled using the Miller equation.**

		Sub → Amb (g/kg <sup>-1</sup> )	Amb → Elev (g/kg <sup>-1</sup> )
NH <sub>4</sub> <sup>+</sup>	TDZ	9.21	8.69
	TDP	2241.92	2264.70
	TAZ	1.76	1.67
NO <sub>3</sub> <sup>-</sup>	TDZ	8.68	9.53
	TDP	2346.00	2275.33
	TAZ	1.64	1.79


The data columns represent the change in TDZ, TDP, and TAZ concentration going from subambient to ambient and ambient to elevated CO<sub>2</sub> concentrations, respectively. The values are calculated as deviations from the mean TDZ, TDP, and TAZ concentrations as produced from FAO and USDA data (Wuehler et al., 2005). Baseline values for TDZ, TDP, and TAZ were 8.90, 2278.00, and 1.70 g kg<sup>-1</sup>, respectively.

et al., 1997). Conflicting results, however, have also been reported (e.g., Havelka et al., 1984). Many of the field and open top chamber studies were grown under natural light and thus received substantially greater photosynthetic flux density than our chamber-grown plants. These higher light conditions would be more favorable to biomass accumulation. Also, these studies typically applied high amounts of mixed N fertilizer (e.g., NH<sub>4</sub>NO<sub>3</sub>), and yields and biomass have been found to be greater under mixed N nutrition than under either NH<sub>4</sub><sup>+</sup> or NO<sub>3</sub><sup>-</sup> alone (Cox and Reisenauer, 1973; Gentry et al., 1989; Heberer and Below, 1989; Wang and Below, 1995). Finally, the wheat cultivar we used (*T. aestivum* cv. Veery 10) is a short-statured variety that has rarely been used in other studies and may have accounted for some of the differences between our study and other published data.

Our results that NH<sub>4</sub><sup>+</sup>-supplied plants had greater yield and yield components than NO<sub>3</sub><sup>-</sup>-supplied plants at ambient CO<sub>2</sub> have been observed previously (Wang and Below, 1996; Chen et al., 1998). Wang and Below (1995) observed greater numbers of kernels head<sup>-1</sup> and KN in plants supplied NO<sub>3</sub><sup>-</sup> that was not observed here. Their study, however, supplied NH<sub>4</sub><sup>+</sup> at relatively high levels (~8.9 vs. 0.2 mM NH<sub>4</sub><sup>+</sup> - N in our study). Several studies (Bennett and Adams, 1970; Cox and Reisenauer, 1973) have found that incipient NH<sub>4</sub><sup>+</sup> toxicity can start appearing at N levels as low as 0.08–0.2 mM NH<sub>4</sub><sup>+</sup>, although the onset of NH<sub>4</sub><sup>+</sup> toxicity depends on light level (Magalhaes and Wilcox, 1984; Britto and Kronzucker, 2002) and solution pH (Findenegg, 1987). The poorer performance of the NH<sub>4</sub><sup>+</sup> treatment in Wang and Below (1995), therefore, might derive from NH<sub>4</sub><sup>+</sup> toxicity. We have previously determined that the 0.2 mM NH<sub>4</sub><sup>+</sup>-supplied to our plants to be sufficiently high for normal growth, but low enough to avoid toxicity problems under our experimental conditions (Bloom et al., 2002).

#### PLANT NUTRIENTS

Our second hypothesis, that nutrient concentrations are differentially affected by the inorganic N form supplied to the plants and CO<sub>2</sub> enrichment, was supported by our data. CO<sub>2</sub> concentration



**FIGURE 5 | The effect of N form and CO<sub>2</sub> concentration on root tissue nutrient concentrations of wheat grown hydroponically to senescence.** Closed (NH<sub>4</sub><sup>+</sup>) and open (NO<sub>3</sub><sup>-</sup>) symbols represent the bulked treatment mean

(*n* = 10). Macro- and micronutrients are listed in the upper left of each frame. The lack of error bars reflects that the root mass for each treatment was bulked and analyzed as a unit.

**Table 4 | Organ nutrient allocation as percentage of the plant total under the CO<sub>2</sub> and N form treatments.**

		Root	Cha	Shoots	Grain	Root	Chaff	Shoots	Grain	Root	Chaff	Shoots	Grain	Root	Chaff	Shoots	Grain
		<b>Total N</b>				<b>P</b>				<b>K</b>				<b>S</b>			
<b>Sub</b>	NH <sub>4</sub> <sup>+</sup>	5.22	9.23	10.35	75.20	4.34	14.68	38.99	41.98	2.24	22.75	58.15	16.86	4.71	16.67	63.11	15.50
<b>Amb</b>		5.02	6.36	9.08	79.54	4.56	12.10	36.78	46.55	2.38	16.30	62.59	18.73	5.18	11.25	61.49	22.09
<b>Elev</b>		4.32	8.36	9.79	77.52	4.13	12.96	40.89	42.02	2.48	16.33	63.12	18.06	4.12	14.88	63.54	17.46
<b>Sub</b>	NO <sub>3</sub> <sup>-</sup>	6.45	4.66	11.89	77.00	6.46	10.21	41.24	42.10	10.05	10.50	64.04	15.41	8.14	5.21	66.52	20.12
<b>Amb</b>		7.96	4.95	10.48	76.61	9.03	10.91	35.97	44.09	17.87	10.54	56.04	15.56	13.60	7.87	56.96	21.57
<b>Elev</b>		6.67	6.88	9.82	76.64	7.30	12.78	36.04	43.88	12.74	12.44	59.35	15.48	10.93	7.19	61.52	20.35
		<b>B</b>				<b>Ca</b>				<b>Mg</b>				<b>Zn</b>			
<b>Sub</b>	NH <sub>4</sub> <sup>+</sup>	3.53	23.50	69.11	3.86	10.74	14.26	69.31	5.69	4.25	12.40	57.24	26.11	4.57	9.23	24.98	61.22
<b>Amb</b>		2.03	23.61	70.72	3.63	12.45	11.71	68.88	6.95	4.13	12.12	53.64	30.11	5.57	11.31	20.97	62.15
<b>Elev</b>		2.07	25.21	68.62	4.10	11.44	13.49	68.80	6.27	3.80	14.35	54.20	27.66	5.12	12.16	25.49	57.23
<b>Sub</b>	NO <sub>3</sub> <sup>-</sup>	5.25	17.24	74.74	2.77	14.82	7.12	73.89	4.17	4.34	9.17	66.09	20.40	4.17	5.75	27.20	62.88
<b>Amb</b>		9.11	21.42	67.10	2.37	18.83	10.06	66.59	4.53	5.19	13.73	58.29	22.78	9.56	8.51	16.76	65.16
<b>Elev</b>		6.38	23.09	67.86	2.67	21.56	9.90	63.65	4.89	6.12	13.90	54.40	25.58	7.14	8.96	18.65	65.25
		<b>Mn</b>				<b>Fe</b>				<b>Cu</b>							
<b>Sub</b>	NH <sub>4</sub> <sup>+</sup>	38.53	11.97	35.41	14.09	72.81	3.39	18.44	5.36	15.77	10.71	28.01	45.52				
<b>Amb</b>		30.72	14.50	39.80	14.98	81.28	1.23	12.21	5.29	19.64	8.48	24.49	47.39				
<b>Elev</b>		23.01	18.98	43.24	14.78	74.85	2.01	19.47	3.67	18.00	9.19	24.97	47.84				
<b>Sub</b>	NO <sub>3</sub> <sup>-</sup>	7.72	16.35	62.77	13.16	78.58	0.92	16.79	3.70	18.96	8.27	32.51	40.26				
<b>Amb</b>		5.47	20.25	60.01	14.26	87.37	0.74	8.39	3.50	40.79	6.79	19.64	32.78				
<b>Elev</b>		10.15	23.71	52.07	14.07	87.06	0.70	9.46	2.78	21.70	8.50	24.48	45.32				

and N form interactions may alter tissue demands for nutrients. For many nutrients, ratios between different elements are typically maintained within a narrow range (Garten, 1976; Bloom et al., 1985; Loladze, 2002). CO<sub>2</sub> concentration and N form may disturb the balance between different nutrients, leading to a cascade of changes in demand, accumulation, and allocation among the different plant tissues (e.g., Loladze, 2002; Högy and Fangmeier, 2008; Natali et al., 2009). Nitrate-supplied plants accumulated the greatest amounts of nutrients at ambient CO<sub>2</sub> (Table 2). Some portion of the greater response of NH<sub>4</sub><sup>+</sup>-supplied plants to CO<sub>2</sub> derived from a dilution effect from the greater biomass at ambient CO<sub>2</sub> concentrations (Figures 2 and 3). Total amounts of nutrients tended to decline with CO<sub>2</sub> enrichment for NH<sub>4</sub><sup>+</sup>-supplied plants, which had the greatest amounts of macro/micronutrients at subambient CO<sub>2</sub> (Table 2). These results have not been observed in other published studies (e.g., Fangmeier et al., 1997; Wu et al., 2004). Growth chamber studies, however, tend to have more exaggerated differences among treatments than field and greenhouse experiments (Högy and Fangmeier, 2008), and N source cannot be well-controlled in field and greenhouse experiments.

The observed increase in NO<sub>3</sub><sup>-</sup>-N concentration with CO<sub>2</sub> concentration in NO<sub>3</sub><sup>-</sup>-supplied plants has been reported previously (Bloom et al., 2002), and adds further support to the hypothesis that elevated CO<sub>2</sub> concentrations and the resulting decrease in photorespiration inhibit shoot NO<sub>3</sub><sup>-</sup> photoassimilation. Nevertheless, tissue NO<sub>3</sub><sup>-</sup>-N concentrations observed here were substantially lower than those in the earlier study (Bloom et al., 2002). Again, this may derive from difference in life stages in

the two studies. Most of the N available to the plant for grain filling comes from N translocation rather than uptake from the substrate (Simpson et al., 1983). Probably, the plants continued to assimilate plant NO<sub>3</sub><sup>-</sup> using a non-photorespiratory dependent process such as root assimilation after root N uptake slowed or stopped. Loss of NO<sub>3</sub><sup>-</sup> through root efflux to the nutrient solution also may have contributed to the lower concentration of NO<sub>3</sub><sup>-</sup>-N.

The partitioning and accumulation of all mineral elements was affected in some manner by the CO<sub>2</sub> treatment and N form supplied to the plants. Observations that cation concentrations decrease under NH<sub>4</sub><sup>+</sup> supply (e.g., Cox and Reisenauer, 1973; Gashaw and Mugwira, 1981; Wang and Below, 1998) relative to NO<sub>3</sub><sup>-</sup> supply were not apparent in this study. Again, this could be partly due to the relatively low concentration of NH<sub>4</sub><sup>+</sup>-supplied in our study, the age of the plants at harvest, and differences among wheat cultivars.

Allocation of nutrients within the plant followed similar trends for both N forms, with the exceptions of Mn and Cu (Table 2). Interestingly, in NO<sub>3</sub><sup>-</sup>-supplied plants, shoot Mn concentrations increased slightly with CO<sub>2</sub>, and these plants allocated far more Mn to the shoots than NH<sub>4</sub><sup>+</sup>-supplied plants at all CO<sub>2</sub> concentrations. Manganese (Mn<sup>2+</sup>) has been found to activate Rubisco in place of Mg<sup>2+</sup> and the Rubisco-Mn complex has been observed to decrease Rubisco carboxylase activity while minimally affecting or even enhancing oxygenase activity (Jordan and Ogren, 1983). The slight increase in shoot Mn with CO<sub>2</sub> corresponded to a large 23% decrease in Mg concentration. Manganese, which can act as a cofactor for glutamine synthetase (Smirnoff and Stewart, 1987), was also the only nutrient that NH<sub>4</sub><sup>+</sup>-supplied plants allocated a

greater percentage to the roots at the expense of the shoots. NO<sub>3</sub><sup>-</sup>-supplied plants typically allocated a higher percentage of most nutrients to the roots, as has been reported previously (Wang and Below, 1995, 1998).

Phytate, which forms complexes with divalent cations, has been found to hinder human Zn and Fe absorption during digestion and thus has been labeled an “anti-nutrient.” It may serve a number of valuable functions, however, including roles as an anti-oxidant and anti-cancer agent (Raboy, 2009). Phytate is also the major repository of grain P, and variation in P supply to the developing seed is the major determinant of net seed phytate accumulation (Raboy, 1997, 2009; Cakmak et al., 2010). To our knowledge, no published studies have explicitly looked at how phytate is affected by CO<sub>2</sub> concentration. Elevated CO<sub>2</sub> has been found to have a much larger negative impact on Zn and Fe concentrations than on P in wheat (Loladze, 2002; Cakmak et al., 2010). Several studies (e.g., Fangmeier et al., 1999; Högy and Fangmeier, 2008) have observed that P increases slightly with CO<sub>2</sub> concentration, and because the majority of P is tied up in phytate, this may cause increases in grain phytate concentrations as atmospheric CO<sub>2</sub> rises. As a result, bioavailable Zn and Fe—Zn and Fe not bound to phytate—is expected to decrease even further (Loladze, 2002).

Nonetheless, we did not observe such trends in macro- and micronutrient concentrations in this study. The mechanism behind these contrasting results is not clear, although the environmental conditions and nutrient solution in which the plants were grown likely had some role. The modeled data demonstrated only a small negative impact of CO<sub>2</sub> concentration on bioavailable Zn concentrations (Table 4), which was unexpected. Indeed, the grain from NO<sub>3</sub><sup>-</sup>-supplied plants actually showed a slight increase in bioavailable Zn between ambient and elevated CO<sub>2</sub>. These results combined with the differences in grain bioavailable Zn between NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup>-supplied plants demonstrates that N form may differentially affect the nutritional status of this important nutrient, especially in less developed countries that might be more dependent on phytate-rich grains for their Zn nutrition (Table 3). The milling process removes some, if not most, of the phytate and grain mineral content with the bran fraction of the grain (Gutteri et al., 2006). Regardless, with over 50% of the human population suffering from Zn deficiencies, even small increases in

bioavailable Zn would be beneficial (Loladze, 2002). This modeling exercise, however, is not a prediction of how increasing CO<sub>2</sub> will affect wheat nutrition so much as illustrates that N source may mediate, to some extent, the effects of CO<sub>2</sub> on phytate and bioavailable Zn, and that N source will become an even more important agricultural consideration in the future.

In summary, both CO<sub>2</sub> concentration and N form strongly affect biomass and yield in hydroponically grown wheat, as well as nutrient concentrations in above- and belowground tissues. Interactions among plant nutrient concentrations, CO<sub>2</sub> concentrations, and N form are complex and non-linear. The impact of N form and CO<sub>2</sub> concentration on the mechanisms affecting nutrient accumulation and distribution requires further research and extension to more realistic and agriculturally relevant growing conditions found in greenhouse and field studies. Of course, in greenhouse and field studies, control of N source is limited and control of atmospheric CO<sub>2</sub> concentration is expensive. The effects of CO<sub>2</sub> and N form on agriculture and human nutrition observed here are interesting and suggest a new area of research on mitigating the effects of climate change on agriculture. The supply of fertilizers (e.g., urea, NH<sub>4</sub>NO<sub>3</sub>, anhydrous NH<sub>3</sub>, organic amendments) or addition of nitrification inhibitors that increase the amount of available NH<sub>4</sub><sup>+</sup> may have beneficial effects for human nutrition, particularly in regards to micronutrient deficiencies such as Zn and Fe that currently affect billions of people worldwide. In the face of the potentially negative consequences of climate change on agriculture, all avenues of mitigation must be examined, and even small improvements may prove worthwhile.

## ACKNOWLEDGMENTS

The authors would like to thank Kenneth Brown and Jan Peerson at the International Zinc Collaborative Group for providing access to their food balance sheet database which was used in calculating total dietary zinc and phytate values. We would also like to thank Hsien Easlon for his advice and critical review of the manuscript. Finally we thank two anonymous reviewers for their critical reviews of the manuscript. This work was supported by NSF IOS-08-18435 and the National Research Initiative Competitive Grant no. 2008-35100-04459 from the USDA National Institute of Food and Agriculture.

## REFERENCES

- Amthor, J. S. (2001). Effects of atmospheric CO<sub>2</sub> concentration on wheat yield: a review of results from experiments using various approaches to control CO<sub>2</sub> concentration. *Field Crop. Res.* 73, 1–34.
- Bennett, A. C., and Adams, F. (1970). Concentration of NH<sub>3</sub>(aq) required for incipient NH<sub>3</sub> toxicity to seedlings. *Soil Sci. Soc. Am. Proc.* 34, 259–263.
- Bloom, A. J. (1997). “Nitrogen as a limiting factor: crop acquisition of ammonium and nitrate,” in *Agricultural Ecology*, ed L. E. Jackson (San Diego, CA: Academic Press), 145–172.
- Bloom, A. J., Asensio, J. S. R., Randall, L., Rachmilevitch, S., Cousins, A. B., and Carlisle, E. A. (2012). CO<sub>2</sub> enrichment inhibits shoot nitrate assimilation in C<sub>3</sub> but not C<sub>4</sub> plants and slows growth under nitrate in C<sub>3</sub> plants. *Ecology* 93, 355–377.
- Bloom, A. J., Burger, M., Asensio, J. S. R., and Cousins, A. B. (2010). Carbon dioxide enrichment inhibits nitrate assimilation in wheat and *Arabidopsis*. *Science* 328, 899–903.
- Bloom, A. J., Caldwell, R. M., Finazzo, J., Warner, R. L., and Weissbart, J. (1989). Oxygen and carbon dioxide fluxes from barley shoots dependent on nitrate assimilation. *Plant Physiol.* 91, 352–356.
- Bloom, A. J., Chapin, F. S. III, and Mooney, H. A. (1985). Resource limitation in plants – an economic analogy. *Annu. Rev. Ecol. Syst.* 16, 363–392.
- Bloom, A. J., Smart, D. R., Nguyen, D. T., and Searles, P. S. (2002). Nitrogen assimilation and growth of wheat under elevated carbon dioxide. *Proc. Natl. Acad. Sci. U.S.A.* 99, 1730–1735.
- Bock, B. R. (1987). Increases in maximum yield of spring wheat by maintaining relatively high ammonium/nitrate ratios in soils. *J. Fertil. Issues* 4, 68–72.
- Britto, D. T., and Kronzucker, H. J. (2002). NH<sub>4</sub><sup>+</sup> toxicity in higher plants: a critical review. *J. Plant Physiol.* 159, 567–584.
- Cakmak, I., Pfeiffer, W. H., and McClafferty, B. (2010). Biofortification of durum wheat with zinc and iron. *Cereal Chem.* 87, 10–20.
- Camberato, J. J., and Bock, B. R. (1990). Spring wheat response to enhanced ammonium supply: I dry matter and nitrogen content. *Agron. J.* 82, 463–467.
- Chaudhuri, U. N., Kirkham, M. B., and Kanemasu, E. T. (1990). Root growth of winter wheat under elevated carbon dioxide and drought. *Crop Sci.* 30, 853–857.

- Chen, J.-G., Cheng, S.-H., Cao, W., and Zhou, X. (1998). Involvement of endogenous plant hormones in the effect of mixed nitrogen source on growth and tillering of wheat. *J. Plant Nutr.* 21, 87–97.
- Cousins, A. B., and Bloom, A. J. (2004). Oxygen consumption during leaf nitrate assimilation in a C<sub>3</sub> and C<sub>4</sub> plant: the role of mitochondrial respiration. *Plant Cell Environ.* 27, 1537–1545.
- Cox, W. J., and Reisenauer, H. M. (1973). Growth and ion uptake by wheat supplied nitrogen as nitrate or ammonium or both. *Plant Soil* 38, 363–380.
- Epstein, E., and Bloom, A. J. (2005). *Mineral Nutrition of Plants: Principles and Perspectives*, 2nd Edn. Sunderland, MA: Sinauer Associates.
- Fangmeier, A., De Temmerman, L., Mortensen, L., Kemp, K., Burke, J., Mitchell, R., van Oijen, M., and Weigel, H.-J. (1999). Effects on nutrients and on grain quality in spring wheat crops grown under elevated CO<sub>2</sub> concentrations and stress condition in the European multiple-site experiment 'ESPACe-wheat'. *Eur. J. Agron.* 10, 215–229.
- Fangmeier, A., Grütters, U., Hertstein, U., Sandhage-Hofmann, A., Vermehren, B., and Jäger, H.-J. (1996). Effects of elevated CO<sub>2</sub>, nitrogen supply, and tropospheric ozone on spring wheat. I. Growth and yield. *Environ. Pollut.* 91, 381–390.
- Fangmeier, A., Grütters, U., Högy, P., Vermehren, B., and Jäger, H.-J. (1997). Effects of elevated CO<sub>2</sub>, nitrogen supply, and tropospheric ozone on spring wheat – II. Nutrients (N, P, K, S, Ca, Mg, Fe, Mn, Zn). *Environ. Pollut.* 96, 43–59.
- Findenegg, G. R. (1987). A comparative study of ammonium toxicity at different constant pH of the nutrient solution. *Plant Soil* 103, 239–243.
- Garten, C. T. (1976). Correlations between concentrations of elements in plants. *Nature* 261, 686–688.
- Gashaw, L., and Mugwira, L. M. (1981). Ammonium-N and nitrate-N effects on the growth and mineral compositions of Triticale, wheat, and rye. *Agron. J.* 73, 47–51.
- Gentry, L. E., Wang, X. T., and Below, F. E. (1989). Nutrient uptake by wheat seedlings that differ in response to mixed N nutrition. *J. Plant Nutr.* 12, 363–373.
- Goyal, S. S., and Huffaker, R. C. (1984). "Nitrogen toxicity in plants," in *Nitrogen in Crop Production*, ed. R. D. Hauck (Madison, WI: ASA, CSSA, SSSA), 97–118.
- Guttieri, M. J., Peterson, K. M., and Souza, E. J. (2006). Mineral distributions in milling fractions of low phytic acid wheat. *Crop Sci.* 46, 2692–2698.
- Haug, W., and Lantzsch, H.-J. (1983). Sensitive method for the rapid determination of phytate in cereals and cereal products. *J. Sci. Food Agric.* 34, 1423–1426.
- Havelka, U. D., Wittenbach, V. A., and Boyle, M. G. (1984). CO<sub>2</sub>-enrichment effects on wheat yield and physiology. *Crop Sci.* 24, 1163–1168.
- Haynes, R. J. (1986). "Uptake and assimilation of mineral nitrogen by plants," in *Mineral Nitrogen in the Plant-Soil System*, ed. R. J. Haynes (Orlando, FL: Academic Press), 303–378.
- Heberer, J. A., and Below, F. E. (1989). Mixed nitrogen nutrition and productivity of wheat growing in hydroponics. *Ann. Bot.* 63, 613–619.
- Högy, P., and Fangmeier, A. (2008). Effects of elevated CO<sub>2</sub> on grain quality of wheat. *J. Cereal Sci.* 48, 580–591.
- Högy, P., Wieser, H., Köhler, P., Schwadorf, K., Breuer, J., Franzaring, J., Muntiferung, R., and Fangmeier, A. (2009). Effects of elevated CO<sub>2</sub> on grain yield and quality of wheat: results from a 3-year free-air CO<sub>2</sub> enrichment experiment. *Plant Biol.* 11(Suppl. 1), 60–69.
- Jordan, D. B., and Ogren, W. L. (1983). Species variation in kinetic properties of ribulose 1,5-bisphosphate carboxylase/oxygenase. *Arch. Biochem. Biophys.* 227, 425–433.
- Kruse, J., Hetzger, I., Mai, C., Polle, A., and Rennenberg, H. (2003). Elevated pCO<sub>2</sub> affects N-metabolism of young poplar plants (*Populus tremula* x *P. alba*) differently at deficient and sufficient N-supply. *New Phytol.* 157, 65–81.
- Loladze, I. (2002). Rising atmospheric CO<sub>2</sub> and human nutrition: toward globally imbalanced plant stoichiometry? *Trends Ecol. Evol. (Amst.)* 17, 457–461.
- Magalhaes, J. S., and Wilcox, G. E. (1984). Ammonium toxicity development in tomato plants relative to nitrogen form and light intensity. *J. Plant Nutr.* 7, 1477–1496.
- Manderscheid, R., Bender, J., Jäger, H.-J., and Weigel, H. J. (1995). Effects of season long CO<sub>2</sub> enrichment on cereals. II. Nutrient concentrations and grain quality. *Agric. Ecosyst. Environ.* 54, 175–185.
- Manoj-Kumar, I. (2011). Will rising CO<sub>2</sub> affect Zn and Fe bioavailability in food grains? *Curr. Sci.* 101, 467.
- McKee, I. F., Bullimore, J. F., and Long, S. P. (1997). Will elevated CO<sub>2</sub> concentration protect the yield of wheat from O<sub>3</sub> damage? *Plant Cell Environ.* 20, 77–84.
- Miller, L. V., Krebs, N. F., and Hambidge, K. M. (2007). A mathematical model of zinc absorption in humans as a function of dietary zinc and phytate. *J. Nutr.* 137, 135–141.
- Natali, S. M., Sañudo-Wilhelmy, S. A., and Lerdau, M. T. (2009). Plant and soil mediation of elevated CO<sub>2</sub> impacts on trace metals. *Ecosystems* 12, 715–727.
- Raboy, V. (1997). "Accumulation and storage of phosphate and minerals," in *Cellular and Molecular Biology of Plant Seed Development*, eds B. A. Larkin and I. K. Vasil (Dordrecht: Kluwer Academic Publishers), 441–477.
- Raboy, V. (2009). Approaches and challenges to engineering seed phytate and total phosphorus. *Plant Sci.* 177, 281–296.
- Simpson, R. J., Lambers, H., and Dalling, M. J. (1983). Nitrogen redistribution during grain growth in wheat (*Triticum aestivum* L.). *Plant Physiol.* 71, 7–14.
- Smirnov, N., and Stewart, G. R. (1987). Glutamine synthetase and ammonium assimilation in roots of zinc-tolerant and non-tolerant clones of *Descampsia cespitosa* (L.) Beauv. and *Anthoxanthum odoratum* L. *New Phytol.* 107, 659–670.
- Taub, D. R., Miller, B., and Allen, H. O. (2008). Effects of elevated CO<sub>2</sub> on the protein concentration of food crops: a meta-analysis. *Glob. Change Biol. Bioenergy* 14, 565–575.
- Thompson, G. B., and Woodward, F. I. (1994). Some influences of CO<sub>2</sub> enrichment, nitrogen nutrition, and competition on grain yield and quality in spring wheat and barley. *J. Exp. Bot.* 45, 937–942.
- Wang, X. T., and Below, F. E. (1992). Root growth, nitrogen uptake, and tillering of wheat induced by mixed-nitrogen source. *Crop Sci.* 32, 997–1002.
- Wang, X. T., and Below, F. E. (1995). Tillering, nutrient accumulation, and yield of winter wheat as influenced by nitrogen form. *J. Plant Nutr.* 18, 1177–1189.
- Wang, X. T., and Below, F. E. (1996). Cytokinins in enhanced growth and tillering of wheat induced by mixed-nitrogen source. *Crop Sci.* 36, 121–126.
- Wang, X. T., and Below, F. E. (1998). Accumulation and partitioning of mineral nutrients in wheat as influenced by nitrogen form. *J. Plant Nutr.* 21, 49–61.
- Wechsung, G., Wechsung, F., Wall, G. W., Adamsen, F. J., Kimball, B. A., Garcia, R. L., Pinter, J. R. Jr., and Kartschall, T. (1995). Biomass and growth rate of a spring wheat root system grown in free-air CO<sub>2</sub> enrichment (FACE) and ample soil moisture. *J. Biogeogr.* 22, 623–634.
- Weigel, H. J., Manderscheid, R., Jäger, H.-J., and Mejer, G. J. (1994). Effects of season-long CO<sub>2</sub> enrichment on cereals. I. Growth performance and yield. *Agric. Ecosyst. Environ.* 48, 231–240.
- Welch, R. M., and Graham, R. D. (2004). Breeding for micronutrients in staple food crops from a human nutrition perspective. *J. Exp. Bot.* 55, 353–364.
- Wu, D.-X., Wang, G.-X., Bai, Y.-F., and Liao, J.-X. (2004). Effects of elevated CO<sub>2</sub> concentration on growth, water use, yield, and grain quality of wheat under two soil water levels. *Agric. Ecosyst. Environ.* 104, 493–507.
- Wuehler, S. E., Peerson, J. M., and Brown, K. H. (2005). Use of national food balance data to estimate the adequacy of zinc in national food supplies: methodology and regional estimates. *Public Health Nutr.* 8, 812–819.

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 03 July 2012; paper pending published: 23 July 2012; accepted: 07 August 2012; published online: 03 September 2012.

Citation: Carlisle E, Myers S, Raboy V and Bloom A (2012) The effects of inorganic nitrogen form and CO<sub>2</sub> concentration on wheat yield and nutrient accumulation and distribution. *Front. Plant Sci.* 3:195. doi: 10.3389/fpls.2012.00195

This article was submitted to *Frontiers in Plant Nutrition*, a specialty of *Frontiers in Plant Science*.

Copyright © 2012 Carlisle, Myers, Raboy and Bloom. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in other forums, provided the original authors and source are credited and subject to any copyright notices concerning any third-party graphics etc.