

Photorespiration plays an important role in the regulation of photosynthetic electron flow under fluctuating light in tobacco plants grown under full sunlight

Wei Huang^{1,2}, Hong Hu^{1,2} and Shi-Bao Zhang^{1,2*}

¹ Key Laboratory of Economic Plants and Biotechnology, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming, China, ² Yunnan Key Laboratory for Wild Plant Resources, Kunming, China

OPEN ACCESS

Edited by:

Richard Sayre,
New Mexico Consortium at Los
Alamos National Labs, USA

Reviewed by:

Robert L. Houtz,
University of Kentucky, USA
Nabil I. Elsheery,
Tanta University, Egypt

*Correspondence:

Shi-Bao Zhang,
Key Laboratory of Economic Plants
and Biotechnology, Kunming Institute
of Botany, Chinese Academy of
Sciences, 132# Lanhei Road,
Heilongtan, Kunming 650201,
Yunnan, China
sbzhang@mail.kib.ac.cn

Specialty section:

This article was submitted to
Plant Physiology,
a section of the journal
Frontiers in Plant Science

Received: 14 May 2015

Accepted: 27 July 2015

Published: 11 August 2015

Citation:

Huang W, Hu H and Zhang S-B (2015)
Photorespiration plays an important
role in the regulation of photosynthetic
electron flow under fluctuating light in
tobacco plants grown under full
sunlight. *Front. Plant Sci.* 6:621.
doi: 10.3389/fpls.2015.00621

Plants usually experience dynamic fluctuations of light intensities under natural conditions. However, the responses of mesophyll conductance, CO₂ assimilation, and photorespiration to light fluctuation are not well understood. To address this question, we measured photosynthetic parameters of gas exchange and chlorophyll fluorescence in tobacco leaves at 2-min intervals while irradiance levels alternated between 100 and 1200 μmol photons m⁻² s⁻¹. Compared with leaves exposed to a constant light of 1200 μmol photons m⁻² s⁻¹, both stomatal and mesophyll conductances were significantly restricted in leaves treated with fluctuating light condition. Meanwhile, CO₂ assimilation rate and electron flow devoted to RuBP carboxylation at 1200 μmol photons m⁻² s⁻¹ under fluctuating light were limited by the low chloroplast CO₂ concentration. Analysis based on the C₃ photosynthesis model indicated that, at 1200 μmol photons m⁻² s⁻¹ under fluctuating light, the CO₂ assimilation rate was limited by RuBP carboxylation. Electron flow devoted to RuBP oxygenation at 1200 μmol photons m⁻² s⁻¹ under fluctuating light remained at nearly the maximum level throughout the experimental period. We conclude that fluctuating light restricts CO₂ assimilation by decreasing both stomatal and mesophyll conductances. Under such conditions, photorespiration plays an important role in the regulation of photosynthetic electron flow.

Keywords: CO₂ assimilation, fluctuating light, photorespiration, photosynthetic electron flow, mesophyll conductance

Introduction

In nature, plants grown in open habitats usually experience changes in light intensities because of clouds. Even on clear days, the leaves of understory plants are frequently exposed to short-term fluctuating light levels due to movements by the leaves and stems of other plants above them. To cope with fluctuating light conditions, plants must regulate their photosynthetic processes.

Abbreviations: A_n, CO₂ assimilation rate; C_c, chloroplast CO₂ concentration; F_v/F_m' , the maximum quantum yield of PSII after light adaptation; C_{trans}, the chloroplast CO₂ concentration at which the transition from RuBP carboxylation to RuBP regeneration occurred; Φ_{PSII}, effective quantum yield of PSII; g_m, mesophyll conductance; g_s, stomatal conductance; J_C, electron flow devoted to RuBP carboxylation; J_O, electron flow devoted to RuBP oxygenation; J_T, total electron flow through PSII; J_{max}, the maximum rate of RuBP regeneration; NPQ, non-photochemical quenching; PSI, photosystem I; PSII, photosystem II; qP, coefficient of PSII photochemical quenching; V_{Cmax}, the maximum rate of RuBP carboxylation.

Under constant low light, most of the absorbed light energy can be used to drive photosynthesis, even when stomatal conductance (g_s) is reduced. Under constant high light, the rate of CO_2 assimilation (A_n) is maintained at an elevated level due to high g_s and mesophyll conductance (g_m) (Yamori et al., 2010, 2011). Fluctuating light restricts both g_s and CO_2 assimilation rate (Fay and Knapp, 1993; Kirschbaum et al., 1998). However, it is unclear how g_m and photorespiration respond to those changes in irradiance.

Under natural conditions, g_m is an important determinant of the CO_2 assimilation rate, especially at high light levels (Carriquí et al., 2015). Several environmental factors, such as water status and temperature, can affect g_m (Flexas et al., 2002; Scafaro et al., 2011; Walker et al., 2013). For tobacco (*Nicotiana tabacum*) plants grown with adequate water and optimum temperature, g_m is mainly dependent upon the growth light intensity (Yamori et al., 2010). Plants exposed to strong light have higher values of g_m when compared with those grown under low light. When light levels are constant, g_m does not appear to be dependent upon light intensity (Yamori et al., 2010). However, the effect of fluctuating light condition on g_m is unclear. According to the model of Farquhar et al. (1980), CO_2 assimilation in C_3 plants is limited by either the carboxylation or the regeneration of ribulose-1,5-bisphosphate (RuBP). In tobacco, a model C_3 plant, the rate of CO_2 assimilation under high light is influenced by leaf nitrogen (N) content. CO_2 assimilation rate under high light tends to be limited by RuBP regeneration for plants grown at high N concentration (Yamori et al., 2010, 2011). However, that presumption is based on high values of g_s and g_m . Once g_s and g_m decrease because of environmental stresses such as drought, CO_2 assimilation rate is then partially constrained by RuBP carboxylation (Flexas et al., 2002; Flexas and Medrano, 2002). Therefore, if fluctuating light levels restrict g_m , then A_n likely tends to be limited by RuBP carboxylation.

Photorespiration, an inevitable process in photosynthesis, plays a supporting role in photosynthetic CO_2 assimilation (Timm et al., 2012; Busch et al., 2013; Weber and Bauwe, 2013). This process is initiated by the oxygenation of RuBP, in which one molecule of glycolate-2-phosphate and one molecule of glycerate-3-phosphate are produced (Ogren, 1984). Although glycolate-2-phosphate cannot be used by plants for biosynthetic reactions, and is also a potential inhibitor of chloroplast functioning (Anderson, 1971), it can be converted into glycerate-3-phosphate through the photorespiratory pathway (Leegood et al., 1995). When g_s and g_m are diminished, the decreased chloroplast CO_2 concentration increases the specificity of Rubisco to O_2 and then induces a rise in the rate of RuBP oxygenation (Wingler et al., 1999, 2000).

Plants avoid those detrimental effects of glycolate-2-phosphate and other photorespiratory intermediates by activating the photorespiratory pathway when chloroplast CO_2 concentration is low. In *Arabidopsis thaliana* plants grown under low irradiance, photorespiration plays a minor role in regulating photosynthetic electron flow after exposure to short-term fluctuating light (Kono et al., 2014). The growth light intensity

significantly affects the development of the photorespiratory pathway (Huang et al., 2014). For example, plants such as tobacco grown under bright light have a greater capacity than those under low light (Huang et al., 2014). However, little is known about how the photorespiratory pathway functions in the acclimation to fluctuating light by plants grown under high light. Because this pathway is critical to the control of A_n and photosynthetic electron flow (Takahashi et al., 2007; Timm et al., 2012; Huang et al., 2014), it is important that research focused on photosynthetic regulation under fluctuating light should include growth light intensity as an experimental variable.

In this study, we measured the photosynthetic parameters of gas exchange and chlorophyll fluorescence to investigate the responses of g_m , A_n , and photosynthetic electron flow to fluctuations in light levels. We also examined the limiting step of CO_2 assimilation and the role of the photorespiratory pathway in modulating photosynthetic electron flow under alternating light conditions. Our objective was to improve our understanding of how photosynthesis is regulated when sun-grown plants are exposed to changes in irradiance. The following questions were addressed: (1) Is g_m restricted by fluctuating light? (2) What is the limiting step of A_n under fluctuating light? and (3) Does the photorespiratory pathway play an important role in regulating photosynthetic electron flow under fluctuating light?

Materials and Methods

Plant Materials and Growing Conditions

Following seed germination, seedlings of tobacco cv. y87 were cultivated in phytotron for 7 weeks. Afterwards, they were grown in plastic pots in an open field at Kunming Institute of Botany, Yunnan, China (elevation 1900 m, 102°41'E, 25°01'N). During our experiment period (10 May to 24 June 2013), none of the plants experienced any water or nutrient stresses. The average temperature at Kunming was 20.9°C in May and 20.6°C in June. Fully expanded mature leaves on 13-week-old plants were used for photosynthetic measurements.

Analyses of Gas Exchange, Chlorophyll Fluorescence, and Mesophyll Conductance

Photosynthetic parameters for gas exchange and chlorophyll fluorescence were monitored with an open gas exchange system that incorporated infrared CO_2 and water vapor analyzers (Li-6400XT; Li-Cor Biosciences, Lincoln, NE, USA) and a 2-cm² measuring head (6400-40 Leaf Chamber Fluorometer; Li-Cor Biosciences). Measurements were made in a phytotron where relative air humidity (60%) and air temperature (25°C) were controlled. The atmospheric CO_2 concentration was maintained at 400 $\mu\text{mol mol}^{-1}$ by the Li-6400XT. To generate a light response curve, we initially exposed the mature leaves to strong irradiance (2000 $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$) for 20 min to obtain steady, high levels of g_s and CO_2 assimilation. Afterward, photosynthetic parameters were evaluated at 2-min intervals at photosynthetic photon flux densities (PPFDs) of 2000, 1600, 1200, 800, 500, 300, 200, 100, 50, 20, or 0 $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$. To investigate the responses of g_s , g_m , CO_2 assimilation

rate, and photosynthetic electron flow to fluctuating light, we also evaluated those photosynthetic parameters under light levels that alternated every 2 min between 100 and 1200 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ after dark-adaptation for 30 min. Photosynthetic induction curves were also developed at 1200 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ after 30 min of darkness. Values for those parameters were recorded automatically by the Li-6400XT at 2-min intervals.

The CO_2 assimilation rate vs. chloroplast CO_2 concentration (C_c) was examined at 1200 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ (von Caemmerer and Farquhar, 1981). For each A_n/C_c curve, the photosynthetic rate reached a steady state at 400 $\mu\text{mol mol}^{-1} \text{CO}_2$, then decreased to a lower limit of 50 $\mu\text{mol mol}^{-1}$ before increasing stepwise to an upper limit of 1600 $\mu\text{mol mol}^{-1}$. Each stepwise measurement was completed within 2–3 min. Using those A_n/C_c curves, we calculated the maximum rates of RuBP regeneration (J_{max}) and RuBP carboxylation (V_{cmax}) according to the method of Long and Bernacchi (2003).

The fluorescence parameters F_o' , F_m' , and F_s were evaluated as previously described in Baker and Rosenqvist (2004). Here, F_o' and F_m' represented the minimum and maximum fluorescence after light-adaptation, respectively. F_s indicated the light-adapted steady-state fluorescence. The maximum quantum yield of PSII after light adaptation (F_v'/F_m') was calculated as $(F_m' - F_o')/F_m'$. Coefficient of PSII photochemical quenching (qP) was calculated as $(F_m' - F_s)/(F_m' - F_o')$. Effective quantum yield of PSII (Φ_{PSII}) was calculated as $(F_m' - F_s)/F_m'$ (Genty et al., 1989).

Total photosynthetic electron flow through PSII was calculated as $J_T = \Phi_{\text{PSII}} \times \text{PPFD} \times L_{\text{abs}} \times 0.5$ (Krall and Edwards, 1992), where L_{abs} represented leaf absorbance and was assumed to be 0.85 for sun-grown tobacco leaves that receive high-nitrogen nutrition (Miyake et al., 2005). The constant of 0.5 was applied based on the assumption that photons were equally distributed between photosystem I (PSI) and PSII (Miyake et al., 2005). Following the assumption that the water–water cycle is not a major alternative electron sink when CO_2 assimilation is limited (Driever and Baker, 2011), we allocated the electron flow through PSII to RuBP carboxylation (J_C) and oxygenation (J_O). Values for J_C and J_O were estimated according to the method of Valentini et al. (1995):

$$J_O = 2/3 \times (J_T - 4 \times (A_n + R_d))$$

$$J_C = 1/3 \times (J_T + 8 \times (A_n + R_d))$$

where A_n was the net rate of CO_2 assimilation and R_d represented the rate of mitochondrial respiration as measured after 30 min of dark-adaptation.

We recorded values for mesophyll conductance (g_m) at 1200 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ after plants were exposed to either fluctuating or constant light for 60 min. For our comparisons, g_m was also estimated at 1200 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ in light response curves. Values for g_m were estimated through a combination analysis of gas exchange and chlorophyll fluorescence, and according to the following equation (Harley et al., 1992; Loreto et al., 1992; Warren and Dreyer, 2006; Yamori et al., 2010, 2011):

$$g_m = \frac{A_n}{C_i - \Gamma^*(J_T + 8(A_n + R_d))/(J_T - 4(A_n + R_d))}$$

where A_n was the net rate of CO_2 assimilation, C_i was the intercellular CO_2 concentration, J_T was total photosynthetic electron flow through PSII, R_d was the rate of mitochondrial respiration, and Γ^* was the CO_2 compensation point in the absence of daytime respiration (Farquhar et al., 1980; Brooks and Farquhar, 1985), with the latter assumed to be 32.2 at 25°C (Long and Bernacchi, 2003). Using the estimated g_m , we calculated the chloroplast CO_2 concentration with the following equation (Long and Bernacchi, 2003; Warren and Dreyer, 2006; Yamori et al., 2010, 2011):

$$C_c = C_i - \frac{A_n}{g_m}$$

where C_i was the intercellular CO_2 concentration, A_n was the net rate of CO_2 assimilation, and g_m was mesophyll conductance. To identify the limiting step of CO_2 assimilation under fluctuating light, we applied the method of Yamori et al. (2010, 2011) to determine C_{trans} , the chloroplast CO_2 concentration at which the transition from RuBP carboxylation to RuBP regeneration occurred:

$$C_{\text{trans}} = \frac{K_c(1 + OK_o)J_{\text{max}}/4V_{\text{cmax}} - 2\Gamma^*}{1 - J_{\text{max}}/4V_{\text{cmax}}}$$


where K_c ($\mu\text{mol mol}^{-1}$) and K_o (mmol mol^{-1}) were the Michaelis constants for CO_2 and O_2 , respectively (Farquhar et al., 1980), and were assumed to be 406.7 $\mu\text{mol mol}^{-1}$ and 277 mmol mol^{-1} at 25°C, respectively (Long and Bernacchi, 2003); J_{max} was the maximum rate of RuBP regeneration; V_{cmax} was the maximum rate of RuBP carboxylation; and Γ^* was the CO_2 compensation point in the absence of daytime respiration. The limiting step of CO_2 assimilation was then determined by comparing the values of C_c and C_{trans} .

Statistical Analysis

The results were displayed as mean values of four independent measurements. We used One-Way ANOVA and SPSS 16.0 software (SPSS Inc., Chicago, IL, USA) to examine differences among treatments involving fluctuating vs. constant light. Those differences were considered significant at $P < 0.05$.

Results


Light response curves indicated that g_s was maintained at high levels ($>0.3 \text{ mol m}^{-2} \text{s}^{-1}$) when plants were exposed to light intensities above 100 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ (Figure 1A). When light levels were reduced from 100 to 0 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$, values for g_s decreased sharply from 0.29 to 0.14 $\text{mol m}^{-2} \text{s}^{-1}$ within 6 min (Figure 1A). This indicated that stomatal conductance in sun-grown tobacco leaves is very sensitive to light intensity in sun-grown tobacco leaves. Under strong irradiance, i.e., 1500 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$, A_n was 28.5 $\mu\text{mol CO}_2 \text{ m}^{-2} \text{s}^{-1}$ (Figure 1B). At levels below 1500 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$, values for J_T , J_C , J_O , and J_O/J_C gradually rose with increasing PPFD, peaking at 233 $\mu\text{mol electrons m}^{-2} \text{s}^{-1}$,




164 $\mu\text{mol electrons m}^{-2} \text{s}^{-1}$, 69 $\mu\text{mol electrons m}^{-2} \text{s}^{-1}$, and 0.43, respectively (Figures 1C,D).

Fluctuating light conditions significantly restricted the opening of stomata. After plants were alternately exposed to 100 and 1200 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ every 2 min for 60 min, g_s was 0.18 $\text{mol m}^{-2} \text{s}^{-1}$ (Figure 2A). However, when plants were illuminated at a constant 1200 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ for 60 min, g_s was 0.30 $\text{mol m}^{-2} \text{s}^{-1}$. After 60 min of fluctuating light, the CO_2 assimilation rate at 1200 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ was 18.1 $\mu\text{mol CO}_2 \text{ m}^{-2} \text{s}^{-1}$ vs. 25.4 $\mu\text{mol CO}_2 \text{ m}^{-2} \text{s}^{-1}$ after exposure to 1200 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ for 60 min (Figure 2B). Those values for g_s and A_n differed significantly between constant and fluctuating-light treatments, demonstrating that the latter condition inhibited g_s as well as CO_2 assimilation. This finding was consistent with those reported previously (Fay and Knapp, 1993; Kirschbaum et al., 1998).

By contrast, values for qP at 1200 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ differed only slightly between the constant and fluctuating light treatments (Figure 3A), while F_v'/F_m' and Φ_{PSII} at 1200 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ were significantly lower under fluctuating light ($P < 0.001$; Figures 3B,C). The parameter F_v'/F_m' represents the maximum efficiency of PSII when all reaction centers are “open,” and qP is the factor that relates maximum PSII efficiency to the operating PSII efficiency (Farage et al., 2006). Because Φ_{PSII} is the product of qP and F_v'/F_m' , the difference in Φ_{PSII} that we found between fluctuating light and constant light resulted from the change in F_v'/F_m' . These results suggested that although fluctuating light had little effect on the coefficient of PSII photochemical quenching, it induced a significant decline in the maximum efficiency of PSII.




After 60 min of treatment, total electron flow through PSII (J_T) at 1200 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ was significantly higher under constant illumination than under fluctuating light, i.e.,




225 vs. 185 $\mu\text{mol electrons m}^{-2} \text{s}^{-1}$, respectively (**Figure 4A**). During that time period, the value for electron flow devoted to RuBP oxygenation (J_O) at 1200 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ changed only slightly between constant- and fluctuating-light treatments (**Figure 4B**). By contrast, electron flow devoted to RuBP carboxylation (J_C) at 1200 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ was higher under constant light (151 $\mu\text{mol electrons m}^{-2} \text{s}^{-1}$) than under fluctuating light (115 $\mu\text{mol electrons m}^{-2} \text{s}^{-1}$) (**Figure 4C**). Consequently, the ratio J_O/J_C at 1200 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ was higher for plants treated with fluctuating light because of the lower value for J_C (**Figure 4D**). These results indicated that fluctuations in irradiance levels suppressed photosynthetic electron flow, primarily by restricting electron flow devoted to RuBP carboxylation. By comparison, electron flow devoted to RuBP oxygenation was hardly affected by fluctuating light conditions.

After pooling the photosynthesis data collected at 1200 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ under fluctuating light, we determined that g_s was linearly and positively correlated with A_n , J_T , and J_C





(**Figures 5A,B**). We found it interesting that J_O was independent of g_s (**Figure 5B**), which implied that RuBP carboxylation and RuBP oxygenation responded differently to g_s . Under fluctuating light, J_O remained at nearly the maximum level throughout the experimental period. In the initial stage of fluctuating light treatment, electron flow attributed to RuBP oxygenation contributed largely to the total electron transport through PSII (**Figure 5C**).

To analyze the limiting step of CO_2 assimilation under fluctuating light, we examined the relationship between photosynthesis and chloroplast CO_2 concentration. Here, the



ratio of the maximum rate of RuBP regeneration (J_{\max}) to that of RuBP carboxylation (V_{cmax}) was 0.92, and the chloroplast CO_2 concentration at which the transition from RuBP carboxylation to RuBP regeneration occurred (C_{trans}) was $135 \mu\text{mol mol}^{-1}$ (**Figure 6**). After exposure to fluctuating light conditions for 60 min, g_m at $1200 \mu\text{mol photons m}^{-2} \text{s}^{-1}$ was $0.21 \text{ mol m}^{-2} \text{s}^{-1}$, which was significantly lower than that found with light curves ($0.29 \text{ mol m}^{-2} \text{s}^{-1}$) or under constant light ($0.28 \text{ mol m}^{-2} \text{s}^{-1}$) (**Figure 7**). For the light curves, C_c at $1200 \mu\text{mol photons m}^{-2} \text{s}^{-1}$ was $154 \mu\text{mol mol}^{-1}$. After exposure to fluctuating light or constant light for 60 min, the value for C_c was 105 or $131 \mu\text{mol mol}^{-1}$, respectively. This indicated that fluctuating light not only decreased g_s but also restricted g_m , leading to a decline in C_c . Because C_c was significantly lower than C_{trans} ($P < 0.0001$), the rate of CO_2 assimilation at $1200 \mu\text{mol photons m}^{-2} \text{s}^{-1}$ under fluctuating light was limited by RuBP carboxylation.



Discussion

The limiting step of A_n is mainly determined by the relative values of C_{trans} and C_c . For tobacco plants supplied with high concentrations of nitrogen, photosynthesis tends to be limited by RuBP regeneration because C_c is higher than C_{trans} (Yamori et al., 2010, 2011). Such previous conclusions have been based on experiments that involved high levels of g_s and g_m under constant strong light. Although g_s and A_n can be significantly inhibited under fluctuating light, the limiting step of A_n under such conditions has been unknown. Here, our results indicate that both g_s and g_m are significantly restricted under fluctuating light, leading to the decrease in C_c . After exposure to fluctuating light for 60 min, C_c was 105 vs. $135 \mu\text{mol mol}^{-1}$

for C_{trans} . Those data provided evidence that, at $1200 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$, the photosynthetic process is limited by RuBP carboxylation under fluctuating light. Meanwhile, the high activation of photorespiration contributed largely to the regulation of photosynthetic electron flow.

Carriqui et al. (2015) have demonstrated that g_m plays an important role in determining the CO_2 assimilation rate, especially at high light intensities. Nevertheless, the response of g_m to light intensity remains controversial. For example, in sclerophylls such as *Banksia integrifolia*, *B. serrata*, and *B. paludosa*, the average g_m under ambient CO_2 concentration is 22% lower at 500 than at $1500 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$ (Hassiotou et al., 2009). However, the average g_m calculated for wheat leaves is not affected by light intensity (Tazoe et al., 2009). In tobacco leaves, g_m is significantly lower at 250 than at $1000 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$ (Flexas et al., 2007). By contrast, Yamori et al. (2010) have shown that g_m differs little between constant high light and constant low light in tobacco leaves. Our results indicated that g_m at $1200 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$ under the fluctuating light was 25% lower than the level calculated at constant light of $1200 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$ (Figure 7). We believe that this difference was caused by the use of a low light regime ($100 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$) in fluctuating light. We found that, under fluctuating light, g_m was regulated by both high and low light levels; i.e., although the former induced an increase in g_m , this effect could be partially reversed when plants were then exposed to reduced irradiance.

According to the photosynthesis model of Farquhar et al. (1980), CO_2 assimilation in C_3 plants is constrained by RuBP carboxylation and/or RuBP regeneration. Therefore, based on that model, the limiting step can be altered in two ways: (1) adjustments in the balance between the maximum rates of RuBP regeneration and RuBP carboxylation, or (2) changes in the chloroplast CO_2 concentration (Hikosaka et al., 2006; Yamori et al., 2011). For example, in research with tobacco plants, Yamori et al. (2011) have reported that the CO_2 assimilation rate at $380 \mu\text{mol mol}^{-1} \text{ CO}_2$ and $1500 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$ (A_{380}) depends upon the leaf-N content and is mainly determined by J_{max}/V_{cmax} . Furthermore, at high leaf-N content, A_{380} is limited by RuBP regeneration due to the low ratio of J_{max}/V_{cmax} (Yamori et al., 2010, 2011). However, those conclusions have been drawn from experiments with plants that had high values for both g_s and g_m , and which did not consider the effects of fluctuating light levels. By comparison, our photosynthetic data for g_s , A_n , J_T , and J_{max}/V_{cmax} ratio are very similar to those that describe the performance of plants grown with a high nitrogen supply (Yamori et al., 2011), indicating that plants grown with high N concentration were used in the present study. Furthermore, CO_2 assimilation rate at $1200 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$ under constant light was limited by RuBP regeneration. When plants were exposed to fluctuating light, the declines in g_s and g_m resulted in a decrease in C_c . The low light regimes under fluctuating light decreased the Rubisco activation state (Yamori et al., 2012), which further restricted the Calvin cycle. The rate of CO_2 assimilation under high light during the fluctuating-light treatment tended to be limited by RuBP carboxylation. Fluctuating light has altered the limiting step of CO_2 assimilation in tobacco plants with high leaf-N content.

Previous studies with *A. thaliana* have investigated the roles of cyclic electron flow (CEF) and O_2 -dependent alternative electron sinks in regulating photosynthetic electron flow under fluctuating light (Suorsa et al., 2012; Kono et al., 2014). It is believed that CEF is essential for proper acclimation of PSI to such light condition (Suorsa et al., 2012). However, the contribution of photorespiration to photodamage under fluctuating light is small in *Arabidopsis* leaves sampled from plants exposed to low light (Kono et al., 2014). In tobacco, the capacity of the photorespiratory pathway is strongly influenced by the growth light intensity, with sun leaves up-regulating this pathway to control CO_2 assimilation and photosynthetic electron flow (Huang et al., 2014). However, it is unknown what role the photorespiratory pathway has in enabling plants normally grown under high light to adapt to fluctuating light conditions. Our data demonstrated that, when plants were exposed to fluctuating light, the reduction in C_c meant that less electron flow could be devoted to RuBP carboxylation. However, we found that the flow devoted to RuBP oxygenation was completely and highly activated under such conditions.

Suppression of CO_2 fixation can cause over-acidification of lumen in the thylakoid membrane, which then activates non-photochemical quenching (NPQ) to dissipate excess light energy harmlessly as heat (Flexas and Medrano, 2002; Takahashi et al., 2007; Huang et al., 2012). At $1200 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$, F_v'/F_m' were lower under fluctuating light than under constant light. Because F_v'/F_m' is inversely related to NPQ, this result was evidence of the higher activation of NPQ under fluctuating light. Furthermore, an increase in the proton gradient across the thylakoid membrane can limit linear electron flow (LEF) via cytochrome b6/f (Tikkanen and Aro, 2014). Consumption of photochemical energy, such as ATP and NADPH, through the photorespiratory pathway is thought to alleviate such over-acidification. Especially in the initial stage of our fluctuating-light period, electron flow that was consumed by the photorespiratory pathway largely contributed to the operation of LEF. Therefore, for tobacco plants grown under full sunlight, photorespiratory pathway would be essential for regulating photosynthetic electron flow under fluctuating light, even though our findings contradict a previous report concerning low-light-grown *A. thaliana* (Kono et al., 2014).

Although photorespiratory intermediates such as glycine and glycerate inhibit the Calvin cycle (Chastain and Ogren, 1989; Eisenhut et al., 2007; Timm et al., 2012), they can be converted to glycerate-3-phosphate through the photorespiratory pathway (Peterhansel and Maurino, 2011). This process is critical for photosynthesis and photoprotection (Takahashi et al., 2007). Under fluctuating light, a reduction in C_c will accelerate RuBP oxygenation and, ultimately, the production of those intermediates. If the photorespiratory pathway is maintained at a low level under such conditions, the accumulation of those intermediates inhibits CO_2 assimilation as well as photosynthetic electron flow, causing acceleration of photodamage (Chastain and Ogren, 1989; Eisenhut et al., 2007; Takahashi et al., 2007). In plants with a high rate of CO_2 assimilation, rapid acceleration of photorespiratory pathway results in low glycine and glycerate contents (Timm et al., 2012). Therefore, to overcome those

detrimental effects of photorespiratory intermediates, this pathway is highly activated under fluctuating light, which then benefits photosynthetic CO₂ assimilation and photosynthetic electron flow. In addition, the operation of this pathway is necessary for the regeneration of RuBP (Takahashi et al., 2007). To optimize photosynthetic CO₂ fixation, the rates of RuBP oxygenation and RuBP regeneration through photorespiratory pathway must be balanced. Therefore, under fluctuating light conditions, strong activation of the photorespiratory pathway accelerates RuBP regeneration, preventing a decrease in the RuBP pool and favoring the Calvin cycle.

In summary, our results provide evidence that, for sun-grown tobacco leaves, fluctuating light conditions significantly decrease both stomatal and mesophyll conductances, as well as chloroplast CO₂ concentration. Consequently, the rate of

CO₂ assimilation is limited by RuBP carboxylation under such conditions. Meanwhile, the photorespiratory pathway is highly activated to regulate photosynthetic electron flow and benefit photosynthetic CO₂ fixation. Thus, strong activation of this pathway is an important strategy by which sun-grown plants adapt to fluctuating light.

Acknowledgments

This study was supported by the National Natural Science Foundation of China (Grant 31300332), funding from the China Postdoctoral Science Foundation to WH (2014T70892), and the following scientific foundations from the Yunnan Tobacco Academy of Agriculture: 110201101003 (TS-03), 2011YN02, 2011YN03.

References

- Anderson, L. E. (1971). Chloroplast and cytoplasmic enzymes. II. Pea leaf triose phosphate isomerases. *Biochim. Biophys. Acta* 235, 237–244. doi: 10.1016/0005-2744(71)90051-9
- Baker, N. R., and Rosenqvist, E. (2004). Applications of chlorophyll fluorescence can improve crop production strategies: an examination of future possibilities. *J. Exp. Bot.* 55, 1607–1621. doi: 10.1093/jxb/erh196
- Brooks, A., and Farquhar, G. D. (1985). Effect of temperature on the CO₂/O₂ specificity of ribulose-1,5-bisphosphate carboxylase/oxygenase and the rate of respiration in the light. *Planta* 165, 397–406. doi: 10.1007/BF00392238
- Busch, F. A., Sage, T. L., Cousins, A. B., and Sage, R. F. (2013). C₃ plants enhance rates of photosynthesis by re-assimilating photorespired and respired CO₂. *Plant Cell Environ.* 36, 200–212. doi: 10.1111/j.1365-3040.2012.02567.x
- Carriqui, M., Cabrera, M., Conesa, M. A., Coopman, R. E., Douthé, C., Gago, J., et al. (2015). Diffusional limitations explain the lower photosynthetic capacity of ferns as compared with angiosperms in a common garden study. *Plant Cell Environ.* 38, 448–460. doi: 10.1111/pce.12402
- Chastain, C. J., and Ogren, W. L. (1989). Glyoxylate inhibition of ribulosebisphosphate carboxylase/oxygenase activation state *in vivo*. *Plant Cell Physiol.* 30, 937–944.
- Driever, S. M., and Baker, N. R. (2011). The water-water cycle in leaves is not a major alternative electron sink for dissipation of excess excitation energy when CO₂ assimilation is restricted. *Plant Cell Environ.* 34, 837–846. doi: 10.1111/j.1365-3040.2011.02288.x
- Eisenhut, M., Bauwe, H., and Hagemann, M. (2007). Glycine accumulation is toxic for the cyanobacterium *Synechocystis* sp. strain PCC 6803, but can be compensated by supplementation with magnesium ions. *FEMS Microbiol. Lett.* 277, 232–237. doi: 10.1111/j.1574-6968.2007.00960.x
- Farage, P. K., Blowers, D., Long, S. P., and Baker, N. R. (2006). Low growth temperatures modify the efficiency of light use by photosystem II for CO₂ assimilation in leaves of two chilling-tolerant C₄ species, *Cyperus longus* L. and *Miscanthus × giganteus*. *Plant Cell Environ.* 29, 720–728. doi: 10.1111/j.1365-3040.2005.01460.x
- Farquhar, G. D., von Caemmerer, S., and Berry, J. A. (1980). A biochemical model of photosynthetic CO₂ assimilation in leaves of C₃ species. *Planta* 149, 78–90. doi: 10.1007/BF00386231
- Fay, P. A., and Knapp, A. K. (1993). Photosynthetic and stomatal responses of *Avena sativa* (Poaceae) to a variable light environment. *Am. J. Bot.* 80, 1369–1373. doi: 10.2307/2445664
- Flexas, J., Bota, J., Escalona, J. M., Sampol, B., and Medrano, H. (2002). Effects of drought on photosynthesis in grapevines under field conditions: an evaluation of stomatal and mesophyll limitations. *Funct. Plant Biol.* 29, 461–471. doi: 10.1071/PP01119
- Flexas, J., Diaz-Espejo, A., Galmés, J., Kaldenhoff, R., Medrano, H., and Ribas-Carbo, M. (2007). Rapid variations of mesophyll conductance in response to changes in CO₂ concentration around leaves. *Plant Cell Environ.* 30, 1284–1298. doi: 10.1111/j.1365-3040.2007.01700.x
- Flexas, J., and Medrano, H. (2002). Energy dissipation in C₃ plants under drought. *Funct. Plant Biol.* 29, 1209–1215. doi: 10.1071/FP02015
- Genty, B., Briantais, J. M., and Baker, N. R. (1989). The relationship between the quantum yield of photosynthetic electron transport and quenching of chlorophyll fluorescence. *Biochim. Biophys. Acta* 99, 87–92. doi: 10.1016/S0304-4165(89)80016-9
- Harley, P. C., Loreto, F., Marco, G. D., and Sharkey, T. D. (1992). Theoretical considerations when estimating the mesophyll conductance to CO₂ flux by analysis of the response of photosynthesis to CO₂. *Plant Physiol.* 98, 1429–1436. doi: 10.1104/pp.98.4.1429
- Hassiotou, F., Ludwig, M., Renton, M., Veneklaas, E. J., and Evans, J. R. (2009). Influence of leaf dry mass per area, CO₂, and irradiance on mesophyll conductance in sclerophylls. *J. Exp. Bot.* 60, 2303–2314. doi: 10.1093/jxb/erp021
- Hikosaka, K., Ishikawa, K., Borjigidai, A., Muller, O., and Onoda, Y. (2006). Temperature acclimation of photosynthesis: mechanisms involved in the changes in temperature dependence of photosynthetic rate. *J. Exp. Bot.* 57, 291–302. doi: 10.1093/jxb/erj049
- Huang, W., Yang, S. J., Zhang, S. B., Zhang, J. L., and Cao, K. F. (2012). Cyclic electron flow plays an important role in photoprotection for the resurrection plant *Paraboea rufescens* under drought stress. *Planta* 235, 819–828. doi: 10.1007/s00425-011-1544-3
- Huang, W., Zhang, S. B., and Hu, H. (2014). Sun leaves up-regulate the photorespiratory pathway to maintain a high rate of CO₂ assimilation in tobacco. *Front. Plant Sci.* 5:688. doi: 10.3389/fpls.2014.00688
- Kirschbaum, M. U. F., Küppers, M., Schneider, H., Giersch, C., and Noe, S. (1998). Modelling photosynthesis in fluctuating light with inclusion of stomatal conductance, biochemical activation and pools of key photosynthetic intermediates. *Planta* 204, 16–26. doi: 10.1007/s004250050225
- Kono, M., Noguchi, K., and Terashima, I. (2014). Roles of the cyclic electron flow around PSI (CEF-PSI) and O₂-dependent alternative pathways in regulation of the photosynthetic electron flow in short-term fluctuating light in *Arabidopsis thaliana*. *Plant Cell Physiol.* 55, 990–1004. doi: 10.1093/pcp/pcu033
- Krall, J. P., and Edwards, G. E. (1992). Relationship between photosystem II activity and CO₂ fixation in leaves. *Physiol. Plant.* 86, 180–187. doi: 10.1111/j.1399-3054.1992.tb01328.x
- Leegood, R. C., Lea, P. J., Adcock, M. D., and Hausler, R. E. (1995). The regulation and control of photorespiration. *J. Exp. Bot.* 46, 1397–1414. doi: 10.1093/jxb/46.special_issue.1397
- Long, S. P., and Bernacchi, C. J. (2003). Gas exchange measurements, what can they tell us about the underlying limitations to photosynthesis? Procedures and sources of error. *J. Exp. Bot.* 54, 2393–2401. doi: 10.1093/jxb/erg262
- Loreto, F., Harley, P. C., Marco, G. D., and Sharkey, T. D. (1992). Estimation of mesophyll conductance to CO₂ flux by three different methods. *Plant Physiol.* 98, 1437–1443. doi: 10.1104/pp.98.4.1437

- Miyake, C., Horiguchi, S., Makino, A., Shinzaki, Y., Yamamoto, H., and Tomizawa, K. (2005). Effects of light intensity on cyclic electron flow around PSI and its relationship to non-photochemical quenching of chl fluorescence in tobacco leaves. *Plant Cell Physiol.* 46, 1819–1830. doi: 10.1093/pcp/pci197
- Ogren, W. L. (1984). Photorespiration: pathways, regulation, and modification. *Annu. Rev. Plant Physiol.* 35, 415–442. doi: 10.1146/annurev.pp.35.060184.002215
- Peterhansel, C., and Maurino, V. G. (2011). Photorespiration redesigned. *Plant Physiol.* 155, 49–55. doi: 10.1104/pp.110.165019
- Scafaro, A. P., Von Caemmerer, S., Evans, J. R., and Atwell, B. J. (2011). Temperature response of mesophyll conductance in cultivated and wild *Oryza* species with contrasting mesophyll cell wall thickness. *Plant Cell Environ.* 34, 1999–2008. doi: 10.1111/j.1365-3040.2011.02398.x
- Suorsa, M., Järvi, S., Grieco, M., Nurmi, M., Pietrzykowska, M., Rantala, M., et al. (2012). PROTON GRADIENT REGULATION5 is essential for proper acclimation of *Arabidopsis* photosystem I to naturally and artificially fluctuating light conditions. *Plant Cell* 24, 2934–2948. doi: 10.1105/tpc.112.097162
- Takahashi, S., Bauwe, H., and Badger, M. R. (2007). Impairment of the photorespiratory pathway accelerates photoinhibition of photosystem II by suppression of repair but not acceleration of damage processes in *Arabidopsis*. *Plant Physiol.* 144, 487–494. doi: 10.1104/pp.107.097253
- Tazoe, Y., von Caemmerer, S., Badger, M. R., and Evans, J. R. (2009). Light and CO₂ do not affect the mesophyll conductance to CO₂ diffusion in wheat leaves. *J. Exp. Bot.* 60, 2291–2301. doi: 10.1093/jxb/erp035
- Tikkanen, M., and Aro, E. M. (2014). Integrative regulatory network of plant thylakoid energy transduction. *Trends. Plant Sci.* 19, 10–17. doi: 10.1016/j.tplants.2013.09.003
- Timm, S., Florian, A., Arrivault, S., Stitt, M., Fernie, A. R., and Bauwe, H. (2012). Glycine decarboxylase controls photosynthesis and plant growth. *FEBS Lett.* 586, 3692–3697. doi: 10.1016/j.febslet.2012.08.027
- Valentini, R., Epron, D., De Angelis, P., Matteucci, G., and Dreyer, E. (1995). *In situ* estimation of net CO₂ assimilation, photosynthetic electron flow and photorespiration in Turkey oak (*Q. cerris* L.) leaves: diurnal cycles under different levels of water supply. *Plant Cell Environ.* 18, 631–640. doi: 10.1111/j.1365-3040.1995.tb00564.x
- von Caemmerer, S., and Farquhar, G. D. (1981). Some relationships between the biochemistry of photosynthesis and the gas exchange of leaves. *Planta* 153, 376–387. doi: 10.1007/BF00384257
- Walker, B., Ariza, L. S., Kaines, S., Badger, M. R., and Cousins, A. B. (2013). Temperature response of *in vivo* Rubisco kinetics and mesophyll conductance in *Arabidopsis thaliana*: comparisons to *Nicotiana tabacum*. *Plant Cell Environ.* 36, 2108–2119. doi: 10.1111/pce.12166
- Warren, C. R., and Dreyer, E. (2006). Temperature response of photosynthesis and internal conductance to CO₂: results from two independent approaches. *J. Exp. Bot.* 57, 3057–3067. doi: 10.1093/jxb/erl067
- Weber, A. P., and Bauwe, H. (2013). Photorespiration – a driver for evolutionary innovations and key to better crops. *Plant Biol.* 15, 621–623. doi: 10.1111/plb.12036
- Wingler, A., Lea, P. J., Quick, W. P., and Leegood, R. C. (2000). Photorespiration: metabolic pathways and their role in stress protection. *Philos. Trans. R. Soc. B Biol. Sci.* 355, 1517–1529. doi: 10.1098/rstb.2000.0712
- Wingler, A., Quick, W. P., Bungard, R. A., Bailey, K. J., Lea, P. J., and Leegood, R. C. (1999). The role of photorespiration during drought stress: an analysis utilising barley mutants with reduced activities of photorespiratory enzymes. *Plant Cell Environ.* 22, 361–373. doi: 10.1046/j.1365-3040.1999.00410.x
- Yamori, W., Evans, J. R., and von Caemmerer, S. (2010). Effects of growth and measurement light intensities on temperature dependence of CO₂ assimilation rate in tobacco leaves. *Plant Cell Environ.* 33, 332–343. doi: 10.1111/j.1365-3040.2009.02067.x
- Yamori, W., Masumoto, C., Fukayama, H., and Makino, A. (2012). Rubisco activase is a key regulator of non-steady-state photosynthesis at any leaf temperature and, to a lesser extent, of steady-state photosynthesis at high temperature. *Plant J.* 71, 871–880. doi: 10.1111/j.1365-313X.2012.05041.x
- Yamori, W., Nagai, T., and Makino, A. (2011). The rate-limiting step for CO₂ assimilation at different temperatures is influenced by the leaf nitrogen content in several C₃ crop species. *Plant Cell Environ.* 34, 764–777. doi: 10.1111/j.1365-3040.2011.02280.x

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2015 Huang, Hu and Zhang. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.