



Induced Systemic Resistance (ISR) and Fe Deficiency Responses in Dicot Plants

Francisco J. Romera^{1*}, María J. García², Carlos Lucena², Ainhoa Martínez-Medina³, Miguel A. Aparicio⁴, José Ramos⁴, Esteban Alcántara¹, Macarena Angulo¹ and Rafael Pérez-Vicente²

¹ Department of Agronomy, Campus de Excelencia Internacional Agroalimentario CeIA3, Universidad de Córdoba, Córdoba, Spain, ² Department of Botany, Ecology and Plant Physiology, Campus de Excelencia Internacional Agroalimentario CeIA3, Universidad de Córdoba, Córdoba, Spain, ³ Molecular Interaction Ecology, German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Leipzig, Germany, ⁴ Department of Microbiology, Campus de Excelencia Internacional Agroalimentario CeIA3, Universidad de Córdoba, Córdoba, Spain

OPEN ACCESS

Edited by:

Thomas J. Buckhout,
Humboldt-Universität zu Berlin,
Germany

Reviewed by:

Christos Zamioudis,
Democritus University of Thrace,
Greece
Louis Grillet,
Academia Sinica, Taiwan

*Correspondence:

Francisco J. Romera
ag1roruf@uco.es

Specialty section:

This article was submitted to
Plant Nutrition,
a section of the journal
Frontiers in Plant Science

Received: 19 December 2018

Accepted: 21 February 2019

Published: 11 March 2019

Citation:

Romera FJ, García MJ, Lucena C, Martínez-Medina A, Aparicio MA, Ramos J, Alcántara E, Angulo M and Pérez-Vicente R (2019) Induced Systemic Resistance (ISR) and Fe Deficiency Responses in Dicot Plants. *Front. Plant Sci.* 10:287. doi: 10.3389/fpls.2019.00287

Plants develop responses to abiotic stresses, like Fe deficiency. Similarly, plants also develop responses to cope with biotic stresses provoked by biological agents, like pathogens and insects. Some of these responses are limited to the infested damaged organ, but other responses systemically spread far from the infested organ and affect the whole plant. These latter responses include the Systemic Acquired Resistance (SAR) and the Induced Systemic Resistance (ISR). SAR is induced by pathogens and insects while ISR is mediated by beneficial microbes living in the rhizosphere, like bacteria and fungi. These root-associated mutualistic microbes, besides impacting on plant nutrition and growth, can further boost plant defenses, rendering the entire plant more resistant to pathogens and pests. In the last years, it has been found that ISR-eliciting microbes can induce both physiological and morphological responses to Fe deficiency in dicot plants. These results suggest that the regulation of both ISR and Fe deficiency responses overlap, at least partially. Indeed, several hormones and signaling molecules, like ethylene (ET), auxin, and nitric oxide (NO), and the transcription factor MYB72, emerged as key regulators of both processes. This convergence between ISR and Fe deficiency responses opens the way to the use of ISR-eliciting microbes as Fe biofertilizers as well as biopesticides. This review summarizes the progress in the understanding of the molecular overlap in the regulation of ISR and Fe deficiency responses in dicot plants. Root-associated mutualistic microbes, rhizobacteria and rhizofungi species, known for their ability to induce morphological and/or physiological responses to Fe deficiency in dicot plant species are also reviewed herein.

Keywords: dicotyledons, ethylene, iron, ISR, rhizobacteria, rhizofungi, rhizosphere, stress responses

INTRODUCTION

In the last decades, crop productivity has been mainly based on the use of high-yielding varieties and in the application of high amounts of fertilizers and pesticides. Despite crop protection measures, current losses are estimated at 20–40% for the major food crops world-wide (Savary et al., 2012). Hence, novel strategies for crop production, with less reliance on chemical products need to be developed. In relation to plant mineral nutrition, two strategies that can contribute to

this goal are the development of crop varieties more efficient in nutrient acquisition and better management of the rhizosphere (Shen et al., 2013). The rhizosphere, the soil volume influenced by the root system, is one of the most energy-rich habitats on Earth, allowing the life of a myriad of microbes (Pieterse et al., 2014; Pii et al., 2015). Many of them are pathogenic and threaten plant growth. However, there are also many others that are beneficial for plants, like rhizobacteria (“PGPB or PGPR: Plant Growth-Promoting Bacteria or Rhizobacteria”) and fungi (“PGPF: Plant Growth-Promoting Fungi”), which can improve plant growth and benefit the adaptation of plants to adverse conditions (Yang et al., 2008; de Zelicourt et al., 2013; Pieterse et al., 2014; Pii et al., 2015; Verbon and Liberman, 2016). Some rhizosphere microbes can have negative effects on plant mineral nutrition, for example, by competing with plants for some nutrients. However, several genera of the rhizosphere microbiota can facilitate nutrient acquisition by plants, thus having positive effects. These beneficial microbes include, among others, mycorrhizal fungi and *Rhizobium*, which establish mutualistic symbiosis with plant roots that improve phosphorus (P) or nitrogen (N) nutrition, respectively (Guinel, 2015; Wang W. et al., 2017). Additionally, there are free-living mutualistic microbes that can improve plant nutrition through different mechanisms, such as the release of nutrient solubilizing compounds or the modification of root physiology and architecture (Jin et al., 2014; Mimmo et al., 2014; Zamioudis et al., 2014, 2015; Contreras-Cornejo et al., 2015; Li et al., 2015; Pii et al., 2015; García-López et al., 2016; Garnica-Vergara et al., 2016; Verbon et al., 2017).

Among the essential mineral nutrients required by plants, iron (Fe), along with P and N, represent the major constraints for crop productivity worldwide (Pii et al., 2015; Scagliola et al., 2016; Tsai and Schmidt, 2017b). Iron deficiency is widely distributed, mainly in calcareous soils (approximately one third of cultivated lands) which are abundant in arid and semiarid regions (Briat et al., 2015). To cope with Fe deficiency, plants develop morphological and physiological responses, mainly in their roots, aimed to facilitate its acquisition (see following Section; Kobayashi and Nishizawa, 2012; Brumbarova et al., 2015; Lucena et al., 2015). Despite these responses, in many cases it is necessary to apply Fe fertilizers to correct Fe deficiency. For Fe supply in the field, the most common practice is the application of Fe chelates to soils, which are generally expensive and therefore restricted to high added-value field-grown crops (Briat et al., 2015). An alternative is the use of more Fe efficient plant genotypes. However, different results obtained with sterile soils have shown that, even with these genotypes, the cooperation of rhizosphere microbes is necessary for an adequate Fe acquisition (Jin et al., 2014; Pii et al., 2015).

Several studies demonstrated that the application of some beneficial microbes to soils can improve the Fe nutrition of plants (de Santiago et al., 2009, 2013; Zhang et al., 2009; Freitas et al., 2015; Li et al., 2015; Ipek et al., 2017; Sonbarse et al., 2017; Aras et al., 2018; Arıkan et al., 2018). However, the main mechanisms driving such effects are complex and not fully understood. One possible mechanism is the release of Fe solubilizing compounds to soils (Jin et al., 2014; Mimmo et al., 2014; Pii et al., 2015). Moreover, the rhizosphere mutualistic microbiota can also improve plant Fe uptake by the alteration

of the root physiology and architecture (Zamioudis et al., 2014, 2015; Contreras-Cornejo et al., 2015; Garnica-Vergara et al., 2016; Scagliola et al., 2016; Verbon et al., 2017). In the last years it has been found that some rhizosphere microbes can induce physiological and morphological responses in roots of dicot plants similar to the ones induced by plants under Fe deficiency (Zhang et al., 2009; Orozco-Mosqueda et al., 2013; Jin et al., 2014; Pieterse et al., 2014; Zamioudis et al., 2014, 2015; Zhao et al., 2014; Pii et al., 2016b; Zhou et al., 2016a; Martínez-Medina et al., 2017; Verbon et al., 2017). It is remarkable that these rhizosphere microbes are also capable of eliciting the Induced Systemic Resistance (ISR) against pathogens and insects. This observation suggests that both processes (ISR and Fe deficiency responses) might be closely interconnected, and opens new possibilities for optimizing the management of the rhizosphere microbiota for improving Fe nutrition and health (Pieterse et al., 2014; Zamioudis et al., 2014, 2015; Verbon et al., 2017). However, the nodes of convergence between the two processes remain unclear.

Elucidating the main nodes of interconnection between the pathways regulating microbe-elicited ISR and Fe uptake is critical for optimizing the use of plant mutualistic microbes in agriculture. This review summarizes the progress in the understanding of the molecular overlap in the regulation of ISR and Fe deficiency responses in dicot plants. We further describe and evaluate rhizobacteria and rhizofungi species, known for their ability to induce morphological and/or physiological responses to Fe deficiency in dicot plants and with potential for a future use as Fe biofertilizers.

Fe DEFICIENCY RESPONSES IN DICOT PLANTS

Iron (Fe) is abundant in most soils, mainly as Fe^{3+} , although its availability to plants is low, especially in calcareous soils (Briat et al., 2015). Based on the mechanisms used by plant roots to facilitate mobilization and uptake of Fe, plants are classified into Strategy I species (dicots and non-grass monocots) and Strategy II species (grasses; Kobayashi and Nishizawa, 2012; Ivanov et al., 2012). Dicots, such as *Arabidopsis* and tomato, are Strategy I species which have to reduce Fe^{3+} to Fe^{2+} at the root surface, by means of a ferric reductase (encoded by *FRO2* in *Arabidopsis*), prior to its subsequent uptake through a Fe^{2+} transporter (encoded by *IRT1* in *Arabidopsis*; Ivanov et al., 2012; Kobayashi and Nishizawa, 2012). This review is devoted to dicots, where ISR mechanisms have been more extensively studied (Balmer et al., 2013). Consequently, the mechanisms described thereafter correspond to Strategy I plant species. For details about the Strategy II plant species readers are referred to other articles in this special issue.

When grown under Fe deficiency, Strategy I species develop several physiological and morphological responses, mainly in roots, known as Fe deficiency responses. Those responses are aimed at facilitating Fe mobilization and uptake (Ivanov et al., 2012; Kobayashi and Nishizawa, 2012; Brumbarova et al., 2015; Lucena et al., 2015). Among the physiological responses are: an enhanced ferric reductase activity due to upregulation of the *FRO*

genes; an enhanced Fe²⁺ uptake capacity due to upregulation of the *IRT1* genes; the acidification of the rhizosphere due to upregulation of *AHA* or *HA* (H⁺-ATPase) genes (Waters et al., 2007; Brumbarova et al., 2015; Lucena et al., 2015); an increase of the synthesis and release of organic acids, like citrate and malate, to the medium (Kabir et al., 2012; Schmidt et al., 2014); an increase of the synthesis and release of phenolic compounds to the medium due to upregulation of genes like *F6'H1*, *S8H*, *BGLU42*, and *ABCG37* (Schmid et al., 2014; Schmidt et al., 2014; Zamioudis et al., 2014; Tsai and Schmidt, 2017a; Siwinska et al., 2018; Tsai et al., 2018); and an increase of the synthesis and release of flavins to the medium (Rodríguez-Celma and Schmidt, 2013). The acidification facilitates the solubilisation of Fe and the functioning of the ferric reductase which has an optimum pH around 5.0 (Lucena et al., 2007; Waters et al., 2007). Organic acids can act as chelating agents for Fe in the soil and also inside the plant (Durrett et al., 2007; Schmidt et al., 2014). In fact, Fe is moved through the xylem chelated with citrate (Durrett et al., 2007; Schmidt et al., 2014). Phenolic compounds, like coumarins, and flavins can act as chelating and reducing agents of Fe³⁺, thus facilitating its mobilization in the rhizosphere (Rodríguez-Celma and Schmidt, 2013; Tsai and Schmidt, 2017a; Rajniak et al., 2018). The *F6'H1* ("Feruloyl-CoA 6'-Hydroxylase1") and *S8H* ("Scopoletin 8-Hydroxylase") genes encode enzymes involved in the last steps of the synthesis of the coumarins scopoletin and fraxetin (Schmid et al., 2014; Schmidt et al., 2014; Tsai and Schmidt, 2017a; Siwinska et al., 2018; Tsai et al., 2018). The *ABCG37* gene (also named *PDR9*) encodes an ABC transporter involved in the release of coumarins to the medium (Fourcroy et al., 2014, 2016; Zamioudis et al., 2014) while the *BGLU42* gene encodes a β-glucosidase, possibly required for the processing of glycosylated phenolic compounds as an essential step for their secretion in the root vicinity (Zamioudis et al., 2014; Stringlis et al., 2018b). Among the morphological responses are: development of subapical root hairs, cluster roots, and transfer cells, all of which are aimed to increase the surface of contact with the soil (Römheld and Marschner, 1986; Lucena et al., 2015; Romera et al., 2017). Both physiological and morphological responses are mainly located in the subapical regions of the roots (Römheld and Marschner, 1986).

The regulation of the physiological and morphological responses described above is not fully understood but in the last years several transcription factors (TFs) that participate in the activation of most of their associated genes have been described (Ivanov et al., 2012; Kobayashi and Nishizawa, 2012; Brumbarova et al., 2015; Zhang et al., 2015; Li et al., 2016; Liang et al., 2017). In *Arabidopsis*, the master regulator of most of these genes is FIT (bHLH29), homolog of the tomato FER (Bauer et al., 2007 and references therein). The FIT regulatory network comprises other bHLH TFs of the Ib subgroup, such as bHLH38, bHLH39, bHLH100, and bHLH101. All of them have redundant functions and can interact with FIT to form heterodimers that activate the expression of the Fe acquisition genes *FRO2* and *IRT1* (Yuan et al., 2008; Wang N. et al., 2013; Brumbarova et al., 2015). *FIT/FER* is induced in roots in response to Fe deficiency while the other Ib bHLH genes cited above are induced in both roots and leaves in response to Fe deficiency (Brumbarova et al., 2015 and

references therein). FIT also controls MYB10 and MYB72, two other TFs essential for plant growth on low Fe conditions (Palmer et al., 2013; Zamioudis et al., 2014, 2015). Besides the FIT/Ib bHLH regulatory network, there is another regulatory network related to the POPEYE (PYE; bHLH47) TF and associated with the vasculature (Brumbarova et al., 2015). In the last years, it has been found that, under Fe-deficiency conditions, IVc subgroup bHLH TFs [bHLH34, bHLH104, bHLH105(ILR3), and bHLH115] activate *FIT/bHLH38/39/100/101* and *PYE* expression (Zhang et al., 2015; Li et al., 2016; Liang et al., 2017). Upstream of the IVc subgroup bHLH TFs is the BRUTUS (BTS) protein, which possesses Fe-binding domains and that interacts with IVc bHLH TFs, targeting them for proteasomal degradation (Zhang et al., 2015; Liang et al., 2017). Since the IVc bHLH TFs act as positive regulators of Fe deficiency responses, the current data suggests that BTS is a negative regulator of Fe deficiency responses (Zhang et al., 2015; Hindt et al., 2017).

The mechanisms by which plants perceive Fe deficiency and how this perception is transmitted to the activation of the responses is not fully understood. Several studies support a role for hormones and other plant signaling molecules in the activation of FIT and other TFs and, consequently, in the upregulation of the ferric reductase, the Fe²⁺ transporter and other Fe-related genes. Within them, the plant hormone ethylene (ET) has been found to play a key role in the regulation of most of the physiological and morphological responses to Fe deficiency (Figure 1; reviewed in Lucena et al., 2015; Li and Lan, 2017; Romera et al., 2017). Besides ET, auxin, nitric oxide (NO), sucrose, and glutathione (GSH) have also been involved in the regulation of Fe deficiency responses; all of them increase in Fe-deficient roots although their specific roles are not fully understood (Romera et al., 1999, 2011, 2017; Lucena et al., 2006, 2015; Graziano and Lamattina, 2007; Waters et al., 2007; Baciaicoa et al., 2009, 2011; García et al., 2010, 2011; Chen et al., 2010; Lingam et al., 2011; Meiser et al., 2011; Koen et al., 2012; Yang et al., 2014; Shanmugam et al., 2015; Lin et al., 2016; Li and Lan, 2017; Kailasam et al., 2018). By contrast to these activating signals, other ones have been implicated in the suppression of Fe deficiency responses, like cytokinins (Séguéla et al., 2008), jasmonic acid (JA; Maurer et al., 2011), brassinosteroids (Wang et al., 2012), and some phloem Fe-related signals (García et al., 2013, 2018). To integrate both positive and negative signals in the regulation of Fe acquisition genes in roots, a model has been proposed where auxin/ET/NO would act as activators of their expression, while LODIS ("LOng Distance Iron Signal": a phloem Fe-related signal) would act to repress them (Lucena et al., 2006; García et al., 2011, 2018; Romera et al., 2011, 2017).


INDUCED SYSTEMIC RESISTANCE (ISR)

Besides responses to abiotic stresses, plants also respond to biotic stresses provoked by biological agents, like pathogens or insects (Pieterse et al., 2014; Martínez-Medina et al., 2017; Verbon et al., 2017). Some of these responses are localized but others are systemic, spreading far from the attacked organ and inducing defensive responses in the entire

plant (Pieterse et al., 2014; Verbon et al., 2017). Within this second possibility, induced resistance is a physiological state of enhanced defensive capacity of the plant triggered by biological or chemical inducers, which protects plant tissues not exposed to the initial attack against future attack by pathogens and herbivorous insects (Van Loon et al., 1998). Induced resistance can be triggered in plants by the infection of pathogens, in response to insect herbivory, or upon root colonization by certain rhizosphere mutualistic microbes. Two of the most studied forms of induced resistance are SAR (Systemic Acquired Resistance), triggered by plant pathogens, and ISR, triggered by root-colonizing mutualistic microbes, like *Pseudomonas simiae* (syn. *Pseudomonas fluorescens*), *Paenibacillus polymyxa*, or *Trichoderma* spp. (Table 1; Zhang et al., 2009; Pieterse et al., 2012, 2014; Alizadeh et al., 2013; Zamioudis et al., 2014, 2015; Zhao et al., 2014; Martínez-Medina et al., 2017; Verbon et al., 2017). SAR and ISR are mainly differentiated on the basis of the elicitor and the regulatory pathways involved, though the signaling pathways that regulate SAR and ISR share some components (Pieterse et al., 1998, 2002, 2012, 2014; Van Loon et al., 1998; Choudhary et al., 2007).

Over the last years, SAR and ISR have been extensively reviewed (Van Loon et al., 1998; Choudhary et al., 2007; Pieterse et al., 2012, 2014), so here we only discuss major principles in both responses. In systemic tissues, SAR is characterized by increased levels of the hormone salicylic acid (SA) which, through the redox-regulated protein NON-EXPRESSION OF PR GENES1 (NPR1), activates the expression of a large set of PATHOGENESIS-RELATED (PR) genes, involved in defense responses (Figure 1; Pieterse et al., 1998, 2002, 2012, 2014; Van Loon et al., 1998; Choudhary et al., 2007). By contrast to SAR, ISR is generally mediated by an SA-independent pathway where JA and ET are the central players, and typically functions without PR gene activation (Figure 1; Pieterse et al., 1998, 2002, 2012, 2014; Van Loon et al., 1998; Choudhary et al., 2007). Despite these differences, it has been shown that NPR1 is also required for the JA/ET-dependent ISR triggered by rhizosphere microbes although its role seems to be different in both processes (Figure 1; Pieterse et al., 2014; Nie et al., 2017). In SA signaling, NPR1 is related to a function in the nucleus while in JA/ET signaling it is related to a cytosolic function (Pieterse et al., 2014). Despite these general differences, in some particular cases, ISR can require SA accumulation (Ryu et al., 2003; Alizadeh et al., 2013). Moreover, the signaling pathways involved in the induction of ISR can be different depending on the microbial species and the plant species (Ryu et al., 2003; Jankiewicz and Koltonowicz, 2012; Alizadeh et al., 2013).

The discovery of ISR occurred around 1991, when several researchers showed that root colonization by certain non-pathogenic bacterial races promoted the health of plants upon the stimulation of their defense responses (reviewed in Pieterse et al., 2014). After these pioneering works with bacteria, ISR was further extended to rhizosphere fungi, like *Trichoderma* spp. or *Piriformospora indica* (Table 1; Segarra et al., 2009; Alizadeh et al., 2013; Pieterse et al., 2014).



How Is ISR Triggered by Beneficial Rhizosphere Microbes?

The ways beneficial rhizosphere microbes elicit ISR are not totally understood but several microbial elicitors have been proposed to be responsible for its onset. These elicitors, upon perception, would trigger the ISR through the action of diverse plant hormones (Figure 1; Pieterse et al., 2012, 2014; Sharifi and Ryu, 2018; Tyagi et al., 2018). Among these elicitors,

there are Microbe-Associated Molecular Patterns (MAMPs) and other elicitors, like Volatile Organic Compounds (VOCs) or siderophores (Figure 1; Zhang et al., 2007, 2009; Jankiewicz and Koltonowicz, 2012; Orozco-Mosqueda et al., 2013; Pieterse et al., 2014; Zamioudis et al., 2015; Garnica-Vergara et al., 2016; Martínez-Medina et al., 2017; Sharifi and Ryu, 2018; Tyagi et al., 2018; Villena et al., 2018). MAMPs (when produced by pathogens are named Pathogen-Associated Molecular Patterns: PAMPs) are conserved microbial molecules released by the microbes, like flagellin, chitin, and lipopolysaccharides (LPS; Zeidler et al., 2004; Pieterse et al., 2014; Villena et al., 2018). VOCs are low molecular weight compounds derived from different biosynthetic pathways, with high vapor pressure and that can evaporate and disperse easily (Sharifi and Ryu, 2018; Tyagi et al., 2018). At present, over 1000 volatile compounds (including alkanes, alcohols, esters, ketones, sulfides, terpenoids, and sesquiterpenes) have been identified (Tyagi et al., 2018). Those derived from beneficial microbes can trigger drastic changes in plant growth patterns, generally by altering hormone signaling (Garnica-Vergara et al., 2016; Martínez-Medina et al., 2017; Sharifi and Ryu, 2018; Tyagi et al., 2018). Siderophores are Fe chelating agents released by the bacteria to further acquire Fe from the medium (Lemanceau et al., 2009; Aznar and Dellagi, 2015; Aznar et al., 2014, 2015).

Microbe-Associated Molecular Patterns are perceived by Pattern Recognition Receptors (PRRs) while other elicitors could be perceived by other Receptors (Rs), not known in all cases (Figure 1; Jankiewicz and Koltonowicz, 2012; Pieterse et al., 2014; Aznar and Dellagi, 2015; Aznar et al., 2015; Sharifi and Ryu, 2018; Tyagi et al., 2018; Villena et al., 2018). Upon perception, the elicitors trigger the ISR by affecting diverse plant hormones that act as central players in the plant immune signaling network leading to the activation of the defense responses (Figure 1; Pieterse et al., 2012, 2014; Sharifi and Ryu, 2018; Tyagi et al., 2018). In some cases, the microbes themselves can also produce different hormones, like auxin or cytokinins, that upon perception by the plant hormonal receptors (HRs) can cause changes in the root physiology and morphology (Grady et al., 2016; Scagliola et al., 2016; Asari et al., 2017; Kudoyarova et al., 2017; Patel and Saraf, 2017). Among the hormones implicated in the ISR, JA, ET, auxin, and NO play a key role (Knoester et al., 1999; Ton et al., 2001; Shoresh et al., 2005; Zhang et al., 2007; Van der Ent et al., 2008; Camehl et al., 2010; Acharya et al., 2011; Pieterse et al., 2014; Garnica-Vergara et al., 2016; Hossain et al., 2017; Martínez-Medina et al., 2017; Nie et al., 2017; Nascimento et al., 2018; Stringlis et al., 2018a).

ISR Characteristics

One general characteristic of the microbial elicitors that induce ISR is their redundancy. This redundancy implies that microbial mutants defective in one elicitor can induce ISR through other elicitors (Meziane et al., 2005; Pieterse et al., 2014; Zamioudis et al., 2015). For example, the siderophore pseudobactin was as effective in inducing ISR as live bacteria but a mutant defective in pseudobactin biosynthesis was equally effective (Meziane et al., 2005). Beneficial ISR-eliciting microbes do not directly activate defense responses but sensitize the whole plant (a phenomenon called priming) for a faster and stronger activation of defense

responses upon invasion by pathogens (Choudhary et al., 2007; Berendsen et al., 2012; Jung et al., 2012; Pieterse et al., 2014; Martínez-Medina et al., 2016). A high percentage of the genes, predominantly associated with defense responses, induced by the elicitors, like flagellin, are suppressed by the ISR-eliciting microbes to allow the establishment of a mutually beneficial interaction with the host root (Stringlis et al., 2018a). There is increasing evidence that beneficial soil-borne microbes hijack plant hormone signaling pathways to suppress the host defenses (Pieterse et al., 2012). This is also the case for the symbiotic relationship between legumes and rhizobia where the defense reactions set up by the plant are quickly suppressed, allowing microbial entry and the potential successful rhizobial establishment in plant roots (Guinel, 2015).

To elicit ISR, beneficial rhizobacteria must reach a minimal concentration equal to 10^5 – 10^7 colony forming unit (CFU) per gram of root for several days (Jankiewicz and Koltonowicz, 2012; Bakker et al., 2013; Pieterse et al., 2014). It should be noted that in the rhizosphere, the microbial density can range from 10^8 to 10^9 bacteria per gram and that its diversity is generally less than in the bulk soil since plant exudates specifically stimulate or repress members of the microbial community shaping the root microbiome (Figure 1; Berendsen et al., 2012; Bakker et al., 2013; Pii et al., 2016a; Stringlis et al., 2018b). In this sense, very recently it has been found that the release of the antimicrobial coumarin scopoletin by roots of *Arabidopsis* plants inoculated with the rhizobacterium *P. simiae* inhibits some soil-borne pathogens but not the rhizobacterium (Stringlis et al., 2018b). Coumarins are phenolic compounds that are also released by Fe-deficient roots to favor the Fe acquisition of plants (Schmid et al., 2014; Schmidt et al., 2014; Tsai and Schmidt, 2017a; Siwinska et al., 2018; Tsai et al., 2018; see also Section “Fe deficiency responses in dicot plants”). Consequently, the ISR-eliciting microbes, by inducing the release of coumarins and other Fe deficiency responses in plants, can improve the Fe nutrition of plants but, at the same time, they can benefit from a niche where their competitors are eliminated or restricted (Stringlis et al., 2018b).

INTERRELATIONSHIP BETWEEN ISR AND Fe DEFICIENCY RESPONSES IN DICOT PLANTS

Since Fe acquisition is a limiting factor in most soils, Fe is a central player in the tripartite interaction among beneficial microbes, pathogens, and plants (López-Berges et al., 2013; Naranjo-Arcos and Bauer, 2016; Verbon et al., 2017). This close interrelation is in good agreement with the already described relationship between Fe homeostasis and defense responses against pathogens in plants (Lemanceau et al., 2009; Aznar et al., 2015; Verbon et al., 2017) and with the crosstalk between ISR and Fe deficiency responses (Pieterse et al., 2014; Verbon et al., 2017). The relationship between plant defense responses and Fe deficiency is complex and depends on several factors, like the plant genotype, the kind of pathogens and the intensity and duration of the deficiency. In some cases, plants are more tolerant to pathogens under conditions of Fe deficiency,

TABLE 1 | Microbial species that induce Fe deficiency responses when applied to dicot plants.


Microbial species	Plant species	Mode appl.	Signals	Fe def. resp.	Fe genes	Fe Gr.	Refs
Rhizobacteria							
<i>Azospirillum brasilense</i>	<i>Solanum lycopersicum</i>	gm(a)	ET Auxin	Root hairs	nd	^	Ribaudo et al., 2006
<i>Azospirillum brasilense</i>	<i>Cucumis sativus</i>	gm(ns)	nd	FCR, pH	<i>FIT FRO1 IRT1 HA1</i>	Fe	Pii et al., 2016b
<i>Bacillus subtilis</i>	<i>Arabidopsis thaliana</i>	gm(a)	VOCs	FCR, pH	<i>FIT FRO2 IRT1</i>	Fe	Zhang et al., 2009
<i>Arthrobacter agilis*</i>	<i>Medicago truncatula</i>	gm(a)	VOCs	FCR, pH	nd	Fe ^	Orozco-Mosqueda et al., 2013
<i>Pseudomonas simiae</i>	<i>Arabidopsis thaliana</i>	gm(a)	VOCs Auxin NO	FCR phenolics	<i>FIT FRO2 bHLH38 bHLH39 IRT1 F6'H1 MYB72 BGLU42 ABCG37</i>		Zamioudis et al., 2014, 2015; Stringlis et al., 2018b
<i>Enterobacter*</i>	<i>Cucumis sativus</i>	gm(ns)	Auxin	FCR	nd	nd	Scagliola et al., 2016
<i>Pseudomonas</i>							
<i>Paenibacillus polymyxa</i>	<i>Arabidopsis thaliana</i>	gm(a)	Auxin	FCR, pH, phenolics	<i>FIT FRO2 IRT1 MYB72 F6'H1</i>	Fe ^	Zhou et al., 2016a
<i>Bacillus amyloliquefaciens</i>	<i>Arabidopsis thaliana</i>	gm(a)	VOCs Auxin NO	FCR	<i>FIT FRO2 IRT1</i>	Fe ^	Wang J. et al., 2017; Zhou et al., 2017
<i>Burkholderia cepacia</i>	<i>Astragalus sinicus</i>	ri(s)	Auxin	FCR, pH, flavins	<i>FIT FRO2 IRT1 AHA2</i>	Fe ^	Zhou et al., 2018
Rhizofungi							
<i>Trichoderma asperellum</i>	<i>Cucumis sativus</i>	gm(s)	nd	FCR	nd	Fe ^	Zhao et al., 2014
<i>Trichoderma asperellum,</i> <i>T. harzianum</i>	<i>Arabidopsis thaliana</i>	gm(a)	VOCs	FCR, root hairs	<i>FIT FRO2 bHLH38 bHLH39 IRT1 MYB72</i>	nd	Martínez-Medina et al., 2017
<i>Trichoderma asperellum,</i> <i>T. harzianum</i>	<i>Solanum lycopersicum</i>	gm(a)	VOCs	FCR, root hairs	<i>FER FRO1 IRT1</i>	nd	Martínez-Medina et al., 2017

Mode appl., mode of application of the microbes (gm, application to the growth medium; ri, root immersion; a, agar; ns, nutrient solution; s, soil); Signals, elicitors and hormones involved; Fe def. resp., Fe deficiency responses; FCR, ferric chelate reductase activity; pH, acidification; Fe genes, Fe acquisition genes; Fe Gr., increased shoot Fe concentration (Fe) and increased shoot growth (^); nd, not determined; *, microbial species whose association with ISR is not yet clear.

probably because pathogens require an adequate quantity of Fe for full virulence (Kieu et al., 2012; López-Berges et al., 2013). However, in other cases, plants are more susceptible to pathogens under Fe-deficient conditions (Verbon et al., 2017 and references therein). The competition for Fe between soil-borne pathogens and their antagonistic microorganisms has been related to disease suppression; siderophores produced in the rhizosphere by PGPR can inhibit growth of the pathogens by depriving them of Fe (Verbon et al., 2017). In contrast to the negative effect of some soil-borne pathogens on Fe acquisition, there are several recent reviews showing an important role of beneficial rhizosphere microbes on the Fe nutrition of plants (Jin et al., 2014; Mimmo et al., 2014; Pii et al., 2015; İpek and Esitken, 2017). These microbes can directly improve Fe nutrition through the release of H⁺ and/or Fe-solubilizing compounds to soils, like siderophores and organic acids, or by inducing changes in root physiology and architecture, which can improve the acquisition of Fe and also of other nutrients (Orozco-Mosqueda et al., 2013; Jin et al., 2014; Mimmo et al., 2014; Zhao et al., 2014; Contreras-Cornejo et al., 2015; Pii et al., 2015, 2016b; Garnica-Vergara et al., 2016; Scagliola et al., 2016; Verbon and Liberman, 2016; Zhou et al., 2016a,b; Martínez-Medina et al., 2017; Sonbarse et al., 2017; Sharifi and Ryu, 2018; Stringlis et al., 2018a). In this way, it has been demonstrated that ISR-eliciting microbes can induce Fe deficiency responses in their host roots, such as enhanced ferric reductase activity, acidification of the rhizosphere, release of phenolics and flavins,

and development of root hairs; and the expression of the genes associated with these responses, such as *FIT*, *bHLH38*, *bHLH39*, *MYB72*, *MYB10*, *FRO2*, *IRT1*, *AHA*, *F6'H1*, *BGLU42*, *ABCG37*, and others (Figures 1, 2 and Table 1; Ribaudo et al., 2006; Zhang et al., 2009; Zamioudis et al., 2014, 2015; Zhao et al., 2014; Pii et al., 2016b; Scagliola et al., 2016; Verbon and Liberman, 2016; Zhou et al., 2016a,b, 2018; Martínez-Medina et al., 2017; Verbon et al., 2017; see also Section “Fe deficiency responses in dicot plants” and Section “Rhizosphere microbial species that induce Fe deficiency responses and improve Fe acquisition”).

Since bacteria that elicit ISR can release siderophores to the medium, it has been speculated that perhaps these Fe chelating agents could deprive plants of Fe and in this way cause the induction of Fe deficiency responses (Van der Ent, 2008; Aznar et al., 2014; Pieterse et al., 2014; Aznar and Dellagi, 2015; Zamioudis et al., 2015). However, mutants defective in siderophore biosynthesis also induce Fe deficiency responses, which suggests that they could induce these responses through other mechanisms (Meziane et al., 2005; Pieterse et al., 2014; Zamioudis et al., 2015). A possibility could be through alteration of hormone biosynthesis and signaling in the plants. In this sense, the plant hormone ET has been implicated in both the activation of ISR (Knoester et al., 1999; Ton et al., 2001; Shores et al., 2005; Camehl et al., 2010; Pieterse et al., 2012; Garnica-Vergara et al., 2016; Hossain et al., 2017; Nie et al., 2017) and the activation of Fe deficiency responses (Figures 1, 2; reviewed in Lucena et al., 2015; Li and Lan, 2017; Romera et al., 2017). Besides




ET, other hormones and signaling molecules, like auxin, and NO, have also been implicated in both processes (García et al., 2010; Acharya et al., 2011; Romera et al., 2011, 2017; Zamioudis, 2012; Jin et al., 2014; Contreras-Cornejo et al., 2015; Garnica-Vergara et al., 2016; Poupin et al., 2016; Stringlis et al., 2018a). Moreover, the root-specific MYB72 TF, that plays a key role in the onset of ISR (Van der Ent et al., 2008; Segarra et al., 2009; Zamioudis et al., 2014; Verbon et al., 2017; Stringlis et al., 2018b), is also essential for plant growth on low Fe conditions (**Figure 2**; García et al., 2011; Palmer et al., 2013; Zamioudis et al., 2014, 2015; Lucena et al., 2015; Verbon et al., 2017; see also Section “Fe deficiency responses in dicot plants”). This indicates that MYB72 is a node of convergence between ISR and Fe deficiency responses (Van der Ent et al., 2008; Segarra et al., 2009; Pieterse et al., 2014; Zamioudis et al., 2014, 2015; Verbon et al., 2017). All these results suggest that the regulatory pathways of ISR and Fe deficiency responses overlap. In the following paragraphs the main common components shared by ISR and Fe deficiency responses are described and analyzed.

Ethylene and Other Hormones and Signaling Molecules

As previously stated (see Section “Induced Systemic Resistance”), ISR is generally mediated by a pathway where JA and ET are the central players (**Figure 1**; Pieterse et al., 1998, 2002,

2012, 2014; Van Loon et al., 1998; Choudhary et al., 2007). In this pathway, JA acts upstream of ET, and ET upstream of the NPR1 protein (**Figure 1**; Pieterse et al., 1998, 2002, 2012, 2014; Van Loon et al., 1998; Choudhary et al., 2007). NPR1 functions in the nucleus as a transcriptional coactivator of SA-responsive genes during the SAR pathway, but NPR1 also plays a cytosolic function in the JA/ET signaling during the ISR pathway (**Figure 1**; Pieterse et al., 2014). In relation to these components (ET, JA, SA, and NPR1), ET has been clearly implicated in the regulation of Fe deficiency responses in Strategy I plants and in rice, that presents characteristics of both Strategy I and II plant species (reviewed in Lucena et al., 2015; Li and Lan, 2017; Romera et al., 2017). However, the roles of the other components (JA, SA, and NPR1) on Fe deficiency responses are not yet clear. For instance, JA has been implicated in the regulation of Fe deficiency responses in Strategy I plants but as a suppressor of these responses (Maurer et al., 2011). In rice, JA has been shown to activate the expression of some Fe deficiency responses at the very early stages of the Fe deficiency (0.5–1 h) but strongly suppresses them later on (3–6 h; Kobayashi et al., 2016). In relation to the role of SA on the regulation of Fe deficiency responses, the results are contrasting. It was shown that Arabidopsis lines overexpressing the SA-inducible transcription factor OBF–BINDING PROTEIN 3 (OBP3) present upregulation of *bHLH038* and *bHLH039*, encoding two TFs that play a key role in the activation of Fe acquisition genes (Kang et al., 2003; see Section “Fe deficiency responses in dicot plants”). In the same way, exogenous application of SA to Arabidopsis plants upregulates the expression of *YELLOW STRIPE-LIKE1* (*YSL1*) and *YSL3*, which are involved in Fe translocation and homeostasis (Chen et al., 2014; Kumar et al., 2017). Shen et al. (2016) further found that Fe deficiency increases SA contents in shoots and roots of Arabidopsis plants, and that the SA biosynthesis defective mutant *phytoalexin deficient 4* (*pad4*) presents altered Fe deficiency responses, suggesting a link between SA and Fe deficiency. However, Maurer et al. (2014) found that SA and SA signaling through NPR1 (**Figure 1**) do not affect Fe deficiency responses. To complete this network of interactions, some Fe-related TFs, such as ILR3 (bHLH105; belonging to the Ivc bHLH subgroup; Zhang et al., 2015; see also Section “Fe deficiency responses in dicot plants”), can affect JA and SA biosynthesis (Aparicio and Pallás, 2017). After analyzing all these results, it is clear that the role of JA and SA on the regulation of Fe deficiency responses in dicots deserves further research.

In relation to ET, this plant hormone is synthesized from the amino acid methionine, through a pathway requiring SAMS (S -adenosyl methionine synthetase), ACS [1-aminocyclopropane-1-carboxylic acid (ACC) synthase], and ACO (ACC oxidase; Sauter et al., 2013; Wang F. et al., 2013; Dubois et al., 2018):



Although ET’s mode of action is not fully understood, a linear signaling pathway has been proposed in Arabidopsis

(Shakeel et al., 2013; Wang F. et al., 2013; Dubois et al., 2018):

ET \rightarrow ET receptors \rightarrow CTR1 \rightarrow EIN2 \rightarrow EIN3/EILs
 \rightarrow ERFs \rightarrow ET responses

In this signaling pathway, all the components are proteins and, within them, EIN3/EILs and ERFs are TFs (for more details, see Shakeel et al., 2013; Wang F. et al., 2013; Lucena et al., 2015; Dubois et al., 2018).

The participation of ET in the regulation of Fe deficiency responses was first proposed by Romera and Alcántara (1994) and has been further supported by many experimental data (recently reviewed in Lucena et al., 2015; Li and Lan, 2017; Romera et al., 2017). As more recent evidence, it should be mentioned that FIT (master regulator of Fe acquisition genes in Arabidopsis; see Section “Fe deficiency responses in dicot plants”) interacts with the EIN3 and EIL1 TFs, associated with ET signaling, and with MED16 and MED25, Mediators, to form a complex implicated in the transcription of Fe acquisition genes (Figure 2; Lingam et al., 2011; Yang et al., 2014; Brumbarova et al., 2015). In the same way, recently it has been found that the ERF4 and ERF72 TFs, that are also associated with ET signaling, are induced under Fe deficiency and participate in the regulation of Fe deficiency responses (Liu et al., 2017a,b). Fe deficiency causes the upregulation of genes involved in both ET synthesis, like *SAM*, *ACS*, *ACO*, *MTK*, *MTI1*, *MPK3*, and *MPK6*, and signaling, like *EIN2*, *EIN3*, *EIL1*, *EIL3*, *ERF4*, and *ERF72*, in roots of different dicot plants (García et al., 2010; Ye et al., 2015; Li and Lan, 2017; Romera et al., 2017). *MTK* (5-methylthioribose kinase) and *MTI1* (5-methylthioribose-1-phosphate isomerase1) participate in the Yang cycle and are necessary for ET biosynthesis (Pommerrenig et al., 2011; Sauter et al., 2013). Both *MTK* and *MTI1*, besides their upregulation under Fe deficiency, are also regulated by ET, probably through FIT (Figure 2; Colangelo and Guerinot, 2004; García et al., 2010). *MTI1* (At2g05830) was previously identified as eIF-2B, a putative eukaryotic translation initiation factor (García et al., 2010; Pommerrenig et al., 2011). The mitogen-activated protein kinases 3 and 6 (*MPK3*/*MPK6*) are related to ET (Li et al., 2012; Contreras-Cornejo et al., 2015; Dubois et al., 2018) and can activate *ACS2/6* in Fe deficiency-induced ET production (Ye et al., 2015).

Ethylene has been involved in the regulation of both morphological and physiological responses to Fe deficiency in Strategy I plants (Romera and Alcántara, 1994, 2004; Lucena et al., 2015; Li and Lan, 2017; Romera et al., 2017). In relation to the morphological responses, ET has been implicated in the formation of subapical root hairs, cluster roots, and transfer cells (reviewed in Lucena et al., 2015). In relation to the physiological responses, ET has been implicated in the upregulation of *FIT* (or its tomato homolog *FER*), *bHLH38*, *bHLH39*, and *MYB72*, encoding key TFs (Figure 2; Lucena et al., 2006; García et al., 2010; Lingam et al., 2011). *bHLH38* and *bHLH39* interact with *FIT* to form heterodimers that activate the expression of the Fe acquisition genes *FRO2* (ferric reductase) and *IRT1* (iron transporter) (Figure 2; Yuan et al., 2008; Wang N. et al., 2013). Similarly, the acidification capacity,

depending on *AHA*-like genes (Colangelo and Guerinot, 2004), is also activated by *FIT* and consequently by ET (Figure 2; Waters et al., 2007; Lucena et al., 2015). In relation to the excretion of phenolics, it has been found that the expression of *F6'H1* and *S8H*, involved in their synthesis, is dependent on *FIT* (Figure 2; García et al., 2010; Schmid et al., 2014; Tsai and Schmidt, 2017a; Tsai et al., 2018; see also “Fe deficiency responses in dicots”). Consequently, both of them would also be dependent on ET (Figure 2). In supporting this view, *S8H* (At3g12900) is greatly induced in Fe deficient roots and drastically inhibited by ethylene inhibitors (García et al., 2010). Besides all the above genes related to Fe acquisition, ET also participates in the activation of *NAS1* (nicotianamine synthase1), *NAS2*, and *FRD3* (ferric reductase defective3), that are very important for internal Fe mobilization and homeostasis (García et al., 2010).

Besides its involvement in Fe deficiency responses, ET has also been implicated in the development of ISR triggered by root-colonizing microbes, acting downstream of JA (Figure 1; Knoester et al., 1999; Ton et al., 2001; Shores et al., 2005; Ribaud et al., 2006; Van Loon et al., 2006; Van der Ent, 2008; Camehl et al., 2010; Pieterse et al., 2012; Zamioudis, 2012; Contreras-Cornejo et al., 2015; Garnica-Vergara et al., 2016; Hossain et al., 2017; Nie et al., 2017; Nascimento et al., 2018). In general, root colonization by ISR-eliciting microbes does not induce a direct enhancement of ET and JA biosynthesis (Knoester et al., 1999; Pieterse et al., 2014) except in some cases (Ribaud et al., 2006; Contreras-Cornejo et al., 2015). However, the expression of genes involved in ET biosynthesis, like *ACS*s and *ACOs*, and signaling, like *ETR1*, *EIL3*, *CTR1*, and *ERFs*, is frequently upregulated in roots by ISR-eliciting microbes (Shores et al., 2005; Ribaud et al., 2006; Van der Ent et al., 2008; Velivelli et al., 2015; Zamioudis et al., 2015; Poupin et al., 2016). For example, root colonization by *P. simiae* WCS417 induced the upregulation of *ACS2*, *ACS6*, and *EIL3* in Arabidopsis roots (Van der Ent et al., 2008; Zamioudis et al., 2015), which are also upregulated under Fe deficiency (García et al., 2010; Ye et al., 2015).

In both processes, ISR and Fe deficiency responses, ET can have a dual role. It is necessary for the activation of Fe deficiency responses (Lucena et al., 2015; Li and Lan, 2017; Romera et al., 2017) and for the onset of ISR (Knoester et al., 1999; Ton et al., 2001; Shores et al., 2005; Ribaud et al., 2006; Pieterse et al., 2012, 2014; Garnica-Vergara et al., 2016; Hossain et al., 2017; Nie et al., 2017; Nascimento et al., 2018). However, when accumulated in excess, ET can have negative effects on the responses to Fe deficiency (Romera et al., 1999), on the growth of plants and on the mutualistic interactions with beneficial microbes (Pierik et al., 2007; Camehl et al., 2010; Gamalero and Glick, 2015; Nascimento et al., 2018). This dual role also occurs for the nodulation between legumes and rhizobia, where ET is crucial for the proper development of the rhizobial colonization but also acts as a negative regulator to limit the number of rhizobial infections (Zamioudis, 2012; Guinel, 2015). To avoid the detrimental effects of ET, some beneficial microbes and plant species possess the enzyme ACC deaminase, that eliminates the ET

precursor ACC (Gamalero and Glick, 2015; Singh et al., 2015; Nascimento et al., 2018).

The participation of ET on morphological and physiological responses to Fe deficiency and on ISR probably follows different ET signaling pathways. While several Arabidopsis ethylene insensitive mutants, like *etr1*, *ein2*, *ein3*, and *eir1*, are blocked in their capacity to mount the ISR (Knoester et al., 1999; Ton et al., 2001; Van Loon et al., 2006; Camehl et al., 2010; Zamioudis, 2012; Alizadeh et al., 2013; Contreras-Cornejo et al., 2015; Hossain et al., 2017; Nie et al., 2017), and to develop morphological responses to Fe deficiency (Romera and Alcántara, 2004), they can induce most of the physiological responses to Fe deficiency (Romera and Alcántara, 2004; Lucena et al., 2006; García et al., 2010). These differences are perhaps related to the existence of an alternate route for ethylene signaling, besides the conventional one that includes EIN2 (see above; Shakeel et al., 2013). At this point, it has been suggested that for several physiological responses, ET could act through a pathway where EIN2 is not strictly required (Lucena et al., 2015).

Besides ET, other hormones and signaling molecules, such as auxin, GSH and NO have also been involved in both ISR and Fe deficiency responses (Ribaudo et al., 2006; Graziano and Lamattina, 2007; Bacaicoa et al., 2009, 2011; Chen et al., 2010; García et al., 2010, 2011, 2018; Acharya et al., 2011; Romera et al., 2011, 2017; Jin et al., 2014; Contreras-Cornejo et al., 2015; Shanmugam et al., 2015; Zamioudis et al., 2015; Garnica-Vergara et al., 2016; Poupin et al., 2016; Zhou et al., 2016a, 2017, 2018; Wang J. et al., 2017; Gullner et al., 2018; Kailasam et al., 2018; Sharifi and Ryu, 2018; Stringlis et al., 2018a; Sumayo et al., 2018; Tyagi et al., 2018). All of them increase in roots under Fe deficiency and frequently upon colonization of roots by ISR-eliciting microbes (Romera et al., 1999; Ribaudo et al., 2006; Graziano and Lamattina, 2007; Bacaicoa et al., 2009, 2011; Chen et al., 2010; Contreras-Cornejo et al., 2015; Shanmugam et al., 2015; Zamioudis et al., 2015; Zhou et al., 2016a, 2017; Wang J. et al., 2017; Kailasam et al., 2018). Some microbial elicitors, like VOCs or LPS, can affect ET, auxin or NO production and/or signaling, and in this way upregulate Fe-related genes (Zeidler et al., 2004; Zhang et al., 2007, 2009; Kwon et al., 2010; Liu and Zhang, 2015; Zamioudis et al., 2015; Garnica-Vergara et al., 2016; Zhou et al., 2017; Wang J. et al., 2017; Sharifi and Ryu, 2018; Tyagi et al., 2018). As examples, VOCs from *Bacillus subtilis* GB03 upregulated the expression of several ET biosynthesis genes (Kwon et al., 2010) and VOCs from *P. simiae* WCS417 or *Bacillus amyloliquefaciens* BF06 caused NO accumulation in Arabidopsis roots and upregulated several Fe-related genes (Zamioudis et al., 2015; Wang J. et al., 2017). Moreover, some of these hormones and signaling molecules can affect the perception of microbial elicitors. For example, the PRR for flagellin (FLAGELLIN-SENSING 2-FLS2) is regulated by ET (Boutrot et al., 2010).

Ethylene, auxin, and NO are closely interrelated since each one can affect the synthesis and/or action of the others (Figure 2; Ribaudo et al., 2006; Chen et al., 2010; García et al., 2011, 2018; Romera et al., 2011, 2017; Contreras-Cornejo et al., 2015; Garnica-Vergara et al., 2016; Poupin et al., 2016; Zhou et al., 2017). As a probe of their close interrelationship, *FIT*,

MYB72, and other Fe- and ISR-related genes are similarly affected by ET, auxin, or NO treatments. They are upregulated by these molecules, or their precursors, and downregulated by their inhibitors (Graziano and Lamattina, 2007; Chen et al., 2010; García et al., 2010, 2011; Zamioudis et al., 2015; Wang J. et al., 2017; Zhou et al., 2017; Stringlis et al., 2018a; Sumayo et al., 2018).

In the last years, the roles of GSH and NO in the activation of responses to Fe deficiency are becoming more complex since several experimental results have shown that S-nitrosoglutathione (GSNO), derived from GSH and NO, specifically works in such an activation having a different role than the one of NO (García et al., 2018; Kailasam et al., 2018). NO, GSH, and GSNO have also been implicated in plant defense responses against pathogens, and NO and GSH in ISR (Zamioudis et al., 2015; Yun et al., 2016; Gullner et al., 2017, 2018). Moreover, in plant defense responses, NO and GSNO exhibit additive functions and, by extension, may have distinct or overlapping molecular targets (Yun et al., 2016). The roles of GSH and GSNO in these processes have been related to their capacity to detoxify toxins (by their conjugation with GSH), to their interconnection with reactive oxygen species and SA, and to their capacity to modulate the redox state of NPR1 and to S-nitrosylate defense-related TFs and transcriptional coregulators (Yun et al., 2016; Gullner et al., 2017, 2018). However, to our knowledge, only NO has been implicated in the signaling processes leading to the activation of Fe acquisition genes by ISR-eliciting microbes (Zamioudis et al., 2015) while GSH and GSNO have not yet been studied in relation to this activation, which deserves further research.

MYB72 and Other Transcription Factors

Some years ago, it was found that the *MYB72* gene, encoding a TF, was greatly induced in Arabidopsis roots under Fe deficiency (Colangelo and Guerinot, 2004) and also upon their colonization with the ISR-eliciting rhizobacterium *P. simiae* WCS417 (Verhagen et al., 2004; Van der Ent, 2008; Van der Ent et al., 2008). Later on, *MYB72* upregulation has also been demonstrated with other ISR-eliciting microbes (Segarra et al., 2009; Alizadeh et al., 2013; Pieterse et al., 2014; Martínez-Medina et al., 2017; Verbon et al., 2017). Furthermore, Arabidopsis *myb72* knockout mutants are defective in the activation of ISR which suggests that *MYB72* plays a key role in the early signaling steps of this process (Figure 2; Van der Ent, 2008; Van der Ent et al., 2008; Segarra et al., 2009; Alizadeh et al., 2013; Zamioudis et al., 2014, 2015; Stringlis et al., 2018b). However, overexpression of *MYB72* did not result in enhanced resistance against any of the pathogens tested, demonstrating that *MYB72* is not sufficient for the expression of ISR (Van der Ent et al., 2008). In both Fe deficiency and ISR, *MYB72* is upregulated along with *MYB10*, also encoding a TF (Colangelo and Guerinot, 2004; Zamioudis, 2012; Palmer et al., 2013; Zamioudis et al., 2015). *MYB72* and *MYB10* physically interact *in vivo* and function redundantly in regulating the expression of genes involved in the shikimate, the phenylpropanoid and the nicotianamine (NA) biosynthesis pathways (Zamioudis, 2012; Zamioudis et al., 2014, 2015; Stringlis et al., 2018b). Under

Fe deficiency, both MYB72 and MYB10 act early in the Fe deficiency regulatory cascade to drive gene expression of *NAS2* and *NAS4*, two NA synthase genes (Palmer et al., 2013). An important difference of the participation of MYB72 and MYB10 in ISR and Fe deficiency responses is that the Arabidopsis *myb72* mutants are defective in the activation of ISR (see above) while they behave apparently normal when grown on alkaline soil, a condition that favors Fe deficiency (Palmer et al., 2013). However, the *myb10myb72* double mutant displays seedling lethality when grown on alkaline soil (Palmer et al., 2013). This suggests that MYB10 and MYB72 have overlapping roles in relation to Fe deficiency (Palmer et al., 2013). In relation to ISR, MYB72, and MYB10 coordinately suppress the expression of a large group of defense-related genes upon root colonization by *P. simiae* WCS417, enabling the bacteria to establish long-term associations with host roots (Zamioudis, 2012). The bacteria can also colonize roots of the *myb72* mutant (Van der Ent et al., 2008), suggesting that MYB10 may compensate in defense suppression (Zamioudis, 2012).

MYB72 is consequently a node of convergence between ISR and Fe deficiency responses in dicots. This convergence is further supported when considering that *MYB72* expression is controlled by FIT either under Fe deficiency or upon colonization of roots by ISR-eliciting microbes (Figure 2; Colangelo and Gueriot, 2004; Sivitz et al., 2012; Palmer et al., 2013; Zamioudis et al., 2015). Moreover, FIT interacts with the bHLH38 TF to control *MYB72* expression upon colonization of roots by ISR-eliciting microbes (Zamioudis et al., 2015), as occurred with *FRO2* and *IRT1* expression under Fe deficiency (Yuan et al., 2008). *FIT*, *bHLH38*, and *MYB72* expression is activated by ET (García et al., 2010, 2011), which suggests a connection between the regulation of Fe deficiency responses and ISR through this plant hormone. In supporting this view, several *P. simiae* WCS417-inducible *MYB72* target genes, like *BGLU42* (β -glucosidase), *CYP71B5* (cytochrome P450), *At5g55620* (unknown function), and *bHLH39* (TF), are all induced by Fe deficiency and also activated by ET (García et al., 2010; Zamioudis et al., 2014). Moreover, *MYB72* and *BGLU42* present ET-responsive elements in their promoters (García et al., 2010). Similarly, there are genes associated with the biosynthesis and release of coumarins in the rhizosphere, like *F6'H1*, *S8H*, and *ABCG37*, that are upregulated both under Fe deficiency and upon colonization of roots by ISR-eliciting microbes. These genes are dependent on FIT (*F6'H1*, *S8H*) or *MYB72* (*ABCG37*) and, consequently, on ET (Figure 2; García et al., 2010; Schmid et al., 2014; Schmidt et al., 2014; Zamioudis et al., 2014; Tsai and Schmidt, 2017a; Siwinska et al., 2018; Tsai et al., 2018; see also Section “Fe deficiency responses in dicot plants”).

The results about the relationship of *MYB72* and ET are controversial. For example, Van der Ent et al. (2006) showed that *MYB72* transcript levels accumulated after treatment with the ET precursor ACC and that they did not accumulate in the Arabidopsis ethylene insensitive mutant *ein2-1* upon root colonization with the ISR-eliciting bacterium *P. simiae* WCS417. However, later on, these authors found that *MYB72* transcript levels did not accumulate after treatment with ACC while they

accumulated in the *ein2-1* mutant upon treatment with *P. simiae* WCS417 (Van der Ent et al., 2008). After these latter results, they concluded that *MYB72* expression was not regulated by ET (Van der Ent et al., 2008). Similarly, it was found, by using yeast two-hybrid screening, that *MYB72* physically interacted *in vitro* with EIL3, a TF associated with ET signaling (Van der Ent et al., 2008) while later on it was found, by using Bimolecular Fluorescence Complementation, that *MYB72* and EIL3 did not interact *in vivo* (Zamioudis, 2012). Curiously, *EIL3* expression is upregulated both under Fe deficiency (García et al., 2010) and upon colonization of roots with *P. simiae* WCS417 (Van der Ent et al., 2008). Today, there is enough evidence to support the regulation of *FIT*, *bHLH38*, and *bHLH39* by ET and, consequently, the one of *MYB72*. For example, both *FIT* and *MYB72* are upregulated by ET treatment (García et al., 2010). In relation to FIT regulation, it has been shown that the ET-signaling TFs EIN3 and EIL1 interact with FIT to favor its stability and activity (Lingam et al., 2011; Yang et al., 2014; Brumbarova et al., 2015; see Subsection “Ethylene and other hormones and signaling molecules”). The upregulation of *MYB72* in the *ein2-1* mutant could be explained by taking into account the upregulation of *FIT*, *bHLH38*, and *bHLH39* in this mutant (García et al., 2010). This could be related to the existence of an alternate route for ethylene signaling, besides the conventional one including EIN2 (Shakeel et al., 2013; Lucena et al., 2015). The complexity of the relationship between *MYB72* and ET is also manifested when considering that *MYB72* can also affect ET biosynthesis: the *SAM1* gene, encoding a SAM synthetase enzyme involved in ET synthesis, is upregulated by *MYB72* (Figure 2; Zamioudis et al., 2014) and also is upregulated under Fe deficiency in Arabidopsis roots (García et al., 2010). In accordance with the influence of *MYB72* on ET, the exogenous application of ACC induced wild-type levels of resistance in the Arabidopsis *myb72-1* mutant, suggesting that *MYB72* acts upstream of ET in the ISR pathway (Van der Ent et al., 2008).

Besides the participation of the *MYB72*, *MYB10*, *FIT*, *bHLH38*, and *bHLH39* TFs in both ISR and Fe deficiency responses, there are other TFs that also play a key role in both processes. Among them, the EIN3/EIL1 (related to ET signaling) and MED16 (Mediator) TFs can be highlighted (see Subsection “Ethylene and other hormones and signaling molecules”). These TFs are required to activate the expression of Fe acquisition genes by interacting with FIT (Figure 2; Lingam et al., 2011; Yang et al., 2014; Zhang et al., 2014; Brumbarova et al., 2015; Lucena et al., 2015). In relation to ISR, the Arabidopsis *ein3-1* mutant did not express ISR in response to treatment with the bacterium *P. simiae* WCS417, which suggests that the EIN3 TF also plays a key role in this process (Knoester et al., 1999). MED16 is a key component in the JA/ET-mediated immunity against necrotrophic pathogens (Wang et al., 2015). In addition, and besides affecting the expression of Fe acquisition genes, it greatly influences the expression of *MYB72* and *MYB10* (Zhang et al., 2014), two important components in both Fe deficiency responses and ISR (see above). Probably, the physical interaction of MED16 with FIT is necessary for the activation of *MYB72* and *MYB10* expression (Figure 2).

TABLE 2 | Microbial species that improve Fe nutrition when applied to dicot plants grown in calcareous soils (or in artificial calcareous soils).

Microbial species	Plant species	Mode appl.	Fe def. resp.	Fe Gr.	Refs
Rhizobacteria					
<i>Bacillus subtilis</i>	<i>Manihot esculenta</i>	ri	nd	Fe \wedge	Freitas et al., 2015
<i>Paenibacillus polymyxa</i>	<i>Arabidopsis thaliana</i>	ri	nd	Fe	Zhou et al., 2016a
<i>Bacillus</i> sp.	<i>Pyrus communis</i>	ri	FCR organic acids	Fe	Ipek et al., 2017
<i>Agrobacterium</i> sp.*					
<i>Alcaligenes</i> sp.*					
<i>Pantoea</i> sp.*					
<i>Alcaligenes</i> sp.*	<i>Malus domestica</i>	i	FCR organic acids	Fe	Aras et al., 2018
<i>Pantoea</i> sp.*					
<i>Bacillus</i> sp.	<i>Prunus persica</i>	ri	FCR organic acids	Fe	Ankan et al., 2018
<i>Agrobacterium</i> sp.*					
<i>Alcaligenes</i> sp.*					
<i>Staphylococcus</i> sp.*					
Rhizofungi					
<i>Trichoderma asperellum</i>	<i>Lupinus albus</i>	gm and ri	nd	Fe	de Santiago et al., 2009
<i>Trichoderma asperellum</i>	<i>Cucumis sativus</i>	ri	nd	Fe	de Santiago et al., 2013

Mode appl., mode of application of the microbes (gm, application to the growth medium; i, through the irrigation system; ri, root immersion). FCR, ferric chelate reductase activity; Fe Gr., increased shoot Fe concentration (Fe) and increased shoot growth (\wedge); nd, not determined; *, microbial species whose association with ISR is not yet clear.

Internal Fe Content

Ethylene, auxin, and NO greatly activate the expression of Fe acquisition genes in plants grown with low levels of Fe (or without Fe), but have much less effect in plants grown with high levels of Fe (Lucena et al., 2006; Graziano and Lamattina, 2007; Chen et al., 2010; García et al., 2011). This suggests that the upregulation of Fe acquisition genes does not solely depend on hormones and signaling molecules (such as ET, auxin, or NO), that would act as activators, but also on the internal Fe content of plants, that would act as a repressor (Lucena et al., 2006; García et al., 2011, 2013, 2018; Romera et al., 2011, 2017). However, different results suggest that total Fe in roots is not the repressor of Fe acquisition genes but instead it is a Fe-related signal moving from shoots to roots through the phloem (García et al., 2013, 2018). Very recently, it has been found that this shoot Fe-related signal can affect the synthesis of ET on roots (García et al., 2018). To integrate all these regulatory components, a model has been proposed where ET/auxin/NO act as activators of Fe acquisition genes while a phloem Fe-related signal acts as repressor (García et al., 2011, 2013, 2018; Lucena et al., 2015; Romera et al., 2017).

In relation to ISR, there are also several results showing that the effects of ISR-eliciting microbes on the induction of Fe deficiency responses could also be dependent on the Fe concentration of the medium. For example, the expression of two Fe-related genes, like *MYB72*, and *FRO2*, was induced by *P. simiae* WCS417 independently of the Fe concentration in the medium but their absolute values were decreased when the Fe concentration increased (Zamioudis et al., 2015). Other examples, the expression of *FIT*, *IRT1* and *FRO2*, and the ferric reductase activity, decreased in *Arabidopsis* plants inoculated with *P. polymyxa* BFKC01 when the Fe concentration in the medium increased (Zhou et al., 2016a). The expression of *FIT*, *IRT1*, *FRO1*, and *HA1*, and the ferric reductase activity, in cucumber plants inoculated with *Azospirillum brasilense*

Cd(DSM-1843) or other rhizobacterial species, decreased when the Fe concentration in the medium increased (Pii et al., 2016b; Scagliola et al., 2016). The acidification capacity induced by the rhizobacterium *Arthrobacter agilis* in *Medicago truncatula* roots was lower in plants grown with high levels of Fe than in those grown with low levels of Fe (Orozco-Mosqueda et al., 2013). Moreover, the expression of several Fe-related genes induced by *P. simiae* WCS417, like *MYB72*, *FRO2*, and *IRT1*, is transitory (Zamioudis et al., 2015). This suggests that, after the induction of the Fe acquisition genes by ISR-eliciting microbes, plants acquire enough Fe and turn off these genes, to avoid toxicity and to conserve energy. The same occurs when plants induce the Fe acquisition genes under Fe deficiency and, as a consequence, they get sufficient Fe (Vert et al., 2003; Lucena et al., 2015).

RHIZOSPHERE MICROBIAL SPECIES THAT INDUCE Fe DEFICIENCY RESPONSES AND IMPROVE Fe ACQUISITION

As previously stated, beneficial rhizosphere microbes can contribute to improve Fe acquisition. This is most likely due to their capacity to induce Fe deficiency responses, such as enhanced ferric reductase activity, acidification of the rhizosphere, release of phenolics and flavins, and development of root hairs (see Section “Interrelationship between ISR and Fe deficiency responses in dicot plants”). These Fe deficiency responses are induced in a similar way as they are induced under Fe deficiency conditions. For example, ISR-eliciting microbes induce the upregulation of the genes associated with the Fe deficiency responses, like *FRO2*, *IRT1*, *AHA*, *F6'H1*, *BGLU42*, *ABCG37*, and others, and these genes are activated by the

TFs that activate them under Fe deficiency, like FIT(FER), bHLH38, bHLH39, MYB72, and MYB10 (Figure 2; Zhang et al., 2009; Palmer et al., 2013; Zamioudis et al., 2014, 2015; Zhao et al., 2014; Pii et al., 2016b; Scagliola et al., 2016; Verbon and Liberman, 2016; Zhou et al., 2016a, 2017, 2018; Martínez-Medina et al., 2017; Verbon et al., 2017; Wang J. et al., 2017; Stringlis et al., 2018a,b). Moreover, the hormones and signaling molecules related to the activation of these TFs, like ET, auxin, and NO, are similar in both ISR and Fe deficiency responses (see Subsection “Ethylene and other hormones and signaling molecules”).

Among the beneficial rhizosphere microbes that can activate the ISR are rhizobacteria, like *P. simiae* (syn. *P. fluorescens*), *B. subtilis*, *P. polymyxa* and *A. brasilense*; rhizofungi, like *Trichoderma* spp.; mycorrhizal fungi, like *Rhizophagus irregularis* (syn. *Glomus intraradices*) and *P. indica*; and non-pathogenic races of *Fusarium oxysporum* (Segarra et al., 2009; Zhang et al., 2009; Patil et al., 2011; Pieterse et al., 2012, 2014; Alizadeh et al., 2013; Zamioudis et al., 2014, 2015; Zhao et al., 2014; Pii et al., 2016b; Martínez-Medina et al., 2017; Verbon et al., 2017). In the case of mycorrhizal fungi, the enhanced defensive capacity provoked by them is also named MIR (“Mycorrhiza-Induced Resistance”) and can favor P acquisition (Jung et al., 2012; Cameron et al., 2013). Furthermore, it has been suggested that MIR can involve an ISR component elicited by bacteria in the mycorrhizosphere (Cameron et al., 2013). This is supported by the synergistic effects in defense responses when both arbuscular mycorrhiza and rhizobacteria are simultaneously applied (Pérez de Luque et al., 2017). The synergy between both kind of microbes paves the way to study the consortia of mycorrhiza and rhizobacteria in relation to the acquisition of Fe and P, and perhaps of other nutrients.

In Table 1 are summarized several rhizobacteria and rhizofungi species (most of them trigger ISR) that have been shown to induce Fe deficiency responses and to improve Fe acquisition and/or growth when applied to dicot plants. In Table 2 are summarized the ones that have been shown to cause similar effects when applied to dicot plants grown in calcareous soils (or in artificial calcareous soils).

CONCLUDING REMARKS AND FUTURE PERSPECTIVES

The ability of ISR-eliciting microbes to trigger both defense responses and Fe deficiency responses opens the way to use them as biopesticides and also as Fe biofertilizers. This represents a very important opportunity to diminish the application of fertilizers and pesticides in a more sustainable agriculture. However, and in relation to Fe nutrition, the use of ISR-eliciting microbes is in its infancy since it is not sufficiently known the behavior of these microbes on crops grown in calcareous soils. Most research works about the relationship of these microbes with the Fe nutrition of plants have been carried out with Arabidopsis plants grown on agar plates. For their application to crop plants in the field it would be also necessary to study the behavior of these microbes with plant species growing in calcareous

soils, including their capacity to thrive in these soils and to compete with wild soil microbes. More research is also needed to know the best ways for their application, by analyzing and comparing, both biologically and economically, the different possibilities, like direct application to soil, root immersion of plantlets before transplanting them (in the case of crop trees), application to seeds, and application into the irrigation systems (probably as spores). In the same way, it is necessary to study whether it is better to apply individual microbial species or consortia of different microbial species (Alizadeh et al., 2013; Sonbarse et al., 2017). In this latter case, and since ET can play a dual role in both ISR and Fe deficiency responses, it would be interesting to analyze the interactions between plant species and microbial species possessing the ACC deaminase enzyme and those that do not. Anyway, the research about ISR-eliciting microbes and Fe nutrition is a very fascinating topic for the near future.

AUTHOR'S NOTE

We apologize to authors whose works were not cited in this review due to our ignorance and to manuscript length restrictions. We encourage authors with papers relating ISR and Fe nutrition in dicot plants to send them to us: we are preparing a website (<http://www.uco.es/rhizoferrum>) to keep updated all the microbial species that elicit ISR, induce Fe deficiency responses, and improve Fe acquisition.

AUTHOR CONTRIBUTIONS

FR, MG, and CL revised the information related to ISR and Fe deficiency signaling. RP-V revised the information related to volatile compounds. EA, JR, MAA, and MA revised the information related to the different microbial species and their effects on Fe nutrition. FR wrote a first draft of the manuscript. JR, RP-V, and AM-M corrected and improved the manuscript.

FUNDING

This work was supported by “Plan Propio” Universidad de Córdoba (Spain), the “Junta de Andalucía” (Research Groups AGR115, BIO159, and BIO202) and “LABORATORIOS JAER S.A.” (company related to the production of micronutrient fertilizers which is starting to work on micronutrient biofertilizers). The funder played no role in the study design, the collection, analysis or interpretation of data, the writing of this paper or the decision to submit it for publication.

ACKNOWLEDGMENTS

We thank Dr. Brian M. Waters of the University of Nebraska (Lincoln, United States), for the English correction and valuable suggestions in the editing of the manuscript, and Gabriel Martí (“LABORATORIOS JAER S.A.”) for funding the publication of this Review.

REFERENCES

- Acharya, K., Chandra, S., Chakraborty, N., and Acharya, R. (2011). Nitric oxide functions as a signal in induced systemic resistance. *Arch. Phytopathol. Plant Protect.* 44, 1335–1342. doi: 10.1080/03235408.2010.496552
- Alizadeh, H., Behboudi, K., Ahmadzadeh, M., Javan-Nikkhah, M., Zamioudis, C., Pieterse, C. M. J., et al. (2013). Induced systemic resistance in cucumber and *Arabidopsis thaliana* by the combination of *Trichoderma harzianum* Tr6 and *Pseudomonas* sp. Ps14. *Biol. Control* 65, 14–23. doi: 10.1016/j.biocontrol.2013.01.009
- Aparicio, F., and Pallás, V. (2017). The coat protein of Alfalfa mosaic virus interacts and interferes with the transcriptional activity of the bHLH transcription factor ILR3 promoting salicylic acid-dependent defence signaling response. *Mol. Plant Pathol.* 18, 173–186. doi: 10.1111/mpp.12388
- Aras, S., Arıkan, S., İpek, M., Esitken, A., Pırlak, L., Dönmez, M. F., et al. (2018). Plant growth promoting rhizobacteria enhanced leaf organic acids, FC-R activity and Fe nutrition of apple under lime soil conditions. *Acta Physiol. Plant.* 40:120. doi: 10.1007/s11738-018-2693-9
- Arıkan, S., Esitken, A., İpek, M., Aras, S., Sahin, M., Pırlak, L., et al. (2018). Effect of plant growth promoting rhizobacteria on Fe acquisition in peach (*Prunus persica* L.) under calcareous soil conditions. *J. Plant Nutr.* 41, 2141–2150. doi: 10.1080/01904167.2018.1482910
- Asari, S., Tarkowská, D., Rolčík, J., Novák, O., Velázquez-Palmero, D., Bejai, S., et al. (2017). Analysis of plant growth-promoting properties of *Bacillus amyloliquefaciens* UCMB5113 using *Arabidopsis thaliana* as host plant. *Planta* 245, 15–30. doi: 10.1007/s00425-016-2580-9
- Aznar, A., Chen, N. W., Rigault, M., Riache, N., Joseph, D., Desmaële, D., et al. (2014). Scavenging iron: a novel mechanism of plant immunity activation by microbial siderophores. *Plant Physiol.* 164, 2167–2183. doi: 10.1104/pp.113.233585
- Aznar, A., Chen, N. W. G., Thomine, S., and Dellagi, A. (2015). Immunity to plant pathogens and iron homeostasis. *Plant Sci.* 240, 90–97. doi: 10.1016/j.plantsci.2015.08.022
- Aznar, A., and Dellagi, A. (2015). New insights into the role of siderophores as triggers of plant immunity: what can we learn from animals? *J. Exp. Bot.* 66, 3001–3010. doi: 10.1093/jxb/erv155
- Bacaicoa, E., Mora, V., Zamarreño, A. M., Fuentes, M., Casanova, E., and García-Mina, J. M. (2011). Auxin: a major player in the shoot-to-root regulation of root Fe-stress physiological responses to Fe deficiency in cucumber plants. *Plant Physiol. Biochem.* 49, 545–556. doi: 10.1016/j.plaphy.2011.02.018
- Bacaicoa, E., Zamarreño, A. M., Leménager, D., Baigorri, R., and García-Mina, J. M. (2009). Relationship between the hormonal balance and the regulation of iron deficiency stress responses in cucumber. *J. Am. Soc. Hortic. Sci.* 134, 589–601. doi: 10.21273/JASHS.134.6.589
- Bakker, P. A. H. M., Doornbos, R. F., Zamioudis, C., Berendsen, R. L., and Pieterse, C. M. J. (2013). Induced systemic resistance and the rhizosphere microbiome. *Plant Pathol. J.* 29, 136–143. doi: 10.5423/PPJ.SI.07.2012.0111
- Balmer, D., Planchamp, C., and Mauch-Mani, B. (2013). On the move: induced resistance in monocots. *J. Exp. Bot.* 64, 1249–1261. doi: 10.1093/jxb/ers248
- Bauer, P., Ling, H. Q., and Guerinot, M. L. (2007). FIT, the FER-LIKE IRON DEFICIENCY INDUCED TRANSCRIPTION FACTOR in *Arabidopsis*. *Plant Physiol. Biochem.* 45, 260–261. doi: 10.1016/j.plaphy.2007.03.006
- Berendsen, R. L., Pieterse, C. M., and Bakker, P. A. (2012). The rhizosphere microbiome and plant health. *Trends Plant Sci.* 17, 478–486. doi: 10.1016/j.tplants.2012.04.001
- Boutrot, F., Segonzac, C., Chang, K. N., Qiao, H., Ecker, J. R., Zipfel, C., et al. (2010). Direct transcriptional control of the *Arabidopsis* immune receptor FLS2 by the ethylene-dependent transcription factors EIN3 and EIL1. *Proc. Natl. Acad. Sci. U.S.A.* 107, 14502–14507. doi: 10.1073/pnas.1003347107
- Briat, J. F., Dubos, C., and Gaymard, F. (2015). Iron nutrition, biomass production, and plant product quality. *Trends Plant Sci.* 20, 33–40. doi: 10.1016/j.tplants.2014.07.005
- Brumbarova, T., Bauer, P., and Ivanov, R. (2015). Molecular mechanisms governing *Arabidopsis* iron uptake. *Trends Plant Sci.* 20, 124–133. doi: 10.1016/j.tplants.2014.11.004
- Camehl, I., Sherameti, I., Venus, Y., Bethke, G., Varma, A., Lee, J., et al. (2010). Ethylene signalling and ethylene-targeted transcription factors are required to balance beneficial and nonbeneficial traits in the symbiosis between the endophytic fungus *Piriformospora indica* and *Arabidopsis thaliana*. *New Phytol.* 185, 1062–1073. doi: 10.1111/j.1469-8137.2009.03149.x
- Cameron, D. D., Neal, A. L., van Wees, S. C. M., and Ton, J. (2013). Mycorrhiza-induced resistance: more than the sum of its parts? *Trends Plant Sci.* 18, 539–545. doi: 10.1016/j.tplants.2013.06.004
- Chen, C. C., Chien, W. F., Lin, N. C., and Yeh, K. C. (2014). Alternative functions of *Arabidopsis* YELLOW STRIPE-LIKE3: from metal translocation to pathogen defense. *PLoS One* 9:e98008. doi: 10.1371/journal.pone.0098008
- Chen, W. W., Yang, J. L., Qin, C., Jin, C. W., Mo, J. H., Ye, T., et al. (2010). Nitric oxide acts downstream of auxin to trigger root ferric-chelate reductase activity in response to iron deficiency in *Arabidopsis thaliana*. *Plant Physiol.* 154, 810–819. doi: 10.1104/pp.110.161109
- Choudhary, D. K., Prakash, A., and Johri, B. N. (2007). Induced systemic resistance (ISR) in plants: mechanism of action. *Indian J. Microbiol.* 47, 289–297. doi: 10.1007/s12088-007-0054-2
- Colangelo, E. P., and Guerinot, M. L. (2004). The essential basic helix-loop-helix protein FIT1 is required for the iron deficiency response. *Plant Cell* 16, 3400–3412. doi: 10.1105/tpc.104.024315
- Contreras-Cornejo, H. A., López-Bucio, J. S., Méndez-Bravo, A., Macías-Rodríguez, L. I., Ramos-Vega, M., Guevara-García, Á. A., et al. (2015). Mitogen-activated protein kinase 6 and ethylene and auxin signaling pathways are involved in *Arabidopsis* root-system architecture alterations by *Trichoderma atroviride*. *Mol. Plant Microbe Interact.* 28, 701–710. doi: 10.1094/MPMI-01-15-0005-R
- de Santiago, A., García-López, A. M., Quintero, J. M., Avilés, M., and Delgado, A. (2013). Effect of *Trichoderma asperellum* strain T34 and glucose addition on iron nutrition in cucumber grown on calcareous soils. *Soil Biol. Biochem.* 57, 598–605. doi: 10.1016/j.soilbio.2012.06.020
- de Santiago, A., Quintero, J. M., Avilés, M., and Delgado, A. (2009). Effect of *Trichoderma asperellum* strain T34 on iron nutrition in white lupin. *Soil Biol. Biochem.* 41, 2453–2459. doi: 10.1016/j.soilbio.2009.07.033
- de Zelicourt, A., Al-Yousif, M., and Hirt, H. (2013). Rhizosphere microbes as essential partners for plant stress tolerance. *Mol. Plant* 6, 242–245. doi: 10.1093/mp/sst028
- Dubois, M., Van den Broeck, L., and Inzé, D. (2018). The pivotal role of ethylene in plant growth. *Trends Plant Sci.* 23, 311–323. doi: 10.1016/j.tplants.2018.01.003
- Durrett, T. P., Gassmann, W., and Rogers, E. E. (2007). The FRD3-mediated efflux of citrate into the root vasculature is necessary for efficient iron translocation. *Plant Physiol.* 144, 197–205. doi: 10.1104/pp.107.097162
- Fourcroy, P., Sisó-Terraza, P., Sudre, D., Saviróon, M., Reyt, G., Gaymard, F., et al. (2014). Involvement of the ABCG37 transporter in secretion of scopoletin and derivatives by *Arabidopsis* roots in response to iron deficiency. *New Phytol.* 201, 155–167. doi: 10.1111/nph.12471
- Fourcroy, P., Tissot, N., Reyt, G., Gaymard, F., Briat, J. F., and Dubos, C. (2016). Facilitated Fe nutrition by phenolic compounds excreted by the *Arabidopsis* ABCG37/PDR9 transporter requires the IRT1/FRO2 high-affinity root Fe²⁺ transport system. *Mol. Plant* 9, 485–488. doi: 10.1016/j.molp.2015.09.010
- Freitas, M. A., Medeiros, F. H. V., Carvalho, S. P., Guilherme, L. R. G., Teixeira, W. D., Zhang, H., et al. (2015). Augmenting iron accumulation in cassava by the beneficial soil bacterium *Bacillus subtilis* (GBO3). *Front. Plant Sci.* 6:596. doi: 10.3389/fpls.2015.00596
- Gamalerio, E., and Glick, B. R. (2015). Bacterial modulation of plant ethylene levels. *Plant Physiol.* 169, 13–22. doi: 10.1104/pp.15.00284
- García, M. J., Corpas, F. J., Lucena, C., Alcántara, E., Pérez-Vicente, R., Zamarreño, Á. M., et al. (2018). A shoot Fe signaling pathway requiring the OPT3 transporter controls GSNO Reductase and ethylene in *Arabidopsis thaliana* roots. *Front. Plant Sci.* 9:1325. doi: 10.3389/fpls.2018.01325
- García, M. J., Lucena, C., Romera, F. J., Alcántara, E., and Pérez-Vicente, R. (2010). Ethylene and nitric oxide involvement in the up-regulation of key genes related to iron acquisition and homeostasis in *Arabidopsis*. *J. Exp. Bot.* 61, 3885–3899. doi: 10.1093/jxb/erl189
- García, M. J., Romera, F. J., Stacey, M. G., Stacey, G., Villar, E., Alcántara, E., et al. (2013). Shoot to root communication is necessary to control the expression of iron-acquisition genes in Strategy I plants. *Planta* 237, 65–75. doi: 10.1007/s00425-012-1757-0
- García, M. J., Suárez, V., Romera, F. J., Alcántara, E., and Pérez-Vicente, R. (2011). A new model involving ethylene, nitric oxide and Fe to explain the regulation

- of Fe-acquisition genes in Strategy I plants. *Plant Physiol. Biochem.* 49, 537–544. doi: 10.1016/j.plaphy.2011.01.019
- García-López, A. M., Avilés, M., and Delgado, A. (2016). Effect of various microorganisms on phosphorus uptake from insoluble Ca-phosphates by cucumber plants. *J. Plant Nutr. Soil Sci.* 179, 454–465. doi: 10.1002/jpln.201500024
- Garnica-Vergara, A., Barrera-Ortiz, S., Muñoz-Parra, E., Raya-González, J., Méndez-Bravo, A., Macías-Rodríguez, L., et al. (2016). The volatile 6-pentyl-2H-pyran-2-one from *Trichoderma atroviride* regulates *Arabidopsis thaliana* root morphogenesis via auxin signaling and ETHYLENE INSENSITIVE 2 functioning. *New Phytol.* 209, 1496–1512. doi: 10.1111/nph.13725
- Grady, E. N., MacDonald, J., Liu, L., Richman, A., and Yuan, Z. C. (2016). Current knowledge and perspectives of *Paenibacillus*: a review. *Microb. Cell Fact.* 15:203. doi: 10.1186/s12934-016-0603-7
- Graziano, M., and Lamattina, L. (2007). Nitric oxide accumulation is required for molecular and physiological responses to iron deficiency in tomato roots. *Plant J.* 52, 949–960. doi: 10.1111/j.1365-313X.2007.03283.x
- Guinel, F. C. (2015). Ethylene, a hormone at the center-stage of nodulation. *Front. Plant Sci.* 6:1121. doi: 10.3389/fpls.2015.01121
- Gullner, G., Komives, T., Király, L., and Schröder, P. (2018). Glutathione S-transferase enzymes in plant-pathogen interactions. *Front. Plant Sci.* 9:1836. doi: 10.3389/fpls.2018.01836
- Gullner, G., Zechmann, B., Künstler, A., and Király, L. (2017). “The signaling roles of glutathione in plant disease resistance,” in *Glutathione in Plant Growth, Development, and Stress Tolerance*, eds M. A. Hossain, M. G. Mostofa, P. D. Vivancos, D. J. Burritt, M. Fujita, and L. S. P. Tran (Cham: Springer International Publishing), 331–357. doi: 10.1007/978-3-319-66682-2_15
- Hindt, M. N., Akmakjian, G. Z., Pivarski, K. L., Punshon, T., Baxter, I., Salt, D. E., et al. (2017). BRUTUS and its paralogs, BTS LIKE1 and BTS LIKE2, encode important negative regulators of the iron deficiency response in *Arabidopsis thaliana*. *Metalomics* 9, 876–890. doi: 10.1039/c7mt00152e
- Hossain, M. M., Sultana, F., and Hyakumachi, M. (2017). Role of ethylene signalling in growth and systemic resistance induction by the plant growth-promoting fungus *Penicillium viridicatum* in *Arabidopsis*. *J. Phytopathol.* 165, 432–441. doi: 10.1111/jph.12577
- İpek, M., Aras, S., Arıkan, S., Esitken, A., Pırlak, L., Dönmez, M. F., et al. (2017). Root plant growth promoting rhizobacteria inoculations increase ferric chelate reductase (FC-R) activity and Fe nutrition in pear under calcareous soil conditions. *Sci. Hortic.* 219, 144–151. doi: 10.1016/j.scienta.2017.02.043
- İpek, M., and Esitken, A. (2017). “The actions of PGPR on micronutrient availability in soil and plant under calcareous soil conditions: an evaluation over Fe nutrition,” in *Plant-Microbe Interactions in Agro-Ecological Perspectives*, Vol. 2, eds D. Singh, H. Singh, and R. Prabha (Singapore: Springer), 81–100.
- Ivanov, R., Brumbarova, T., and Bauer, P. (2012). Fitting into the harsh reality: regulation of iron-deficiency responses in dicotyledonous plants. *Mol. Plant* 5, 27–42. doi: 10.1093/mp/psr065
- Jankiewicz, U., and Koltonowicz, M. (2012). The involvement of *Pseudomonas* bacteria in induced systemic resistance in plants (Review). *Appl. Biochem. Microbiol.* 48, 244–249. doi: 10.1134/S0003683812030052
- Jin, C. W., Ye, Y. Q., and Zheng, S. J. (2014). An underground tale: contribution of microbial activity to plant iron acquisition via ecological processes. *Ann. Bot.* 113, 7–18. doi: 10.1093/aob/mct249
- Jung, S. C., Martínez-Medina, A., Lopez-Raez, J. A., and Pozo, M. J. (2012). Mycorrhiza-Induced Resistance and priming of plant defenses. *J. Chem. Ecol.* 38, 651–664. doi: 10.1007/s10886-012-0134-6
- Kabir, A. H., Paltridge, N. G., Able, A. J., Paull, J. G., and Stangoulis, J. C. R. (2012). Natural variation for Fe-efficiency is associated with up-regulation of Strategy I mechanisms and enhanced citrate and ethylene synthesis in *Pisum sativum* L. *Planta* 235, 1409–1419. doi: 10.1007/s00425-011-1583-9
- Kailasam, S., Wang, Y., Lo, J. C., Chang, H. F., and Yeh, K. C. (2018). S-nitrosoglutathione works downstream of nitric oxide to mediate iron deficiency signaling in *Arabidopsis*. *Plant J.* 94, 157–168. doi: 10.1111/tj.13850
- Kang, H. G., Foley, R. C., Onate-Sanchez, L., Lin, C., and Singh, K. B. (2003). Target genes for OBP3, a Dof transcription factor, include novel basic helix-loop-helix domain proteins inducible by salicylic acid. *Plant J.* 35, 362–372. doi: 10.1046/j.1365-313X.2003.01812.x
- Kieu, N. P., Aznar, A., Segond, D., Rigault, M., Simond-Côte, E., Kunz, C., et al. (2012). Iron deficiency affects plant defence responses and confers resistance to *Dickeya dadantii* and *Botrytis cinerea*. *Mol. Plant Pathol.* 13, 816–827. doi: 10.1111/J.1364-3703.2012.00790.X
- Knoester, M., Pieterse, C. M. J., Bol, J. F., and Van Loon, L. C. (1999). Systemic resistance in *Arabidopsis* induced by rhizobacteria requires ethylene-dependent signaling at the site of application. *Mol. Plant Microbe Interact.* 12, 720–727. doi: 10.1094/MPMI.1999.12.8.720
- Kobayashi, T., Itai, R. N., Senoura, T., Oikawa, T., Ishimaru, Y., Ueda, M., et al. (2016). Jasmonate signaling is activated in the very early stages of iron deficiency responses in rice roots. *Plant Mol. Biol.* 91, 533–547. doi: 10.1007/s11103-016-0486-3
- Kobayashi, T., and Nishizawa, N. K. (2012). Iron uptake, translocation, and regulation in higher plants. *Annu. Rev. Plant Biol.* 63, 131–152. doi: 10.1146/annurev-arplant-042811-10552
- Koen, E., Szymańska, K., Klinguer, A., Dobrowolska, G., Besson-Bard, A., and Wendehenne, D. (2012). Nitric oxide and glutathione impact the expression of iron uptake- and iron transport-related genes as well as the content of metals in *A. thaliana* plants grown under iron deficiency. *Plant Signal. Behav.* 7, 1246–1250. doi: 10.4161/psb.21548
- Kudoyarova, G. R., Vysotskaya, L. B., Arkhipova, T. N., Kuzmina, L. Y., Galimsyanova, N. F., Sidorova, L. V., et al. (2017). Effect of auxin producing and phosphate solubilizing bacteria on mobility of soil phosphorus, growth rate, and P acquisition by wheat plants. *Acta Physiol. Plant.* 39:253. doi: 10.1007/s11738-017-2556-9
- Kumar, R. K., Chu, H. H., Abundis, C., Vasques, K., Rodriguez, D. C., Chia, J. C., et al. (2017). Iron-nicotianamine transporters are required for proper long distance iron signaling. *Plant Physiol.* 175, 1254–1268. doi: 10.1104/pp.17.00821
- Kwon, Y. S., Ryu, C.-M., Lee, S., Park, H. B., Han, K. S., Lee, J. H., et al. (2010). Proteome analysis of *Arabidopsis* seedlings exposed to bacterial volatiles. *Planta* 232, 1355–1370. doi: 10.1007/s00425-010-1259-x
- Lemanceau, P., Expert, D., Gaymard, F., Bakker, P. A. H. M., and Briat, J. F. (2009). Role of iron in plant-microbe interactions. *Adv. Bot. Res.* 51, 491–549. doi: 10.1016/S0065-2296(09)51012-9
- Li, G., Meng, X., Wang, R., Mao, G., Han, L., Liu, Y., et al. (2012). Dual-level regulation of ACC synthase activity by MPK3/MPK6 cascade and its downstream WRKY transcription factor during ethylene induction in *Arabidopsis*. *PLoS Genet.* 8:e1002767. doi: 10.1371/journal.pgen.1002767
- Li, R.-X., Cai, F., Pang, G., Shen, Q.-R., Li, R., and Chen, W. (2015). Solubilisation of phosphate and micronutrients by *Trichoderma harzianum* and its relationship with the promotion of tomato plant growth. *PLoS One* 10:e0130081. doi: 10.1371/journal.pone.0130081
- Li, W., and Lan, P. (2017). The understanding of the plant iron deficiency responses in Strategy I plants and the role of ethylene in this process by omic approaches. *Front. Plant Sci.* 8:40. doi: 10.3389/fpls.2017.00040
- Li, X., Zhang, H., Ai, Q., Liang, G., and Yu, D. (2016). Two bHLH transcription factors, bHLH34 and bHLH104, regulate iron homeostasis in *Arabidopsis thaliana*. *Plant Physiol.* 170, 2478–2493. doi: 10.1104/pp.15.01827
- Liang, G., Zhang, H., Li, X., Ai, Q., and Yu, D. (2017). bHLH transcription factor bHLH115 regulates iron homeostasis in *Arabidopsis thaliana*. *J. Exp. Bot.* 68, 1743–1755. doi: 10.1093/jxb/erx043
- Lin, X. Y., Ye, Y. Q., Fan, S. K., Jin, C. W., and Zheng, S. J. (2016). Increased sucrose accumulation regulates iron-deficiency responses by promoting auxin signaling in *Arabidopsis* plants. *Plant Physiol.* 170, 907–920. doi: 10.1104/pp.15.01598
- Lingam, S., Mohrbacher, J., Brumbarova, T., Potuschak, T., Fink-Straube, C., Blondet, E., et al. (2011). Interaction between the bHLH transcription factor FIT and the ETHYLENE INSENSITIVE3/ETHYLENE INSENSITIVE3-LIKE1 reveals molecular linkage between the regulation of iron acquisition and ethylene signaling in *Arabidopsis*. *Plant Cell* 23, 1815–1829. doi: 10.1105/tpc.111.084715
- Liu, W., Karemera, N. J. U., Wu, T., Yang, Y., Zhang, X., Xu, X., et al. (2017a). The ethylene response factor ATERF4 negatively regulates the iron deficiency response in *Arabidopsis thaliana*. *PLoS One* 12:e0186580. doi: 10.1371/journal.pone.0186580
- Liu, W., Li, Q., Wang, Y., Wu, T., Yang, Y., Zhang, X., et al. (2017b). Ethylene response factor ATERF72 negatively regulates *Arabidopsis thaliana* response to iron deficiency. *Biochem. Biophys. Res. Commun.* 491, 862–868. doi: 10.1016/j.bbrc.2017.04.014
- Liu, X. M., and Zhang, H. (2015). The effects of bacterial volatile emissions on plant abiotic stress tolerance. *Front. Plant Sci.* 6:774. doi: 10.3389/fpls.2015.00774

- López-Berges, M., Turra, D., Capilla, J., Schaffner, L., Matthijs, S., Jöchl, C., et al. (2013). Iron competition in fungus-plant interactions: the battle takes place in the rhizosphere. *Plant Signal. Behav.* 8:e23012. doi: 10.4161/psb.23012
- Lucena, C., Romera, F. J., García, M. J., Alcántara, E., and Pérez-Vicente, R. (2015). Ethylene participates in the regulation of Fe deficiency responses in Strategy I plants and in rice. *Front. Plant Sci.* 6:1056. doi: 10.3389/fpls.2015.01056
- Lucena, C., Romera, F. J., Rojas, C. L., García, M. J., Alcántara, E., and Pérez-Vicente, R. (2007). Bicarbonate blocks the expression of several genes involved in the physiological responses to Fe deficiency of Strategy I plants. *Funct. Plant Biol.* 34, 1002–1009. doi: 10.1071/FP07136
- Lucena, C., Waters, B. M., Romera, F. J., García, M. J., Morales, M., Alcántara, E., et al. (2006). Ethylene could influence ferric reductase, iron transporter and H⁺-ATPase gene expression by affecting FER (or FER-like) gene activity. *J. Exp. Bot.* 57, 4145–4154. doi: 10.1093/jxb/erl189
- Martínez-Medina, A., Flors, V., Heil, M., Mauch-Mani, B., Pieterse, C. M. J., Pozo, M. J., et al. (2016). Recognizing plant defense priming. *Trends Plant Sci.* 21, 818–822. doi: 10.1016/j.tplants.2016.07.009
- Martínez-Medina, A., Van Wees, S. C. M., and Pieterse, C. M. J. (2017). Airborne signals from *Trichoderma* fungi stimulate iron uptake responses in roots resulting in priming of jasmonic acid dependent defences in shoots of *Arabidopsis thaliana* and *Solanum lycopersicum*. *Plant Cell Environ.* 40, 2691–2705. doi: 10.1111/pce.13016
- Maurer, F., Müller, S., and Bauer, P. (2011). Suppression of Fe deficiency gene expression by jasmonate. *Plant Physiol. Biochem.* 49, 530–536. doi: 10.1016/j.plaphy.2011.01.025
- Maurer, F., Naranjo Arcos, M. A., and Bauer, P. (2014). Responses of a triple mutant defective in three iron deficiency-induced Basic Helix-Loop-Helix genes of the subgroup Ib(2) to iron deficiency and salicylic acid. *PLoS One* 9:e99234. doi: 10.1371/journal.pone.0099234
- Meiser, J., Lingam, S., and Bauer, P. (2011). Post-transcriptional regulation of the Fe deficiency bHLH transcription factor FIT is affected by iron and nitric oxide. *Plant Physiol.* 157, 2154–2166. doi: 10.1104/pp.111.183285
- Meziane, H., Van der Sluis, I., Van Loon, L. C., Höfte, M., and Bakker, P. A. (2005). Determinants of *Pseudomonas putida* WCS358 involved in inducing systemic resistance in plants. *Mol. Plant Pathol.* 6, 177–185. doi: 10.1111/j.1364-3703.2005.00276.x
- Mimmo, T., Del Buono, D., Terzano, R., Tomasi, N., Vigani, G., Crecchio, C., et al. (2014). Rhizospheric organic compounds in the soil-microorganism-plant system: their role in iron availability. *Eur. J. Soil Sci.* 65, 629–642. doi: 10.1111/ejss.12158
- Naranjo-Arcos, M. A., and Bauer, P. (2016). “Iron nutrition, oxidative stress, and pathogen defense,” in *Nutritional Deficiency*, eds P. Erkekoglu and B. Kocer-Gumusel (Rijeka: InTechOpen), 63–98. doi: 10.5772/63204
- Nascimento, F. X., Rossi, M. J., and Glick, B. R. (2018). Ethylene and 1-aminocyclopropane-1-carboxylate (ACC) in plant-bacterial interactions. *Front. Plant Sci.* 9:114. doi: 10.3389/fpls.2018.00114
- Nie, P., Li, X., Wang, S., Guo, J., Zhao, H., and Niu, D. (2017). Induced systemic resistance against *Botrytis cinerea* by *Bacillus cereus* AR156 through a JA/ET- and NPR1-dependent signaling pathway and activates PAMP-triggered immunity in *Arabidopsis*. *Front. Plant Sci.* 8:238. doi: 10.3389/fpls.2017.00238
- Orozco-Mosqueda, M. C., Velázquez-Becerra, C., Macías-Rodríguez, L. I., Santoyo, G., Flores-Cortez, I., Alfaro-Cuevas, R., et al. (2013). *Arthrobacter agilis* UMCV2 induces iron acquisition in *Medicago truncatula* (Strategy I plant) in vitro via dimethylhexadecylamine emission. *Plant Soil* 362, 51–66. doi: 10.1007/s11104-012-1263-y
- Palmer, C. M., Hind, M. N., Schmidt, H., Clemens, S., and Guerinot, M. L. (2013). MYB10 and MYB72 are required for growth under iron-limiting conditions. *PLoS Genet.* 9:e1003953. doi: 10.1371/journal.pgen.1003953
- Patel, T., and Saraf, M. (2017). Biosynthesis of phytohormones from novel rhizobacterial isolates and their in vitro plant growth-promoting efficacy. *J. Plant Interact.* 12, 480–487. doi: 10.1080/17429145.2017.1392625
- Patil, S., Sriram, S., Savitha, M. J., and Arulmani, N. (2011). Induced systemic resistance in tomato by non-pathogenic *Fusarium* species for the management of *Fusarium* wilt. *Arch. Phytopathol. Plant Protect.* 44, 1621–1634. doi: 10.1080/03235408.2010.526774
- Pérez de Luque, A., Tille, S., Johnson, I., Pascual-Pardo, D., Ton, J., and Cameron, D. D. (2017). The interactive effects of arbuscular mycorrhiza and plant growth-promoting rhizobacteria synergistically enhance host plant defences against pathogens. *Sci. Rep.* 7:16409. doi: 10.1038/s41598-017-16697-4
- Pierik, R., Sasidharan, R., and Voesenek, L. A. C. J. (2007). Growth control by ethylene: adjusting phenotypes to the environment. *J. Plant Growth Regul.* 26, 188–200. doi: 10.1007/s00344-006-0124-4
- Pieterse, C. M. J., Van der Does, D., Zamioudis, C., Leon-Reyes, A., and Van Wees, S. C. M. (2012). Hormonal modulation of plant immunity. *Annu. Rev. Cell Dev. Biol.* 28, 489–521. doi: 10.1146/annurev-cellbio-092910-154055
- Pieterse, C. M. J., Van Wees, S. C. M., Ton, J., Van Pelt, J. A., and Van Loon, L. C. (2002). Signalling in rhizobacteria-Induced systemic resistance in *Arabidopsis thaliana*. *Plant Biol.* 4, 535–544. doi: 10.1055/s-2002-354411
- Pieterse, C. M. J., Van Wees, S. C. M., van Pelt, J. A., Knoester, M., Laan, R., Gerrits, H., et al. (1998). A novel signaling pathway controlling induced systemic resistance in *Arabidopsis*. *Plant Cell* 10, 1571–1580. doi: 10.1105/tpc.10.9.1571
- Pieterse, C. M. J., Zamioudis, C., Berendsen, R. L., Weller, D. M., Van Wees, S. C. M., and Bakker, P. A. H. M. (2014). Induced systemic resistance by beneficial microbes. *Annu. Rev. Phytopathol.* 52, 347–375. doi: 10.1146/annurev-phyto-082712-102340
- Pii, Y., Borruso, L., Brusetti, L., Crecchio, C., Cesco, S., and Mimmo, T. (2016a). The interaction between iron nutrition, plant species and soil type shapes the rhizosphere microbiome. *Plant Physiol. Biochem.* 99, 39–48. doi: 10.1016/j.plaphy.2015.12.002
- Pii, Y., Marastoni, L., Springeth, C., Fontanella, M. C., Beone, G. M., Cesco, S., et al. (2016b). Modulation of Fe acquisition process by *Azospirillum brasilense* in cucumber plants. *Environ. Exp. Bot.* 130, 216–225. doi: 10.1016/j.envexpbot.2016.06.011
- Pii, Y., Mimmo, T., Tomasi, N., Terzano, R., Cesco, S., and Crecchio, C. (2015). Microbial interactions in the rhizosphere: beneficial influences of plant growth-promoting rhizobacteria on nutrient acquisition process. A review. *Biol. Fertil. Soils* 51, 403–415. doi: 10.1007/s00374-015-0996-1
- Pommerrenig, B., Feussner, K., Zierer, W., Rabinovych, V., Klebl, F., Feussner, I., et al. (2011). Phloem-specific expression of Yang cycle genes and identification of novel Yang cycle enzymes in *Plantago* and *Arabidopsis*. *Plant Cell* 23, 1904–1919. doi: 10.1105/tpc.110.079657
- Poupin, M. J., Greve, M., Carmona, V., and Pinedo, I. (2016). A complex molecular interplay of auxin and ethylene signaling pathways is involved in *Arabidopsis* growth promotion by *Burkholderia phytofirmans* PsJN. *Front. Plant Sci.* 7:492. doi: 10.3389/fpls.2016.00492
- Rajniak, J., Giehl, R. F. H., Chang, E., Murgia, I., von Wirén, N., and Sattely, E. S. (2018). Biosynthesis of redox-active metabolites in response to iron deficiency in plants. *Nat. Chem. Biol.* 14, 442–450. doi: 10.1038/s41589-018-0019-2
- Ribaudo, C. M., Krumpal, E. M., Cassán, F. D., Bottini, R., Cantore, M. L., and Curá, J. A. (2006). *Azospirillum* sp. promotes root hair development in tomato plants through a mechanism that involves ethylene. *J. Plant Growth Regul.* 24, 175–185. doi: 10.1007/s00344-005-0128-5
- Rodríguez-Celma, J., and Schmidt, W. (2013). Reduction-based iron uptake revisited. On the role of secreted iron-binding compounds. *Plant Signal. Behav.* 8:e26116. doi: 10.4161/psb.26116
- Romera, F. J., and Alcántara, E. (1994). Iron-deficiency stress responses in cucumber (*Cucumis sativus* L.) roots. A possible role for ethylene? *Plant Physiol.* 105, 1133–1138.
- Romera, F. J., and Alcántara, E. (2004). Ethylene involvement in the regulation of Fe-deficiency stress responses by Strategy I plants. *Funct. Plant Biol.* 31, 315–328. doi: 10.1071/FP03165
- Romera, F. J., Alcántara, E., and De la Guardia, M. D. (1999). Ethylene production by Fe-deficient roots and its involvement in the regulation of Fe-deficiency stress responses by Strategy I plants. *Ann. Bot.* 83, 51–55. doi: 10.1006/anbo.1998.0793
- Romera, F. J., García, M. J., Alcántara, E., and Pérez-Vicente, R. (2011). Latest findings about the interplay of auxin, ethylene and nitric oxide in the regulation of Fe deficiency responses by Strategy I plants. *Plant Signal. Behav.* 6, 167–170. doi: 10.4161/psb.6.1.14111
- Romera, F. J., Lucena, C., García, M. J., Alcántara, E., and Pérez-Vicente, R. (2017). “The role of ethylene and other signals in the regulation of Fe deficiency responses by dicot plants,” in *Stress Signaling in Plants: Genomics and Proteomics Perspectives*, Vol. 2, eds M. Sarwat, A. Ahmad, M. Z. Abidin, and M. Ibrahim (Cham: Springer), 277–300.

- Römheld, V., and Marschner, H. (1986). Mobilization of iron in the rhizosphere of different plant species. *Adv. Plant Nutr.* 2, 155–204. doi: 10.1007/BF02220801
- Ryu, C.-M., Hu, C.-H., Reddy, M. S., and Kloepper, J. W. (2003). Different signaling pathways of induced resistance by rhizobacteria in *Arabidopsis thaliana* against two pathogens of *Pseudomonas syringae*. *New Phytol.* 160, 413–420. doi: 10.1046/j.1469-8137.2003.00883.x
- Sauter, M., Moffatt, B., Saechao, M. C., Hell, R., and Wirtz, M. (2013). Methionine salvage and S-adenosylmethionine: essential links between sulfur, ethylene and polyamine biosynthesis. *Biochem. J.* 451, 145–154. doi: 10.1042/BJ20121744
- Savary, S., Ficke, A., Aubertot, J. N., and Hollier, C. (2012). Crop losses due to diseases and their implications for global food production losses and food security. *Food Secur.* 4, 519–537. doi: 10.1007/s12571-012-0200-5
- Scagliola, M., Pii, Y., Mimmo, T., Cesco, S., Ricciuti, P., and Crecchio, C. (2016). Characterization of plant growth promoting traits of bacterial isolates from the rhizosphere of barley (*Hordeum vulgare* L.) and tomato (*Solanum lycopersicon* L.) grown under Fe sufficiency and deficiency. *Plant Physiol. Biochem.* 107, 187–196. doi: 10.1016/j.plaphy.2016.06.002
- Schmid, N. B., Giehl, R. F. H., Döll, S., Mock, H. P., Strehmel, N., Scheel, D., et al. (2014). Feruloyl-CoA 6'-Hydroxylase1-dependent coumarins mediate iron acquisition from alkaline substrates in *Arabidopsis*. *Plant Physiol.* 164, 160–172. doi: 10.1104/pp.113.228544
- Schmidt, H., Günther, C., Weber, M., Spörlein, C., Loscher, S., Böttcher, C., et al. (2014). Metabolome analysis of *Arabidopsis thaliana* roots identifies a key metabolic pathway for iron acquisition. *PLoS One* 9:e102444. doi: 10.1371/journal.pone.0102444
- Segarra, G., Van der Ent, S., Trillas, I., and Pieterse, C. M. J. (2009). MYB72, a node of convergence in induced systemic resistance triggered by a fungal and a bacterial beneficial microbe. *Plant Biol.* 11, 90–96. doi: 10.1111/j.1438-8677.2008.00162.x
- Séguéla, M., Briat, J. F., Vert, G., and Curie, C. (2008). Cytokinins negatively regulate the root iron uptake machinery in *Arabidopsis* through a growth-dependent pathway. *Plant J.* 55, 289–300. doi: 10.1111/j.1365-313X.2008.03502.x
- Shakeel, S. N., Wang, X., Binder, B. M., and Schaller, G. E. (2013). Mechanisms of signal transduction by ethylene: overlapping and non-overlapping signalling roles in a receptor family. *AoB Plants* 5:lt010. doi: 10.1093/aobpla/plt010
- Shanmugam, V., Wang, Y. W., Tsendee, M., Karunakaran, K., and Yeh, K. C. (2015). Glutathione plays an essential role in nitric oxide-mediated iron-deficiency signaling and iron-deficiency tolerance in *Arabidopsis*. *Plant J.* 84, 464–477. doi: 10.1111/tpj.13011
- Sharifi, R., and Ryu, C. M. (2018). Sniffing bacterial volatile compounds for healthier plants. *Curr. Opin. Plant Biol.* 44, 88–97. doi: 10.1016/j.pbi.2018.03.004
- Shen, C., Yang, Y., Liu, K., Zhang, L., Guo, H., Sun, T., et al. (2016). Involvement of endogenous salicylic acid in iron-deficiency responses in *Arabidopsis*. *J. Exp. Bot.* 67, 4179–4193. doi: 10.1093/jxb/erw196
- Shen, J., Li, C., Mi, G., Li, L., Yuan, L., Jiang, R., et al. (2013). Maximizing root/rhizosphere efficiency to improve crop productivity and nutrient use efficiency in intensive agriculture of China. *J. Exp. Bot.* 64, 1181–1192. doi: 10.1093/jxb/ers342
- Shoresh, M., Yedidia, I., and Chet, I. (2005). Involvement of jasmonic acid/ethylene signaling pathway in the systemic resistance induced in cucumber by *Trichoderma asperellum* T203. *Phytopathology* 95, 76–84. doi: 10.1094/PHYTO-95-0076
- Singh, R. P., Shelke, G. M., Kumar, A., and Jha, P. N. (2015). Biochemistry and genetics of ACC deaminase: a weapon to “stress ethylene” produced in plants. *Front. Microbiol.* 6:937. doi: 10.3389/fmicb.2015.00937
- Sivitz, A. B., Hermand, V., Curie, C., and Vert, G. (2012). *Arabidopsis* bHLH100 and bHLH101 control iron homeostasis via a FIT-independent pathway. *PLoS One* 7:e44843. doi: 10.1371/journal.pone.0044843
- Siwinska, J., Siatkowska, K., Olry, A., Grosjean, J., Hehn, A., Bourgaud, F., et al. (2018). Scopoletin 8-hydroxylase: a novel enzyme involved in coumarin biosynthesis and iron-deficiency responses in *Arabidopsis*. *J. Exp. Bot.* 69, 1735–1748. doi: 10.1093/jxb/ery005
- Sonbarse, P. P., Sharma, P., and Parvatam, G. (2017). PGPR's mix treatment to *Moringa* improved plant growth and iron content in foliage as substantiated by biochemical and molecular methods. *J. Plant Interact.* 12, 526–532. doi: 10.1080/17429145.2017.1400125
- Stringlis, I. A., Proietti, S., Hickman, R., Van Verk, M. C., Zamioudis, C., and Pieterse, C. M. J. (2018a). Root transcriptional dynamics induced by beneficial rhizobacteria and microbial immune elicitors reveal signatures of adaptation to mutualists. *Plant J.* 93, 166–180. doi: 10.1111/tpj.13741
- Stringlis, I. A., Yua, K., Feussner, K., de Jonge, R., Van Bentum, S., Van Verk, M. C., et al. (2018b). MYB72-dependent coumarin exudation shapes root microbiome assembly to promote plant health. *Proc. Natl. Acad. Sci. U.S.A.* 115, E5213–E5222. doi: 10.1073/pnas.1722335115
- Sumayo, M. M., Son, J.-S., and Ghim, S.-Y. (2018). Exogenous application of phenylacetic acid promotes root hair growth and induces the systemic resistance of tobacco against bacterial soft-rot pathogen *Pectobacterium carotovorum* subsp. *carotovorum*. *Funct. Plant Biol.* 45, 1119–1127. doi: 10.1071/FP17332
- Ton, J., Davison, S., Van Wees, S. C. M., Van Loon, L. C., and Pieterse, C. M. J. (2001). The *Arabidopsis* ISR1 locus controlling rhizobacteria-mediated induced systemic resistance is involved in ethylene signaling. *Plant Physiol.* 125, 652–661. doi: 10.1104/pp.125.2.652
- Tsai, H. H., Rodríguez-Celma, J., Lan, P., Wu, Y. C., Vélez-Bermúdez, I. C., and Schmidt, W. (2018). Scopoletin 8-Hydroxylase-mediated fraxetin production is crucial for iron mobilization. *Plant Physiol.* 177, 194–207. doi: 10.1104/pp.18.00178
- Tsai, H. H., and Schmidt, W. (2017a). Mobilization of iron by plant-borne coumarins. *Trends Plant Sci.* 22, 538–548. doi: 10.1016/j.tplants.2017.03.008
- Tsai, H. H., and Schmidt, W. (2017b). One way. Or another? Iron uptake in plants. *New Phytol.* 214, 500–505. doi: 10.1111/nph.14477
- Tyagi, S., Mulla, S. I., Lee, K. J., Chae, J. C., and Shukla, P. (2018). VOCs-mediated hormonal signaling and crosstalk with plant growth promoting microbes. *Crit. Rev. Biotechnol.* 38, 1277–1296. doi: 10.1080/07388551.2018.1472551
- Van der Ent, S. (2008). *Transcriptional Regulators of Rhizobacteria Induced Systemic Resistance*. Ph.D. Thesis, Utrecht University Repository, Utrecht.
- Van der Ent, S., Pozo, M. J., Verhagen, B. W. M., Bakker, D., Van Loon, L. C., and Pieterse, C. M. J. (2006). Transcription factors in roots and shoots of *Arabidopsis* involved in rhizobacteria-induced systemic resistance. *IOBC/WPRS Bull.* 29, 157–161.
- Van der Ent, S., Verhagen, B. W. M., Van Doorn, R., Bakker, D., Verlaan, M. G., Pel, M. J. C., et al. (2008). MYB72 is required in early signaling steps of rhizobacteria induced systemic resistance in *Arabidopsis*. *Plant Physiol.* 146, 1293–1304. doi: 10.1104/pp.107.113829
- Van Loon, L. C., Bakker, P. A. H. M., and Pieterse, C. M. J. (1998). Systemic resistance induced by rhizosphere bacteria. *Annu. Rev. Phytopathol.* 36, 453–483. doi: 10.1146/annurev.phyto.36.1.453
- Van Loon, L. C., Geraats, B. P. J., and Linthorst, H. J. M. (2006). Ethylene as a modulator of disease resistance in plants. *Trends Plant Sci.* 11, 184–191. doi: 10.1016/j.tplants.2006.02.005
- Velivelli, S. L. S., Lojan, P., Cranenbrouck, S., Dupré de Boulois, H., Suarez, J. P., Declerck, S., et al. (2015). The induction of Ethylene Response Factor 3 (ERF3) in potato as a result of co-inoculation with *Pseudomonas* sp. R41805 and *Rhizophagus irregularis* MUCL 41833 - a possible role in plant defense. *Plant Signal. Behav.* 10:e988076. doi: 10.4161/15592324.2014.988076
- Verbon, E. H., and Liberman, L. M. (2016). Beneficial microbes affect endogenous mechanisms controlling root development. *Trends Plant Sci.* 21, 218–229. doi: 10.1016/j.tplants.2016.01.013
- Verbon, E. H., Trapet, P. L., Stringlis, I. A., Kruijs, S., Bakker, P. A. H. M., and Pieterse, C. M. J. (2017). Iron and immunity. *Annu. Rev. Phytopathol.* 55, 355–375. doi: 10.1146/annurev-phyto-080516-035537
- Verhagen, B. W. M., Glazebrook, J., Zhu, T., Chang, H.-S., Van Loon, L. C., and Pieterse, C. M. J. (2004). The transcriptome of rhizobacteria-induced systemic resistance in *Arabidopsis*. *Mol. Plant Microbe Interact.* 17, 895–908. doi: 10.1094/MPMI.2004.17.8.895
- Vert, G. A., Briat, J. F., and Curie, C. (2003). Dual regulation of the *Arabidopsis* high-affinity root iron uptake system by local and long-distance signals. *Plant Physiol.* 132, 796–804. doi: 10.1104/pp.102.016089
- Villena, J., Kitazawa, H., Van Wees, S. C. M., Pieterse, C. M. J., and Takahashi, H. (2018). Receptors and signaling pathways for recognition of bacteria in livestock and crops: prospects for beneficial microbes in healthy growth strategies. *Front. Immunol.* 9:2223. doi: 10.3389/fimmu.2018.02223

- Wang, B., Li, Y., and Zhang, W. H. (2012). Brassinosteroids are involved in response of cucumber (*Cucumis sativus*) to iron deficiency. *Ann. Bot.* 110, 681–688. doi: 10.1093/aob/mcs126
- Wang, C., Yao, J., Du, X., Zhang, Y., Sun, Y., Rollins, J. A., et al. (2015). The *Arabidopsis* Mediator Complex Subunit16 is a key component of basal resistance against the necrotrophic fungal pathogen *Sclerotinia sclerotiorum*. *Plant Physiol.* 169, 856–872. doi: 10.1104/pp.15.00351
- Wang, F., Cui, X., Sun, Y., and Dong, C. H. (2013). Ethylene signaling and regulation in plant growth and stress responses. *Plant Cell Rep.* 32, 1099–1109. doi: 10.1007/s00299-013-1421-6
- Wang, N., Cui, Y., Liu, Y., Fan, H., Du, J., Huang, Z., et al. (2013). Requirement and functional redundancy of Ib subgroup bHLH proteins for iron deficiency responses and uptake in *Arabidopsis thaliana*. *Mol. Plant* 6, 503–513. doi: 10.1093/mp/sss089
- Wang, J., Zhou, C., Xiao, X., Xie, Y., Zhu, L., and Ma, Z. (2017). Enhanced iron and selenium uptake in plants by volatile emissions of *Bacillus amyloliquefaciens* (BF06). *Appl. Sci.* 7:85. doi: 10.3390/app7010085
- Wang, W., Shi, J., Xie, Q., Jiang, Y., Yu, N., and Wang, E. (2017). Nutrient exchange and regulation in arbuscular mycorrhizal symbiosis. *Mol. Plant* 10, 1147–1158. doi: 10.1016/j.molp.2017.07.012
- Waters, B. M., Lucena, C., Romera, F. J., Jester, G. G., Wynn, A. N., Rojas, C. L., et al. (2007). Ethylene involvement in the regulation of the H⁺-ATPase *CsHA1* gene and of the new isolated ferric reductase *CsFRO1* and iron transporter *CsIRT1* genes in cucumber plants. *Plant Physiol. Biochem.* 45, 293–301. doi: 10.1016/j.plaphy.2007.03.011
- Yang, J., Kloepper, J. W., and Ryu, C. M. (2008). Rhizosphere bacteria help plants tolerate abiotic stress. *Trends Plant Sci.* 14, 1–4. doi: 10.1016/j.tplants.2008.10.004
- Yang, Y., Ou, B., Zhang, J., Si, W., Gu, H., Qin, G., et al. (2014). The *Arabidopsis* Mediator subunit MED16 regulates iron homeostasis by associating with EIN3/EIL1 through subunit MED25. *Plant J.* 77, 838–851. doi: 10.1111/tpj.12440
- Ye, L., Li, L., Wang, L., Wang, S., Li, S., Du, J., et al. (2015). MPK3/MPK6 are involved in iron deficiency-induced ethylene production in *Arabidopsis*. *Front. Plant Sci.* 6:953. doi: 10.3389/fpls.2015.00953
- Yuan, Y. X., Wu, H. L., Wang, N., Li, J., Zhao, W. N., Du, J., et al. (2008). FIT interacts with AtbHLH038 and AtbHLH039 in regulating iron uptake gene expression for iron homeostasis in *Arabidopsis*. *Cell Res.* 18, 385–397. doi: 10.1038/cr.2008.26
- Yun, B. W., Skelly, M. J., Yin, M., Yu, M., Mun, B. G., Lee, S. U., et al. (2016). Nitric oxide and S-nitrosoglutathione function additively during plant immunity. *New Phytol.* 211, 516–526. doi: 10.1111/nph.13903
- Zamioudis, C. (2012). *Signaling in Arabidopsis Roots in Response to Beneficial Rhizobacteria*. Ph.D. Thesis, Utrecht University, Utrecht.
- Zamioudis, C., Hanson, J., and Pieterse, C. M. J. (2014). β -Glucosidase BGLU42 is a MYB72-dependent key regulator of rhizobacteria-induced systemic resistance and modulates iron deficiency responses in *Arabidopsis* roots. *New Phytol.* 204, 368–379. doi: 10.1111/nph.12980
- Zamioudis, C., Korteland, J., Van Pelt, J. A., van Hamersveld, M., Dombrowski, N., Bai, Y., et al. (2015). Rhizobacterial volatiles and photosynthesis-related signals coordinate MYB72 expression in *Arabidopsis* roots during onset of induced systemic resistance and iron-deficiency responses. *Plant J.* 84, 309–322. doi: 10.1111/tpj.12995
- Zaidler, D., Zähringer, U., Gerber, I., Dubery, I., Hartung, T., Bors, W., et al. (2004). Innate immunity in *Arabidopsis thaliana*: lipopolysaccharides activate nitric oxide synthase (NOS) and induce defense genes. *Proc. Natl. Acad. Sci. U.S.A.* 101, 15811–15816. doi: 10.1073/pnas.0404536101
- Zhang, H., Kim, M. S., Krishnamachari, V., Payton, P., Sun, Y., Grimson, M., et al. (2007). Rhizobacterial volatile emissions regulate auxin homeostasis and cell expansion in *Arabidopsis*. *Planta* 226, 839–851. doi: 10.1007/s00425-007-0530-2
- Zhang, H., Sun, Y., Xie, X., Kim, M. S., Dowd, S. E., and Paré, P. W. (2009). A soil bacterium regulates plant acquisition of iron via deficiency inducible mechanisms. *Plant J.* 58, 568–577. doi: 10.1111/j.1365-313X.2009.03803.x
- Zhang, J., Liu, B., Li, M., Feng, D., Jin, H., Wang, P., et al. (2015). The bHLH transcription factor bHLH104 interacts with IAA-LEUCINE RESISTANT3 and modulates iron homeostasis in *Arabidopsis*. *Plant Cell* 27, 787–805. doi: 10.1105/tpc.114.132704
- Zhang, Y., Wu, H., Wang, N., Fan, H., Chen, C., Cui, Y., et al. (2014). Mediator subunit 16 functions in the regulation of iron uptake gene expression in *Arabidopsis*. *New Phytol.* 203, 770–783. doi: 10.1111/nph.12860
- Zhao, L., Wang, F., Zhang, Y., and Zhang, J. (2014). Involvement of *Trichoderma asperellum* strain T6 in regulating iron acquisition in plants. *J. Basic Microbiol.* 54, S115–S124. doi: 10.1002/jobm.201400148
- Zhou, C., Guo, J., Zhu, L., Xiao, X., Xie, Y., Zhu, J., et al. (2016a). *Paenibacillus polymyxa* BFKC01 enhances plant iron absorption via improved root systems and activated iron acquisition mechanisms. *Plant Physiol. Biochem.* 105, 162–173. doi: 10.1016/j.plaphy.2016.04.025
- Zhou, C., Ma, Z., Xiao, X., Xie, Y., Zhu, J., and Wang, J. (2016b). Potential enhancement of plant iron assimilation by microbial-induced root exudation of phenolic compounds. *Res. Rev. J. Bot. Sci.* 5, 34–37.
- Zhou, C., Zhu, L., Ma, Z., and Wang, J. (2017). *Bacillus amyloliquefaciens* SAY09 increases cadmium resistance in plants by activation of auxin-mediated signaling pathways. *Genes* 8:E173. doi: 10.3390/genes8070173
- Zhou, C., Zhu, L., Ma, Z., and Wang, J. (2018). Improved iron acquisition of *Astragalus sinicus* under low iron-availability conditions by soil-borne bacteria *Burkholderia cepacia*. *J. Plant Interact.* 13, 9–20. doi: 10.1080/17429145.2017.1407000

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2019 Romera, García, Lucena, Martínez-Medina, Aparicio, Ramos, Alcántara, Angulo and Pérez-Vicente. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.