



Corrigendum: Depletion of Arabidopsis ACYL-COA-BINDING PROTEIN3 Affects Fatty Acid Composition in the Phloem

OPEN ACCESS

Edited and reviewed by:

Kent D. Chapman,
University of North Texas,
United States

*Correspondence:

Mee-Len Chye
mlchye@hku.hk

Specialty section:

This article was submitted to
Plant Physiology,
a section of the journal
Frontiers in Plant Science

Received: 23 November 2020

Accepted: 26 January 2021

Published: 18 February 2021

Citation:

Hu T-H, Lung S-C, Ye Z-W and
Chye M-L (2021) Corrigendum:
Depletion of Arabidopsis
ACYL-COA-BINDING PROTEIN3
Affects Fatty Acid Composition in the
Phloem. *Front. Plant Sci.* 12:632503.
doi: 10.3389/fpls.2021.632503

Tai-Hua Hu, Shiu-Cheung Lung, Zi-Wei Ye and Mee-Len Chye*

School of Biological Sciences, The University of Hong Kong, Pokfulam, Hong Kong

Keywords: acyl-CoA esters, fatty acids, linolenic acid, jasmonate, oxylipins, wounding

A Corrigendum on

Depletion of Arabidopsis ACYL-COA-BINDING PROTEIN3 Affects Fatty Acid Composition in the Phloem

by Hu, T.-H., Lung, S.-C., Ye, Z.-W., and Chye, M.-L. (2018) *Front. Plant Sci.* 9:2.
doi: 10.3389/fpls.2018.00002

In the original article, there was a mistake in **Figure 3** as published. The representative images of the unwounded VC at 0 hpw and the unwounded VC at 3 hpw were duplicated. **Figure 3B** has now been updated by replacing the two duplicated images with new representative images previously taken from the same VC batch in the experiment. The corrected **Figure 3** appears below.

The authors apologize for this error and please note that the replaced images do not change the scientific conclusions of the article in any way. The original article has been updated.

Copyright © 2021 Hu, Lung, Ye and Chye. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.




FIGURE 3 | GUS activity assays of transgenic Arabidopsis *AtACBP3pro::GUS* plants following wounding. **(A)** Four-week-old *AtACBP3pro::GUS* transgenic plants were wounded with a pair of forceps and harvested at 1, 2, 3, and 4 h post wounding (hpw) for quantitative GUS activity measurements. Vector (pBI101.3)-transformed plants were analysed at 1 hpw and served as a negative control, while unwounded *AtACBP3pro::GUS* plants were used as a baseline control. *(Continued)*

FIGURE 3 | “***” indicates statistically significant difference ($P < 0.01$, $n = 3$ by Student’s t -test) in comparison to unwounded samples collected at the same time point. Error bars represent standard deviations. **(B)** A representation of histochemical staining of wounded 4-week-old leaves from *AtACBP3pro::GUS* and vector (pBI101.3)-transformed (VC) *Arabidopsis*. These experiments were repeated twice with consistent results. Scale bars = 0.8 cm.