



Erratum: Floral Color Diversity: How Are Signals Shaped by Elevational Gradient on the Tropical–Subtropical Mountainous Island of Taiwan?

OPEN ACCESS

Approved by:

Frontiers Editorial Office, Frontiers Media SA, Switzerland

*Correspondence:

Frontiers Production Office production.office@frontiersin.org

Specialty section:

This article was submitted to Functional Plant Ecology, a section of the journal Frontiers in Plant Science

Received: 01 February 2021 Accepted: 01 February 2021 Published: 18 February 2021

Citation:

Frontiers Production Office (2021)
Erratum: Floral Color Diversity: How
Are Signals Shaped by Elevational
Gradient on the Tropical–Subtropical
Mountainous Island of Taiwan?
Front. Plant Sci. 12:662738.
doi: 10.3389/fpls.2021.662738

Frontiers Production Office*

Frontiers Media SA, Lausanne, Switzerland

Keywords: flowers, bee vision, phylogeny, community, altitude, tropical-subtropical, island

An Erratum on

Floral Color Diversity: How Are Signals Shaped by Elevational Gradient on the Tropical–Subtropical Mountainous Island of Taiwan?

by Tai, K.-C., Shrestha, M., Dyer, A. G., Yang, E.-C., and Wang, C.-N. (2020). Front. Plant Sci. 11:582784. doi: 10.3389/fpls.2020.582784

Due to a production error, there was a mistake in **Table 2** as published. The heading mistakenly contained "B =" which should have been removed. The corrected heading and **Table 2** appear below.

The publisher apologizes for this mistake. The original article has been updated.

1

Copyright © 2021 Frontiers Production Office. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

TABLE 2 | Summary of the phylogenetic structure of low, middle and high-altitude species in Taiwan.

$\mathsf{MPD}_{\mathsf{phylo}}$

	Altitude	N	Actual	Nulls (mean \pm sd)	P
	Low	399	268.1	267.0 ± 3.0	0.639
	Middle	186	267.8	267.1 ± 5.7	0.54
	High	142	260.6	267.0 ± 6.8	0.174
MNTD _{phylo}					
	Low	399	94	93.3 ± 2.1	0.645
	Middle	186	106.4	111.8 ± 4.2	0.098
	High	142	105.3	118.9 ± 5.0	0.004

The P-value is determined by the rank of the actual MPD_{phylo} (or MNTD_{phylo}) in 1000 nulls MPD_{phylo} (or MNTD_{phylo}). Larger P-value indicates stronger phylogenetic over-dispersion, while lower P-value indicates stronger phylogenetic cluster. Phylogenetic distance calculated from the phylogenetic tree was the distance analyzed here.