



Editorial: Lipid Metabolism in Development and Environmental Stress Tolerance for Engineering Agronomic Traits

Zhi-Yan Du^{1*}, Susanne Hoffmann-Benning², Shiwen Wang³, Lina Yin³, Agnieszka Zienkiewicz⁴ and Krzysztof Zienkiewicz⁴

¹ Department of Molecular Biosciences and Bioengineering, University of Hawaii at Manoa, Honolulu, HI, United States, ² Department of Biochemistry and Molecular Biology, Michigan State University, East Lansing, MI, United States, ³ Institute of Soil and Water Conservation, Northwest A&F University, Yangling, China, ⁴ Centre for Modern Interdisciplinary Technologies, Nicolaus Copernicus University in Toruń, Toruń, Poland

Keywords: lipid biosynthesis and turnover, membrane and storage lipids, development, biotic and abiotic stress, genetic engineering

Editorial on the Research Topic

Lipid Metabolism in Development and Environmental Stress Tolerance for Engineering Agronomic Traits

OPEN ACCESS

Edited and reviewed by:

Xue-Rong Zhou,
Agriculture and Food, Commonwealth
Scientific and Industrial Research
Organisation (CSIRO), Australia

*Correspondence:

Zhi-Yan Du
duz@hawaii.edu

Specialty section:

This article was submitted to
Plant Metabolism and Chemodiversity,
a section of the journal
Frontiers in Plant Science

Received: 12 July 2021

Accepted: 26 July 2021


Published: 24 August 2021

Citation:

Du Z-Y, Hoffmann-Benning S,
Wang S, Yin L, Zienkiewicz A and
Zienkiewicz K (2021) Editorial: Lipid
Metabolism in Development and
Environmental Stress Tolerance for
Engineering Agronomic Traits.
Front. Plant Sci. 12:739786.
doi: 10.3389/fpls.2021.739786

Lipids are the essential building blocks of cellular membranes, and they play fundamental roles in numerous biological activities such as photosynthesis, protection, environmental and cellular communication, and storage of carbon and energy. Lipid metabolism is a dynamic and complicated process that includes lipid biosynthesis, transport, accumulation, turnover, and excretion, acting to regulate plant development and tolerance to various environmental stresses. Understanding how lipid metabolism regulates the development and growth in response to adverse conditions raises many interesting questions. Communication between membrane and storage lipids has been observed during the progress, for example, membrane degradation accompanied by lipid droplet (LD) accumulation. Recently, autophagy has been connected with lipid metabolism in plants and algae under stress conditions (Zienkiewicz and Zienkiewicz). Because of the vital role of lipid metabolism in photosynthetic organisms, exploring lipid processes is very important for engineering crops to obtain better agronomic traits including productivity, nutrition, stress tolerance and resilience.

This Research Topic includes 10 original review and research articles, with a special focus on lipid metabolism in crops. The review by Zienkiewicz and Zienkiewicz summarizes the most recent advances on LD degradation, which is a key process for the release of triacylglycerol (TAG). As the major reservoir of cellular carbon and energy, TAG is packed in LD, a dynamic organelle at the center of cellular metabolism. The review also compares the LD degradation in plants and algae (Figure 1A). Looking at lipid metabolism in the context of membrane composition and transport, Salvaing et al. report the function of the Arabidopsis ALA10, a phospholipid flippase of the P₄ type-ATPase family, in reducing the 18:2 desaturation of phosphatidylcholine in the endoplasmic reticulum (ER), while Liao et al. show the roles of two rice acyl-CoA-binding proteins, OsACBP4 (plasma membrane and ER) and OsACBP5 (apoplast), in the intra- and extracellular transport of acyl-CoA esters, respectively (Figure 1B). Other contributions directly address the role of lipid



metabolism during development, *EgMIXTA1*, an R2R3 MYB-type transcription factor, promotes the cuticular wax formation in *Eustoma grandiflorum* (lisianthus) leaves (Wang et al.); together with the specificities of extraplastidial acyltransferase enzymes, the olive *OeFAD2-2* and *OeFAD2-5* regulate the oleic/linoleic acid ratio in the mesocarp of olive cultivars (Figure 1C) (Hernández et al.). The Research Topic also highlights the significance of lipid metabolism in environmental stress response and tolerance (Figure 1D). In *Arabidopsis*, expression of *CsKCS6*, a gene coding the 3-Ketoacyl-CoA synthase (KCS) that catalyzes the biosynthesis of very-long-chain fatty acid (VLCFA) in *Citrus sinensis* (Newhall navel orange), reduces water loss and ion leakage and increases tolerance to drought and salt stress (Guo et al.); *Arabidopsis* phospholipid:diacylglycerol acyltransferase 1 (*AtPDAT1*) promotes the activities of acyl-CoA:lysophosphatidylcholine acyltransferase (LPCAT)

and acylCoA:lysophosphatidylethanolamine acyltransferase (LPEAT) and improves resilience to heat and cold exposure (Demski et al.). In *Vaccinium corymbosum* L., cv. Duke (Duke blueberry), intermittent warming treatment of the fruit regulates its lipid metabolism such as increases in phosphatidylcholine, linoleic acid, and oleic acid, which enhances the low-temperature tolerance in the fruit (Dai et al.). In the roots of maize, integrated lipidomic and transcriptomic analyses reveal the membrane lipid remodeling and gene regulations under saline-alkaline (Xu et al.) or cold stress (Zhao et al.), which are applicable for engineering the crop for better performance under these stress conditions.

The collection of articles in this Research Topic demonstrates the significance of lipid metabolism in various biological processes and the findings will contribute to engineering agronomic traits of crops for the increasing demand for food and bioproducts.

AUTHOR CONTRIBUTIONS

All authors listed have made a substantial, direct and intellectual contribution to the work, and approved it for publication.

ACKNOWLEDGMENTS

The editors would like to thank all reviewers who evaluated manuscripts and contributors for this Research Topic. Research of the Topic Editors was supported by the funds from CTAHR (2303491) and OVCR, UHM (Z-YD), Natural Science Foundation of Shaanxi Province, China (2021JM-604) (LY), NSF (1841251) and USDA-NIMSS (NC-1203) (SH-B).

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2021 Du, Hoffmann-Benning, Wang, Yin, Zienkiewicz and Zienkiewicz. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.