



Corrigendum: Involvement of Arabidopsis Multi-Copper Oxidase-Encoding *LACCASE12* in Root-to-Shoot Iron Partitioning: A Novel Example of Copper-Iron Crosstalk

OPEN ACCESS

Edited and reviewed by:

Francisco Javier Romera,
University of Cordoba, Spain

*Correspondence:

María Bernal
maria.bernal@ruhr-uni-bochum.de
Ute Krämer
ute.kraemer@ruhr-uni-bochum.de

Specialty section:

This article was submitted to
Plant Nutrition,
a section of the journal
Frontiers in Plant Science

Received: 11 November 2021

Accepted: 16 November 2021

Published: 03 December 2021

Citation:

Bernal M and Krämer U (2021)
Corrigendum: Involvement of
Arabidopsis Multi-Copper
Oxidase-Encoding *LACCASE12* in
Root-to-Shoot Iron Partitioning: A
Novel Example of Copper-Iron
Crosstalk.
Front. Plant Sci. 12:813380.
doi: 10.3389/fpls.2021.813380

María Bernal^{1,2*} and Ute Krämer^{1*}

¹ Department of Molecular Genetics and Physiology of Plants, Faculty of Biology and Biotechnology, Ruhr University Bochum, Bochum, Germany, ² Department of Plant Nutrition, Estación Experimental de Aula Dei-CSIC, Zaragoza, Spain

Keywords: copper, iron, multicopper oxidase, homeostasis, deficiency

A Corrigendum on

Involvement of Arabidopsis Multi-Copper Oxidase-Encoding *LACCASE12* in Root-to-Shoot Iron Partitioning: A Novel Example of Copper-Iron Crosstalk

by Bernal, M., and Krämer, U. (2021). *Front. Plant Sci.* 12:688318. doi: 10.3389/fpls.2021.688318

In the original article, there was a mistake in **Figure 2B** as published. The panel B showed the positive control of panel A and it was accidentally taken from a different experiment when the figure was prepared. The corrected **Figure 2** with the correct positive control plate appears below.

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2021 Bernal and Krämer. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.




FIGURE 2 | Heterologous expression of the *AtLAC12* cDNA complements a Fe uptake-defective *fet3fet4* mutant of *Saccharomyces cerevisiae*. Wild-type and the Fe uptake-defective *fet3fet4* mutant of *Saccharomyces cerevisiae* transformed with the empty vector pFL61 Gateway (eV) or expressing *IRT1* or *LAC12* cDNAs of *A. thaliana*, respectively. Aliquots of 10 μL of 10-fold serial dilutions (starting from $\text{OD}_{600} = 0.3$, ca. 10^7 cells ml^{-1}) were spotted on **(A)** SD-URA medium (pH 5.7) and **(B)** SD-URA medium (pH 5.7) supplemented with 0.5 mM FeSO_4 . Images are representative of three independent transformant colonies from each of two independent experiments.