



OPEN ACCESS

EDITED BY

Paromik Bhattacharyya,
Institute of Himalayan Bioresource
Technology (CSIR), India

REVIEWED BY

Ajay Madhusudan Sorty,
Aarhus University, Denmark
Radha Prasanna,
Indian Agricultural Research Institute
(ICAR), India

*CORRESPONDENCE

Rafael Jorge León Morcillo
✉ rafael.morcillo@csic.es
Eduarne Baroja-Fernández
✉ e.baroja@csic.es
Javier Pozueta-Romero
✉ javier.pozueta@csic.es

[†]These authors have contributed
equally to this work

SPECIALTY SECTION

This article was submitted to
Plant Physiology,
a section of the journal
Frontiers in Plant Science

RECEIVED 09 September 2022

ACCEPTED 05 December 2022

PUBLISHED 23 December 2022

CITATION

Morcillo RJL, Baroja-Fernández E,
López-Serrano L, Leal-López J,
Muñoz FJ, Bahaji A, Férrez-Gómez A
and Pozueta-Romero J (2022) Cell-
free microbial culture filtrates as
candidate biostimulants to enhance
plant growth and yield and activate
soil- and plant-associated
beneficial microbiota.
Front. Plant Sci. 13:1040515.
doi: 10.3389/fpls.2022.1040515

COPYRIGHT

© 2022 Morcillo, Baroja-Fernández,
López-Serrano, Leal-López, Muñoz,
Bahaji, Férrez-Gómez and Pozueta-
Romero. This is an open-access article
distributed under the terms of the
[Creative Commons Attribution License
\(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or
reproduction in other forums is
permitted, provided the original
author(s) and the copyright owner(s)
are credited and that the original
publication in this journal is cited, in
accordance with accepted academic
practice. No use, distribution or
reproduction is permitted which does
not comply with these terms.

Cell-free microbial culture filtrates as candidate biostimulants to enhance plant growth and yield and activate soil- and plant-associated beneficial microbiota

Rafael Jorge León Morcillo^{1*†}, Eduarne Baroja-Fernández^{2*†},
Lidia López-Serrano¹, Jesús Leal-López¹,
Francisco José Muñoz², Abdellatif Bahaji²,
Alberto Férrez-Gómez¹ and Javier Pozueta-Romero^{1*}

¹Institute for Mediterranean and Subtropical Horticulture "La Mayora" (IHSM), Consejo Superior de Investigaciones Científicas-Universidad de Málaga, Málaga, Spain, ²Instituto de Agrobiotecnología (IdAB), Consejo Superior de Investigaciones Científicas-Gobierno de Navarra, Nafarroa, Spain

In this work we compiled information on current and emerging microbial-based fertilization practices, especially the use of cell-free microbial culture filtrates (CFs), to promote plant growth, yield and stress tolerance, and their effects on plant-associated beneficial microbiota. In addition, we identified limitations to bring microbial CFs to the market as biostimulants. In nature, plants act as metaorganisms, hosting microorganisms that communicate with the plants by exchanging semiochemicals through the phytosphere. Such symbiotic interactions are of high importance not only for plant yield and quality, but also for functioning of the soil microbiota. One environmentally sustainable practice to increasing crop productivity and/or protecting plants from (a)biotic stresses while reducing the excessive and inappropriate application of agrochemicals is based on the use of inoculants of beneficial microorganisms. However, this technology has a number of limitations, including inconsistencies in the field, specific growth requirements and host compatibility. Beneficial microorganisms release diffusible substances that promote plant growth and enhance yield and stress tolerance. Recently, evidence has been provided that this capacity also extends to phytopathogens. Consistently, soil application of microbial cell-free culture filtrates (CFs) has been found to promote growth and enhance the yield of horticultural crops. Recent studies have shown that the response of plants to soil application of microbial CFs is associated with strong proliferation of the resident beneficial soil microbiota. Therefore, the use of microbial CFs to enhance both crop yield and stress tolerance, and to activate beneficial soil microbiota could be a safe, efficient and environmentally friendly approach to minimize shortfalls related to the technology of microbial inoculation. In this review, we compile information on microbial CFs and the main constituents

(especially volatile compounds) that promote plant growth, yield and stress tolerance, and their effects on plant-associated beneficial microbiota. In addition, we identify challenges and limitations for their use as biostimulants to bring them to the market and we propose remedial actions and give suggestions for future work.

KEYWORDS

plant-microbe interaction, biostimulants, crop yield, stress tolerance, volatile organic compounds, soil- and plant-associated microbiota

Introduction

Plant' growth and development are influenced by microorganisms occurring in the phytosphere that communicate with plants by exchanging chemical signals (Hartmann et al., 2014). Some of these microorganisms can benefit host plants in a variety of ways, a scenario of utmost interest when searching for new and efficient agricultural approaches based on manipulation of plant-associated microbiota. Beneficial microorganisms can directly promote plant growth through mechanisms involving production of bioactive compounds (e.g. phytohormones, volatile compounds, peptides, etc.), dinitrogen fixation, solubilization of minerals and organic material and enhancement of water and nutrient uptake and use (Tsavkelova et al., 2006a; Rodríguez et al., 2007; Francis et al., 2010). These microorganisms can also indirectly promote plant growth by antagonism/antibiosis against pathogens, alleviation of stress caused by environmental pollutants or other stressful abiotic conditions (e.g. drought and salinity), or by triggering in the host plant enhanced defense capacities against pathogen attack.

A decline in natural resources and the environmental damage caused by practices relying on the excessive and inappropriate application of fertilizers and depletion of soil and water resources have become major limitations in conventional agriculture. A more sustainable and eco-friendly agriculture requires increases in product yield quality, while reducing the negative environmental impact of agrochemicals on soil fertility and biodiversity; potential solutions may be fostered by microbial-based approaches (Calvo et al., 2014). The aim of this review was to compile information on current and emerging microbial-based fertilization practices, particularly the use of microbial inoculants, microbial-derived compounds and microbial culture filtrates (CFs), to promote plant growth, yield and stress tolerance, and their effects on plant-associated beneficial microbiota. In addition, we identify challenges and limitations to bring microbial CFs to the market as biostimulants compliant with scientific requirements of the official regulations for fertilizer products.

Soil inoculation of beneficial microorganisms: limitations of a widely used practice to enhance crop yield and/or protect plants from (a)biotic stresses in an eco-friendly manner

One environmentally safe and sustainable practice to promote plant growth, increase crop yield and/or enhance stress tolerance is based on the inoculation of soil with plant growth promoting microorganisms (PGPM) (Miransari, 2011; Calvo et al., 2014; Ahmad et al., 2018; Backer et al., 2018; Fiorentino et al., 2018; Zhong et al., 2019; Noceto et al., 2021). Microbial inoculants consist of one or a reduced number of microbial strains which are grown separately or in mixed culture fermentation, concentrated and then formulated with an appropriate carrier into the final product form. Despite its undisputable success (Li et al., 2022), the technology based on PGPM inoculation has faced a number of limitations and inconsistencies that are summarized in Table 1. First, the efficiency of inoculation depends on soil pH, temperature and nutrient content, interaction with the crop species (host specificity), competition with native strains and compatibility between the microbial inoculants (Svenningsen et al., 2018; Emmanuel and Babalola, 2020). Second, development of multi-strain bioinoculants on a large-scale level is difficult since each co-inoculant requires specific culture media and physical-chemical conditions (Reddy and Saravanan, 2013). Third, inoculation with beneficial microorganisms without a proper carrier or formulation may result in a rapid decline in the applied microbial population, as the inoculated microbes must compete with the often better-adapted native microbial community (Bashan et al., 2014; Cardinale et al., 2015; Parnell et al., 2016). For instance, the abundance in soil of some well-known beneficial microbes such as *Bacillus amyloliquefaciens* FZB42 and *Trichoderma harzianum* falls below detection limits just a few weeks after application (Papavizas, 1982; Kröber et al., 2014; Oskiera et al., 2017). Although seed coating with beneficial

TABLE 1 Limitations of soil inoculation of PGPM and application of microbial-derived compounds and microbial culture filtrates.

	PGPM inoculation	Application of microbial-derived compounds	Application of microbial culture filtrates
Efficiency depends on environmental factors, host compatibility and competition with native microbes	x		
Efficiency depends on microbial culture composition and age	x	x	x
Difficulty to develop multi-strain bioinoculants	x		
Manner of application: necessity for a proper carrier or formulation	x	x	x
Pathogenicity risk to indigenous microbial communities and/or plants	x		
Biostimulant effect depends on plant nutritional status	x		
Scaling-up: difficulty to culture plant-associated microbes in reactors	x	x	x
Long and difficult process to isolate, identify and purify the beneficial compound		x	
Dose-dependence response		x	x
Complex synergic and cooperative interactions between different compounds to promote plant growth		x	
Contradictory effect on different plant species		x	
Antagonistic effects on native microbiota		x	
High production cost		x	x
"X" highlights the limitation of each microbial-derived method.			

microorganisms may be a suitable option to maintain microbial survival in soil over a longer period, an appropriate coating requires a delicate balance between coating materials, microbe and compatible chemistry, which is not always easy to obtain (Glare et al., 2012; Parnell et al., 2016). Moreover, an adequate delivery system for microorganisms in the soil is also required, which represents a major challenge to industry since it involves mass production, formulation, and application of the beneficial microbes (Ravensberg, 2011; Glare et al., 2012; Vassilev et al., 2020). Fourth, some PGPM including some *Pseudomonas* spp. are opportunistic pathogens (Belimov et al., 2007; Sitaraman, 2015) and thus their use is associated with a pathogenicity risk. Fifth, the growth-promoting effect of inoculating beneficial microorganisms strongly depends on the nutritional status of the plant in relation to the extent that they can be rendered ineffective in promoting plant growth (Hoeksema et al., 2010; Pineda et al., 2013). Thus, plants respond better to mycorrhizal inoculation when grown in soils with high microbial diversity and when subjected to phosphorous limitation (Hoeksema et al., 2010). For instance, *Colletotrichum tofieldiae* promotes growth in *Arabidopsis thaliana* only under phosphorus deficiency conditions (Hiruma et al., 2016). Inoculation with the growth-promoting rhizobacterium *B. amyloliquefaciens* GB03 can have deleterious effects on plant growth under phosphate deficiency conditions, due to an

activation of the phytohormone-mediated immune response modulated by a phosphate-starvation response (Morcillo et al., 2020). Sixth, the PGPM inoculation efficiency largely depends upon production of bioactive compounds by the inoculated microbes, which in turn strongly depends on abiotic and biotic environmental contexts. Therefore, it can never be guaranteed that inoculation of a particular microbe will result in the production of compounds with plant growth-promoting or stress tolerance-conferring properties. Seventh, most of soil and plant-associated microorganisms cannot be cultured in reactors. Eighth, inoculation of non-native, allochthonous microorganisms is known to produce strong shifts in microbial communities (Schmidt et al., 2014; Diagne et al., 2018; Berg et al., 2021), with unpredictable and unwanted effects (Hart et al., 2018). For instance, non-native mycorrhizal fungal commercial inoculants may lead to undesirable promotion of exotic over native plant species (Burkle and Belote, 2015; Middleton et al., 2015; Hart et al., 2018).

Some of the constraints of the classic single and multi-strain bioinoculation approach can be circumvented by holistic approaches based on the use of SynComs (for **S**ynthetic **m**icrobial **C**ommunities), which has emerged as a new paradigm not only to better understand plant-microbe interactions, but also to benefit from them

(Castrillo et al., 2017; Durán et al., 2018; Kwak et al., 2018; Carrión et al., 2019; Marin et al., 2021). SynComs are based on the use of metagenomic tools to determine the structure and potential function of plant-associated microbial communities, followed by the isolation and co-culturing of multiple locally adapted native microorganisms. Establishment and survival of inoculated SynComs in the field are higher than that of single or multi-strain bioinoculations, as SynComs are capable of competing with the pre-existing microbiota present in the plant or soil (Liu et al., 2022; Shayanthan et al., 2022). However, despite the obvious theoretical advantages of application of SynComs designed “à la carte” to mimic the role of a particular microbiome, this technology still has some constraints, including technical limitations in the correct metagenomic identification of the isolated microorganisms (Liu et al., 2020), development of SynComs inoculants at a large-scale industrial level, dependence upon the nutritional status of the plant, maintenance of the stability and function of SynComs over time under changing environmental conditions in open field, etc.

Application of microbial-derived compounds: A step to minimize shortfalls related to PGPM inoculation technology

Depending on environmental conditions, microorganisms can release diffusible compounds including phytohormones, siderophores, proteins, peptides, sugar-derived molecules, amino acids, exopolysaccharides, organic acids and volatile compounds that alter metabolism, enhance photosynthesis, promote plant growth, confer resistance to (a)biotic stresses and cause massive lateral root formation, thus improving the root's exploratory capacity for nutrients and predisposing plants for colonization and infection by microbes (Ryu et al., 2003; Arkhipova et al., 2005; Tsavkelova et al., 2006b; Spaepen et al., 2007; Berg, 2009; Contreras-Cornejo et al., 2009; Ortíz-Castro et al., 2009; Chanclud and Morel, 2016; Saha et al., 2016; Sánchez-López et al., 2016; Egamberdieva et al., 2017; García-Gómez et al., 2019; Morcillo and Manzanera, 2021). Some of these compounds are capable of activating soil microbial activity (Rodríguez-Morgado et al., 2017; Macías-Benitez et al., 2020). To address limitations related to PGPM inoculation technology, the application of small quantities of microbial bioactive compounds in pure form, either as alternatives, supplements or complements to microbial cells, has been proposed as a possible approach for improving crop productivity and stress tolerance while reducing agrochemical use (Kanchiswamy et al., 2015; Naamala and Smith, 2021). This approach offers reliability and the easy control of the quantity and quality of a compound of interest (Table 1). Furthermore, compared with the PGPM

inoculation technology, the use of pure microbial compounds can benefit a broader range of crops and minimize pathogenicity risk (Table 1). Although the stability of many of these compounds depends on abiotic environmental factors (temperature, salt concentration in soil, pH, etc.) and biotic factors (they could be used by native microorganisms as nutrient source), their rapid perception by plants can prime them for growth promotion. However, there are quite a number of limitations associated with the use of microbe-derived compounds including time-consuming processes of isolation, identification and purification of bioactive compounds, dose-dependence of the response, complex synergic and cooperative interactions between different compounds to promote plant growth, contradictory effects of the same compound on different plants, antagonistic effects on beneficial microbiota, etc. (Naamala and Smith, 2021) (Table 1).

Application of cell-free microbial culture filtrates: A sustainable and environmentally friendly approach to activate the soil- and plant-associated beneficial microbiota and cope with constraints related to PGPM inoculation and application of microbial-derived compounds?

Some of the limitations related to the use of microbe-derived compounds could be circumvented by the use of cell-free filtrates of beneficial bacterial and fungal cultures, which are mixtures of phytohormones, siderophores, proteins, peptides, amino acids, exopolysaccharides, organic acids, volatile compounds, etc. derived from broth cultures processed through centrifugation or filtration (i.e. micro/ultra/nanofiltration and inverse osmosis) for cell removal (Pellegrini et al., 2020). Agronomic studies have provided evidence that application of these complex cocktails is an efficient approach to promote plant growth and enhance yield and stress tolerance in a wide range of crops while reducing the use of agrochemicals. Notably, recent studies have shown that cell-free CFs of phytopathogens can also be used to enhance yield and stress tolerance (Baroja-Fernández et al., 2021 and unpublished results). Table 2 summarizes details of the studies on effects of microbial CF application. Most of these studies indicated that phytohormones occurring in the microbial CFs (especially indole acetic acid (IAA)) are major determinants of the response of plants to these extracts. However, some studies indicated that microbial amino acids, peptides, extracellular proteins, lipopeptides and siderophores could also play important roles in the response of plants to fungal CFs. For instance, Buensateai et al. (2013) showed that application of *Bacillus* sp. strain CaSUT007 CFs enriched in extracellular proteins increased root and shoot

TABLE 2 Studies on effects of microbial CF application on plant.

Microbial species		Culture medium	Plant species	Application manner	Effect on plant	Mechanism/ mode of action	Reference
<i>Bacterial species</i>							
<i>Azotobacter vinelandii</i>	Beneficial	Specific medium	<i>Solanum lycopersicum</i>	Root irrigation with culture supernatants	Increase shoot dry weight and fruit production	CFs contain auxins, gibberellins and cytokinin-like substances	Azcón and Barea (1975)
<i>Azotobacter beijerinckii</i>							
<i>Streptomyces olivaceoviridis</i>	Beneficial	Starch-casein medium	<i>Triticum aestivum</i>	Pretreatment of wheat grain with CFs	Enhance growth vigor and crop yield	CFs contain auxins, gibberellins and cytokinin-like substances	Aldesuquy et al. (1998)
<i>Streptomyces rimosus</i>							
<i>Streptomyces rochei</i>							
<i>Streptomyces atroolivaceus</i>	Beneficial	MBGM	<i>Triticum aestivum</i>	Pretreatment of wheat grain with CFs	Increases the shoot length, fresh and dry mass, root fresh and dry mass but suppresses the depth of the root system	Effects probably caused by activity of plant growth regulators	El-Shanshoury (1989)
<i>Azospirillum brasilense</i>	Beneficial	NFb	<i>Oryza sativa</i>	CFs applied in hydroponic medium	Enhance root growth and development	CFs contain IAA	El-Khawas and Adachi (1999)
<i>Klebsiella pneumoniae</i>		NFDM					
<i>Bacillus amyloliquefaciens</i> (FZB24, FZB42, FZB45)	Beneficial	GNB	<i>Zea mays</i>	Coleoptiles incubated with CFs	Enhance length growth	CFs contain IAA	Idris et al. (2004)
<i>Bacillus subtilis</i> FZB37							
<i>Bacillus amyloliquefaciens</i> KPS46	Beneficial	GNB	<i>Glycine max</i>	Pretreatment of seeds with CFs	Increases root and shoot length and plant biomass	IAA and extracellular proteins	Buensanteai et al (2008)
<i>Streptomyces coelicolor</i>	Beneficial	GYMA broth	<i>Triticum aestivum</i>	Seed coating with CFs	Improve plant growth under water-stress conditions	CFs contain IAA	Yandigeri et al. (2012)
<i>Streptomyces olivaceus</i>							
<i>Streptomyces geysiriensis</i>							
<i>Methylobacterium</i> spp.	Beneficial	AMS	<i>Triticum aestivum</i>	Pretreatment of seeds with CFs	Enhances seed germination and seedling growth	CFs contain CKs	Meena et al. (2012)
<i>Bacillus</i> sp. CaSUT007	Beneficial	GNB	<i>Manihot esculenta</i>	Pretreatment of stakes with CFs	Increases root and shoot lengths	IAA and extracellular proteins	Bulgarelli et al. (2013)
<i>Bacillus subtilis</i> EA-CB0575	Beneficial	TSB, SBM	<i>Musa</i> spp.	Pretreatment of germinated seeds with CFs	Increases shoot length and dry weight	CFs contain Lipopeptides and siderophores	Posada et al. (2016)
<i>Burkholderia seminalis</i>	Beneficial	Specific medium	<i>Solanum lycopersicum</i>	<i>In vitro</i> application of CFs	Increases seed germination	CFs contain IAA	Tallapragada et al. (2015)

(Continued)

TABLE 2 Continued

Microbial species		Culture medium	Plant species	Application manner	Effect on plant	Mechanism/ mode of action	Reference
<i>Enterococcus faecium</i>	Neutral	LB	<i>Cucumis melo</i>	Soil irrigation with CFs	Increases shoot and root lengths, plant fresh weight, and chlorophyll content	CFs contain IAA and GAs	Lee et al. (2015)
<i>Streptomyces</i> sp.	Beneficial	TYB	<i>Solanum lycopersicum</i>	Soil irrigation with CFs	Enhances plant growth	Effects probably caused by IAA production	Kaur et al. (2019)
<i>Azospirillum brasilense</i> (Ab-V5, Ab-V6)	Beneficial	DYGS	<i>Glycine max</i>	Spraying of leaves or seeds	Increase root nodulation and root development	Indolic compounds	Rondina et al. (2020)
<i>Pectobacterium carotovorum</i>	Pathogenic	LB	<i>Arabidopsis thaliana</i>	Soil irrigation with CFs	Increase shoot and root biomass but not production	Unknwon	Ávila and Poveda (2022)
<i>Pseudomonas syringa</i>							
Fungal species							
<i>Piriformospora indica</i>	Beneficial	Specific medium	<i>Zea mays</i>	Root irrigation with CFs	Enhances shoot but not root growth	Unknown	Varma et al. (1999)
<i>Piriformospora indica</i>	Beneficial	M+ medium	<i>Arabidopsis thaliana</i>	<i>In vitro</i> application of CFs	Induces root branching	Production of a diffusible factor that is probably IAA	Sirrenberg et al. (2007)
<i>Sebacina vermifera</i>	Beneficial	MYP	<i>Panicum virgatum</i>	Pretreatment of seeds with CFs	Enhances seed germination and biomass production	Unknown	Ghimire et al. (2009)
<i>Cladosporium</i> sp. MH-6	Beneficial	Czapek's broth	<i>Cucumis sativus</i>	<i>In vitro</i> application of CFs	Increases shoot and root lengths and plant dry and fresh weight	CFs contain GAs	Hamayun et al. (2010)
<i>Penicillium minioluteum</i> LHL09	Beneficial	Czapek's broth	<i>Oryza sativa</i> (waito-C)/ <i>Glycine max</i>	<i>In vitro</i> application of CFs	Promotes plant growth and nitrogen assimilation, with and without sodium chloride-induced salinity	CFs contain GAs	Khan et al. (2011)
<i>Piriformospora indica</i>	Beneficial	<i>Aspergillus</i> minimal medium	<i>Helianthus annuus L</i>	Root irrigation with CFs	Increases plant growth and seed production. Increases oil content in seeds	Unknown	Bagde et al. (2011)
<i>Shimizuomyces paradoxus</i>	Pathogenic	PDB	<i>Brassica napus</i>	Pretreatment of seeds with CF and foliar spray	Increases seed germination and seedling growth	Unknown	Sung et al. (2011)
<i>Penicillium citrinum</i> KACC43900	Beneficial	Czapek's broth	<i>Carex kobomugi</i>	Injection/foliar spray with CFs	Increases leaf blade length, chlorophyll and carotenoids contents and photosynthesis	Unknown	Hwang et al. (2011)
<i>Fusarium oxysporum</i>	Pathogenic	CYA	<i>Cajanus cajan</i>	Pretreatment of seeds with CF	Increases seed germination and promotes plant growth	Unknown	Jalander and Gachande (2012)
<i>Penicillium</i> sp.	Beneficial	Czapek's broth	<i>Suaeda japonica</i>	<i>In vitro</i> application of CFs	Increases plant length	CFs contain GAs	You (2012)
<i>Trichoderma virens</i>	Beneficial	Richard's solution	<i>Capsicum annuum</i>	Seed coating with CFs	Enhance seed germination and plant vigor	Unknown	Rahman et al. (2012)
<i>Trichoderma pseudokoningii</i>							

(Continued)

TABLE 2 Continued

Microbial species		Culture medium	Plant species	Application manner	Effect on plant	Mechanism/ mode of action	Reference
<i>Trichoderma harzianum</i>							
<i>Penicillium</i> sp. PNF2	Beneficial	PDB	<i>Sesamum indicum</i>	Pretreatment of seeds with CFs	Increases shoot length and fresh weight	CFs contain IAA	Radhakrishnan et al. (2013)
<i>Fusarium oxysporum</i>	Pathogenic	LGN	<i>Solanum lycopersicum</i>	Pretreatment of seeds with CFs	Inhibit seed germination and plant growth	CFs contain toxins	Raithak and Gachande (2013)
<i>Alternaria solani</i>							
<i>Penicillium</i> spp.	Beneficial/ Pathogenic	ME	<i>Triticum aestivum</i>	Pretreatment of seeds with CFs	Increases seed germination and plant biomass	Unknown	Khokhar et al. (2013)
<i>Piriformospora indica</i>	Beneficial	<i>Aspergillus</i> minimal medium	<i>Aristolochia elegans</i>	Soil irrigation with CFs	Increases shoot and root length and fresh and dry weight	Unknown	Bagde et al. (2013)
<i>Trichoderma</i> spp.	Beneficial	ME	<i>Cicer arietinum</i>	Pretreatment of seeds with CFs	Increases seed germination and promote plant growth	Unknown	Ali et al. (2014)
<i>Penicillium</i> spp. (NICS01, DFC01)	Beneficial	PDB	<i>Sesamum indicum</i>	Pretreatment of seeds with CFs	Increase shoot and root length and fresh and dry weight	Amino acids	Radhakrishnan et al., 2014
<i>Pseudomonas</i> sp.	Beneficial	King's B medium	<i>Coriandrum sativum</i>	Pretreatment of seeds with CF and foliar spray	Increases fresh and dry weight and oil production	Unknown	Hegazi et al. (2015)
<i>Alternaria alternata</i>	Pathogenic	Czapek's broth	<i>Triticum aestivum</i>	Pretreatment of seeds with CF and foliar spray	Increase seed germination and plant growth	Unknown	Bhajibhujje (2015)
<i>Alternaria solani</i>							
<i>Penicillium nordicum</i>	Pathogenic	CYA	<i>Sorghum bicolor</i>	Pretreatment of seeds with CFs	Inhibit seed germination	CFs contain mycotoxin	Vankudoth et al. (2015)
<i>Penicillium citrinum</i> ,							
<i>Penicillium chrysogenum</i>							
<i>Penicillium commune</i>							
<i>Penicillium verrucosum</i>							
<i>Penicillium camemberti</i>							
<i>Penicillium digitatum</i>							
<i>Penicillium oxalicum</i>	Beneficial	PDB	<i>Pennisetum glaucum</i>	Pretreatment of seeds with CFs	Increases seed germination and seedling vigor	Unknown	Murali and Amruthesh (2015)
<i>Fusarium tricinctum</i> RSF-4L	Pathogenic	Czapek's broth	<i>Oryza sativa</i> L. cv. Dongjin	<i>In vitro</i> application of CFs	Increase shoot and root lengths, plant fresh weight, and chlorophyll content	CFs contain IAA	Khan et al. (2015)
<i>Alternaria alternata</i> RSF-6L							

(Continued)

TABLE 2 Continued

Microbial species		Culture medium	Plant species	Application manner	Effect on plant	Mechanism/ mode of action	Reference
<i>Purpureocillium lilacinum</i>	Beneficial	Specific medium	<i>Solanum lycopersicum</i>	Pretreatment of seeds with CFs/ Soil irrigation with CFs	Increases seed germination and promotes plant growth	CFs contain IAA	Cavello et al. (2015)
<i>Fusarium oxysporum</i>	Pathogenic	Czapek's broth	<i>Triticum aestivum</i> <i>Hordeum vulgare</i> <i>Solanum tuberosum</i>	Pretreatment of seeds with CFs	Inhibit seed germination and plant growth	Unknown	Ogórek (2016)
<i>Fusarium sulphureum</i>							
<i>Gibberella avenacea</i>							
<i>Gibberella intricans</i>							
<i>Trichoderma harzianum</i> WKY1	Beneficial	Czapek's broth	<i>Shorgum</i>	Soil irrigation with CFs	Increase shoot and root lengths, plant fresh weight, and total phenol content	CFs contain IAA	Saber et al. (2017)
<i>Aspergillus fumigatus</i> TS1	Beneficial	Czapek's broth	<i>Oryza sativa</i> (waito-C)	Application to the apical meristem	Enhance chlorophyll content, root-shoot length, and biomass production	CFs contain IAA and GAs	Bilal et al. (2018)
<i>Fusarium proliferatum</i> BRL1							
<i>Fusarium solani</i>	Pathogenic	Richard's solution	<i>Solanum lycopersicum</i> <i>Brassica rapa</i> <i>Raphanus sativus</i> <i>Trigonella melongena</i>	Pretreatment of seeds with CFs	Enhances seed germination	Unknown	Parveen et al. (2019)
<i>Trichothecium roseum</i>					Inhibit seed germination	CFs contain mycotoxins	
<i>Aspergillus niger</i>							
<i>Cladoporium herbarum</i>							
<i>Alternaria alternata</i>							
<i>Penicillium chrysogenum</i>							
<i>Penicillium expansum</i>							
<i>Trichoderma</i> spp.	Beneficial				Enhance seed germination	Unknown	
<i>Trichoderma asperellum</i>							
<i>Trichoderma harzianum</i>							
<i>Piriformospora indica</i>	Beneficial	CM	<i>Cichorium intybus</i>	Foliar spray	Enhances growth and morpho-physiological traits	Unknown	Rashnoo et al. (2020)
<i>Trichoderma harzianum</i> ,	Beneficial	MS	<i>Capsicum annuum</i>	Soil irrigation with CFs and DEs	Stimulate root growth and enhance fruit yield	CFs produce changes in plant-associated microbiota	Baroja-Fernández et al. (2021)
<i>Alternaria alternata</i>	Pathogenic						
<i>Penicillium aurantiogriseum</i>							

(Continued)

TABLE 2 Continued

Microbial species		Culture medium	Plant species	Application manner	Effect on plant	Mechanism/ mode of action	Reference
<i>Gibberella intermedia</i>	Beneficial	Czapek's broth	<i>Oryza sativa</i> (waito-C)	Application to the apical meristem	Increases shoot growth	CFs contain GAs	Khalmuratova et al. (2021)
<i>Fusarium oxysporum</i>	Pathogenic	PDB	<i>Arabidopsis thaliana</i>	Soil irrigation with CFs	Increase shoot and root biomass but not production	Unkwnon	Ávila and Poveda (2022)
<i>Pythium irregulare</i>							
<i>Rhizoctonia solani</i>							
<i>Chaetomium globosum</i>	Beneficial	ME	<i>Cichorium intybus</i>	Soil irrigation with CFs	Increase of biomass, shoots and roots length, and leaf area	Increases phenylalanine pathway and chicoric acid	Spinelli et al. (2022)
<i>Minimedusa polyspora</i>						Increases phenylalanine pathway and 4-OH-benzoate	
<i>Geotrichum candidum</i>	Neutral	PDB	<i>Vigna radiata</i>	Pretreatment of seeds with CFs	Increases seed germination and promotes plant growth	CFs contain IAA	George et al. (2019)
<i>Saccharomyces cerevisiae</i>	Beneficial	Sucrose	<i>Coreandrum sativum</i>	Pretreatment of seeds with CF and foliar spray	Increases fresh and dry weight and oil production	Unknown	Hegazi et al. (2015)

IAA, Indole-3-acetic acid; GAs, Gibberellins; AMS, Ammonium mineral salt medium supplemented with methanol; CM, Complex medium; CYA, Czapek yeast medium; GNB, Glucose-enriched nutrient broth; GYMA broth, Glucose, yeast extract, malt extract; LB, Luria-Bertani broth; LGN, Liquid glucose nitrate medium; MBGM, Modified bouillon glycerol medium; ME, Malt extract broth; MS, Murashige and Skoog medium; MYP broth, Malta Yeast Peptone broth; PDB, Potato Dextrose Broth; NFB, Nitrogen-free malate; NFD, Nitrogen-free dextrose; SBM, Sporulation Bacillus Medium; TSB, Trypticase Soy Broth; TYB, Tryptone-Yeast Extract Broth. Specific medium, see publication for details.

lengths and total biomass of cassava stalks. Furthermore, Posada et al. (2016) showed that application of *B. subtilis* EA-CB0575 CFs enhanced dry weight of banana plants by the action of lipopeptides and siderophores. Moreover, Buensanteai et al. (2008) showed that extracts of *B. amyloliquefaciens* strain KPS46 promoted soybean growth through the actions of the antibiotic surfactin and proteins secreted by the bacterium, including auxin biosynthetic enzymes, proteins related to phosphate solubilization and nitrogen metabolism, antifungal lipopeptides and proteins related to protection against oxidative stress. Also, CFs enriched in amino acids secreted by several *Penicillium* spp. enhanced shoot and root length as well as the biomass of sesame plants under well-irrigated and drought conditions (Radhakrishnan et al., 2014).

Baroja-Fernández et al. (2021) have recently shown that soil application of CFs of beneficial and phytopathogenic fungi cultured in Murashige & Skoog (MS) medium promoted root growth, enhanced fruit yield and altered composition of fruits of pepper plants. In the same study, the authors found that CFs of the different fungal species possessed volatile organic compounds

(VOCs) that, once distilled and applied to soil, promoted responses similar to those triggered by direct application of the fungal CFs. These findings indicated that (i) CFs of both beneficial and phytopathogenic fungi can be used to improve crop yield and (ii) VOCs mediate the crops' responses to fungal CF application. Some bioactive VOCs present in the fungal CFs are shown in Table 3. As further discussed below, it is conceivable that some these compounds (particularly acetic acid) are involved in the crop response to soil application of CFs. Notably, high-throughput sequencing analyses revealed that soil application of fungal CFs and distillates (DEs) promoted similar changes in the soil microbiota, and promoted the proliferation of the same beneficial microbial taxa (Baroja-Fernández et al., 2021) (Table 4). Collectively, the findings of Baroja-Fernández et al. (2021) indicated that (i) CFs of both beneficial and fungal phytopathogens can be used to activate the soil and plant-associated beneficial microbiota, and (ii) microbial VOCs mediate the plants' responses to soil application of fungal CFs through mechanisms involving stimulation of the beneficial soil microbiota as schematically illustrated in Figure 1.

TABLE 3 List of potentially relevant bioactive VOCs present in at least one of the fungal CFs used by Baroja-Fernández et al. (2021) and their effects on plant.

VOCs	Effect on plant	Reference
1-butanol-3-methyl	Enhances fresh shoot and root biomass and chlorophyll content in Arabidopsis and increases both root length and thickness in <i>Agave salmiana</i> .	Camarena-Pozos et al. (2019)
	Increases size, fresh weight and total chlorophyll content in Arabidopsis	Lee et al. (2019a)
1-butanol-2-methyl	Increases size, fresh weight and total chlorophyll content in Arabidopsis	Lee et al. (2019a)
1-hexanol	Promotes growth of Arabidopsis	Blom et al. (2011)
benzaldehyde	Increases the volatile oil accumulation in <i>Atractylodes lancea</i>	Zhou et al. (2016)
2-phenylethyl alcohol	Enhances fresh shoot and root biomass, chlorophyll content, in Arabidopsis and increases both root length and thickness in <i>A. salmiana</i>	Camarena-Pozos et al. (2019)
acetic acid	Enhances drought tolerance in Arabidopsis, rapeseed, maize, rice and wheat plants	Kim et al. (2017)
	Increases root biomass and photosynthesis in mung bean	Rahman et al. (2019)
acetoin	Increases root length, shoot length and dry weight in <i>L. sativa</i> seedlings	Fincheira et al. (2017)
	Induces stomatal closure in Arabidopsis and <i>Nicotiana benthamiana</i>	Wu et al. (2018)
2,3-butanediol	Promotes growth of Arabidopsis	Ryu et al. (2003)
	Induces stomatal closure in Arabidopsis and <i>N. benthamiana</i>	Wu et al. (2018)
	Increases root length, shoot length and dry weight in <i>L. sativa</i> seedlings	Fincheira et al. (2017)
2-heptanone	Promotes the growth of Arabidopsis seedlings	Jiang et al. (2019)
2-nonanone	Increases root length, shoot length and dry weight in <i>L. sativa</i> seedlings	Fincheira et al. (2017)
acetophenone	Elicits the increase of biomass in Arabidopsis	Camarena-Pozos et al. (2019)
cis-thujopsene	Induces lateral root formation of Arabidopsis seedlings and poplar plantlets	Ditengou et al. (2015)

Challenges and limitations of the microbial CF technology

Despite having great potential as a strategy for improving productivity in a sustainable and eco-friendly manner, the technology based on microbial CF application is still at its infancy and faces important challenges and limitations before it can be widely used (Table 1). First, one challenging aspect of the microbial CF-based technology is the manner of application of the extracts, especially in cases in which bioactivity of CFs is based on compounds with high evaporation rates such as VOCs. Most studies on the effect of application of microbial CFs on plants are based on seed coating and soil applications of CFs (Table 2), but other means of delivery, should be explored to develop appropriate and durable methods that can be used in the field. Second, another challenging aspect of the microbial CF-based technology is the scaling up from lab-scale shake flasks to stirred tanks-based pilot-scale production (Figure 2). This also applies to the technologies of soil inoculation of beneficial microorganisms and application of microbial-derived compounds. Overcoming this limitation is not an easy task due to marked differences in hydromechanical properties and nutrients/oxygen gradients between flasks and large


fermentors. For successful scaling up, key parameters affecting heat, momentum and mass transfer should be considered (Trujillo-Roldán et al., 2013). Moreover, some physical parameters should be combined to obtain dimensionless numbers intended to be kept constant during the scaling up process. Third, there are no studies on techno-economic viability of large-scale production of microbial CFs involving (1) propagation of the microbial strain until desired inoculum concentration is reached, (2) fermentation of the microbial strains in large, industrial size fermentors until desired cell concentration is reached and (3) microbial cell removal. For the first two steps, some techno-economic models have been developed for typical liquid biofertilizer production plants (Pérez-Sánchez et al., 2018). These models can be used to investigate the main factors that affect the production process, in order to optimize plant productivity and reliability, and also to reduce costs. Fourth, the limitations to microbial CF use are closely related to the downstream processes for production. However, there are no studies on the formulation and shelf life of microbial CFs and on allocation of fitness costs for resources for the large-scale synthesis of these extracts. Clearly, the formulation of new products ready to be commercialized requires further scientific and industrial up-scaling studies. Fifth, the effects

TABLE 4 List of microbial species whose populations are enriched by the soil application of fungal CFs and DEs used by Baroja-Fernández et al. (2021).

Microbial species	Mechanism/mode of action	Reference
Bacterial species		
<i>Burkholderia arboris</i>	Solubilizes phosphate and produces IAA and siderophores	Zhang et al. (2022)
<i>Burkholderia silvatlantica</i>	Has ACC deaminase activity Fixes N ₂	Onofre-Lemus et al. (2009)
		Perin et al. (2006)
<i>Caballeronia udeis</i>	Solubilizes phosphate and produces siderophores	Puri et al. (2020)
<i>Duganella ginsengisoli</i>	Produces IAA	Goodwin (2022)
<i>Pseudomonas brassicacearum</i>	Has ACC deaminase activity	Belimov et al. (2007)
<i>Pseudomonas mediterranea</i>	Solubilizes organic phosphate and produces siderophores, proteases, ammonia and IAA	Gu et al. (2020)
<i>Pseudomonas Knackmussii</i>	Solubilizes phosphate and produces IAA and siderophores	Rabhi et al. (2018)
<i>Rhodanobacter glycinis</i>	Synthesizes osmolytes and biocontrol-related substances	Lee et al. (2019b)
Fungal species		
<i>Candida subhashii</i>	Biological control of plant pathogenic fungi	Hilber-Bodmer et al. (2017)
<i>Geotrichum candidum</i>	Produces phytohormones and reactive oxygen species	Waqas et al. (2017)
	Solubilizes phosphate	Wu et al. (2012)
	Produces IAA, ammonia and polyamines	Fu et al. (2016)
	Produces IAA and siderophores and has ACC deaminase activity	George et al. (2019)
<i>Pseudogymnoascus spp.</i>	Solubilizes phosphate	Abdel-Ghany et al. (2019)

of CFs on plants and associated microbiota may vary depending on the plant and microbial species and ecotypes as well as on media composition, age and growth conditions of the microbial culture. Thus, whereas Khokhar et al. (2013) reported that application of CFs of several *Penicillium* spp. cultured for 15 days at 20 °C in malt extract broth exerted a positive effect on wheat germination and growth, Vankudoth et al. (2015) reported that application of CFs of the same fungal species cultured for 12 days at 27 °C in CYA broth exerted a negative effect on sorghum germination. CFs of the phytopathogen *Fusarium oxysporum* grown in Czapek-Dox and potato dextrose liquid media exerted a negative effect on seed germination and growth of cucumber and garden cress (*Lepidium sativum* L.) plants (Melo and Piccinin, 1999; Ogórek, 2016), whereas application of CFs of *F. oxysporum* cultured for 5 days at 25 °C in CYA broth enhanced pigeonpea (*Cajanus cajan* L.) seed germination and growth (Jalander and Gachande, 2012). Application of CFs of the fungal phytopathogen *A. alternata* cultured in Czapek broth and MS media promoted growth of rice and wheat plants and enhanced pepper fruit yield (Bhajibhuje, 2015; Khan et al., 2015; Baroja-Fernández et al., 2021), whereas application of CFs of the same species cultured in Richard's solution exerted a negative effect on germination of seeds of

several crop plants (Parveen et al., 2019). In many instances, the growth inhibitory effect of the CFs of phytopathogens was due to toxins released by the microorganism to the culture medium (Raithak and Gachande, 2013; Vankudoth et al., 2015; Parveen et al., 2019). Sixth, above threshold levels, many microbial compounds are toxic to plants. Therefore, excess application of microbial CFs has the potential to exert a negative effect on plants. However, after due assessment of the dose-response effect on specific crops, microbial CFs can be safely managed. Seventh, although Baroja-Fernández et al. (2021) showed that application of CFs of diverse microorganisms resulted in activation of beneficial soil and plant-associated microbiota without significant changes in the relative abundance of populations of pathogenic microbial species, it is important to ensure that these results can be extrapolated to other CFs in different soil types and environmental scenarios. Eighth, CFs based on co-cultivation of various microorganisms might be an efficient approach to obtain widely range of bioactive compounds. Nevertheless, this practice faces similar problems to those of multi-microbial bioinoculants, since each co-inoculant requires specific culture conditions (Reddy and Saravanan, 2013). Ninth, there are few studies on the mechanisms and modes of action of microbial CFs on plants.




The necessity of identifying the mechanisms and modes of action of cell-free microbial CFs for their use as biostimulants

In our opinion, the exploitation of microbial CFs as biostimulants (defined by the European Biostimulant Industry Council (<https://biostimulants.eu/>) as “substances and/or microorganisms whose function when applied to plants or to soil is to stimulate natural processes to enhance or benefit nutrient uptake, nutrient efficiency, tolerance to abiotic stress and crop quality” (du Jardin, 2015)) is only just evolving and its broad potential is now beginning to be demonstrated. The majority of studies describing the positive effect(s) of the application of microbial CFs on plant growth and yield did not identify modes or mechanisms of action of these extracts (Table 1). At most, some of these studies proposed that phytohormones and other growth

promoting compounds released by microbes in their culture media are involved in the response of plants to microbial CFs. This may result in the assumption by many that these complex, multicomponent mixtures are “magic potions” or “snake oil” not compliant with scientific requirements of the official regulations for fertilizer products (Yakhin et al., 2017). The identification of mechanisms and modes of action of microbial CFs on plants and the characterization of the biological functions and ecological roles of their bioactive components, based on reasonable scientific hypotheses rather than a try-it-and-see approach, could be important not only to develop novel products able to increase yields in crops in a sustainable and environmentally benign manner, but also to obtain clues for the biotechnological design of more productive and efficient crops.

Baroja-Fernández et al. (2021) showed that VOCs are important mediators in the response of plants and plant-associated microbiota to soil application of microbial CFs. However, the bioactive VOCs produced by fungal cultures and



their mechanisms and modes of action remain unverified. Furthermore, whether the effect exerted by soil application of VOCs on plants is direct or mediated by changes induced in the composition of the soil microbiota (or both) still needs to be determined, as does whether the effect exerted by soil application of VOCs on the plant-associated microbiota is direct or mediated by changes induced in the root metabolism (Figure 1). That VOCs isolated from CFs of diverse beneficial and phytopathogenic microorganisms promoted similar growth and fruit yield and quality responses in crops indicate that plants respond similarly to a wide range of bioactive microbial VOCs. Alternatively, it is likely that many microorganisms produce the same growth promoting VOCs. One of such compounds could be acetic acid, which is present in soils, where microorganisms produce it in response to biotic and abiotic stresses (Adeleke et al., 2017). Recent studies have shown that soil application of acetic acid, but not other organic acids, enhances drought tolerance in *Arabidopsis* and important crops including maize, wheat, rice and cassava (Kim et al., 2017; Utsumi et al., 2019) and tolerance to bacterial wilt disease (Wang et al., 2021). In *Arabidopsis*, the enhancement of drought resistance promoted by soil application of acetic acid involves JA signaling and an ON/OFF switching epigenetic mechanism dependent on histone deacetylase HDA6 (Kim et al., 2017). However, the biochemical and molecular mechanisms underlying the enhancement of drought tolerance promoted by soil application of acetic acid in crops remain to be elucidated.

Several lines of evidence indicate that the manner of application of microbial volatile compounds and/or plant growth conditions are important determinants of the biochemical and molecular mechanisms triggered in plants. Application of these compounds *via* the air promotes plant

growth and metabolic and developmental changes, enhances photosynthesis and improves nutrient and water acquisition (Ryu et al., 2003; Zhang et al., 2008; Zhang et al., 2009; Ezquer et al., 2010; Gutiérrez-Luna et al., 2010; Garnica-Vergara et al., 2016; Sánchez-López et al., 2016; García-Gómez et al., 2019; García-Gómez et al., 2020). In *Arabidopsis*, these responses are associated with changes in the transcriptome, proteome, metabolome, hormone and redox-proteome through mechanisms involving long-distance communication between roots and the aerial part of the plant and proteostatic regulation of central metabolic pathways in the plastidial compartment (Zhang et al., 2007; Zhang et al., 2008; Zhang et al., 2009; Sánchez-López et al., 2016; Amezttoy et al., 2019; García-Gómez et al., 2020; Amezttoy et al., 2021; Gámez-Arcas et al., 2022). However, Baroja-Fernández et al. (2021) showed that no such mechanisms operate in crops irrigated with VOC-containing microbial CFs and DEs. Needless to say, further work will be necessary to understand how plants respond to microbial volatiles.

The discovery that soil application of VOCs from diverse microorganisms can enhance crop yield and promote similar changes in the soil microbiota extends knowledge on the mechanisms modulating the physiology of the plant and their interaction with plant-associated microbes, raising questions on their ecological significance and biotechnological applications. Because microbes respond to microbial VOCs, it is likely that the fungal VOC-promoted enrichment of the soil beneficial microbiota is due to direct action of these compounds on the microorganisms, which in turn determine plant growth and metabolism through emission of semiochemicals. These compounds trigger the production of molecules by roots that, once transported to the

aerial part of the plant, act as long-distance signals that promote growth and enhance yield (Figure 1A). In line with this presumption, Yuan et al. (2017) showed that exposure of soil to VOCs of *B. amyloliquefaciens* NJN-6 altered the composition of soil microbial communities. Compounds secreted by roots in their exudates attract beneficial microorganisms which, in turn, strongly influence plant fitness and enhance the capacity of plants to adapt to environmental changes and stress (Baudoin and Benizri, 2003; Badri and Vivanco, 2009; Badri et al., 2013; Bulgarelli et al., 2013; Schulz-Bohm et al., 2017; Sasse et al., 2018; Zhalnina et al., 2018; Huang et al., 2019; Chen et al., 2020; Vilchez et al., 2020; He et al., 2022). Therefore, it is conceivable that enrichment in the plant-associated beneficial microbiota triggered by application of fungal CFs and DEs is due to an alteration of composition of root exudates (Figure 1B). This capacity of root exudates to influence the soil microbiota is not static since the composition of root exudates is not uniform over time (Zhalnina et al., 2018) and depends on the plant species or genotype (Bulgarelli et al., 2012; Bodenhausen et al., 2014; Bouffaud et al., 2014; Zhong et al., 2019). For instance, it has been shown that soybean genotype significantly influences the structure and composition of its associated rhizosphere and affects connections in rhizobacterial networks (Zhong et al., 2019).

Additional remarks

According to the Regulation (EU) 2019/1009 of the European Parliament and of the Council of 5 June 2019 laying down rules on the making available on the market of EU fertilising products, a microbial plant biostimulant may contain micro-organisms belonging to the *Azotobacter*, *Rhizobium*, *Azospirillum* genera and mycorrhizal fungi, including dead or empty-cell micro-organisms and non-harmful residual elements of the media on which the microorganisms were cultured (<https://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:32019R1009>). Although regulations are of great importance for guaranteeing food security and preserving environmental integrity, the stringency and exclusivity of the list of microorganisms that can be used for the production of biostimulants may strongly limit the potential benefits of these products. As shown in the review, growing

References

- Abdel-Ghany, T. M., Mohamed, Z. H., Al Abboud, M. A., Helmy, E. A., Al-Rajhi, A. M. H., and Shater, A. R. M. (2019). Solubilization of inorganic phosphate by rhizospheric fungi isolated from soil cultivated with *Sorghum bicolor* L. *BioRes.* 14 (3), 5521–5532. doi: 10.15376/biores.14.3.5521-5532
- Adeleke, R., Nwangburuka, C., and Oboirien, B. (2017). Origins, roles and fate of organic acids in soils: A review. *S. Afr. J. Bot.* 108, 393–406. doi: 10.1016/J.SAJB.2016.09.002
- Ahmad, M., Pataczek, L., Hilger, T. H., Zahir, Z. A., Hussain, A., Rasche, F., et al. (2018). Perspectives of microbial inoculation for sustainable development and environmental management. *Front. Microbiol.* 9. doi: 10.3389/FMICB.2018.02992

evidence has been compiled demonstrating that application of cell-free CFs of beneficial and phytopathogenic microorganisms is an efficient approach to promote plant growth and enhance yield and stress tolerance in a wide range of crops while reducing the use of agrochemicals. Therefore, it may be appropriate to enlarge the list of microorganisms that can be used for the production of cell-free CF-based biostimulants according to EU regulation on fertilising products, assuming scientific evidence can demonstrate that these products are safe for both the environment and consumers.

Authors contributions

RL, EB-F and J-PR conceptualized the topic and wrote the manuscript. All authors contributed in the literature search and discussions, and reviewed and approved the final manuscript.

Funding

This work was supported by the Agencia Estatal de Investigación (AEI) and Fondo Europeo de Desarrollo Regional (Spain) (grant PID2019-104685GB-I00).

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

- Aldequy, H. S., Mansour, F. A., and Abo-Hamed, S. A. (1998). Effect of the culture filtrates of *Streptomyces* on growth and productivity of wheat plants. *Folia Microbiol.* 43 (5), 465–470. doi: 10.1007/BF02820792

- Ali, A., Haider, M. S., Ashfaq, M., and Hanif, S. (2014). Effect of culture filtrates of *Trichoderma* spp. on seed germination and seedling growth in chickpea—an *in vitro* study. *Pakistan J. Phytopathol.* 26 (1), 01–05.

- Ameztoy, K., Baslam, M., Sánchez-López, Á.M., Muñoz, F. J., Bahaji, A., Almagro, G., et al. (2019). Plant responses to fungal volatiles involve global posttranslational thiol redox proteome changes that affect photosynthesis. *Plant Cell Environ.* 42 (9), 2627–2644. doi: 10.1111/pce.13601

- Ameztoy, K., Sánchez-López, Á.M., Muñoz, F. J., Bahaji, A., Almagro, G., Baroja-Fernández, E., et al. (2021). Proteostatic regulation of MEP and shikimate pathways by redox-activated photosynthesis signaling in plants exposed to small fungal volatiles. *Front. Plant Sci.* 12. doi: 10.3389/fpls.2021.637976
- Arkhipova, T. N., Veselov, S. U., Melentiev, A. I., Martynenko, E. V., and Kudoyarova, G. R. (2005). Ability of bacterium *Bacillus subtilis* to produce cytokinins and to influence the growth and endogenous hormone content of lettuce plants. *Plant Soil*. 272 (1), 201–209. doi: 10.1007/s11104-004-5047-x
- Ávila, A. C., and Poveda, J. (2022). Induction of immune response in *Arabidopsis thaliana* treated with phytopathogen filtrates. *Biol. Life Sci. Forum*. 11, 85. doi: 10.3390/IECPS2021-11974
- Azcón, R., and Barea, J. M. (1975). Synthesis of auxins, gibberellins and cytokinins by *Azotobacter vinelandii* and *Azotobacter beijerinckii* related to effects produced on tomato plants. *Plant Soil*. 43 (1), 609–619. doi: 10.1007/BF01928522
- Backer, R., Rokem, J. S., Ilangumaran, G., Lamont, J., Praslickova, D., Ricci, E., et al. (2018). Plant growth-promoting rhizobacteria: Context, mechanisms of action, and roadmap to commercialization of biostimulants for sustainable agriculture. *Front. Plant Sci.* 871. doi: 10.3389/fpls.2018.01473
- Badri, D. V., Chaparro, J. M., Zhang, R., Shen, Q., and Vivanco, J. M. (2013). Application of natural blends of phytochemicals derived from the root exudates of arabis to the soil reveal that phenolic-related compounds predominantly modulate the soil microbiome. *J. Biol. Chem.* 288 (7), 4502–4512. doi: 10.1074/JBC.M112.433300
- Badri, D. V., and Vivanco, J. M. (2009). Regulation and function of root exudates. *Plant Cell Environ.* 32 (6), 666–681. doi: 10.1111/j.1365-3040.2009.01926.x
- Bagde, U., Prasad, R., and Varma, A. (2011). Influence of culture filtrate of *Piriformospora indica* on growth and yield of seed oil in *Helianthus annuus*. *Symbiosis*. 53, 83–88. doi: 10.1007/s13199-011-0114-6
- Bagde, U. S., Prasad, R., and Varma, A. (2013). Impact of culture filtrate of *Piriformospora indica* on biomass and biosynthesis of active ingredient aristolochic acid in *Aristolochia elegans*. *Mart. Int. J. Biol.* 6, 29. doi: 10.5539/ijb.v6n1p29
- Baroja-Fernández, E., Almagro, G., Sánchez-López, Á.M., Bahaji, A., Gámez-Arcas, S., De Diego, N., et al. (2021). Enhanced yield of pepper plants promoted by soil application of volatiles from cell-free fungal culture filtrates is associated with activation of the beneficial soil microbiota. *Front. Plant Sci.* 12. doi: 10.3389/fpls.2021.752653
- Bashan, Y., de-Bashan, L. E., Prabhu, S. R., and Hernandez, J.-P. (2014). Advances in plant growth-promoting bacterial inoculant technology: formulations and practical perspective–2013). *Plant Soil*. 378 (1), 1–33. doi: 10.1007/s11104-013-1956-x
- Baudoin, E., and Benizri, E. (2003). Impact of artificial root exudates on the bacterial community structure in bulk soil and maize rhizosphere. *Soil Biol. Biochem.* 35, 1183–1192. doi: 10.1016/S0038-0717(03)00179-2
- Belimov, A. A., Dodd, I. C., Safronova, V. I., Hontzas, N., and Davies, W. J. (2007). *Pseudomonas brassicacearum* strain Am3 containing 1-aminocyclopropane-1-carboxylate deaminase can show both pathogenic and growth-promoting properties in its interaction with tomato. *J. Exp. Bot.* 58 (6), 1485–1495. doi: 10.1093/jxb/erm010
- Berg, G. (2009). Plant-microbe interactions promoting plant growth and health: perspectives for controlled use of microorganisms in agriculture. *Appl. Microbiol. Biotechnol.* 84 (1), 11–18. doi: 10.1007/s00253-009-2092-7
- Berg, G., Kusstatscher, P., Abdelfattah, A., Cernava, T., and Smalla, K. (2021). Microbiome modulation—toward a better understanding of plant microbiome response to microbial inoculants. *Front. Microbiol.* 12. doi: 10.3389/fmicb.2021.650610
- Bhajibhujje, M. N. (2015). Response of metabolites from culture filtrates of *Alternaria* species against *Triticum aestivum* l. *Int. J. Life Sci.* 3, 55–62.
- Bilal, L., Asaf, S., Hamayun, M., Gul, H., Iqbal, A., Ullah, I., et al. (2018). Plant growth promoting endophytic fungi *Aspergillus fumigatus* T51 and *Fusarium proliferatum* BRL1 produce gibberellins and regulates plant endogenous hormones. *Symbiosis*. 76, 117–127. doi: 10.1007/s13199-018-0545-4
- Blom, D., Fabbri, C., Connor, E. C., Schiestl, F. P., Klausner, D. R., Boller, T., et al. (2011). Production of plant growth modulating volatiles is widespread among rhizosphere bacteria and strongly depends on culture conditions. *Environ. Microbiol.* 13 (11), 3047–3058. doi: 10.1111/j.1462-2920.2011.02582.x
- Bodenhausen, N., Bortfeld-Miller, M., Ackermann, M., and Vorholt, J. A. (2014). A synthetic community approach reveals plant genotypes affecting the phyllosphere microbiota. *PLoS Genet.* 10 (4), e1004283. doi: 10.1371/journal.pgen.1004283
- Bouffaud, M. L., Poirier, M. A., Muller, D., and Moëgne-Locoz, Y. (2014). Root microbiome relates to plant host evolution in maize and other poaceae. *Environ. Microbiol.* 16 (9), 2804–2814. doi: 10.1111/1462-2920.12442
- Buensateai, N., Yuen, G. Y., and Prathuangwong, S. (2008). The biocontrol bacterium *Bacillus amyloliquefaciens* KPS46 produces auxin, surfactin and extracellular proteins for enhanced growth of soybean plant. *Thai J. Agric. Sci.* 41 (3–4), 101–116.
- Buensateai, N., Sompong, M., Thamnu, K., Athinuwat, D., Brauman, A., and Plassard, C. (2013). The plant growth promoting bacterium *Bacillus* sp. CaSUT007 produces phytohormone and extracellular proteins for enhanced growth of cassava. *Afr. J. Microbiol. Res.* 7 (42), 4949–4954. doi: 10.5897/AJMR12.1839
- Bulgarelli, D., Rott, M., Schlaeppli, K., Ver Loren van Themaat, E., Ahmadijnejad, N., Assenza, F., et al. (2012). Revealing structure and assembly cues for arabisopsis root-inhabiting bacterial microbiota. *Nature*. 488 (7409), 91–95. doi: 10.1038/nature11336
- Bulgarelli, D., Schlaeppli, K., Spaepen, S., Van Themaat, E. V. L., and Schulze-Lefert, P. (2013). Structure and functions of the bacterial microbiota of plants. *Annu. Rev. Plant Biol.* 64, 807–838. doi: 10.1146/annurev-arplant-050312-120106
- Burkle, L. A., and Belote, R. T. (2015). Soil mutualists modify priority effects on plant productivity, diversity, and composition. *Appl. Veg. Sci.* 18, 332–342. doi: 10.1111/avsc.12149
- Calvo, P., Nelson, L., and Kloepper, J. W. (2014). Agricultural uses of plant biostimulants. *Plant Soil* 383 (1–2), 3–41. doi: 10.1007/S11104-014-2131-8/TABLES/1
- Camarena-Pozos, D. A., Flores-Núñez, V. M., López, M. G., López-Bucio, J., and Partida-Martínez, L. P. (2019). Smells from the desert: Microbial volatiles that affect plant growth and development of native and non-native plant species. *Plant Cell Environ.* 42 (4), 1368–1380. doi: 10.1111/PCE.13476
- Cardinale, M., Ratering, S., Suarez, C., Zapata Montoya, A. M., Geissler-Plaum, R., and Schnell, S. (2015). Paradox of plant growth promotion potential of rhizobacteria and their actual promotion effect on growth of barley (*Hordeum vulgare* L.) under salt stress. *Microbiol. Res.* 181, 22–32. doi: 10.1016/j.micres.2015.08.002
- Carrión, V. J., Perez-Jaramillo, J., Cordovez, V., Tracanna, V., de Hollander, M., Ruiz-Buck, D., et al. (2019). Pathogen-induced activation of disease-suppressive functions in the endophytic root microbiome. *Science*. 366 (6465), 606–612. doi: 10.1126/science.aaw9285
- Castrillo, G., Teixeira, P. J. P. L., Paredes, S. H., Law, T. F., de Lorenzo, L., Felcher, M. E., et al. (2017). Root microbiota drive direct integration of phosphate stress and immunity. *Nature*. 543 (7646), 513–518. doi: 10.1038/nature21417
- Cavello, I., Crespo, J., García, S., Zapiola, J., Luna, M., and Cavalitto, S. (2015). Plant growth promotion activity of keratinolytic fungi growing on a recalcitrant waste known as “hair waste”. *Biotechnol. Res. Int.* 2015, 1–10. doi: 10.1155/2015/952921
- Chanclud, E., and Morel, J.-B. (2016). Plant hormones: a fungal point of view. *Mol. Plant Pathol.* 17 (8), 1289–1297. doi: 10.1111/mpp.12393
- Chen, Y., Bonkowski, M., Shen, Y., Griffiths, B. S., Jiang, Y., Wang, X., et al. (2020). Root ethylene mediates rhizosphere microbial community reconstruction when chemically detecting cyanide produced by neighbouring plants. *Microbiome*. 8 (1), 4. doi: 10.1186/s40168-019-0775-6
- Contreras-Cornejo, H. A., Macías-Rodríguez, L., Cortes-Penagos, C., and Loípez-Bucio, J. (2009). *Trichoderma virens*, a plant beneficial fungus, enhances biomass production and promotes lateral root growth through an auxin-dependent mechanism in arabisopsis. *Plant Physiol.* 149 (3), 1579–1592. doi: 10.1104/pp.108.130369
- Diagne, N., Baudoin, E., Svistoonoff, S., Ouattara, C., Diouf, D., Kane, A., et al. (2018). Effect of native and allochthonous arbuscular mycorrhizal fungi on *Casuarina equisetifolia* growth and its root bacterial community. *Arid. Land Res. Manage.* 32 (2), 212–228. doi: 10.1080/15324982.2017.1406413
- Ditengou, F. A., Müller, A., Rosenkranz, M., Felten, J., Lasok, H., van Doorn, M. M., et al. (2015). Volatile signalling by sesquiterpenes from ectomycorrhizal fungi reprogrammes root architecture. *Nat. Commun.* 6 (1), 1–9. doi: 10.1038/ncomms7279
- du Jardin, P. (2015). Plant biostimulants: Definition, concept, main categories and regulation. *Sci. Hortic.* 196, 3–14. doi: 10.1016/J.SCIEN.2015.09.021
- Durán, P., Thiergart, T., Garrido-Oter, R., Agler, M., Kemen, E., Schulze-Lefert, P., et al. (2018). Microbial interkingdom interactions in roots promote arabisopsis survival. *Cell*. 175 (4), 973–983.e14. doi: 10.1016/j.cell.2018.10.020
- Egamberdieva, D., Wirth, S. J., Alqarawi, A. A., Abd-Allah, E. F., and Hashem, A. (2017). Phytohormones and beneficial microbes: Essential components for plants to balance stress and fitness. *Front. Microbiol.* 8. doi: 10.3389/fmicb.2017.02104
- El-Khawas, H., and Adachi, K. (1999). Identification and quantification of auxins in culture media of *Azospirillum* and *Klebsiella* and their effect on rice roots. *Biol. Fertil. Soils*. 28 (4), 377–381. doi: 10.1007/s003740050507
- El-Shanshoury, A. R. (1989). Growth promotion of wheat seedlings by *Streptomyces atroolivaceus*. *J. Agron. Crop Sci.* 163 (2), 109–114. doi: 10.1111/j.1439-037X.1989.tb00743.x

- Emmanuel, O. C., and Babalola, O. O. (2020). Productivity and quality of horticultural crops through co-inoculation of arbuscular mycorrhizal fungi and plant growth promoting bacteria. *Microbiol. Res.* 239, 126569. doi: 10.1016/j.micres.2020.126569
- Ezquer, I., Li, J., Ovecka, M., Baroja-Fernández, E., Muñoz, F. J., Montero, M., et al. (2010). Microbial volatile emissions promote accumulation of exceptionally high levels of starch in leaves in mono- and dicotyledonous plants. *Plant Cell Physiol.* 51 (10), 1674–1693. doi: 10.1093/PCP/PCQ126
- Fincchira, P., Parra, L., Mutis, A., Parada, M., and Quiroz, A. (2017). Volatiles emitted by *Bacillus* sp. BCT9 act as growth modulating agents on *Lactuca sativa* seedlings. *Microbiol. Res.* 203, 47–56. doi: 10.1016/j.micres.2017.06.007
- Fiorentino, N., Ventrino, V., Woo, S. L., Pepe, O., De Rosa, A., Gioia, L., et al. (2018). Trichoderma-based biostimulants modulate rhizosphere microbial populations and improve N uptake efficiency, yield and nutritional quality of leafy vegetables. *Front. Plant Sci.* 9. doi: 10.3389/fpls.2018.00743
- Francis, I., Holsters, M., and Vereecke, D. (2010). The gram-positive side of plant-microbe interactions. *Environ. Microbiol.* 12 (1), 1–12. doi: 10.1111/j.1462-2920.2009.01989.X
- Fu, S.-F., Sun, P.-F., Lu, H.-Y., Wei, J.-Y., Xiao, H.-S., Fang, W.-T., et al. (2016). Plant growth-promoting traits of yeasts isolated from the phyllosphere and rhizosphere of *Drosera spatulata* Lab. *Fungal Biol.* 120, 433–448. doi: 10.1016/j.funbio.2015.12.006
- Gámez-Arcas, S., Baroja-Fernández, E., García-Gómez, P., Muñoz, F. J., Almagro, G., Bahaji, A., et al. (2022). Action mechanisms of small microbial volatile compounds in plants. *J. Exp. Bot.* 73 (2), 498–510. doi: 10.1093/jxb/erab463
- García-Gómez, P., Almagro, G., Sánchez-López, Á.M., Bahaji, A., Amezttoy, K., Ricarte-Bermejo, A., et al. (2019). Volatile compounds other than CO₂ emitted by different microorganisms promote distinct posttranscriptionally regulated responses in plants. *Plant Cell Environ.* 42 (5), 1729–1746. doi: 10.1111/PCE.13490
- García-Gómez, P., Bahaji, A., Gámez-Arcas, S., Muñoz, F. J., Sánchez-López, Á.M., Almagro, G., et al. (2020). Volatiles from the fungal phytopathogen *Penicillium aurantiogriseum* modulate root metabolism and architecture through proteome resetting. *Plant Cell Environ.* 43 (10), 2551–2570. doi: 10.1111/PCE.13817
- Garnica-Vergara, A., Barrera-Ortiz, S., Muñoz-Parra, E., Raya-González, J., Méndez-Bravo, A., Macías-Rodríguez, L., et al. (2016). The volatile 6-pentyl-2H-pyran-2-one from *Trichoderma atroviride* regulates *Arabidopsis thaliana* root morphogenesis via auxin signaling and ETHYLENE INSENSITIVE 2 functioning. *New Phytol.* 209 (4), 1496–1512. doi: 10.1111/NPH.13725
- George, T. K., SubaidaBeevi, S., Asok, A. K., and Shaikmoideen, J. M. (2019). Plant growth promoting endophytic yeast *Geotrichum candidum* (jx 477426) from roots of *Bruguiera cylindrica*. *J. Microbiol. Biotechnol. Food Sci.* 9 (2), 267–272. doi: 10.15414/JMBFS.2019.9.2.267-272
- Ghimire, S. R., Charlton, N. D., and Craven, K. D. (2009). The mycorrhizal fungus, *Sebacina vermifera*, enhances seed germination and biomass production in switchgrass (*Panicum virgatum* L.). *Bioenergy Res.* 2 (1), 51–58. doi: 10.1007/s12155-009-9033-2
- Glare, T., Caradus, J., Gelernter, W., Jackson, T., Keyhani, N., Köhl, J., et al. (2012). Have pesticides come of age? *Trends Biotechnol.* 30 (5), 250–258. doi: 10.1016/j.tibtech.2012.01.003
- Goodwin, P. H. (2022). The rhizosphere microbiome of ginseng. *Microorganisms* 10, 1152. doi: 10.3390/microorganisms10061152
- Gutiérrez-Luna, F. M., López-Bucio, J., Altamirano-Hernández, J., and Valencia-Cantero, E. (2010). Reyes De la Cruz, h., and macías-rodríguez, l. plant growth-promoting rhizobacteria modulate root-system architecture in arabidopsis thaliana through volatile organic compound emission. *Symbiosis.* 51, 75–83. doi: 10.1007/s13199-010-0066-2
- Gu, Y., Wang, J., Xia, Z., and Wei, H. L. (2020). Characterization of a versatile plant growth-promoting rhizobacterium *Pseudomonas mediterranea* strain S58. *Microorganisms.* 8 (3), 334. doi: 10.3390/MICROORGANISMS8030334
- Hamayun, M., Khan, S. A., Khan, A. L., Rehman, G., Kim, Y. H., Iqbal, I., et al. (2010). Gibberellin production and plant growth promotion from pure cultures of cladosporium sp. MH-6 isolated from cucumber (*Cucumis sativus* L.). *Mycologia.* 102 (5), 989–995. doi: 10.3852/09-261
- Hart, M. M., Antunes, P. M., Chaudhary, V. B., and Abbott, L. K. (2018). Fungal inoculants in the field: Is the reward greater than the risk? *Funct. Ecol.* 32 (1), 126–135. doi: 10.1111/1365-2435.12976
- Hartmann, A., Rothballer, M., Hense, B. A., and Schröder, P. (2014). Bacterial quorum sensing compounds are important modulators of microbe-plant interactions. *Front. Plant Sci.* 5. doi: 10.3389/FPLS.2014.00131
- Hegazi, M. A., Metwaly, M. M. S., and Belal, E. B. (2015). Influence of plant growth promoting bacteria (PGPB) on coriander (*Coriandrum sativum* L.) and dill (*Anethum graveolens* L.) plants. *J. Plant Production.* 6 (2), 205–218. doi: 10.21608/jpp.2015.49299
- He, D., Singh, S. K., Peng, L., Kaushal, R., Vilchez, J. I., Shao, C., et al. (2022). Flavonoid-attracted aeromonas sp. from the arabidopsis root microbiome enhances plant dehydration resistance. *ISME J.* 16, 2622–2632. doi: 10.1038/s41396-022-01288-7
- Hilber-Bodmer, M., Schmid, M., Ahrens, C. H., and Freimoser, F. M. (2017). Competition assays and physiological experiments of soil and phyllosphere yeasts identify *Candida subhashii* as a novel antagonist of filamentous fungi. *BMC Microbiol.* 17, 4. doi: 10.1186/s12866-016-0908-z
- Hiruma, K., Gerlach, N., Sacristán, S., Nakano, R. T., Hacquard, S., Kracher, B., et al. (2016). Root endophyte *Colletotrichum tofieldiae* confers plant fitness benefits that are phosphate status dependent. *Cell.* 165 (2), 464–474. doi: 10.1016/j.cell.2016.02.028
- Hoeksema, J. D., Chaudhary, V. B., Gehring, C. A., Johnson, N. C., Karst, J., Koide, R. T., et al. (2010). A meta-analysis of context-dependency in plant response to inoculation with mycorrhizal fungi. *Ecol. Lett.* 13 (3), 394–407. doi: 10.1111/j.1461-0248.2009.01430.x
- Huang, A. C., Jiang, T., Liu, Y. X., Bai, Y. C., Reed, J., Qu, B., et al. (2019). A specialized metabolic network selectively modulates arabidopsis root microbiota. *Science.* 364 (6440), eaau6389. doi: 10.1126/SCIENCE.AAU6389
- Hwang, J. S., You, Y. H., Bae, J. J., Khan, S. A., Kim, J. G., and Choo, Y. S. (2011). Effects of endophytic fungal secondary metabolites on the growth and physiological response of *Carex kobomugi* ohwi. *J. Coast. Res.* 27 (3), 544–548. doi: 10.2112/JCOASTRES-D-10-00090.1
- Idris, E. E. S., Bochow, H., Ross, H., and Borriss, R. (2004). Use of *Bacillus subtilis* as biocontrol agent. VI. phytohormone-like action of culture filtrates prepared from plant-growth promoting bacillus amyloliquefaciens FZB24, FZB42, FZB45 and *Bacillus subtilis* FZB37. *J. Plant Dis. Prof.* 111, 583–597.
- Jalander, V., and Gachande, B. D. (2012). Effect of fungal metabolites of some rhizosphere soil fungi on seed germination and seedling growth of some pulses and cereals. *Sci. Res. Repor.* 2 (3), 265–267.
- Jiang, C. H., Xie, Y. S., Zhu, K., Wang, N., Li, Z. J., Yu, G. J., et al. (2019). Volatile organic compounds emitted by bacillus sp. JC03 promote plant growth through the action of auxin and strigolactone. *Plant Growth Regul.* 87 (2), 317–328. doi: 10.1007/S10725-018-00473-Z
- Kanchiswamy, C. N., Malnoy, M., and Maffei, M. E. (2015). Chemical diversity of microbial volatiles and their potential for plant growth and productivity. *Front. Plant Sci.* 6. doi: 10.3389/FPLS.2015.00151
- Kaur, T., Rani, R., and Manhas, R. K. (2019). Biocontrol and plant growth promoting potential of phylogenetically new *Streptomyces* sp. MR14 of rhizospheric origin. *AMB Express.* 9 (1), 125. doi: 10.1186/S13568-019-0849-7
- Khalmuratova, I., Choi, D.-H., Kim, J.-G., and Lee, I.-S. (2021). Endophytic fungi of salt-tolerant plants: diversity and ability to promote plant growth. *J. Microbiol. Biotechnol.* 31, 1526–1532. doi: 10.4014/jmb.2106.06007
- Khan, A. L., Hamayun, M., Ahmad, N., Hussain, J., Kang, S. M., Kim, Y. H., et al. (2011). Salinity stress resistance offered by endophytic fungal interaction between *Penicillium minioluteum* LHL09 and glycine max. I. *World J. Microbiol. Biotechnol.* 21 (9), 893–902. doi: 10.4014/JMB.1103.03012
- Khan, A. R., Ullah, I., Waqas, M., Shahzad, R., Hong, S.-J., Park, G.-S., et al. (2015). Plant growth-promoting potential of endophytic fungi isolated from *Solanum nigrum* leaves. *World J. Microbiol. Biotechnol.* 31, 1461–1466. doi: 10.1007/s11274-015-1888-0
- Khokhar, I., Haider, M. S., Mukhtar, I., Ali, A., Mushtaq, S., and Ashfaq, M. (2013). Effect of *Penicillium* species culture filtrate on seedling growth of wheat. *Int. J. Agric. Res.* 3 (1), 24–29.
- Kim, J. M., To, T. K., Matsui, A., Tanoi, K., Kobayashi, N. I., Matsuda, F., et al. (2017). Acetate-mediated novel survival strategy against drought in plants. *Nat. Plants.* 3, 17097. doi: 10.1038/NPLANTS.2017.97
- Kröber, M., Wibberg, D., Grosch, R., Eikmeyer, F., Verwaaijen, B., Chowdhury, S. P., et al. (2014). Effect of the strain *Bacillus amyloliquefaciens* FZB42 on the microbial community in the rhizosphere of lettuce under field conditions analyzed by whole metagenome sequencing. *Front. Microbiol.* 5. doi: 10.3389/FMICB.2014.00252
- Kwak, M. J., Kong, H. G., Choi, K., Kwon, S. K., Song, J. Y., Lee, J., et al. (2018). Rhizosphere microbiome structure alters to enable wilt resistance in tomato. *Nat. Biotech.* 36 (11), 1100–1116. doi: 10.1038/NBT.4232
- Lee, S., Behringer, G., Hung, R., and Bennett, J. (2019a). Effects of fungal volatile organic compounds on *Arabidopsis thaliana* growth and gene expression. *Fungal Cell.* 37, 1–9. doi: 10.1016/j.funeco.2018.08.004
- Lee, S. A., Kanth, B. K., Kim, H. S., Kim, T. W., Sang, M. K., Song, J., et al. (2019b). Complete genome sequence of the plant growth-promoting endophytic bacterium *Rhodanobacter glycinis* T01E-68 isolated from tomato (*Solanum lycopersicum* L.) plant roots. *Korean J. Microbiol.* 55 (4), 422–424. doi: 10.7845/KJM.2019.9115
- Lee, K. E., Radhakrishnan, R., Kang, S. M., You, Y. H., Joo, G. J., Lee, I. J., et al. (2015). *Enterococcus faecium* LKE12 cell-free extract accelerates host plant growth

- via gibberellin and indole-3-acetic acid secretion. *World J. Microbiol. Biotechnol.* 25 (9), 1467–1475. doi: 10.4014/JMB.1502.02011
- Liu, H., Brettell, L. E., Qiu, Z., and Singh, B. K. (2020). Microbiome-mediated stress resistance in plants. *Trends Plant Sci.* 25 (8), 733–743. doi: 10.1016/j.tplants.2020.03.014
- Liu, H., Qiu, Z., Ye, J., Verma, J. P., Li, J., and Singh, B. K. (2022). Effective colonisation by a bacterial synthetic community promotes plant growth and alters soil microbial community. *J. Sustain. Agric.* 1 (1), 30–42. doi: 10.1002/sae2.12008
- Li, J., Wang, J., Liu, H., Macdonald, C. A., and Singh, B. K. (2022). Application of microbial inoculants significantly enhances crop productivity: A meta-analysis of studies from 2010 to 2020. *J. Sust. Agric. Environ.* 1 (3), 216–225. doi: 10.1002/sae2.12028
- Macias-Benitez, S., Garcia-Martinez, A. M., Caballero Jimenez, P., Gonzalez, J. M., Tejada Moral, M., and Parrado Rubio, J. (2020). Rhizospheric organic acids as biostimulants: Monitoring feedbacks on soil microorganisms and biochemical properties. *Front. Plant Sci.* 11. doi: 10.3389/FPLS.2020.00633
- Marin, O., González, B., and Poupin, M. J. (2021). From microbial dynamics to functionality in the rhizosphere: a systematic review of the opportunities with synthetic microbial communities. *Front. Plant Sci.* 12. doi: 10.3389/fpls.2021.650609
- Meena, K. K., Kumar, M., Kalyuzhnaya, M. G., Yandigeri, M. S., Singh, D. P., Saxena, A. K., et al. (2012). Epiphytic pink-pigmented methylotrophic bacteria enhance germination and seedling growth of wheat (*Triticum aestivum*) by producing phytohormone. *Antonie van Leeuwenhoek.* 101 (4), 777–786. doi: 10.1007/S10482-011-9692-9
- Melo, I. S. D., and Piccinin, E. (1999). Toxic metabolites from culture filtrate of *Fusarium oxysporum* and its effects on cucumber cells and plantlets. *Rev. microbiologia.* 30, 104–106. doi: 10.1590/S0001-37141999002000003
- Middleton, E. L., Rihardson, S., Koziol, L., Palmer, C. E., Yermakov, Z. Y., Henning, J. A., et al. (2015). Locally adapted arbuscular mycorrhizal fungi improve vigor and resistance to herbivory of native prairie plant species. *Ecosphere* 61–, 16. doi: 10.1890/ES15-00152.1
- Miransari, M. (2011). Soil microbes and plant fertilization. *Appl. Microbiol. Biotechnol.* 92 (5), 875–885. doi: 10.1007/S00253-011-3521-Y
- Morcillo, R. J. L., and Manzanera, M. (2021). The effects of plant-associated bacterial exopolysaccharides on plant abiotic stress tolerance. *Metabolites.* 11 (6), 337. doi: 10.3390/METABO11060337
- Morcillo, R. J., Singh, S. K., He, D., An, G., Vilchez, J. I., Tang, K., et al. (2020). Rhizobacterium-derived diacyetyl modulates plant immunity in a phosphate-dependent manner. *EMBO J.* 39 (2), e102602. doi: 10.15252/EMBJ.2019102602
- Murali, M., and Amruthesh, K. N. (2015). Plant growth-promoting fungus *Penicillium oxalicum* enhances plant growth and induces resistance in pearl millet against downy mildew disease. *J. Phytopathol.* 163, 743–754. doi: 10.1111/jph.12371
- Naamala, J., and Smith, D. L. (2021). Microbial derived compounds, a step toward enhancing microbial inoculants technology for sustainable agriculture. *Front. Microbiol.* 12. doi: 10.3389/fmicb.2021.634807
- Noceto, P. A., Bettenfeld, P., Boussageon, R., Hériché, M., Sportes, A., van Tuinen, D., et al. (2021). Arbuscular mycorrhizal fungi, a key symbiosis in the development of quality traits in crop production, alone or combined with plant growth-promoting bacteria. *Mycorrhiza.* 31 (6), 655–669. doi: 10.1007/S00572-021-01054-1
- Ogórek, R. (2016). Enzymatic activity of potential fungal plant pathogens and the effect of their culture filtrates on seed germination and seedling growth of garden cress (*Lepidium sativum* L.). *Eur. J. Plant Pathol.* 145 (2), 469–481. doi: 10.1007/s10658-016-0860-7
- Onofre-Lemus, J., Hernández-Lucas, I., Girard, L., Caballero-Mellado, J.A.C.C. (2009). (1-aminocyclopropane-1-carboxylate) deaminase activity, a widespread trait in burkholderia species and its growth-promoting effect on tomato plants. *Appl. Environ. Microbiol.* 75, 6581–6590. doi: 10.1128/AEM.01240-09
- Ortiz-Castro, R., Contreras-Cornejo, H. A., Macias-Rodriguez, L., and López-Bucio, J. (2009). The role of microbial signals in plant growth and development. *Plant Signal. Behav.* 4 (8), 701–712. doi: 10.4161/PSB.4.8.9047
- Oskiera, M., Szczec, M., Stępowaska, A., Smolińska, U., and Bartoszewski, G. (2017). Monitoring of *Trichoderma* species in agricultural soil in response to application of biopreparations. *Biol. Control.* 113, 65–72. doi: 10.1016/j.biocontrol.2017.07.005
- Papavizas, G. C. (1982). Survival of *Trichoderma harzianum* in soil and in pea and bean rhizospheres. *Phytopathology.* 72 (1), 121. doi: 10.1094/PHYTO-72-121
- Parnell, J. J., Berka, R., Young, H. A., Sturino, J. M., Kang, Y., Barnhart, D. M., et al. (2016). From the lab to the farm: An industrial perspective of plant beneficial microorganisms. *Front. Plant Sci.* 7. doi: 10.3389/FPLS.2016.01110
- Parveen, S., Wani, A. H., and Bhat, M. Y. (2019). Effect of culture filtrates of pathogenic and antagonistic fungi on seed germination of some economically important vegetables. *Braz. J. Biol. Sci.* 6 (12), 133–139. doi: 10.21472/bjbs.061212
- Pellegrini, M., Pagnani, G., Bernardi, M., Mattedi, A., Spera, D. M., and del Gallo, M. (2020). Cell-free supernatants of plant growth-promoting bacteria: a review of their use as biostimulant and microbial biocontrol agents in sustainable agriculture. *Sustainability.* 12 (23), 9917. doi: 10.3390/SU12239917
- Pérez-Sánchez, A., Singh, S., Pérez-Sánchez, E. J., and Segura-Silva, R. M. (2018). Techno-economic evaluation and conceptual design of a liquid biofertilizer plant. *Rev. Colombiana Biotecnología* 20 (2), 6–18. doi: 10.15446/rev.colomb.biote.v20n2.77053
- Perin, L., Martínez-Aguilar, L., Paredes-Valdez, G., Baldani, J. I., Estrada-de los Santos, P., Reis, V. M., et al. (2006). *Burkholderia silvaticola* sp. nov., a diazotrophic bacterium associated with sugar cane and maize. *Int. J. Syst. Evol. Microbiol.* 56 (8), 1931–1937. doi: 10.1099/IJS.0.64362-0
- Pineda, A., Dicke, M., Pieterse, C. M. J., and Pozo, M. J. (2013). Beneficial microbes in a changing environment: are they always helping plants to deal with insects? *Funct. Ecol.* 27 (3), 574–586. doi: 10.1111/1365-2435.12050
- Posada, L. F., Ramírez, M., Ochoa-Gómez, N., Cuellar-Gaviria, T. Z., Argel-Roldan, L. E., Ramirez, C. A., et al. (2016). Bioprospecting of aerobic endospore-forming bacteria with biotechnological potential for growth promotion of banana plants. *Sci. Hortic.* 212, 81–90. doi: 10.1016/j.scienta.2016.09.040
- Puri, A., Padda, K. P., and Chanway, C. P. (2020). Sustaining the growth of Pinaceae trees under nutrient-limited edaphic conditions via plant-beneficial bacteria. *PLoS One* 2615 (8), e0238055. doi: 10.1371/journal.pone.0238055
- Rabhi, N. E. H., Silini, A., Cherif-Silini, H., Yahiaoui, B., Lekired, A., Robineau, M., et al. (2018). *Pseudomonas knackmussii* MLR6, a rhizospheric strain isolated from halophyte, enhances salt tolerance in arabidopsis thaliana. *J. Appl. Microbiol.* 125 (6), 1836–1851. doi: 10.1111/JAM.14082
- Radhakrishnan, R., Kang, S. M., Baek, I. Y., and Lee, I. J. (2014). Characterization of plant growth-promoting traits of *Penicillium* species against the effects of high soil salinity and root disease. *J. Plant Interact.* 9 (1), 754–762. doi: 10.1080/17429145.2014.930524
- Radhakrishnan, R., Shim, K. B., Lee, B. W., Hwang, C. D., Pae, S. B., Park, C. H., et al. (2013). IAA-producing penicillium sp. NICS01 triggers plant growth and suppresses *Fusarium* sp.-induced oxidative stress in sesame (*Sesamum indicum* L.). *World J. Microbiol. Biotechnol.* 23 (6), 856–863. doi: 10.4014/JMB.1209.09045
- Rahman, M. M., Mostofa, M. G., Rahman, M. A., Islam, M. R., Keya, S. S., Das, A. K., et al. (2019). Acetic acid: a cost-effective agent for mitigation of seawater-induced salt toxicity in mung bean. *Sci. Rep.* 9, 1–15. doi: 10.1038/s41598-019-51178-w
- Rahman, M. A., Sultana, R., Begum, M. F., and Alam, M. F. (2012). Effect of culture filtrates of *Trichoderma* on seed germination and seedling growth in chili international journal of biosciences. *Int. J. Biosci.* 2 (4), 46–55.
- Raithak, P. V., and Gachande, B. D. (2013). Effect of culture filtrates of tomato plant pathogenic fungi on seed germination and seedling growth of tomato (*Lycopersicon esculentum* mill.). *Curr. Botany.* 4 (1), 9–11.
- Rashnoo, A., Movahedi, Z., Rostami, M., and Ghabooli, M. (2020). *Piriformospora indica* culture filtrate and biofertilizer (nitrokar) promote chicory (*Cichorium intybus* L.) growth and morpho-physiological traits in an aeroponic system and soil culture. *Int. J. Hortic. Sci. Technol.* 7, 353–363. doi: 10.22059/ijhst.2020.292414.324
- Ravensberg, W. J. (2011). A roadmap to the successful development and commercialization of microbial pest control products for control of arthropods. (Vol. 10). *Springer Sci. Business Media.* doi: 10.1007/978-94-007-0437-4
- Reddy, C. A., and Saravanan, R. S. (2013). Polymicrobial multi-functional approach for enhancement of crop productivity. *Adv. Appl. Microbiol.* 82, 53–113. doi: 10.1016/B978-0-12-407679-2.00003-X
- Rodriguez, H., Fraga, R., Gonzalez, T., and Bashan, Y. (2007). Genetics of phosphate solubilization and its potential applications for improving plant growth-promoting bacteria. *First Int. Meeting Microbial Phosphate Solubilization* 102, 15–21. doi: 10.1007/978-1-4020-5765-6_2
- Rodriguez-Morgado, B., Jiménez, P. C., Moral, M. T., and Rubio, J. P. (2017). Effect of l-lactic acid from whey wastes on enzyme activities and bacterial diversity of soil. *Biol. Fertil. Soils.* 53 (4), 389–396. doi: 10.1007/S00374-017-1187-Z
- Rondina, A. B. L., dos Santos Sanzovo, A. W., Guimarães, G. S., Wendling, J. R., Nogueira, M. A., and Hungria, M. (2020). Changes in root morphological traits in soybean co-inoculated with *Bradyrhizobium* spp. and *Azospirillum brasilense* or treated with *A. brasilense* exudates. *Biol. Fertil. Soils.* 56 (4), 537–549. doi: 10.1007/s00374-020-01453-0
- Ryu, C. M., Faragt, M. A., Hu, C. H., Reddy, M. S., Wei, H. X., Paré, P. W., et al. (2003). Bacterial volatiles promote growth in arabidopsis. *Proc. Natl. Acad. Sci. U.S.A.* 100 (8), 4927–4932. doi: 10.1073/PNAS.0730845100
- Saber, W. I. A., Ghoneem, K. M., Rashad, Y. M., and Al-Askar, A. A. (2017). *Trichoderma harzianum* WKY1: an indole acetic acid producer for growth improvement and anthracnose disease control in sorghum. *Biocontrol Sci. Technol.* 27 (5), 654–676. doi: 10.1080/09583157.2017.1321733

- Saha, M., Sarkar, S., Sarkar, B., Sharma, B. K., Bhattacharjee, S., and Tribedi, P. (2016). Microbial siderophores and their potential applications: a review. *Environ. Sci. Pollut. Res.* 23 (5), 3984–3999. doi: 10.1007/S11356-015-4294-0
- Sánchez-López, Á.M., Baslam, M., de Diego, N., Muñoz, F. J., Bahaji, A., Almagro, G., et al. (2016). Volatile compounds emitted by diverse phytopathogenic microorganisms promote plant growth and flowering through cytokinin action. *Plant Cell Environ.* 39 (12), 2592–2608. doi: 10.1111/PCE.12759
- Sasse, J., Martinoia, E., and Northen, T. (2018). Feed your friends: do plant exudates shape the root microbiome? *Trends Plant Sci.* 23 (1), 25–41. doi: 10.1016/J.TPLANTS.2017.09.003
- Schmidt, R., Köberl, M., Mostafa, A., Ramadan, E. M., Monschein, M., Jensen, K. B., et al. (2014). Effects of bacterial inoculants on the indigenous microbiome and secondary metabolites of chamomile plants. *Front. Microbiol.* 5. doi: 10.3389/fmicb.2014.00064
- Schulz-Bohm, K., Martín-Sánchez, L., and Garbeva, P. (2017). Microbial volatiles: small molecules with an important role in intra- and interkingdom interactions. *Front. Microbiol.* 8. doi: 10.3389/fmicb.2017.02484
- Shayanthan, A., Ordoñez, P. A. C., and Oresnik, I. J. (2022). The role of synthetic microbial communities (SynCom) in sustainable agriculture. *Front. Agron.* 4. doi: 10.3389/fagro.2022.896307
- Sirenberg, A., Göbel, C., Grond, S., Czempinski, N., Ratzinger, A., Karlovsky, P., et al. (2007). *Piriformospora indica* affects plant growth by auxin production. *Physiol. Plant* 131 (4), 581–589. doi: 10.1111/J.1399-3054.2007.00983.X
- Sitaraman, R. (2015). *Pseudomonas* spp. as models for plant-microbe interactions. *Front. Plant Sci.* 6. doi: 10.3389/fpls.2015.00787
- Spaepen, S., Vanderleyden, J., and Remans, R. (2007). Indole-3-acetic acid in microbial and microorganism-plant signaling. *FEMS Microbiol. Rev.* 31 (4), 425–448. doi: 10.1111/J.1574-6976.2007.00072.X
- Spinelli, V., Brasili, E., Sciubba, F., Ceci, A., Giampaoli, O., Miccheli, A., et al. (2022). Biostimulant effects of *Chaetomium globosum* and *Minimedusa polyspora* culture filtrates on *Cichorium intybus* plant: growth performance and metabolomic traits. *Front. Plant Sci.* 13. doi: 10.3389/fpls.2022.879076
- Sung, G. H., Shrestha, B., Park, K. B., Han, S. K., and Sung, J. M. (2011). Enhancing effect of *Shimizuomyces paradoxus* on seed germination and seedling growth of canola, plant growth of cucumber, and harvest of tomato. *Mycobiology.* 39 (1), 7. doi: 10.4489/MYCO.2011.39.1.007
- Svenningsen, N. B., Watts-Williams, S. J., Joner, E. J., Battini, F., Efthymiou, A., Cruz-Paredes, C., et al. (2018). Suppression of the activity of arbuscular mycorrhizal fungi by the soil microbiota. *ISME J.* 12 (5), 1296–1307. doi: 10.1038/s41396-018-0059-3
- Tallapragada, P., Dikshit, R., and Seshagiri, S. (2015). Isolation and optimization of IAA producing *Burkholderia seminalis* and its effect on seedlings of tomato. *Songklanakarinn J. Sci. Technol.* 37 (5), 553–559.
- Trujillo-Roldán, M. A., Valdez-Cruz, N. A., González-Monterrubio, C. F., Acevedo-Sánchez, E. V., Martínez-Salinas, C., García-Cabrera, R. I., et al. (2013). Scale-up from shake flasks to pilot-scale production of the plant growth-promoting bacterium *Azospirillum brasilense* for preparing a liquid inoculant formulation. *Appl. Microbiol. Biotechnol.* 97, 9665–9674. doi: 10.1007/s00253-013-5199-9
- Tsavkelova, E. A., Klimova, S. Y., Cherdynseva, T. A., and Netrusov, A. I. (2006a). Microbial producers of plant growth stimulators and their practical use: A review. *Appl. Biochem. Microbiol.* 42 (2), 117–126. doi: 10.1134/S0003683806020013
- Tsavkelova, E. A., Klimova, S. Y., Cherdynseva, T. A., and Netrusov, A. I. (2006b). Hormones and hormone-like substances of microorganisms: a review. *Appl. Biochem. Microbiol.* 42 (3), 229–235. doi: 10.1134/S000368380603001X
- Utsumi, Y., Utsumi, C., Tanaka, M., Ha, C., Takahashi, S., Matsui, A., et al. (2019). Acetic acid treatment enhances drought avoidance in cassava (*Manihot esculenta* crantz). *Front. Plant Sci.* 10. doi: 10.3389/fpls.2019.00521
- Vankudoth, K. R., Sivadeveuni, G., and Reddy, S. M. (2015). Influence of different species of *Penicillium* and their culture filtrates on seed germination and seedling growth of sorghum. *J. Biochem. Technol.* 5 (4), 832–837.
- Varma, A., Verma, S., Sudha, Sahay, N., Bütehorn, B., and Franken, P. (1999). *Piriformospora indica*, a cultivable plant-growth-promoting root endophyte. *Appl. Environ. Microbiol.* 65 (6), 2741–2744. doi: 10.1128/AEM.65.6.2741-2744.1999
- Vassilev, N., Vassileva, M., Martos, V., Garcia del Moral, L. F., Kowalska, J., Tylkowski, B., et al. (2020). Formulation of microbial inoculants by encapsulation in natural polysaccharides: focus on beneficial properties of carrier additives and derivatives. *Front. Plant Sci.* 11. doi: 10.3389/fpls.2020.00270
- Vilchez, J. I., Yang, Y., He, D., Zi, H., Peng, L., Lv, S., et al. (2020). DNA Demethylases are required for myo-inositol-mediated mutualism between plants and beneficial rhizobacteria. *Nat. Plants.* 6 (8), 983–995. doi: 10.1038/S41477-020-0707-2
- Wang, Y., Zhao, A., Morcillo, R. J. L., Yu, G., Xue, H., Rufian, J. S., et al. (2021). A bacterial effector protein uncovers a plant metabolic pathway involved in tolerance to bacterial wilt disease. *Mol. Plant* 14 (8), 1281–1296. doi: 10.1016/J.MOLP.2021.04.014
- Waqas, M., Kim, Y. H., Khan, A. L., Shahzad, R., Asaf, S., Hamayun, M., et al. (2017). Additive effects due to biochar and endophyte application enable soybean to enhance nutrient uptake and modulate nutritional parameters. *J. Zhejiang Univ. Sci.* 18 (2), 109. doi: 10.1631/JZUS.B1500262
- Wu, Y., He, Y., Yin, H., Chen, W., Wang, Z., Xu, L., et al. (2012). Isolation of phosphate-solubilizing fungus and its application in solubilization of rock phosphates. *Pakistan J. Biol. Sci. Pjbs* 15, 1144–1151. doi: 10.3923/pjbs.2012.1144.1151
- Wu, L., Li, X., Ma, L., Borriss, R., Wu, Z., and Gao, X. (2018). Acetoin and 2,3-butanediol from *Bacillus amyloliquefaciens* induce stomatal closure in *Arabidopsis thaliana* and *Nicotiana benthamiana*. *J. Exp. Bot.* 69 (22), 5625–5635. doi: 10.1093/jxb/ery326
- Yakhin, O. I., Lubyantsev, A. A., Yakhin, I. A., and Brown, P. H. (2017). Biostimulants in plant science: A global perspective. *Front. Plant Sci.* 7. doi: 10.3389/fpls.2016.02049
- Yandigeri, M. S., Meena, K. K., Singh, D., Malviya, N., Singh, D. P., Solanki, M. K., et al. (2012). Drought-tolerant endophytic actinobacteria promote growth of wheat (*Triticum aestivum*) under water stress conditions. *Plant Growth Regul.* 68 (3), 411–420. doi: 10.1007/S10725-012-9730-2
- You, Y.-H. (2012). Fungal diversity and plant growth promotion of endophytic fungi from six halophytes in suncheon bay. *J. Microbiol. Biotechnol.* 22, 1549–1556. doi: 10.4014/jmb.1205.05010
- Yuan, J., Zhao, M., Li, R., Huang, Q., Raza, W., Rensing, C., et al. (2017). Microbial volatile compounds alter the soil microbial community. *Environ. Sci. Pollut. Res.* 24 (28), 22485–22493. doi: 10.1007/s11356-017-9839-y
- Zhalnina, K., Louie, K. B., Hao, Z., Mansoori, N., da Rocha, U. N., Shi, S., et al. (2018). Dynamic root exudate chemistry and microbial substrate preferences drive patterns in rhizosphere microbial community assembly. *Nat. Microbiol.* 3 (4), 470–480. doi: 10.1038/S41564-018-0129-3
- Zhang, H., Kim, M. S., Krishnamachari, V., Payton, P., Sun, Y., Grimson, M., et al. (2007). Rhizobacterial volatile emissions regulate auxin homeostasis and cell expansion in arabidopsis. *Planta.* 226 (4), 839–851. doi: 10.1007/S00425-007-0530-2
- Zhang, R., Ouyang, J., Xu, X., Li, J., Rehman, M., Deng, G., et al. (2022). Nematicidal activity of *Burkholderia arboris* J211 against *Meloidogyne incognita* on tobacco. *Front. Microbiol.* 13, 915546. doi: 10.3389/fmicb.2022.915546
- Zhang, H., Sun, Y., Xie, X., Kim, M. S., Dowd, S. E., and Paré, P. W. (2009). A soil bacterium regulates plant acquisition of iron via deficiency-inducible mechanisms. *Plant J.* 58 (4), 568–577. doi: 10.1111/J.1365-313X.2009.03803.X
- Zhang, H., Xie, X., Kim, M. S., Korniyev, D. A., Holaday, S., and Paré, P. W. (2008). Soil bacteria augment arabidopsis photosynthesis by decreasing glucose sensing and abscisic acid levels in planta. *Plant J.* 56 (2), 264–273. doi: 10.1111/J.1365-313X.2008.03593.X
- Zhong, Y., Yang, Y., Liu, P., Xu, R., Rensing, C., Fu, X., et al. (2019). Genotype and rhizobium inoculation modulate the assembly of soybean rhizobacterial communities. *Plant Cell Environ.* 42 (6), 2028–2044. doi: 10.1111/pce.13519
- Zhou, J.-Y., Li, X., Zheng, J.-Y., and Dai, C.-C. (2016). Volatiles released by endophytic *Pseudomonas fluorescens* promoting the growth and volatile oil accumulation in *Tractylodes lancea*. *Plant Physiol. Biochem.* 101, 132–140. doi: 10.1016/j.plaphy.2016.01.026