



Stable Isotope Fractionation of Metals and Metalloids in Plants: A Review

Matthias Wiggerhauser^{1*†}, Rebekah E. T. Moore^{2*†}, Peng Wang³, Gerd Patrick Bienert⁴, Kristian Holst Laursen⁵ and Simon Blotevogel⁶

¹Group of Plant Nutrition, Department of Environmental System Science, Institute of Agricultural Sciences, ETH Zurich, Zurich, Switzerland, ²MAGIC Group, Department of Earth Science and Engineering, Imperial College London, London, United Kingdom, ³College of Resources and Environmental Sciences, Nanjing Agricultural University, Nanjing, China, ⁴Crop Physiology, Molecular Life Sciences, Technical University of Munich, Freising, Germany, ⁵Plant Nutrients and Food Quality Research Group, Plant and Soil Science Section and Copenhagen Plant Science Centre, Department of Plant and Environmental Sciences, Faculty of Science, University of Copenhagen, Copenhagen, Denmark, ⁶Laboratoire Matériaux et Durabilité des Constructions (LMDC), UPS/INSA, Université Paul Sabatier - Toulouse III, Toulouse, France

OPEN ACCESS

Edited by:

Marc Hanikenne,
University of Liège, Belgium

Reviewed by:

Stephan Clemens,
University of Bayreuth, Germany
Maria Bernal,
Ruhr University Bochum, Germany

*Correspondence:

Matthias Wiggerhauser
matthias.wiggerhauser@usys.ethz.ch
Rebekah E. T. Moore
r.moore13@imperial.ac.uk

[†]These authors share first authorship

Specialty section:

This article was submitted to
Plant Nutrition,
a section of the journal
Frontiers in Plant Science

Received: 21 December 2021

Accepted: 17 February 2022

Published: 19 April 2022

Citation:

Wiggerhauser M, Moore RET, Wang P, Bienert GP, Laursen KH and Blotevogel S (2022) Stable Isotope Fractionation of Metals and Metalloids in Plants: A Review. *Front. Plant Sci.* 13:840941. doi: 10.3389/fpls.2022.840941

This work critically reviews stable isotope fractionation of essential (B, Mg, K, Ca, Fe, Ni, Cu, Zn, Mo), beneficial (Si), and non-essential (Cd, Tl) metals and metalloids in plants. The review (i) provides basic principles and methodologies for non-traditional isotope analyses, (ii) compiles isotope fractionation for uptake and translocation for each element and connects them to physiological processes, and (iii) interlinks knowledge from different elements to identify common and contrasting drivers of isotope fractionation. Different biological and physico-chemical processes drive isotope fractionation in plants. During uptake, Ca and Mg fractionate through root apoplast adsorption, Si through diffusion during membrane passage, Fe and Cu through reduction prior to membrane transport in strategy I plants, and Zn, Cu, and Cd through membrane transport. During translocation and utilization, isotopes fractionate through precipitation into insoluble forms, such as phytoliths (Si) or oxalate (Ca), structural binding to cell walls (Ca), and membrane transport and binding to soluble organic ligands (Zn, Cd). These processes can lead to similar (Cu, Fe) and opposing (Ca vs. Mg, Zn vs. Cd) isotope fractionation patterns of chemically similar elements in plants. Isotope fractionation in plants is influenced by biotic factors, such as phenological stages and plant genetics, as well as abiotic factors. Different nutrient supply induced shifts in isotope fractionation patterns for Mg, Cu, and Zn, suggesting that isotope process tracing can be used as a tool to detect and quantify different uptake pathways in response to abiotic stresses. However, the interpretation of isotope fractionation in plants is challenging because many isotope fractionation factors associated with specific processes are unknown and experiments are often exploratory. To overcome these limitations, fundamental geochemical research should expand the database of isotope fractionation factors and disentangle kinetic and equilibrium fractionation. In addition, plant growth studies should further shift toward hypothesis-driven experiments, for example, by integrating contrasting nutrient supplies, using established model plants, genetic approaches, and by combining isotope analyses

with complementary speciation techniques. To fully exploit the potential of isotope process tracing in plants, the interdisciplinary expertise of plant and isotope geochemical scientists is required.

Keywords: stable isotopes, fractionation, plant uptake, translocation, process tracing, multiple-collector-ICP-MS, metals, metalloids

INTRODUCTION

Stable isotopes are atoms with the same number of protons and electrons but a different number of neutrons and are considered stable if they do not undergo a measurable nuclear decay. Isotope fractionation describes small changes in the isotope composition (i.e., relative isotope abundances) during chemical reactions or physical processes. Isotope compositions of plant organs can provide a record of biological and physico-chemical processes that control an element's journey from their source to different organs, such as roots, leaves, and fruits. This information is complementary to organ-specific concentrations that provide information on element availability and utilization. This complementary information has been widely used for so-called "traditional isotopes" of, for example, H, C, N, O, and S to investigate for example water and fertilizer utilization, as well as nutrient assimilation (O'Leary et al., 1992; Tcherkez and Tea, 2013; Novak et al., 2019, 2021). Analytical advances during the past two decades now also allow the detection of naturally occurring variations of isotope compositions in plants for other, usually heavier elements (Hoefs, 2018), known as "non-traditional isotopes." Here, we critically review the fractionation of non-traditional stable isotopes to identify biological and physico-chemical processes that control the uptake and transport of these elements within plants (i.e., stable isotope process tracing). To this end, we: (i) provide interdisciplinary fundamentals on plant physiology and stable isotope process tracing, (ii) compile available data on isotope process tracing of B, Mg, Si, K, Ca, Fe, Ni, Cu, Zn, Mo, Cd, and Tl, (iii) interlink and discuss findings and approaches from the individual element reviews, and (iv) highlight best practices for data acquisition and interpretation, and define future research priorities. This review does not cover the use of non-traditional stable isotopes to trace the origin of elements (i.e., "source tracing") and to decipher complexation processes of elements onto soil components and thereby isotope fractionation between soil pools. Both topics have been reviewed recently (Komárek et al., 2021; Wang et al., 2021). Also beyond the scope of this review is the use of enriched stable isotopes as tracers to, for example, identify pathways of elements in plants (e.g., Yan et al., 2018). Finally, this review contributes to bridge plant science and isotope geochemistry by harmonizing nomenclature and working models of plant-based isotope studies to encourage interdisciplinary research and collaboration.

Non-traditional Isotopes: The Basics

In plants, isotope fractionation is usually mass-dependent (Table 1). Mass-dependent isotope effects are induced by slight differences of masses of distinct isotopes. Properties affected by the mass

difference are velocities and diffusivities, as well as vibrational and rotational frequencies and thermodynamic energies (Dauphas and Schauble, 2016). Differences in vibrational frequencies are the most important factor, as they determine isotope fractionation during chemical reactions. The general theory of isotope fractionation has been extensively reviewed elsewhere (Fry, 2006; Hoefs, 2018). We use Ca as an example in the following paragraphs to illustrate the basics of non-traditional isotope fractionation in plants. Isotope compositions are governed by equilibrium and kinetic isotope fractionation (Table 1). Equilibrium fractionation can occur when, for example, Ca ions in the soil or nutrient solution sorb to the negatively charged root apoplast (Cobert et al., 2011). In this case, thermodynamic equilibrium is reached when the adsorption and desorption of Ca ions are equally likely and occur at the same rate. Importantly, at isotope equilibrium, the individual Ca isotope ratios (e.g., $^{44}\text{Ca}/^{40}\text{Ca}$) of the adsorbed and desorbed (i.e., dissolved) phase remain the same over time. At equilibrium conditions, the isotope fractionation depends on the mass difference of the isotopes and the stiffness of the bonds involved. Heavier isotopes preferentially go into stiffer bonds (higher spring constant of the bond, Bigeleisen, 1965). This effect is due to the larger mass-dependent difference in zero-point energy in stiffer bonds. In applied sciences, the stiffness of a bond is often approximated by the bond length or the thermodynamic stability of a chemical species.

If a system is not in thermodynamic equilibrium, kinetic processes control the isotope fractionation. For example, during the precipitation of Ca oxalate in plants (Cobert et al., 2011). Precipitation can induce kinetic isotope fractionation as long as the quantity of mineral Ca oxalate is increasing. At this stage, the forward reaction is quicker than the backward reaction (i.e., the dissolution of Ca oxalate) and favorably light isotopes precipitate into Ca oxalate. This is due to the slightly lower activation energy of lighter isotopes. When the amount of Ca oxalate remains constant, an (isotope) equilibrium between mineral and solution will be reached and the resulting isotope fractionation is governed by equilibrium fractionation. Note that chemical equilibrium is usually established faster than isotope equilibrium.

Isotope fractionation differs in closed and open systems, which can be defined using Rayleigh modeling (Fry, 2006). In closed systems, only negligible quantities of reactants or reaction products are added or removed from a system. For instance, a plant that preferentially takes up light Ca isotopes from a nutrient solution will deplete the nutrient solution of Ca and light Ca isotopes if the nutrient solution is not frequently replenished. As the plant will continue to preferentially remove light isotopes from the nutrient solution, the nutrient solution will become enriched with heavy isotopes while the isotopes taken up by the plant

TABLE 1 | Glossary of terms relevant to isotope composition analysis.

Analytical precision	Usually expressed as the 2× standard deviation (2sd) of repeat measurements of a primary/bracketing standard or matrix-matched reference material (Supplementary Tables S1, S2), or the 2sd of repeated sample measurements. Gives a good estimate of the isotope composition resolvability. Often referred to as “external precision.”
Analyte purification	Isolation of analyte from the matrix of a digested sample. Most common method is ion exchange chromatography: percolation of acids of different molarities through resin-filled columns.
Digestion	Breakdown of molecules to allow liberation of individual elements. Uses hotplate or microwave oven-assisted acid digestion using nitric acid (and hydrogen peroxide) /aqua-regia /perchloric acid /hydrofluoric acid. For Si analysis, alkaline fusion (e.g., using NaOH) is more appropriate, since SiF ₆ is volatile after HF digestion.
Double spike	A mass bias correction technique for elements with ≥ 4 stable isotopes. A mixture of 2 enriched isotope solutions with known concentration and isotope composition is equilibrated with samples. An iterative procedure is used to deconvolute the data.
Equilibrium fractionation	Occurs when two or more substrates are in chemical (isotope) equilibrium. Heavier isotopes are preferentially incorporated into stiffer bonds in equilibrium reactions.
External normalization	A mass bias correction technique using a second element that is close in atomic mass and ionization potential to the analyte.
Isobaric interference	When isotopes of different elements have similar mass, mass spectrometry may be unable to distinguish between them, for example, ⁴⁰ Ca, ⁴⁰ Ar.
Isotope fractionation	Changes in proportions of isotopes that occur during chemical reactions or physical processes.
Kinetic (isotope) fractionation	In chemical reactions that are not at equilibrium, light isotopes accumulate in reaction products as they have slightly lower activation energies and therefore react faster.
Mass bias	Isotope fractionation that occurs inside mass spectrometers due to differences in transmission efficiencies between isotopes. This can induce large isotope fractionations, so post-measurement mathematical correction is required. Mass bias is much stronger in ICP than in TIMS.
Mass-dependent fractionation (MDF)	Fractionation scales with the difference in isotopic mass. For example, the mass difference between ⁶⁶ Zn and ⁶⁴ Zn is ~double that of ⁶⁶ Zn and ⁶⁴ Zn, so fractionations of ~2 and ~1‰ are expected (Maréchal et al., 1999).
Molecular interference	Also known as poly-atomic interferences. These occur within mass spectrometers when the mass of a molecule (e.g., ⁴⁰ Ar ¹⁶ O) overlaps with an isotope of the analyte (e.g., ⁵⁶ Fe).
Multiple collector inductively coupled plasma mass spectrometry	Use an Ar ICP source, magnetic sector analyzer, and Faraday collector array to simultaneously measure isotope abundances at precisions higher than attainable by single-collector ICP-MS. Enables analysis of a large variety of elements, including those with high ionization potentials, such as Zn. Samples are introduced as a solution or aerosol and even solid when laser ablation is used.
Procedural blank	A sample taken through every preparation step to enable the quantification of any contamination that can alter a samples' true isotope composition.
Sample-standard bracketing	A mass bias correction technique. Samples are analyzed between standards. Post-measurement, a correction is made relative to the standard before and after the sample.
Thermal ionization mass spectrometry (TIMS)	More traditional devices for isotope analysis. Samples are deposited on a filament that is heated by an electric current to ionize the analyte. Isotopes are separated using a magnetic sector and an array of faraday cups. TIMS measurements involve lower mass bias and interferences than MC-ICP-MS, however they are slower and accurate mass bias correction mostly relies on the double spike technique, drastically limiting the amount of possible analytes.

become successively heavier. Finally, when the entire Ca is taken up from the solution, the plant will have the same Ca isotope ratio as the solution initially had. In contrast, if the solution is frequently replenished, the plant is virtually exposed to an infinite Ca source and the system is considered open. In such a system, the plant will continuously take up light Ca isotopes from the nutrient solution and the Ca isotope ratios in the plant and in the solution do not change over time.

Isotope ratio analyses are more meaningful when mass balances are calculated. To calculate such “isotope mass balances” in plants, the dry weights, the plant organs (e.g., roots, leaves) concentrations, and isotope ratios of both the plant organs and the nutrient source need to be determined. For instance, the weighted mean of Ca isotope ratios in different plant organs provide the isotope ratio of the whole

plant and thereby the isotope fractionation associated with Ca uptake from soil to plant can be calculated. Note, these calculations consider that there is no significant efflux of Ca from the plant into the soil.

Metals and Metalloids in Plants: The Basics

Plants require 17 elements in order to complete their life cycle (**Table 2**). Besides hydrogen (H), carbon (C), and oxygen (O), which are the main building blocks of plant tissues, the remaining 14 elements are the macronutrients: nitrogen (N), magnesium (Mg), phosphorus (P), sulfur (S), potassium (K), calcium (Ca) and the micronutrients: boron (B), chlorine (Cl), manganese (Mn), iron (Fe), nickel (Ni), copper (Cu), zinc (Zn) and molybdenum (Mo) (Marschner and Marschner, 2012).

TABLE 2 | Overview of elements found in plant tissue that have been studied using stable isotope fractionation (listed according to atomic weight)**.

Element	Symbol (atomic weight)	Main plant available form(s)	Normal/sufficient content (in dry matter [§])	Phloem mobility	Functional roles in plants
Boron	B (10.81)	B(OH) ₄ ⁻ B(OH) ₃	5–60 µg/g	Low	Essential plant nutrient: Stabilization of cell walls, nucleotide biosynthesis, translocation, pollen germination
Magnesium	Mg (24.305)	Mg ²⁺	0.1–1%	High	Essential plant nutrient: Photosynthesis (central atom of chlorophyll), enzyme activation
Silicon	Si (28.085)	H ₄ SiO ₄	0.1–10%	Low	Beneficial plant nutrient: Alleviates abiotic and biotic stresses (salt, heavy metal, fungal disease, osmotic, drought)
Potassium	K (39.098)	K ⁺	1–6%	High	Essential plant nutrient: Translocation, osmoregulation, pH homeostasis, enzyme activation
Calcium	Ca (40.078)	Ca ²⁺	0.2–3%	Low	Essential plant nutrient: Cell wall synthesis and stabilization, signal transduction
Iron	Fe (55.845)	Fe ³⁺ Fe ²⁺	30–300 µg/g	Conditional	Essential plant nutrient: Enzyme activation, electron transport, chlorophyll biosynthesis, oxidative stress protection
Nickel	Ni (58.693)	Ni ²⁺	0.1–1 µg/g	Conditional	Essential plant nutrient: Enzyme activation, urea metabolism
Copper	Cu (63.546)	Cu ⁺ Cu ²⁺	4–25 µg/g	Conditional	Essential plant nutrient: Enzyme activation, synthesis of cell wall components, electron transport, oxidative stress protection
Zinc	Zn (65.38)	Zn ²⁺	15–80 µg/g	Conditional	Essential plant nutrient: Gene expression, enzyme activation, hormone and nucleic acid synthesis, oxidative stress protection
Molybdenum	Mo (95.95)	MoO ₄ ²⁻	0.1–1 µg/g	Conditional	Essential plant nutrient: Enzyme activation, nitrogen fixation, nitrate reduction
Cadmium	Cd (112.41)	Cd ²⁺	0.01–1 µg/g	Conditional	Non-essential, non-beneficial
Thallium	Tl (204.38)	Tl ⁺	site dependent	NA	Non-essential, non-beneficial

Main plant available form(s) of each element, normal or sufficient content, phloem mobility, functional roles in plants and selected additional information of relevance for interpretation of element-specific isotope fractionations. NA = information not available. [§]Dependent on the developmental stage of the plant, the plant species, the plant genotype, and the plant organ. ****Adapted from Pilon-Smits et al. (2009); Husted et al. (2011); Marschner and Marschner (2012); de Bang et al. (2020); www.iupac.org. For element-specific information on uptake, functional roles and utilization in plants, we refer to recent reviews for: B (Yoshinari and Takano, 2017), Ca (Thor, 2019), Mg (Chen et al., 2018), K (Wang and Wu, 2017), Si (Ma, 2015), Cu, Fe, Ni, Zn, and Cd (Andresen et al., 2018), Mo (Tejada-Jimenez et al., 2018).**

Besides the essential plant nutrients, many other elements of the periodic table are also found in plants. Some elements, such as silicon (Si), are considered beneficial for certain plant species. Others with no functional role are present in plant organs as a result of their environmental abundance and due to non-specific uptake by plants. Some of these elements represent a threat to plant growth and performance, for example, cadmium (Cd). Most elements are acquired as inorganic ions from the soil solution and are subsequently translocated in plant organs and utilized in a plethora of biochemical functions (Table 2).

Physiological Processes and Isotope Compositions in Plants

Physiological processes, such as root uptake, short and long-distance transport (in xylem or phloem), storage (e.g., vacuolar), metabolic functionalization, and compartmentalization (Figure 1A), are a series of physico-chemical processes that potentially change isotope compositions of a given element. Physico-chemical processes comprise water mass flow, sorption, ligand exchange, redox changes, precipitation, dissolution, and diffusion.

Plant Root Uptake of Elements

Plants take up most elements in specific, phytoavailable, chemical forms from the soil solution (Tables 2, 3). These are transported in roots *via* the apoplastic, symplastic, or transcellular pathways (Figure 1B; Table 3). Elements enter the root apoplast with water mass flow, mainly driven by the transpiration stream or by diffusion along an electrochemical gradient. The apoplastic mobility of elements is determined by the number and affinity of available adsorption, complexation, and exchange sites within cell walls for the specific element (Meychik et al., 2021). When a physico-chemical barrier restricts further apoplastic transport, elements have to cross a cell membrane to take the symplastic and transcellular pathways, which may require metabolic energy (Figure 1C; Table 3). One such barrier during root-to-shoot transport is the endodermis. It forms the boundary between the cortex and the stele and contains a lignin- and suberin-based hydrophobic radial cell wall “impregnation” called the “Casparian strip” (Geldner, 2013). To cross the endodermis, all elements have to enter the intracellular space at least




FIGURE 1 | Overview of physiological processes potentially causing isotope fractionation. **(A)** In shoots, isotope fractionation may occur during short distance transport across membranes, during long-distance transport in xylem (unidirectional transport) or in the phloem (bidirectional transport), due to metabolic functionalization (e.g., enzyme activation, ligand exchange), element storage or compartmentalization. Major physiological processes in roots are nutrient uptake and short distance transport across membranes. Elements are taken up from the rhizosphere and translocated *via* the xylem to sink tissue (young leaves or generative tissue, such as fruits, grains, or pods). Elements may also be re-translocated *via* the phloem from source tissue (e.g., old leaves) (Continued)

FIGURE 1 | to sink tissue. **(B)** Simplified model of symplastic (blue), apoplastic (red) versus transcellular (orange) element transport routes in plant roots. Cell-to-cell transport in the symplastic pathway occurs *via* plasmodesmata. In the apoplastic pathway elements travel in the cell wall and the intercellular space. The plasma membrane must be crossed to exit the root cortex and enter the stele and the vasculature system. **(C)** Elements may be bound to soil particles in the rhizosphere or to the negatively charged cell wall prior to uptake across the plasma membrane mediated by transport proteins (e.g., *via* carriers and channels). In the cytoplasm, elements may form complexes with enzymes and substrates—for example, ATP activation through Mg-ATP complexation (Adapted from Marschner and Marschner, 2012; Taiz et al., 2015). For glossary of terms and processes see **Table 3**.

temporarily (**Figure 1B**). After crossing the endodermis, an element is loaded into the vascular system (**Table 3**) for further root-to-shoot transport.

Cross-Membrane Transport of Elements

Elements are transported across membranes either by passive diffusion through the phospholipid bilayer or *via* an armada of specialized membrane transport proteins (**Figure 1C**; **Table 3**). Passive diffusion is negligible for charged molecules due to their restricted membrane permeability while passive channels, secondary active carriers, and active transporters generate an element-specific and rapidly controllable pathway through various membranes (Stein and Litman, 2015; Tang et al., 2020). Each transport protein has a unique activity. The subcellular localization of each membrane protein is mostly restricted to one cellular membrane system (e.g., plasma membrane, tonoplast, mitochondrial or chloroplasmic membrane) or even a spatially targeted fraction of a membrane system. For instance, certain transporters that control the uptake of nutrients, such as B and Si, are polarly localized solely at the distal side of root rhizodermis cells, while transporters responsible for the further translocation of nutrients into the cortex are localized to the proximal side (Ma and Yamaji, 2015; Yoshinari and Takano, 2017). Similar to animal epithelial cells, where the mechanisms for polar localization of nutrient transporters at the apical or basolateral membrane were intensively studied (Mellman and Nelson, 2008), many plant nutrient transporters that are either responsible for rhizodermal uptake or xylem loading, have been demonstrated to be polarly localized (Langowski et al., 2010 and references therein).

The main driving force for protein-mediated membrane transport is an electrochemical gradient (**Table 3**) which depends on differences in element concentrations and electrical potential inside and outside of the cell. The electrochemical gradient is established by membrane proteins (H⁺-ATPases), which pump protons out of cells against an inward directed proton gradient (**Figure 1C**). This process requires metabolic energy from ATP hydrolysis and generates an electrical gradient (positive outside, negative inside) and a pH gradient (acidic outside vs. neutral/alkaline inside) across the membrane. As a consequence, there is a permanent driving force (proton motive force) for cations into plant cells and anions out of plant cells (Falhof et al., 2016). These energetic downhill transport processes can be used to energize uphill transport processes that are mainly facilitated by carrier-type transport proteins (Stein and Litman, 2015; Tang et al., 2020).

TABLE 3 | Glossary of terms and processes relevant for understanding isotope fractionation in plants (in alphabetical order).

Active transport – cellular influx and efflux – short distance transport	Movement of an ion/molecule across the membrane against the concentration gradient of the transported atom/molecule and/or against the electrochemical gradient. Requires metabolic energy to pump molecules across a membrane (transport direction = uniport; transport rates = $1-10^2$ molecules per sec; energization = ATP/PPi -hydrolysis; protein types mediating the active transport = pumps, ATPases, which demands structural changes of the protein; examples = Ca, Zn, Mn, Cd, Co P-type ATPases, Cd, Pb, As, Hg (II), Sb ATP-Binding-Cassette (ABC)-transporters).
Apoplastic transport (extracellular)	Transport of an ion/molecule in the apoplast, which comprises the entirety of plant cell wall and the intercellular space, without entering the cell. Note, elements that fulfill functions in the apoplast, like Ca (cell wall stabilization), can be considered as taken up by the plant although they have not entered the symplast.
Bulk Soil	Soil outside of the rhizosphere and neither penetrated nor influenced by plant roots.
Compartmentalization	Separation of ion/molecule into isolated subcellular compartments (organelles, cell wall, etc.) or cells.
Dual-affinity transport proteins	Transport proteins which are suggested to transport efficiently at high or low nutrient concentrations due to their ability to switch their structural or biochemical protein properties <i>via</i> post-translational modifications (e.g., the potassium transporter AtKUP1).
Electrochemical Gradient	A difference in 1) charge and 2) the chemical concentration of a compound across the plasma membrane.
High-affinity transport proteins	Transport proteins which are suggested to dominantly transport at low nutrient concentrations; transport can be saturated (e.g., the iron transporter AtIRT1).
Ligand	A group, ion or molecule coordinated to a central atom or molecule in a complex. For example nicotianamine, phytate, citrate and malate.
Long-distance transport (<i>via</i> mass flow)	Transport of ion/molecule within the vasculature at the level of the whole plant linking a multitude of roots with branches and leaves <i>via</i> the specialized tissues called xylem (unidirectional transport, up in a plant ↑) and phloem (bidirectional transport, up and down in a plant ↓). Driven by pressure flow (hydrostatic or osmotic).
Low-affinity transport proteins	Transport proteins which are suggested to dominantly transport at high nutrient concentrations; transport seems non-saturable (e.g., the cation Cd transporter OsLCT1).
Passive transport – cellular influx and efflux – short distance transport	Movement of an ion/molecule across the membrane along the concentration gradient of the transported atom/molecule and/or along the electrochemical gradient. Protein-mediated or <i>via</i> sole diffusion across the membrane (transport direction = uniport; transport rates = 10^6-10^9 molecules per sec; energization = electrochemical potential gradient; protein types mediating the passive transport = channels; examples = K, B, Si, Cl, Ca channels).
Phytoavailability	Plant availability - i.e., elements are available to plants for uptake. Most commonly used when referring to essential plant nutrients that are in the right chemical form at sufficient concentrations in the soil solution.
Rhizosphere	The soil around living roots, which is influenced by root activity. The radial extension of the rhizosphere can range from sub- μ m to supra-cm scales.
Secondary active transport – cellular influx and efflux – short distance transport	Movement of an ion/molecule across the membrane against the concentration gradient of the transported molecule and/or against the electrochemical gradient (transport direction = uniport, symport, antiport; transport rates = 10^2-10^4 molecules per sec; energization = proton (H^+) gradient established by active transport processes; protein types mediating the active transport = carriers, which demands structural changes of the protein; examples = K, Na, Fe, Zn, Mg, B, Cl transporters).
Selectivity of a transporter	Each individual transport protein, irrespective of whether it is a channel (passive transporter), a carrier (a secondary active transporter) or an active transporter (pump) has a specific transport selectivity allowing for selective passage of (a) specific atom(s)/molecule(s). This transporter-intrinsic specificity does neither depend on the type of transport process (active vs. passive) nor on the transported atom/molecule (ionic versus polar molecule).
Short distance transport	Transport that occurs within, into and out of cells including a membrane crossing in the process. Includes diffusion and mass flow.
Symplastic (intracellular) transport	Cell-to-cell transport of an ion/molecule <i>via</i> plasmodesmata which has once entered the symplast <i>via</i> passive diffusion or a transport protein.
Transcellular (intracellular/extracellular) transport	Transport protein-mediated transport of an ion/molecule from one cell to the other, passing repeatedly from symplast to apoplast, independent of the water mass flow but depending on a non-overlapping localization of influx and efflux transporters in the plasma membrane to allow a directional flow.

Specific uptake and efflux transport proteins have been identified for mineral nutrients in the rhizodermis and root hairs at the plant–soil interface (Bienert et al., 2021), as well as in the cortex cells which have to be passed before nutrients enter the vasculature of the stele (**Figure 1B**). Although transport proteins have often developed a high selectivity for their substrate-of-interest, most are not fully element-specific and can accommodate a range of substrates simply due to physico-chemical similarities. The selectivity of transport proteins is mostly determined by 3D

physico-electrochemical properties of the amino acid residues constituting the surface of and the transport pathway through the protein (Chan and Shukla, 2021). While some transporters are ubiquitously expressed, others are only found at certain developmental stages, times of day, or only in response to stimuli (e.g., nutrient deficiency, mineral toxicity).

In the absence of knowledge on transport proteins, nutrient uptake fluxes have been historically differentiated into low- and high-affinity uptake systems (**Table 3**). Low-affinity uptake systems

act at higher phytoavailable nutrient concentrations while high-affinity uptake systems act at low phytoavailable nutrient concentrations. In addition, dual-affinity transport systems act under a wide concentration range of phytoavailable nutrient concentrations. The high-low-dual-affinity terminology is sometimes confusingly used as it is often extended to the binding affinity of a substrate to a transport protein that also determines nutrient uptake kinetics. However, underlying experimental transporter characteristics have not been determined and environmental parameters might determine transport rates by the respective protein (Dreyer and Michard, 2020).

Long-Distance Transport of Elements

From the root cortex, elements are transported into the stele and the vasculature system for long-distance, unidirectional transport toward the shoot (translocation, **Figure 1A**; **Table 3**). The xylem transports elements with the transpirational flow of water toward the leaves and generative tissues. Re-translocation of elements and bidirectional transport occurs in the phloem where high concentrations of assimilates (mainly sucrose) are loaded from source tissues (e.g., leaves) and are unloaded in sink tissues (e.g., fruits) by transport proteins. This generates a diffusion or osmotic gradient which induces a flow in the phloem and thereby long-distance transport of elements. The unloading of elements from the xylem and phloem as well as the distribution to physiological sink cells and organs again requires membrane transport proteins (see Cross-Membrane Transport of Elements). All these transport processes may cause isotope fractionation and change depending on physiological status, developmental stage, nutritional status, and the environmental condition at which a plant grows.

Binding of Elements to Organic Molecules

While elements are transported to their physiological destination they are exposed to a variety of non-transport related reactions. For instance, B readily forms ester bonds in many metabolites of primary and secondary metabolism while Cu can be retained in prosthetic bonds or as a cofactor in enzymes (Brđar-Jokanović, 2020; Li et al., 2020; Mir et al., 2021). Sorption, chelation, and complexation of elements to and with various organic and inorganic molecules is ubiquitous in plants. Some of these interactions are developmentally or physiologically targeted, others occur solely due to their chemical reactivity (Hanikenne et al., 2021). Noteworthy, in a quantitative sense, is the binding of, for example, Mg in chlorophyll, Fe in ferritin or heme, Zn in zinc finger proteins or nicotianamine, and trace metals to a plethora of thiols. In addition, the affinity of Ca to cell wall components contributes to the stability of cell walls (Thor, 2019).

METHODS FOR ISOTOPE ANALYSES OF PLANTS

Analyzing non-traditional stable isotope compositions of plants requires cautious sample handling, compared to traditional isotope analyses, and additional preparation steps, such as sample purification. In the following sections, the analytical procedures are listed chronologically.

Sample Preparation

For all post-harvest processes, such as plant sample cleaning, drying, and homogenization, sample contamination should be kept to a minimum since it can significantly bias isotope compositions. For guidelines on post-harvest sample preparation, we recommend Hansen et al. (2013). A crucial step for isotope analyses in plants is the cleaning of the roots of hydroponically and soil grown plants. To remove soil particles, weak salt solutions, ultrapure, and ice-cold water (18.2 MΩ.cm) are used (Tang et al., 2016). Salt solutions or weak acids can be further used to extract the apoplastic root pool prior to isotope ratio analysis (e.g., Tang et al., 2016; Garnier et al., 2017). However, both the washing procedure and the extractant used may significantly impact the isotope ratios of the different root pools.

Wet Chemistry

Wet chemistry sample preparation comprises two steps: sample digestion and analyte purification. To minimize sample contamination, these steps need to be conducted in “clean labs” (i.e., laboratories with non-metal fixtures, filtered air, ideally certified), ultra-clean reagents (e.g., distilled acids), and acid-cleaned PTFE/PFA containers (see also Section Quality Control). For most elements, digestion of plants requires strong acids (e.g., HNO₃) and a microwave oven or a hotplate (**Table 1**). Silicon analyses represent a special case, where alkaline fusion procedures are more common than acid digestions (Frick et al., 2020). After digestion, the samples are “purified” by anion/cation exchange and extraction chromatography (**Table 1**; **Supplementary Table S1**) which is the most common technique to separate pure analyte fractions. Often more than one of these “column chemistry” procedures are required to separate residual matrix elements. It is important that the chromatographic yields are close to 100%, as any loss of element on the column can result in isotope fractionation and thereby bias results. For example, Cu isotopes can significantly fractionate between the first and last milliliter of elution from a commonly used column chemistry, and even not collecting the final 5% of Cu can induce a significant isotope fractionation (Maréchal and Albarède, 2002). For some elements, quantitative yields are less important if the double spike technique is used (**Table 1**; **Supplementary Table S1**).

Isotope Analyses and Notation

To resolve differences in stable isotope composition in the per mil range, the use of multiple collector inductively coupled plasma mass spectrometry (MC-ICP-MS) or thermal ionization mass spectrometry (TIMS) is required (**Table 1**; Hoefs, 2018). Sample purification is necessary to avoid isobaric or poly-atomic/molecular interferences on the target element (**Table 1**). However, due to residual impurities or molecular interferences with elements present in the solution or the plasma (like N, O, H, Ar), it is often still challenging to perform precise isotope ratio analysis. Common problems include the formation of argides and those associated with analyzing isotopes that have very low natural abundances. For instance, ⁵⁶Fe overlaps with the argide ⁴⁰Ar¹⁶O. This problem is resolved by, for example, performing analyses at high-resolution modes, which is able to distinguish between Fe and argon oxides (Schoenberg and von Blanckenburg, 2005).

Mass bias during analyses needs to be corrected (Table 1). The simplest method adopted is “sample-standard bracketing” (Table 1). If a mass bias is non-linear, this method will not produce accurate results but can be improved by the addition of external normalization and mass fractionation laws (Table 1; Rehkämper et al., 2001). Such normalization, or “doping” uses the assumption that a dopant will undergo similar instrumental mass bias to the analyte, based on mass and ionization potential (e.g., Ni or Zn as doping element for Cu analyses). For elements with at least four stable isotopes, mass bias can be corrected by the double spike technique (Table 1; Rudge et al., 2009).

Stable isotope compositions are usually reported relative to an internationally recognized primary standard material and as a δ notation, exemplified for Ca (Hoefs, 2018):

$$\delta^{44/40}\text{Ca} (\text{‰}) = \left[\frac{\left(\frac{^{44}\text{Ca}}{^{40}\text{Ca}} \right)_{\text{sample}}}{\left(\frac{^{44}\text{Ca}}{^{40}\text{Ca}} \right)_{\text{standard}}} - 1 \right] 1000 \quad (1)$$

Values are most often presented in per mil, but are also occasionally given as ϵ (10,000 times instead of 1,000 times multiplication). The isotopes used for reporting the isotope composition and the internationally accepted primary reference standard of each reviewed element are listed in Supplementary Table S1. For several elements, it is conventional that only the heavier isotope is included in the δ notation (e.g., $\delta^{44/40}\text{Ca}$ is written as $\delta^{44}\text{Ca}$).

The isotope fractionation between two compartments of soil–plant system is expressed as:

$$\Delta^{44}\text{Ca}_{a-b} (\text{‰}) = \delta^{44}\text{Ca}_a - \delta^{44}\text{Ca}_b \quad (2)$$

In plants, Δ can refer to the isotope fractionation (i) between the whole plant, a, and its source, b, resulting in $\Delta^{44}\text{Ca}_{\text{plant-source}}$ to determine the isotope fractionation during the root uptake of an element or (ii) within a plant (e.g., $\Delta^{44}\text{Ca}_{\text{shoot-root}}$). In isotope studies where full mass balances are difficult to calculate (e.g., for trees), often $\Delta^{44}\text{Ca}_{\text{root-source}}$ is given. This notation is not equal to the isotope fractionation during uptake as the root isotope ratios result from both uptake and further transport of the element.

Quality Control

To assure the quality of the non-traditional stable isotope ratio measurements, two principle quality control measures are necessary. First, the concentration of a “procedural blank” (Table 1) needs to be determined to ensure that the unintended addition of an analyte, for example, Zn, from acids, plastics, and dust, to a sample is kept to a minimum. As a general rule, a procedural blank that contains less than 1% of the analyte mass in a sample creates no significant artificial effect on isotope ratios. Second, reference samples need to be analyzed to show that the sample preparation and analyses are robust and reproducible. The quality of the isotope ratio measurement is commonly reported using the reproducibility of the primary reference standards

(Supplementary Table S1). Furthermore, interlaboratory comparisons of certified reference materials (CRMs) enable the recovery of an element during sample digestion and purification to be calculated (Supplementary Table S2). There are published (although not certified) isotope ratios for some of these materials and we highly recommend their use, since they better match the composition of plant samples and enable the validation of the whole analytical process from sample preparation to isotope analyses.

ISOTOPE FRACTIONATION IN PLANTS

In this section, isotope fractionation in plants is reviewed for individual elements. Where element mass balances were available mass balances were available, isotope fractionation upon uptake by plants (e.g., $\Delta^{44}\text{Ca}_{\text{plant-source}}$) is reported, along with any within-plant isotope fractionation (e.g., $\Delta^{44}\text{Ca}_{\text{shoot-root}}$, $\Delta^{44}\text{Ca}_{\text{grains-leaves}}$). Interpretations on distinct isotope compositions of plants are reviewed when data-driven and well-substantiated hypotheses were provided. Elements that have been analyzed in exploratory studies, without detailed process tracing discussions (K, Mo, and Tl) are reviewed briefly at the end.

Boron

Only a few studies have systematically investigated B isotope fractionation in plants. As full mass balances were not available, only semi-quantitative information on isotope fractionation between source and plant ($\Delta^{11}\text{B}_{\text{plant-source}}$), and within the plant ($\Delta^{11}\text{B}_{\text{within,plant}}$) can be extracted from these data. Nevertheless, B isotope analyses in B sources, such as hydroponic nutrient solutions and plants indicated that $\Delta^{11}\text{B}_{\text{plant-source}}$, may differ between plant species or between cultivars of the same species, as demonstrated for different grapevine cultivars (Marentes et al., 1997; Coetzee et al., 2010; Figure 2). These differences are possibly linked to distinct B demand at similar B supply. In bell pepper, heavy B isotopes were preferentially transported from roots to leaves ($\Delta^{11}\text{B}_{\text{leaves-roots}}$ +0.8 to +27‰, Geilert et al., 2015, 2019). Within the shoots, leaves were isotopically heavier than stems and fruits were in most cases lighter than leaves.

A main driver of B isotope fractionation within the plant may be changes of the B species boric acid ($\text{B}(\text{OH})_3$) and borate ($\text{B}(\text{OH})_4^-$, Geilert et al., 2019; Figure 2). At equilibrium, boric acid ($\text{B}(\text{OH})_3$) is enriched in heavy isotopes compared to borate ($\text{B}(\text{OH})_4^-$, Zeebe, 2005). Boron is transported into the root cytoplasm under B-deficient growth conditions as boric acid by the so-called Nodulin26-like Intrinsic Protein (NIP)-type membrane channels (Yoshinari and Takano, 2017). In the cytoplasm, boric acid is partially transformed to borate, which is effluxed by borate transporter proteins (BORs) membrane transporters to the xylem (Parker and Boron, 2013). Hence, the transport of B from root-to-shoot may favor the transport of light B isotopes. However, this was not the case (Geilert et al., 2019), indicating that isotope fractionation between boric acid and borate was masked by other processes. The B species are also relevant to structural binding of B into cell walls. Borate is thought to function as the structural ester “bridge” to interlink Rhamnolacturonan-II monomers within




FIGURE 2 | A few studies have investigated B isotope fractionation in plants. Bell pepper has been most systematically investigated. (1) Due to lacking B mass balances, $\Delta^{11}\text{B}_{\text{plant-source}}$ is not known, but it can be semi-quantitatively shown that (2) shoots are enriched in heavy isotopes compared to roots. (3) Within plants, leaves are enriched in heavy B isotopes compared to stems while leaves can be heavier and lighter than reproductive organs. (A) At equilibrium, boric acid ($\text{B}(\text{OH})_3$) is isotopically heavier than borate ($\text{B}(\text{OH})_4^-$) which may contribute to the isotope fractionation between root and shoot. (B) The integration of borate into cell walls may cause the enrichment of heavy isotopes in leaves compared to stems.

the pectin fraction of primary cell walls (Yoshinari and Takano, 2017). Therefore, light B isotopes should be integrated into cell walls, if the integration itself does not fractionate B isotopes, while heavy B isotopes remain soluble and mobile (Geilert et al., 2015, 2019). Hence, the integration of B into cell walls could explain the enrichment of heavy isotopes in leaves compared to stems (Geilert et al., 2019).

Magnesium

Plants preferentially take up heavy isotopes from their Mg sources, such as hydroponic nutrient solutions (Black et al., 2008; Bolou-Bi et al., 2010; Wrobel et al., 2020), phlogopite (Bolou-Bi et al., 2010), soil pore water (Tipper et al., 2010; Kimmig et al., 2018), and phytoavailable soil Mg pools (Bolou-Bi et al., 2012; Opfergelt et al., 2014; Kimmig et al., 2018; Wang et al., 2020; **Figure 3**). In ryegrass and clover, root extracts (CaCl_2) of Mg revealed that heavier Mg is preferentially adsorbed to the apoplast (Bolou-Bi et al., 2010). This Mg adsorption likely contributes to the preferential uptake of heavy Mg isotopes into plants (Bolou-Bi et al., 2010; Wang et al., 2020). In wheat, $\Delta^{26}\text{Mg}_{\text{plant-source}}$ was more positive under low compared to regular Mg supply (Wang et al., 2020). This difference was ascribed to distinct cross-membrane transport systems. At low supply,




FIGURE 3 | Mg isotope fractionation in plants has been studied in trees and cereals. Exemplified for cereals: (1) Plants are enriched in heavy isotopes compared to the Mg source while (2) shoots tend to be enriched in light isotopes compared to roots. (3) Grains are enriched in heavy isotopes compared to stems and leaves. (A) Heavy Mg isotopes preferentially bind onto negatively charged surfaces in the root apoplast. (B) Binding of Mg to membrane transporters that are active at low Mg supply may induce a shift towards heavy isotopes during plant uptake and root-to-shoot transport compared to Mg transport at regular (or high) Mg supply. (C) It is assumed that the root contains a heavy Mg pool that is bound to organic ligands (Mg-L) while ionic Mg^{2+} is enriched in light isotopes that may be preferentially transported toward the shoots. (D) During grain filling, organic ligands that contain Mg degrade (e.g., chlorophyll) and release heavy Mg which may lead to a preferential re-translocation of heavy isotopes from senescent tissues.

active Mg-specific proteins facilitate Mg uptake, while at high supply passive channels facilitate Mg uptake. The former may shift the plant Mg toward a heavier isotope composition through binding to the membrane protein (Bolou-Bi et al., 2010; Wang et al., 2020).

Shoots of grasses and clover tended to be lighter than their roots (Black et al., 2008; Bolou-Bi et al., 2010; Gao et al., 2018; Wang et al., 2020; **Figure 3**). Bolou-Bi et al. (2010) ascribed the negative $\Delta^{26}\text{Mg}_{\text{shoot-root}}$ values to two different Mg pools in roots: ionic Mg (favoring light isotopes) and Mg bound to organic ligands (favoring heavy isotopes), such as ATP and proteins. Ionic Mg is thought to be preferentially transported from roots to shoots, leading to an enrichment of light isotopes in shoots compared to the roots. In wheat, the $\Delta^{26}\text{Mg}_{\text{shoot-root}}$ was more negative in the initial growth phases ($\Delta^{26}\text{Mg}_{\text{shoot-root}} -0.85\%$) than in the final growth stages ($\Delta^{26}\text{Mg}_{\text{shoot-root}} -0.25\%$) at regular Mg supply (Wang et al., 2020). The inverse was the case for wheat grown with low Mg supply

($\Delta^{26}\text{Mg}_{\text{shoot-root}}$ initial -0.25% , final -0.55%). Similar as for root uptake, the distinct $\Delta^{26}\text{Mg}_{\text{shoot-root}}$ was ascribed to distinct cross-membrane transport modes at low and high Mg supply.

Mg isotope ratios also vary between different organs of plant shoots (**Figure 3**). In wheat and rice, grains were heavier than the remaining shoot (Black et al., 2008; Gao et al., 2018). However, Mg in wheat ears was lighter than the shoots at regular Mg supply, but heavier than shoots at low Mg supply (Wang et al., 2020). In spruce trees, Mg in older needles was lighter than in young needles ($\Delta^{26}\text{Mg}_{\text{old-young}}$ of -0.30% , Bolou-Bi et al., 2012) while Mg in old leaves of sugar maple tended to be heavier than young leaves (maximum $\Delta^{26}\text{Mg}_{\text{old-young}}$ of 0.20% , Kimmig et al., 2018). These studies indicated that Mg isotope fractionation within shoots may be related to Mg storage during vegetative growth and the subsequent remobilization from these Mg pools during reproductive growth stages. The remobilization of Mg would require a dissociation of Mg from its major organic ligands, such as chlorophyll (Kleczkowski and Igamberdiev, 2021). Theoretical and experimental studies revealed that Mg bound to chlorophyll is isotopically heavier than the bulk leaf or ionic Mg (Black et al., 2007; Moynier and Fujii, 2017; Pokharel et al., 2018; Wrobel et al., 2020). Only one study found that Mg in chlorophyll is lighter than the bulk leaf Mg (Black et al., 2008). Based on these findings, it was suggested that isotopically distinguishable pools of Mg exist in leaves, such as a light ionic Mg pool and a heavy Mg pool stored in chlorophyll (Pokharel et al., 2018). The latter may become a phloem source of Mg during leaf senescence and thereby an export of the heavier chlorophyll-bound Mg pool toward phloem sinks. However, Mg pools other than ionic and chlorophyll-bound exist in leaves and Mg isotopes bound to different organic ligands may be enriched in heavy and light isotopes compared to aqueous ionic Mg (Schott et al., 2016).

Silicon

Most plants take up light Si isotopes from hydroponic nutrient solutions and phytoavailable soil pools (**Figure 4**). Frick et al. (2020) showed that light Si was preferentially taken up by Si-accumulating wheat and non-Si-accumulating tomato and mustard plants. To identify the role of membrane proteins on Si isotope fractionation, Sun et al. (2016a) reduced the biologically mediated Si uptake of rice plants by metabolic inhibitors and by cooling the hydroponic nutrient solution. These treatments reduced the Si uptake 3–4 times and induced a shift toward light isotope uptake compared to the control plants (-0.29%). In mutant rice plants with non-functional Si membrane channel proteins (Lsi1), the Si concentration in rice shoots was 25 times reduced while $\Delta^{30}\text{Si}$ in the shoots did not change (Köster et al., 2009). The enrichment of light isotopes in Si-accumulating and non-Si-accumulating plants was ascribed to diffusion during the membrane passage of Si with or without transport by membrane proteins, respectively (Frick et al., 2020). Note that Si is taken up as uncharged $\text{Si}(\text{OH})_4$ and can therefore diffuse through the cell membrane (Ma, 2015).

No consistent pattern for $\Delta^{30}\text{Si}_{\text{shoot-root}}$ has been found, for example, for the same plant species (e.g., rice, -0.23 to 1.3% , Ding et al., 2008; Sun et al., 2016a,b; **Figure 4**) or among




FIGURE 4 | Si isotope fractionation in plants exemplified for cereals. (1) Root uptake of Si leads in most cases to an enrichment of light isotopes while (2) no consistent pattern of Si isotope fractionation has been observed between root and shoot. (3) Within the shoot, plants become significantly enriched in heavy isotopes along the transpiration stream leading to a strong enrichment of heavy isotopes in husks and grains of rice. **(A)** The preferential uptake of light isotopes has been ascribed to a non-membrane protein-mediated transfer of Si (non-Si accumulators) and a membrane protein-mediated transfer of Si (Lsi1) in Si-accumulating plants. Both processes may be driven by Si diffusion that favors light Si isotopes. **(B)** Precipitation of light Si isotopes into phytoliths (SiO_2) results in an enrichment of heavy isotopes in the Si pool ($\text{H}_4^{30}\text{SiO}_4$) and thereby in an enrichment of heavy isotopes along the transpiration stream.

non-Si-accumulating plants (-0.37 to $+0.72\%$, Frick et al., 2020). The processes that may govern the isotope fractionation between root and shoot are the precipitation of Si as phytoliths in roots and xylem loading (Sun et al., 2016b; Frick et al., 2020). The precipitation of Si as phytoliths favors light isotopes (Sun et al., 2016b). For xylem loading, Si needs to cross a membrane either by membrane diffusion (favors light isotopes) or by the membrane transporter Lsi2. Unlike Lsi1, Lsi2 is located at the proximal side of root exodermis and endodermis, effluxes Si out of the cell and, as a carrier protein, requires energy which is provided by the electrochemical proton gradient (Ma, 2015). A knockout of Lsi2 in rice led to a shift toward light isotopes (of $\sim 0.5\%$) in the shoots compared to the wild-type plant (Köster et al., 2009) indicating that Lsi2 may favor heavy isotopes.


Shoot organs of rice including stems, leaves, and husks become systematically enriched in heavy isotopes along the transpiration stream (Ding et al., 2005, 2008; Sun et al., 2016a; **Figure 4**). Similar patterns were observed for different shoot organs in banana, bamboo, and cucumber (Opfergelt et al., 2006; Ding et al., 2008; Sun et al., 2016b) and within leaves of banana

and rice (Opfergelt et al., 2006; Ding et al., 2008). There, the lightest $\delta^{30}\text{Si}$ were found in petioles, followed by a successive enrichment of heavy Si toward leaf tips. The systematic Si isotope fractionation within shoots was explained by a successive precipitation of light isotopes into phytoliths (Dupuis et al., 2015). The precipitation likely leads to an enrichment of heavy isotopes in the soluble Si fraction that is then further transported along the transpiration stream. In the end points of the transpiration stream, such as rice husk and grains, only a small fraction of the total Si is stored but this fraction is most enriched in heavy isotopes (Sun et al., 2016b). This successive enrichment of heavy isotopes through Si precipitation lead to a Rayleigh-like fractionation (Sun et al., 2008, 2016b). Deviations from the Rayleigh fit could be attributed to membrane protein-facilitated transport (Lsi6) within the shoot (Sun et al., 2016b).

Calcium

In studies that focused on Ca cycling in catchments, Ca in trees (spruce, beech, pines) was lighter compared to its sources (Wiegand et al., 2005; Page et al., 2008; Cenki-Tok et al., 2009; Holmden and Bélanger, 2010; Bullen and Chadwick, 2016; Figure 5). These observations were confirmed in hydroponic studies on soybean, wheat, rice, and beans (Cobert et al., 2011; Schmitt et al., 2013; Christensen et al., 2018). In studies that provided full mass balances, Ca in beans and alpine plants was lighter than Ca in nutrient solutions or bulk soils, respectively ($\Delta^{44/40}\text{Ca}_{\text{plant-source}} -0.23$ to -1.03‰ , Hindshaw et al., 2013; Schmitt et al., 2013). The enrichment of light Ca isotopes during root uptake was assigned to preferential binding of light Ca to negatively charged surfaces, such as pectines in the root apoplast (Cobert et al., 2011; Schmitt et al., 2017, 2018). The fractionation between nutrient solution and beans could be modeled using a constant equilibrium fractionation factor between free ionic Ca and Ca sorbed to pectin groups ($\Delta^{44/40}\text{Ca}_{\text{pectine-ionic}} -0.12\text{‰}$, Schmitt et al., 2013). In addition, Ca channel-mediated membrane transport may fractionate Ca isotopes as the transport rate of Ca ions may be size specific. The effective size of Ca ions depends on the number of water molecules in the first hydration shell and less hydrated Ca ions tend to be isotopically light (Moynier and Fujii, 2017).

There is no systematic fractionation of Ca isotopes between roots and shoots of different species (Hindshaw et al., 2013; Schmitt et al., 2013; Christensen et al., 2018; Figure 5). By employing a sequential extraction procedure for roots, Schmitt et al. (2018) showed that 90% of Ca in the roots of beech trees stabilizes cell walls and membranes as Ca-pectate. This structural Ca was lighter than the water soluble Ca fraction, suggesting that light Ca isotopes were retained in the roots while heavy isotopes were transported to the shoot. Hindshaw et al. (2013) explained the retention of light Ca isotopes in roots with Ca precipitation as oxalate in the root cortex of alpine herb. The assumption that Ca oxalate is enriched in light Ca isotopes was confirmed by extraction of Ca oxalate from leaves (Cobert et al., 2011; Schmitt et al., 2018). However, no profound explanation exists yet for the favorable transport of light isotopes from roots to shoots in, for example, soybean (Christensen et al., 2018).



Leaves tended to be heavier than stems in cereals, beans, and alpine plants (Page et al., 2008; Cobert et al., 2011; Hindshaw et al., 2013; Schmitt et al., 2013, 2017; Christensen et al., 2018; Figure 5). This fractionation might be explained by successive removal of light Ca from the xylem sap into cell walls of the xylem tissues (Cobert et al., 2011; Schmitt et al., 2018). Therefore, heavy Ca is further transported along the transpiration stream toward the leaves. Furthermore, pods of beans were lighter than leaves but not distinguishable from stems at regular Ca supply (Cobert et al., 2011; Schmitt et al., 2013). However, the isotope fractionation between leaves and pods was negligible at low Ca supply which strongly suggests that different processes controlled the Ca transfer to the pods at distinct Ca supply. Despite the fact that Ca is usually considered as phloem immobile, Ca mass balances revealed that Ca can be re-translocated from roots and leaves of several plant species at low Ca supply (Maillard et al., 2015). Hence, the degree of Ca re-translocation in plants could determine the Ca isotope composition in phloem sinks and sources.


Iron

Plants preferentially take up heavy Fe from phytoavailable soil pools ($\Delta^{56}\text{Fe}_{\text{plant-phytoavailable}} -0.11$ to $+1.36\text{‰}$, Garnier et al., 2017;

Wu et al., 2019; Liu et al., 2019; Chen et al., 2021; **Figure 6**). Plant Fe acquisition from soils includes the dissolution of Fe(III) minerals and root membrane transport, but the Fe acquisition strategy differs among plant species (Connorton et al., 2017). Strategy I plants (e.g., tomato) are equipped with root surface reductases to reduce insoluble Fe(III) to Fe(II) prior to cross-membrane transport. Strategy II plants (e.g., oat) release phytosiderophores that mobilize Fe(III) from the mineral lattice. The Fe(III)-phytosiderophore complex is then transported across the plasma membrane. The isotope fractionation $\Delta^{56}\text{Fe}_{\text{plant-bulk.soil}}$ was 1.8‰ lighter in strategy I (e.g., bean) compared to strategy II (e.g., oat) plants (Guelke and von Blanckenburg, 2007). In strategy I plants, kinetically controlled reduction of Fe(III) to Fe(II) prior to membrane transport could yield large isotope fractionation (Johnson et al., 2020). In strategy II plants, Fe binding to chelating molecules released by the plant may induce comparably small equilibrium Fe isotope fractionation (Dideriksen et al., 2008). These results were confirmed in hydroponics where Fe was supplied as Fe(III)-EDTA (Guelke-Stelling and von Blanckenburg, 2012). The negative isotope fractionation during uptake was more pronounced in strategy I (bean) compared to strategy II (oat) plants. Strategy I plants may have reduced Fe(III)-EDTA to Fe(II) while no reduction was required during the ligand exchange from Fe(III)-EDTA to Fe(III)-phytosiderophores in

strategy II plants (Guelke-Stelling and von Blanckenburg, 2012; Liu et al., 2019). Hence, the absence and presence of a Fe reduction step might be the major factor that causes distinct Fe isotope fractionation in strategy I and II plants during Fe acquisition.

Several studies indicated that the distinct $\Delta^{56}\text{Fe}_{\text{plant-source}}$ values of strategy I and II plants are not universally applicable (Kiczka et al., 2010; Moynier et al., 2013; Liu et al., 2019; Wu et al., 2021; **Figure 6**). In field studies, the $\Delta^{56}\text{Fe}_{\text{plant-source}}$ values of strategy I and II plants differed less. This is probably due to less controlled environmental conditions in the field and/or the use of a variety of non-model plants for which Fe acquisition strategies are less known. In addition, there might be other processes than root surface reduction and chelation by phytosiderophores that cause Fe fractionation during plant acquisition. In paddy soil-rice systems, root plaque is formed in the oxidized rhizosphere where Fe precipitates as Fe(III) oxides in the root apoplast (Garnier et al., 2017; Chen et al., 2021). This precipitation induced a strong isotope fractionation between pore water and root plaque ($\Delta^{56}\text{Fe}_{\text{root.plaque-pore.water}} +1.41$ to $+2.24$ ‰). Moreover, Fe isotope composition of roots differed with and without Fe plaque. Similar findings were also reported for plants grown in aerated soils for different Fe pools in the roots of alpine plants (Kiczka et al., 2010). These results highlight that Fe (re-)precipitation processes in the rhizosphere and



apoplast need to be considered to approach the “true” $\Delta^{56}\text{Fe}_{\text{plant-source}}$ value.


Iron isotope fractionation between root and shoot ($\Delta^{56}\text{Fe}_{\text{shoot-root}}$) ranged from -0.60 to $+2.02\text{‰}$ (Kiczka et al., 2010; Guelke-Stelling and von Blanckenburg, 2012; Wu et al., 2019, 2021; Chen et al., 2021; **Figure 6**). There was no clear preference for light or heavy Fe isotope transport from root to shoot while different methodologies to separate Fe root pools limit the comparability of the $\Delta^{56}\text{Fe}_{\text{shoot-root}}$ data (see previous paragraph).

Fe isotopes fractionate significantly between different shoot organs (Guelke and Von Blanckenburg, 2007; Guelke-Stelling and von Blanckenburg, 2012; Arnold et al., 2015; Chen et al., 2021; **Figure 6**). $\Delta^{56}\text{Fe}_{\text{within.shoot}}$ differed between strategy I (e.g., beans) and II plants, similar as for $\Delta^{56}\text{Fe}_{\text{plant-source}}$ (e.g., oat, Guelke-Stelling and von Blanckenburg, 2012). Seeds of strategy I plants were lighter than the remaining shoot while no such fractionation occurred in strategy II plants or rice (Guelke and Von Blanckenburg, 2007; Guelke-Stelling and von Blanckenburg, 2012; Arnold et al., 2015; Chen et al., 2021). The preferential transport of light Fe isotopes into seeds in strategy I was ascribed to Fe reduction (Guelke-Stelling and von Blanckenburg, 2012). Within the leaf cells, heavier isotopes are thought to be preferentially stored as Fe(III) in the vacuoles or in proteins, such as ferritin. The remobilization of Fe requires a reduction of Fe(III) to Fe(II) for transport as, for example, Fe(II)-nicotianamine. Likewise, the negligible Fe isotope fractionation within shoots of strategy II plants indicates that Fe(III) is either quantitatively reduced to Fe(II) or such redox changes are absent.

Nickel

Plants that grew on soils with high Ni concentrations were isotopically lighter than phytoavailable soil pools ($\Delta^{60}\text{Ni}_{\text{plant-extract}}$ -0.51 to -0.06‰ , Estrade et al., 2015; Zelano et al., 2020; **Figure 7**). In hydroponics, light isotopes were also preferentially taken up from the nutrient solution ($\Delta^{60}\text{Ni}_{\text{plant-solution}}$ -0.90 to -0.07‰ , Deng et al., 2014). Nickel hyperaccumulator plants took up lighter isotopes compared to Ni non-hyperaccumulator plants (Deng et al., 2014). In hyperaccumulators, the Ni uptake may be kinetically controlled due to the high metal transport rate of low-affinity transporters that leads to depletion of Ni in the rhizosphere. The Ni flow from bulk solution toward the roots may be diffusion driven which favors light isotopes (Rodushkin et al., 2004). Besides, Ni speciation in the hydroponic nutrient solution may have induced isotope fractionation between the free ionic Ni^{2+} and the complexed Ni pool. Particularly at low Ni supply, only a small fraction of Ni was present as free ionic Ni, while most of the Ni was complexed to EDTA. The EDTA may have enriched the free ionic Ni fraction in light isotopes and thereby contributed to the uptake of light Ni isotopes in plants. Heavy Ni isotope complexation to organic ligands was corroborated by theoretical and experimental studies on Ni-citrate complexes (Fujii et al., 2014; Zelano et al., 2020).


The $\Delta^{60}\text{Ni}_{\text{shoot-root}}$ and $\Delta^{60}\text{Ni}_{\text{within.shoot}}$ values were positive, negative or zero, depending on the plant and growth conditions (Deng et al., 2014; Estrade et al., 2015; Ratié et al., 2019; Zelano



et al., 2020; **Figure 7**). In hydroponically grown plants, a much higher fraction of Ni was transported from roots to shoots in hyperaccumulators, compared to the non-hyperaccumulator. In these experiments, the isotope fractionation followed distinct patterns: hyperaccumulator plants had negative $\Delta^{60}\text{Ni}_{\text{shoot-root}}$ values, whereas those of non-hyperaccumulators were positive (Deng et al., 2014). Both $\Delta^{60}\text{Ni}_{\text{shoot-root}}$ and $\Delta^{60}\text{Ni}_{\text{leaves-stem}}$ varied with plant age and phenological stage of plants (Estrade et al., 2015; Zelano et al., 2020). The biochemical processes that generate the variation in $\Delta^{60}\text{Ni}_{\text{shoot-root}}$ and $\Delta^{60}\text{Ni}_{\text{leaves-stem}}$ are not well understood except that experimentally derived equilibrium fractionation factors as well as combined speciation and isotopes studies suggest that Ni complexation to organic ligands in plants is not a main driver of Ni isotope fractionation in plants (Zelano et al., 2018, 2020).

Copper

Cu isotope fractionation between the Cu source and plants ($\Delta^{65}\text{Cu}_{\text{root-source}}$) and the Cu source and the root can be positive, negative, or zero (Jouvin et al., 2012; Ryan et al., 2013; Li et al., 2016; Blotevogel et al., 2022; **Figure 8**). Results on Cu isotope fractionation during plant uptake are difficult to



compare as in most cases $\Delta^{65}\text{Cu}_{\text{roots-source}}$ values were reported instead of $\Delta^{65}\text{Cu}_{\text{plant-source}}$. Furthermore, plants were exposed to distinct Cu sources in hydroponics (e.g., ionic Cu^{2+} vs. chelated Cu) and Cu bound to the root apoplast was desorbed in one study prior to root analyzes (Ryan et al., 2013). Nonetheless, preferential uptake of light isotopes seems to be controlled by the reduction of Cu(II) to Cu(I) prior to root membrane transport into tomato (Jouvin et al., 2012; Ryan et al., 2013). The Cu reduction may explain why strategy I plants (according to their Fe acquisition strategy, see section Iron) take up lighter isotopes than strategy II plants. For the latter, the Cu reduction seems to be not a requirement for Cu uptake (Ryan et al., 2013). However, studies on yeast mutants showed that specific Cu membrane transporters favor light isotopes without a preceding Cu reduction indicating that the preferential uptake of light Cu isotopes is not solely controlled by Cu reduction (Cadiou et al., 2017). Moreover, distinct Fe supply neither affected $\Delta^{65}\text{Cu}_{\text{plant-source}}$ nor $\Delta^{65}\text{Cu}_{\text{roots-source}}$ (Jouvin et al., 2012; Ryan et al., 2013) while a higher Cu supply shifted $\Delta^{65}\text{Cu}_{\text{roots-source}}$ from light to heavy in grapevine plants (Blotevogel et al.,

2022). This shift was assigned to a switch from active toward passive uptake pathways due to increased Cu availability. Together, the identified processes that control $\Delta^{65}\text{Cu}_{\text{plant-source}}$ are Cu reductase activity at the root surface, membrane transport, and adsorption to root surfaces that may all vary with Cu supply.

Copper isotope fractionation between root and shoot ($\Delta^{65}\text{Cu}_{\text{shoot-root}}$) can be positive, negative or zero (Weinstein et al., 2011; Jouvin et al., 2012; Ryan et al., 2013; Li et al., 2016; Blotevogel et al., 2022; **Figure 8**). Similarly, as for the $\Delta^{65}\text{Cu}_{\text{plants-source}}$ values, the comparison of $\Delta^{65}\text{Cu}_{\text{shoot-root}}$ values may be limited particularly due to different Cu root desorption strategies (Jouvin et al., 2012; Ryan et al., 2013). Enrichment of light isotopes in shoots compared to roots was ascribed to Cu diffusion and membrane transport that were categorized as kinetic fractionation processes (Weinstein et al., 2011; Jouvin et al., 2012). Light isotope enrichment in leaves was accentuated in grapevine at high Cu supply (Blotevogel et al., 2022). Exposed to high Cu concentrations and after root Cu desorption, $\Delta^{65}\text{Cu}_{\text{shoot-root}}$ was neutral in oat (strategy II) while tomato (strategy I) shoots were enriched in heavy isotopes compared to their roots (Ryan et al., 2013). The contrasting $\Delta^{65}\text{Cu}_{\text{shoot-root}}$ in Fe strategy I and II plants were ascribed to the reoxidation of Cu(I)-S to Cu(II) prior to xylem loading (Ryan et al., 2013). Similar to Fe, this reoxidation step may only occur in strategy I plants which results in a strong enrichment of heavy isotopes in the shoots compared to the roots. However, in strategy II plants, the reoxidation step might be absent or all Cu(I) is reoxidized to Cu(II).


Within plant shoots, an enrichment of light isotopes in leaves appears to increase with plant height in hairy-leaved sedge (Weinstein et al., 2011) and/or with Cu concentration in grapevine leaves (Blotevogel et al., 2019, 2022; **Figure 8**). The enrichment of light Cu isotopes was ascribed to diffusion and membrane transport (Weinstein et al., 2011) as well as Cu immobilization by Cu complexation to S (Cu(I)-S) that favors light isotopes (Cadiou et al., 2017; Blotevogel et al., 2019). However, detailed analyses of the Cu tolerant plant *E. splendens* revealed that redistribution of Cu from senescent to younger leaves and reproductive organs can cancel the correlations between leaf height and Cu concentration with Cu isotope ratios (Li et al., 2016). Ryan et al. (2013) found a more pronounced Cu isotope fractionation between stems and leaves in Fe strategy I than in II plants. Similarly to Fe, the stronger isotope fractionation in strategy I plants is likely driven by Cu redox cycles while these cycles may not control the Cu isotope fractionation in shoots of strategy II plants.

Zinc

A previously published review summarized that adsorption, type of membrane transport (low vs. high-affinity transport), speciation, compartmentalization, and diffusion control the Zn isotope fractionation in plants (Caldelas and Weiss, 2017). This conclusion was based on studies that investigated a diversity of plant species and applied methods, such as (i) Zn uptake studies with unicellular organisms

(Gélabert et al., 2006; John et al., 2007), (ii) root extraction techniques (Tang et al., 2016), (iii) elaborated hydroponic (e.g., Smolders et al., 2013) and pot studies (Houben et al., 2014; Couder et al., 2015), (iv) combined Zn speciation and isotope analyses (Aucour et al., 2015), and (v) a set of theoretically and experimentally determined isotope fractionation factors for Zn binding to organic ligands (Jouvin et al., 2012; Fujii et al., 2014; Marković et al., 2017).

After the review of Caldelas and Weiss (2017); a field study on paddy soil-rice systems showed more positive $\Delta^{66}\text{Zn}_{\text{rice-source}}$ values at low Zn than at regular Zn supply (Weiss et al., 2021; **Figure 9**). This shift toward heavy Zn isotopes at low Zn supply can be explained by chelators (e.g., the phytosiderophores 2-deoxymugineic acids) that are secreted by roots to strip Zn




from the soil matrix and subsequent uptake of the entire Zn-phytosiderophore complex by a membrane transporter (Smolders et al., 2013; Arnold et al., 2015; Weiss et al., 2021). Since 2-deoxymugineic acids bind heavy Zn isotopes at equilibrium (Marković et al., 2017), the uptake of Zn-phytosiderophore complexes may have induced the shift toward heavy isotopes. Furthermore, a unicellular organism study showed a dependence of $\Delta^{66}\text{Zn}_{\text{cell-source}}$ on Zn uptake rates (Köbberich and Vance, 2017). At low uptake rates, heavier Zn isotopes were taken up than at high uptake rates. This study is discussed in detail in section Heavy but More Mobile and the Question of Kinetic vs. Equilibrium Fractionation that focuses on unicell studies.

Changes to Zn isotope composition in the shoot of wheat ($\Delta^{66}\text{Zn}_{\text{within.shoot}}$) were measured during the grain filling period (Wiggenhauser et al., 2018). In this period, leaves and stems showed a net loss of Zn and a depletion of light isotopes, while grains became enriched in light isotopes compared to stems and leaves. These results strongly suggest that light Zn isotopes were transported within the phloem toward the grains while heavy isotopes were retained in senescing shoot organs. The retention of heavy Zn isotopes in senescing shoot organs was ascribed to Zn binding to O containing ligands in the apoplast, such as pectins (Aucour et al., 2017).

Cadmium

Plants were isotopically lighter than their phytoavailable Cd source, in hydroponic and soil studies (Wei et al., 2018; Imseng et al., 2019; Moore et al., 2020; Wiggenhauser et al., 2021a; Zhang et al., 2021; Zhong et al., 2021, 2022; **Figure 10**). An exception was that a hyperaccumulator plant grown on Cd contaminated soil had a heavier isotope composition than the phytoavailable pool (+0.02 to +0.18‰, Zhou et al., 2020). Two rice studies found that the enrichment in light isotopes was slightly enhanced in non-flooded compared to flooded soils (Wiggenhauser et al., 2021a; Zhong et al., 2022) while one found the opposite (Zhang et al., 2021). Possible factors causing this difference may be distinct initial soil properties (e.g., pH and soil Cd concentration) or changes in the phytoavailability of elements, such as Mn and Fe upon flooding. Several studies investigated the role of NRAMP5 on Cd isotope fractionation during uptake. In rice (*Oryza sativa*), both a small upregulation and knock-out of *OsNRAMP5* were associated with uptake of lighter Cd isotopes compared to control treatments (Zhang et al., 2021; Zhong et al., 2022). This apparent discrepancy is discussed in section Genetic Approaches for Isotope Fractionation Factors. Yeasts expressing cacao (*Theobroma cacao*, Tc) *TcNRAMP5* preferentially took up lighter Cd isotopes compared to the control yeast (i.e., with empty vector, Moore et al., 2020). In addition to membrane transport, light Cd uptake may be partly due to adsorption of preferentially light Cd onto the root apoplast ($\Delta^{114/110}\text{Cd}_{\text{root absorbed-adsorbed}}$ of -0.17‰, Zhang et al., 2021).

In most cases, shoots were isotopically heavier than roots in wheat, barley, rice, cacao, and a Cd accumulator plant (Wiggenhauser et al., 2016, 2021a,b; Wei et al., 2018; Imseng et al., 2019; Moore et al., 2020; Wang et al., 2021; Zhong et al., 2021; **Figure 10**). The retention of light Cd



isotopes in roots of cereals and numerous different cacao genotypes was ascribed to vacuolar sequestration. This explanation is supported by Rayleigh isotope fractionation models (Imseng et al., 2019; Moore et al., 2020). Furthermore, while fitting the same isotope model, substantial differences in translocation of Cd were observed for distinct cacao clones, which was ascribed to differences in the expression of genes that encode dominant Cd transporters, rather than distinct biochemical processes (Moore et al., 2020). The role of vacuolar sequestration on Cd isotope fractionation was further supported by results of wild-type mutant experiments on rice (Zhang et al., 2021). These results suggested that the tonoplast transporter heavy metal ATPase 3 (HMA3) preferentially transports light Cd isotopes into the vacuole. In the vacuole, Cd is then sequestered by strong binding to thiols that may contribute further to the retention of light Cd (Wei et al., 2018). This hypothesis is supported by synchrotron X-ray absorption spectroscopy (XAS) results that showed the majority of Cd in rice roots can be bound to S-containing ligands, such as glutathione or phytochelatin, which are expected to preferentially bind light Cd isotopes (Yan et al., 2016; Wiggenhauser et al., 2021a; Zhao et al., 2021). Alternatively, Zhong et al. (2022) showed that $\Delta^{114/110}\text{Cd}_{\text{shoot-root}}$ in rice can be further impacted

by the protein CAL1. This protein complexes Cd in xylem parenchyma cells and Cd-CAL1 complexes are then transported into the xylem. Furthermore, the higher expression of *CAL1* coincided with a strong shift toward light isotopes during root to shoot translocation.

Cd isotope composition in reproductive and shoot organs can strongly differ ($\Delta^{114/110}\text{Cd}_{\text{reproductive-shoot.organ}}$ -0.30 to +0.50‰, **Figure 10**). While Rayleigh fractionation modeling for different plants suggests dominantly unidirectional xylem flow from roots to shoots (Wiggenhauser et al., 2016; Wei et al., 2018; Moore et al., 2020), it has been shown that cereals likely use phloem redistribution on grain filling (Wiggenhauser et al., 2021b; Zhong et al., 2021). In rice, an important hub for the transfer of Cd from the xylem to the phloem are the nodes (Yamaji and Ma, 2014). First data on Cd isotopes in nodes indicated that the xylem to phloem transfer of Cd contributes to the enrichment of heavy isotopes in grains (Wiggenhauser et al., 2021b; Zhong et al., 2021). Additionally, combined isotope and gene expression analyses strongly suggested that membrane proteins (OsHMA2, OsLCT1) that transfer Cd from the xylem to the phloem in the node contribute to the enrichment of heavy Cd in grains (Zhong et al., 2021). The role of Cd speciation in the nodes (mainly Cd-S) on Cd isotope fractionation

between nodes, leaves, and grains is not understood yet (Wiggenhauser et al., 2021b). In contrast to cereals, Cd loading into cacao beans may use a different mechanism (Barraza et al., 2019) since Cd in cacao beans was found to be isotopically lighter than in the leaves. The identical isotope compositions between cacao leaves and leaf litter support this hypothesis, because isotope fractionation would be expected if there was phloem redistribution on senescence.

Emerging Elements

Potassium

Soybeans and grasses preferentially took up light isotopes from hydroponics ($\Delta^{41/39}\text{K}_{\text{plant-solution}}$ of -0.60%) and also from phytoavailable soil pools (Christensen et al., 2018; Li et al., 2021). $\Delta^{41/39}\text{K}_{\text{shoots-roots}}$ were reported to be positive and negative, however, full mass balances were not provided. Within shoots, dead leaves were isotopically lighter than living leaves (Li et al., 2021). During plant uptake and translocation, membrane transport may strongly control K isotope fractionation (Christensen et al., 2018). The selectivity of an ion channel is at least partly controlled by the ionic radius and could explain the preferential uptake of ^{39}K over ^{40}K due to the smaller ionic radius of ^{39}K (Lockless et al., 2007). Additionally, K transport through an ion channel requires dehydration which would favor ^{39}K for membrane transport (MacKinnon, 2003; Hofmann et al., 2012; Christensen et al., 2018). Within the plant, diffusion may only play a minor role for K isotope fractionation in moving saps, such as in the xylem and phloem (Christensen et al., 2018). Binding of heavy isotopes to pectate may be an additional factor that controls the K isotope fractionation in plants, as deduced from combined K isotope and XAS analysis (Li et al., 2021).

Molybdenum

The first Mo isotope analyses in plants revealed that in three out of four plant species (lingonberry, common juniper, and rosebay willowherb), heavy isotopes were preferentially transported from roots to stems ($\Delta^{98}\text{Mo}_{\text{stem-root}}$ -0.00 to $+0.70\%$) and from stems to leaves ($\Delta^{98}\text{Mo}_{\text{leaves-root}}$ -0.00 to $+0.40\%$), while in blueberry, Mo isotopes were not fractionated (Malinovsky and Kashulin, 2018). No processes have been yet ascribed to the systematic Mo isotope fractionation patterns.

Thallium

White mustard grown in hydroponics preferentially took up light isotopes ($\Delta^{205}\text{Tl}_{\text{plant-source}}$ -0.20 to -0.09%) while heavy isotopes were transported from roots to shoots, particularly at high Tl supply ($\Delta^{205}\text{Tl}_{\text{shoot-root}}$ -0.07 to $+0.72\%$, Vaněk et al., 2019). Within shoots, Tl isotope fractionation reached $\Delta^{205}\text{Tl}_{\text{shoot-root}}$ of 0.43% (Kersten et al., 2014; Rader et al., 2019; Vaněk et al., 2019). Although there is no known biological function for Tl, it has been posited that the isotope fractionation between roots and shoots is due to physiologically controlled translocation which may be related to membrane transport and Tl speciation. Since Tl^+ and K^+ have similar ionic radii (1.76Å and 1.60Å), it has also been suggested that Tl uses K channels to translocate

(Vaněk et al., 2019). In addition, Tl in plants appears to be mostly present as Tl(I)aq and Tl-acetate (O-ligand) in roots, stems, and leaves.

DISCUSSION

Linkages of biological and physico-chemical processes that are known to fractionate isotopes in plants, and highlights knowledge gaps that could be effectively filled, are summarized in **Supplementary Table S3**. In the next sections, we interlink the information collected for the individual elements in the results section to advance the use of non-traditional isotopes for process tracing in plants.

Similar Elements, Similar Isotope Fractionation?

Chemically similar elements can undergo similar or opposite isotope fractionation in plants. For example, for the redox sensitive transition metals Fe and Cu, reduction can be required prior to membrane transport (Cadiou et al., 2017; Johnson et al., 2020, **Figures 6; 8**). The reduction enriches the reduced Fe and Cu fractions in light isotopes and may largely control the shift toward light isotopes during acquisition in strategy I plants (Guelke-Stelling and von Blanckenburg, 2012; Ryan et al., 2013).

In contrast, for the non-redox sensitive transition metals Zn and Cd, the within-plant isotope fractionation differs as light Zn but heavy Cd isotopes are more mobile in plants (Arnold et al., 2015; Wiggenhauser et al., 2018; **Figures 9, 10**). Currently, two hypotheses are discussed that explain the opposing isotope fractionation between Zn and Cd. The first is that the distinct isotope fractionation is caused by the higher affinity of Cd to S donors of organic molecules (e.g., cysteine) compared to Zn, which in turn has a higher affinity to O and N donors (e.g., histidine) compared to Cd (Maret and Moulis, 2013). Thiol chelators (e.g., phytochelatin) can strongly bind Cd and contribute to the sequestration of light Cd isotopes in vacuoles (Nocito et al., 2011). Based on *ab initio* calculations, preferentially light Zn and Cd isotopes bind to S chelators at equilibrium (e.g., Fujii et al., 2014; Zhao et al., 2021). Given that Cd has a higher affinity to thiols than Zn, competition for thiol binding could control the opposing isotope fractionation of these trace metals in cereals. The second hypothesis is based on Cd sorption experiments on humic acid and *ab initio* calculations (Ratié et al., 2021; Zhao et al., 2021). Binding of ionic Cd to carboxyl groups (Cd-O) leads to a small shift toward light isotopes (Ratié et al., 2021). The isotope shift was ascribed to the dehydration of Cd from $\text{Cd}(\text{H}_2\text{O})_6$ to $\text{Cd}(\text{H}_2\text{O})_5$ prior to binding of Cd to carboxyl groups. The dehydration of Cd favors light isotopes while the opposite is the case for Zn (Fujii et al., 2014; Zhao et al., 2021). Hence, the opposing isotope fractionation of Zn and Cd could be induced by dehydration processes that may take place prior to membrane transport or during ligand exchange and/or by the high affinity of Cd to thiols.

Similarly, the alkaline earth metals Ca and Mg fractionate in an opposite manner during plant uptake (**Figures 3, 5**).

This is partly due to preferential sorption of light Ca and heavy Mg isotopes onto negatively charged root surfaces (Bolou-Bi et al., 2010; Schmitt et al., 2017). Therefore, plants preferentially take up lighter Ca but heavier Mg isotopes compared to their Ca and Mg source. In a study on goethite, which like pectins provides negatively charged O donors, Mg had a tendency to form inner-sphere complexes, resulting in the loss of its water coordination shell. Calcium, in contrast, formed outer-sphere complexes, keeping its water coordination shell (Rahnamaie et al., 2006). Hence, the opposing isotope fractionation of Ca and Mg can likely be explained by a difference in the stability of their hydration sphere (Essington, 2015), as also hypothesized for Zn and Cd. The within-plant isotope fractionation of Ca and Mg does not clearly differ (Figures 3, 5). Hence, processes, such as structural binding to cell walls or oxalate precipitation for Ca, and cross-membrane transport and partitioning in functional molecules for Mg, may mask isotope effects induced by dehydration and sorption.

These examples illustrate that specific chemical properties can induce marked differences in isotope fractionation in plants, even if elements belong to the same group in the periodic table. Hence, the use of different elements to explain isotope fractionation patterns by analogy is limited.

Heavy but More Mobile and the Question of Kinetic vs. Equilibrium Fractionation

For elements, such as Mg, Si, and Cd, heavy isotopes are more mobile than light isotopes in plants (Figures 3, 4, 10). These findings contradict the established idea that biological processes in plants favor the transport of light isotopes due to enzymatic reactions that are kinetically controlled (Dawson et al., 2002). Preferential uptake of heavy Mg isotopes seems to be controlled by the sorption to root surfaces (Figure 4). In this case, equilibrium isotope fractionation seems to outcompete membrane transport, which may be kinetically controlled. Light Si isotopes precipitate in roots as phytoliths and light Cd isotopes are likely sequestered in vacuoles, leading to higher mobility of heavy isotopes within the plant (Figures 4, 10; Ding et al., 2009; Wiggenhauser et al., 2021a). Precipitation of Si in shoots is likely kinetically controlled because it takes place at the end of the transpiration stream, where water evaporation leads to constant supersaturation and new Si influx. In contrast, vacuolar sequestration may be caused by a mix of kinetic and equilibrium fractionation (membrane transport and Cd chelation). These examples illustrate that kinetically controlled processes do not necessarily lead to a higher mobility of light isotopes in plants.

The importance to distinguish between kinetic and equilibrium fractionation was highlighted for Zn uptake of marine diatoms. These unicell organisms preferentially took up heavier Zn isotopes at low supply but light isotopes at high Zn uptake rates (John et al., 2008; Köbberich and Vance, 2017). At high Zn uptake rates, the negative $\Delta^{66}\text{Zn}_{\text{organism-source}}$ was ascribed to kinetic effects during membrane transport and to Zn speciation in the nutrient solution. At low Zn uptake rates, a pseudo-equilibrium may have been established between the ionic Zn

in solution and the binding site of the membrane transporter, leading to a shift toward heavy isotopes (Jouvin et al., 2009; Fujii et al., 2014). These interpretations suggest that isotope fractionation of one biological process may not in all cases be kinetically or equilibrium controlled. Instead, there might be environmental conditions in which only kinetic, only equilibrium or both, determine the isotope fractionation in biological organisms. Deciphering whether isotope fractionation is kinetic or equilibrium is of high priority to advance process tracing by non-traditional isotope systems in plants.

The general rule that biological processes in plants favor the transport of light isotopes due to enzymatic reactions is based on studies on traditional isotopes, such as C, N, and S. For instance, the enzyme rubisco that catalyzes CO₂ fixation in plants induces a kinetically controlled enrichment of light C isotopes during C assimilation (O'Leary, 1993). Similarly, enzymatic processes involving N assimilation, such as nitrate reductase or glutamine synthetase, favor light N isotopes (Tcherkez, 2011). However, metals and metalloids act as cofactors for enzymes rather than being the target element of complex metabolism pathways as with C, N, and S. Yet, enzyme controlled processes can be crucial for the homeostasis of redox sensitive metals, such as Fe and Cu or the Mg-dechelatase mediated decomposition of chlorophyll (Connorton et al., 2017; Chen et al., 2018; Zandi et al., 2020). Hence, the impact of enzymatically controlled processes on the fractionation for non-traditional isotopes in plants might differ from traditional isotope systems.

Unicellular Organisms: A Suitable Model System for Isotope Studies?

Studies of unicellular organisms (unicell) have proven to be an effective tool to gain a detailed understanding of the biological uptake of nutrients and pollutants. Yeast mutants provided evidence that the reduction of Cu(II) to Cu(I), controlled by the reductase FRE1 and FRE2, leads to light Cu isotope enrichment during uptake (Cadiou et al., 2017). However, the mutants with dysfunctional reductases were still enriched in light isotopes, indicating that the high-affinity Cu transporters CTR1 and CTR3 preferentially transport light Cu isotopes. This illustrates that the negative $\Delta^{65}\text{Cu}_{\text{cell-source}}$ is driven by both reduction and membrane transport. The yeast study confirmed previously stated hypotheses that Cu reduction leads to a shift toward light isotopes during root uptake into strategy I plants, such as tomato (Jouvin et al., 2012; Ryan et al., 2013).

Isotope fractionation during uptake and compartmentalization of Cd was investigated using transgenic yeasts and *E. coli* (Horner et al., 2013; Moore et al., 2020). In yeast transformed with cacao (Tc) genes, expression of *TcNRAMP5* was associated with uptake of light Cd isotopes. These findings supported the hypothesis that the preferential binding of light Cd to the membrane transporter contributes to the overall enrichment of light Cd isotopes in cacao plants (Figure 10). Although the growth conditions were similar for both organisms, the magnitude of the fractionation was larger for the yeast suggesting that other transporters are contributing significantly to $\Delta_{\text{organism-source}}$ in cacao. Horner et al. (2013) measured Cd isotopes in cell

walls and other compartments of *E. coli*. The cell walls were enriched in light isotopes which was ascribed to Cd binding to cell wall thiols. Similarly to *E. coli*, Cd isotopes in cereal roots were also enriched in light isotopes possibly due to vacuolar sequestration and detoxification through binding of Cd to chelating thiols in root vacuoles of cereals (Wiggenhauser et al., 2021a).

Cyanobacteria were employed to determine Mg isotope fractionation during uptake and compartmentalization (Pokharel et al., 2018). The positive $\Delta_{\text{organism-source}}$ value for cyanobacteria was related to Mg binding to functional molecules, such as chlorophyll or ATP in the cells. This binding generated intracellular Mg pools that were enriched in heavy (bound Mg) and light Mg isotopes (ionic Mg). Mass balances showed that the light ionic Mg pool was likely transported out of the cell, leading to a net enrichment of heavy Mg isotopes in cyanobacterium and thereby to a positive $\Delta_{\text{organism-source}}$. This intracellular Mg isotope fractionation may explain negative $\text{Mg}^{26}\Delta_{\text{shoot-root}}$ found in plants (Figure 3) where Mg bound to functional molecules (heavy) is retained in roots while ionic Mg (light) is transported through the xylem into the shoot. Likewise, heavy Mg isotopes stored in functional molecules in leaves could be re-translocated from senescing plant organs *via* phloem toward reproductive plant organs. However, such a relation was not found in wheat sampled at different growth stages (Wang et al., 2020). Hence, Mg binding to functional molecules alone does not explain the changing isotope compositions in leaves, which may be affected by Mg import and export in leaves and/or by long-distance transport pathways *via* xylem and phloem.

Together, unicell studies can provide a powerful method to test specific hypotheses on potential fractionation mechanisms in plants and to identify specific isotope fractionation factors. However, results of unicell studies need to be carefully extrapolated to plants as plants have developed several strategies to acquire elements (e.g., regulation of several membrane transporters, root exudation) and are complex organisms that exchange elements between different cells, tissues, and organs.

Genetic Approaches for Isotope Fractionation Factors

A variety of genetic approaches have been used to determine isotope fractionation factors associated with individual membrane proteins. For example, three different techniques have been used to investigate how NRAMP5 impacts the fractionation of Cd during uptake in cacao and rice (unicell transgenics, gene knockout, and gene expression monitoring). A yeast experiment using a gene from cacao (TcNRAMP5) supported the hypothesis that TcNRAMP5 preferentially transports light Cd isotopes (see section Heavy but More Mobile and the Question of Kinetic vs. Equilibrium Fractionation) and thereby contributes to the enrichment of light isotopes in cacao plants. Conversely, the knockout experiment in rice found that rice without *OsNRAMP5* preferentially took up light Cd compared to its wild type (Zhang et al., 2021). Since plants have a suite of different membrane proteins that can also transport Cd, the result favoring the interpretation that *OsNRAMP5* takes up preferentially heavy Cd may reflect the upregulation of genes encoding other membrane

proteins that induce an even larger isotopic shift than *OsNRAMP5*. The two gene expression experiments found that *OsNRAMP5* in rice was moderately upregulated in non-flooded soil (Zhong et al., 2021, 2022). In Zhong et al. (2022), there was no significant difference in $\Delta^{114/110}\text{Cd}_{\text{rice-source}}$ between flooded and non-flooded conditions, while Zhong et al. (2021) found an enrichment in light isotopes in rice that grew on the non-flooded conditions where *OsNRAMP5* was upregulated. Although the experimental conditions and the soil-rice systems of these two experiments were identical, the causes of these distinct observations were not discussed. Nevertheless, the studies of Zhong et al. (2021, 2022) are in agreement with the transgenic experiment results, which strongly suggest that NRAMP5 can contribute to preferential uptake of light isotopes in plants. The comparisons show that complementary techniques (e.g., knockout and transgenics), at similar environmental conditions, are necessary to cross-validate results and interpretations. The distinct outcomes of the NRAMP5 studies highlight that carefully obtained isotope fractionation factors with controlled model systems may still be masked by environmental factors in complex soil-plant systems.

FUTURE PERSPECTIVES

Isotope fractionation in plants is usually very systematic and can differ, for example, among plant ages, species, and nutrient supply. Although biotic and abiotic factors induce systematic isotope fractionation in plants, its interpretation is often challenging and relies on basic knowledge in plant sciences. Here we provide recommendations to better exploit the information provided by varying isotope compositions in plants to further advance non-traditional isotope process tracing in plants. Once these recommendations are implemented, isotope process tracing could be applied to detect and quantify changes in uptake and translocation pathways in response to changing nutrient supply as suggested for Mg (Wang et al., 2020), Cu (Blotevogel et al., 2022), and Zn (Weiss et al., 2021).

Basic Requirements for Plant Isotope Experiments

An indispensable requirement for the interpretation of isotope fractionation patterns in plants is the establishment of fractionation factors for specific physico-chemical processes. This requires experiments to quantify isotope fractionation for individual processes, such as precipitation (e.g., Guinoiseau et al., 2018) or binding to organic ligands (e.g., Ratié et al., 2021). In addition, there is a set of minimum requirements that must be taken into account to ensure a meaningful data set and discussion.

Planning phase:

- design experiments to test specific hypotheses and determine (co-)variables. These can range from environmental conditions, such as (i) pH (Cobert et al., 2011), (ii) varying nutrient sources (Rader et al., 2019), or (iii) nutrient limitations (Smolders et al., 2013), to biological changes, such as wild-type-mutant experiments (Zhang et al., 2021),

- include biological replicates to account for experimental variability (Wang et al., 2020),
- conduct plant growth experiments at physiologically relevant conditions to ensure nutrient levels are not additional variables impacting the isotope signatures of the targeted element,
- thoroughly document growth conditions (e.g., nutrient levels, light, temperature, plant protection regime), specific tissue type, and phenological growth stages at harvest (Meier, 2001).

Sample preparation:

- consider root treatments at harvest, such as root desorption by weak salt solutions (Tang et al., 2016). This separates apoplastic and symplastic element pools and enables the determination of isotope fractionation for apoplastic sorption and cross-membrane transport,
- take into account that the phytoavailability of elements and isotope composition in bulk and rhizosphere soil can be very different (e.g., Imseng et al., 2019).

Data analysis and evaluation:

- calculate the average concentration and isotopic mass balance of target elements. This is needed to elucidate fractionation upon root uptake from growth media (the $\Delta_{\text{root-solution}}$ does not inform on the fractionation during root uptake),
- once isotope compositions in different plant organs are measured, geochemical modeling approaches (e.g., Moore et al., 2020) can be used and incorporated with existing plant science knowledge to interpret what biological and physico-chemical processes are responsible for the systematic isotope fractionation patterns,
- analyze concentrations of other elements that may compete with the element of interest for, for example, membrane transporter and organic ligands.

Innovative Research Ideas and Methods

Provided that the requirements that are listed in section Basic Requirements for Plant Isotope Experiments are met, the following ideas can be included to expand the scope of plant stable isotope analyses. The ideas were selected from the individual element reviews and studies beyond those on stable isotope fractionation in plants.

- Isotope analyses in different plant organs and tissues (e.g., cortex, epidermis, rhizodermis) that are sampled at different growth stages (e.g., flowering, plant maturity) can provide useful information on internal transport processes and pathways of elements within plants (e.g., Wang et al., 2020).
- Measuring the isotope composition of different elements in the soil–plant system (e.g., Wiggenhauser et al., 2018).
- Further compartmentalization, for example, analysis of xylem sap, would provide complementary information to $\Delta_{\text{shoot-root}}$ and $\Delta_{\text{shoot-grain/bean}}$ values to understand the processes that control the transfer of elements from the roots to shoot organs including reproductive organs (Álvarez-Fernández et al., 2014).

- To aid such detailed and comprehensive experiments, improvements to sample throughput should be made. The efficiency of sample purification could be improved by (i) automated ion chromatography or sample purification systems (Husson et al., 2015; Morgan et al., 2018; Kidder et al., 2020; Zhou et al., 2021) or (ii) using new generation MC-ICP-MS instruments that are equipped with high-resolution modes, quadrupole pre-mass filters or collision/reaction cells to reduce interferences (e.g., Christensen et al., 2018; Moynier et al., 2021).
- Wild-type mutant comparisons (e.g., Zhang et al., 2021) as well as monitoring the gene expression of plants in response to different environmental conditions (Zhong et al., 2021) can be used to investigate the role of specific proteins in element storage and transport. Determining the isotope fractionation factor for specific proteins using these techniques is challenging as, for example, compensatory mechanisms, such as the abundance alteration of other proteins, may contribute to the signal (Chang et al., 2020).
- Further constraint of isotope fractionation factors associated with individual transporters can be achieved by characterization of their transport features following heterologous expression in simple systems; such as unicellular organisms like yeasts (e.g., *Saccharomyces cerevisiae*, Cadiou et al., 2017; Moore et al., 2020), insect cells or frog eggs (*Xenopus oocytes*, Larsen et al., 2017), model plant species, or the reconstitution of purified transporters in defined lipid bilayers (Xie, 2008).
- Speciation analyses can provide complementary information to isotope analyses (e.g., Li et al., 2021). Synchrotron XAS provides information on the chemical environment of the target element in plant organs (e.g., bound to O, N, or S donors or specific molecules, such as citrate).
- Combining liquid chromatography with MC-ICP-MS. This has been used to measure Mg and Cu isotope ratios in chlorophyll and proteins (Larner et al., 2019; Wrobel et al., 2020). For structural elements, such as B and Ca, extraction techniques can provide crucial insights into isotope fractionation within plants (Schmitt et al., 2018).
- Analysis of organelles, such as vacuoles and cell walls, could improve the understanding of isotope fractionation on a subcellular level in plants (Horner et al., 2013; Song et al., 2014). Such approaches require small volumes of analytes that could be facilitated by bespoke sample introduction systems (Murphy et al., 2020).
- Coupling laser ablation with MC-ICP-MS could provide *in-situ* isotope ratio analyses at tissue level (Lobo et al., 2018). This method has recently been tested to measure Si isotope ratios of phytoliths in leaves (Frick et al., 2019) and can potentially be applied for other elements.
- Finally, it would be interesting to combine isotope analyses for traditional and non-traditional elements. Nitrogen isotopes may be highly complementary to non-traditional isotopes (e.g., Mg due to its role in chlorophyll, Laursen et al., 2013). Another example is C isotope data, which can provide information on water use efficiency (Newton, 2016). Linking, for example, C and K isotope fractionation may also be highly relevant due to the role of K in controlling stomatal conductance.

CONCLUSION

This review illustrates that non-traditional isotopes usually fractionate significantly and systematically in plants. The current knowledge suggests that fractionation is driven by: diffusion for Si uptake, root apoplast adsorption for Ca and Mg, membrane transport and chemical speciation for Zn and Cd, reduction of Fe and Cu, precipitation of elements into insoluble forms, such as phytoliths (Si) or oxalate (Ca), and structural binding to cell walls as shown for Ca and probably also B. These processes can induce similar (Cu, Fe) or distinct (Ca vs. Mg, Zn vs. Cd) isotope fractionation patterns for chemically similar elements. In addition, isotope compositions of the covered elements vary between plant species/varieties and in distinct environmental conditions. Hence, isotope process tracing in plants can potentially provide unique information on dominant biological and physico-chemical processes that control uptake and transport of elements in distinct soil-plant systems. For instance, it can be used to understand changes in uptake and translocation mechanisms that occur in response to changing nutrient supply, as shown for, for example, Mg, Zn, and Cu. Therefore, isotope process tracing could be a valuable tool to identify how plants cope with distinct environmental stresses and soil management strategies. Further exploitation of the scope of isotope process tracing requires the joint expertise of plant scientists and geochemists to develop hypothesis-driven experimental designs for plant growth trials, conduct robust and efficient isotope analyses, employ biological model systems, integrate contrasting environmental conditions, and to ultimately link isotope fractionation in plants to physiological processes. This expertise can only be covered by research collaborations that include (at least) plant and isotope geochemical scientists. Such interdisciplinary

research may have great potential to overcome current thinking boundaries and thereby advance isotope and plant physiological knowledge.

AUTHOR CONTRIBUTIONS

The interdisciplinary team of plant and geo scientists including MW, RM, GB, KHL, and SB contributed to conceptualization, methodology, investigation, data curation, writing of original draft, and visualization. PW contributed to writing and editing. MW was responsible for the project administration. All authors contributed to the article and approved the submitted version.

ACKNOWLEDGMENTS

We thank Vaněk Aleš, Christoph Cloquet, Daniel Frick, Xiao-Ming Liu, Dmitriy Malinovskiy, Gildas Ratié, Bei Wu, Nuria Basdediós Prieto, Franziska Stamm, and John Christensen for the technical inputs on isotope analyses. We thank Heinz Wiggenhauser for the grand support for the visualization and Meryl Meyer for editing and literature management.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fpls.2022.840941/full#supplementary-material>

REFERENCES

- Álvarez-Fernández, A., Di-az-Benito, P., AbadiA, A., López-Millán, A.-F., and AbadiA, J. (2014). Metal species involved in long distance metal transport in plants. *Front. Plant Sci.* 5:105. doi: 10.3389/fpls.2014.00105
- Andresen, E., Peiter, E., and Küpper, H. (2018). Trace metal metabolism in plants. *J. Exp. Bot.* 69, 909–954. doi: 10.1093/jxb/erx465
- Arnold, T., Markovic, T., Kirk, G. J. D., Schönbacher, M., Rehkämper, M., Zhao, F. J., et al. (2015). Iron and zinc isotope fractionation during uptake and translocation in rice (*Oryza sativa*) grown in oxic and anoxic soils. *C. R. Geosci.* 347, 397–404. doi: 10.1016/j.crte.2015.05.005
- Aucour, A. M., Bedell, J. P., Queyron, M., Magnin, V., Testemale, D., and Sarret, G. (2015). Dynamics of Zn in an urban wetland soil-plant system: coupling isotopic and EXAFS approaches. *Geochim. Cosmochim. Acta* 160, 55–69. doi: 10.1016/j.gca.2015.03.040
- Aucour, A. M., Bedell, J. P., Queyron, M., Tholé, R., Lamboux, A., and Sarret, G. (2017). Zn speciation and stable isotope fractionation in a contaminated urban wetland soil-*Typha latifolia* system. *Environ. Sci. Technol.* 51, 8350–8358. doi: 10.1021/acs.est.6b02734
- Barraza, F., Moore, R. E. T., Rehkämper, M., Schreck, E., Lefeuvre, G., Kreissig, K., et al. (2019). Cadmium isotope fractionation in the soil-cacao systems of Ecuador: a pilot field study. *RSC Adv.* 9, 34011–34022. doi: 10.1039/C9RA05516A
- Bienert, M. D., Werner, L. M., Wimmer, M. A., and Bienert, G. P. (2021). Root hairs: the villi of plants. *Biochem. Soc. Trans.* 49, 1133–1146. doi: 10.1042/BST20200716
- Bigeleisen, J. (1965). Chemistry of isotopes: isotope chemistry has opened new areas of chemical physics, geochemistry, and molecular biology. *Science* 147, 463–471. doi: 10.1126/science.147.3657.463
- Black, J. R., Epstein, E., Rains, W. D., Yin, Q. Z., and Casey, W. H. (2008). Magnesium-isotope fractionation during plant growth. *Environ. Sci. Technol.* 42, 7831–7836. doi: 10.1021/es8012722
- Black, J. R., Yin, Q. Z., Rustad, J. R., and Casey, W. H. (2007). Magnesium isotopic equilibrium in chlorophylls. *J. Am. Chem. Soc.* 129, 8690–8691. doi: 10.1021/ja072573i
- Blotevogel, S., Oliva, P., Denaix, L., Audry, S., Viers, J., and Schreck, E. (2022). Stable cu isotope ratios show changes in cu uptake and transport mechanisms in vitis vinifera due to high cu exposure. *Front. Plant Sci.* 12:755944. doi: 10.3389/fpls.2021.755944
- Blotevogel, S., Schreck, E., Audry, S., Saldi, G. D., Viers, J., Courjault-Radé, P., et al. (2019). Contribution of soil elemental contents and cu and Sr isotope ratios to the understanding of pedogenetic processes and mechanisms involved in the soil-to-grape transfer (soave vineyard, Italy). *Geoderma* 343, 72–85. doi: 10.1016/j.geoderma.2019.02.015
- Bolou-Bi, E. B., Poszwa, A., Leyval, C., and Vigier, N. (2010). Experimental determination of magnesium isotope fractionation during higher plant growth. *Geochim. Cosmochim. Acta* 74, 2523–2537. doi: 10.1016/j.gca.2010.02.010
- Bolou-Bi, E. B., Vigier, N., Poszwa, A., Boudot, J. P., and Dambrine, E. (2012). Effects of biogeochemical processes on magnesium isotope variations in a forested catchment in the Vosges Mountains (France). *Geochim. Cosmochim. Acta* 87, 341–355. doi: 10.1016/j.gca.2012.04.005
- Brdar-Jokanović, M. (2020). Boron toxicity and deficiency in agricultural plants. *Int. J. Mol. Sci.* 21:1424. doi: 10.3390/ijms21041424
- Bullen, T., and Chadwick, O. (2016). Ca, Sr and Ba stable isotopes reveal the fate of soil nutrients along a tropical climosequence in Hawaii. *Chem. Geol.* 422, 25–45. doi: 10.1016/j.chemgeo.2015.12.008

- Cadiou, J. L., Pichat, S., Bondanese, V. P., Soulard, A., Fujii, T., Albarède, F., et al. (2017). Copper transporters are responsible for copper isotopic fractionation in eukaryotic cells. *Sci. Rep.* 7, 1–10. doi: 10.1038/srep44533
- Caldelas, C., and Weiss, D. J. (2017). Zinc homeostasis and isotopic fractionation in plants: a review. *Plant Soil* 411, 17–46. doi: 10.1007/s11104-016-3146-0
- Centi-Tok, B., Chabaux, F., Lemarchand, D., Schmitt, A. D., Pierret, M. C., Viville, D., et al. (2009). The impact of water-rock interaction and vegetation on calcium isotope fractionation in soil- and stream waters of a small, forested catchment (the Strengbach case). *Geochim. Cosmochim. Acta* 73, 2215–2228. doi: 10.1016/j.gca.2009.01.023
- Chan, M. C., and Shukla, D. (2021). Markov state modeling of membrane transport proteins. *J. Struct. Biol.* 213:107800. doi: 10.1016/j.jsb.2021.107800
- Chang, J., Huang, S., Yamaji, N., Zhang, W., Ma, J. F., and Zhao, F. (2020). OSNRAMP1 transporter contributes to cadmium and manganese uptake in rice. *Plant Cell Environ.* 12:755944. doi: 10.1111/pce.13843
- Chen, G., Liu, T., Li, Y., Gao, T., Huang, F., Li, X., et al. (2021). New insight into iron biogeochemical cycling in soil-rice plant system using iron isotope fractionation. *Fundam. Res.* 1, 277–284. doi: 10.1016/j.fmre.2021.04.006
- Chen, Z. C., Peng, W. T., Li, J., and Liao, H. (2018). Functional dissection and transport mechanism of magnesium in plants. *Semin. Cell Dev. Biol.* 74, 142–152. doi: 10.1016/j.semcdb.2017.08.005
- Christensen, J. N., Qin, L., Brown, S. T., and Depaolo, D. J. (2018). Potassium and calcium isotopic fractionation by plants (soybean [*Glycine max*], rice [*Oryza sativa*], and wheat [*Triticum aestivum*]). *ACS Earth Space Chem.* 2, 745–752. doi: 10.1021/acsearthspacechem.8b00035
- Cobert, F., Schmitt, A.-D., Bourgeade, P., Labolle, F., Badot, P.-M., Chabaux, F., et al. (2011). Experimental identification of Ca isotopic fractionations in higher plants. *Geochim. Cosmochim. Acta* 75, 5467–5482. doi: 10.1016/j.gca.2011.06.032
- Coetzee, P. P., Greeff, L., and Vanhaecke, F. (2010). ICP-MS Measurement of ¹¹B/¹⁰B Isotope Ratios in Grapevine Leaves and the Investigation of Possible Boron Isotope Fractionation in Grapevine Plants. *South Afr. J. Enol. Vitic.* 32. doi: 10.21548/32-1-1363
- Connorton, J. M., Balk, J., and Rodríguez-Celma, J. (2017). Iron homeostasis in plants – a brief overview. *Metallomics* 9, 813–823. doi: 10.1039/C7MT00136C
- Couder, E., Mattioli, N., Drouet, T., Smolders, E., Delvaux, B., Iserentant, A., et al. (2015). Transpiration flow controls Zn transport in *Brassica napus* and *Lolium multiflorum* under toxic levels as evidenced from isotopic fractionation. *Compt. Rendus Geosci.* 347, 386–396. doi: 10.1016/j.crte.2015.05.004
- Dauphas, N., and Schauble, E. A. (2016). Mass Fractionation Laws, Mass-Independent Effects, and Isotopic Anomalies. *Annu. Rev. Earth Planet. Sci.* 44, 709–783. doi: 10.1146/annurev-earth-060115-012157
- Dawson, T. E., Mambelli, S., Plamboeck, A. H., Templer, P. H., and Tu, K. P. (2002). Stable isotopes in plant ecology. *Annu. Rev. Ecol. Syst.* 33, 507–559. doi: 10.1146/annurev.ecolsys.33.020602.095451
- de Bang, T. C., Husted, S., Laursen, K. H., Persson, D. P., and Schjoerring, J. K. (2021). The molecular-physiological functions of mineral macronutrients and their consequences for deficiency symptoms in plants. *New Phytol.* 229, 2446–2469. doi: 10.1111/nph.17074
- Deng, T. H. B., Cloquet, C., Tang, Y. T., Sterckeman, T., Echevarria, G., Estrade, N., et al. (2014). Nickel and zinc isotope fractionation in hyperaccumulating and nonaccumulating plants. *Environ. Sci. Technol.* 48, 11926–11933. doi: 10.1021/es5020955
- Dideriksen, K., Baker, J. A., and Stipp, S. L. S. (2008). Fe isotope fractionation between inorganic aqueous Fe(III) and a Fe siderophore complex. *Mineral. Mag.* 72, 313–316. doi: 10.1180/minmag.2008.072.1.313
- Ding, T. P., Ma, G. R., Shui, M. X., Wan, D. F., and Li, R. H. (2005). Silicon isotope study on rice plants from the Zhejiang province, China. *Chem. Geol.* 218, 41–50. doi: 10.1016/j.chemgeo.2005.01.018
- Ding, T. P., Zhou, J. X., Wan, D. F., Chen, Z. Y., Wang, C. Y., and Zhang, F. (2008). Silicon isotope fractionation in bamboo and its significance to the biogeochemical cycle of silicon. *Geochim. Cosmochim. Acta* 72, 1381–1395. doi: 10.1016/j.gca.2008.01.008
- Dreyer, I., and Michard, E. (2020). High- and Low-Affinity Transport in Plants From a Thermodynamic Point of View. *Front. Plant Sci.* 10:1797. doi: 10.3389/fpls.2019.01797
- Dupuis, R., Benoit, M., Nardin, E., and Méheut, M. (2015). Fractionation of silicon isotopes in liquids: The importance of configurational disorder. *Chem. Geol.* 396, 239–254. doi: 10.1016/j.chemgeo.2014.12.027
- Essington, M. E. (2015). *Soil and Water Chemistry: An Integrative Approach. Second edition.* Boca Raton, Florida: CRC Press, Taylor & Francis Group.
- Estrade, N., Cloquet, C., Echevarria, G., Sterckeman, T., Deng, T., Tang, Y. T., et al. (2015). Weathering and vegetation controls on nickel isotope fractionation in surface ultramafic environments (Albania). *Earth Planet. Sci. Lett.* 423, 24–35. doi: 10.1016/j.epsl.2015.04.018
- Falhof, J., Pedersen, J. T., Fuglsang, A. T., and Palmgren, M. (2016). Plasma membrane H⁺-ATPase regulation in the Center of Plant Physiology. *Mol. Plant* 9, 323–337. doi: 10.1016/j.molp.2015.11.002
- Frick, D. A., Remus, R., Sommer, M., Augustin, J., Kaczorek, D., and von Blanckenburg, F. (2020). Silicon uptake and isotope fractionation dynamics by crop species. *Biogeosciences* 17, 6475–6490. doi: 10.5194/bg-17-6475-2020
- Frick, D. A., Schuessler, J. A., Sommer, M., and von Blanckenburg, F. (2019). Laser ablation In situ silicon stable isotope analysis of phytoliths. *Geostand. Geoanal. Res.* 43, 77–91. doi: 10.1111/ggr.12243
- Fry, B. (2006). *Stable Isotope Ecology.* New York: Springer-Verlag.
- Fujii, T., Moynier, F., Blichert-Toft, J., and Albarède, F. (2014). Density functional theory estimation of isotope fractionation of Fe, Ni, Cu, and Zn among species relevant to geochemical and biological environments. *Geochim. Cosmochim. Acta* 140, 553–576. doi: 10.1016/j.gca.2014.05.051
- Gao, T., Ke, S., Wang, S.-J., Li, F., Liu, C., Lei, J., et al. (2018). Contrasting mg isotopic compositions between Fe-Mn nodules and surrounding soils: accumulation of light Mg isotopes by Mg-depleted clay minerals and Fe oxides. *Geochim. Cosmochim. Acta* 237, 205–222. doi: 10.1016/j.gca.2018.06.028
- Garnier, J., Garnier, J. M., Vieira, C. L., Akerman, A., Chmieleff, J., Ruiz, R. I., et al. (2017). Iron isotope fingerprints of redox and biogeochemical cycling in the soil-water-rice plant system of a paddy field. *Sci. Total Environ.* 574, 1622–1632. doi: 10.1016/j.scitotenv.2016.08.202
- Geilert, S., Vogl, J., and Rosner, M. (2015). Boron isotope fractionation in bell pepper. *Mass Spec. Purif. Tech.* 01, 1–6. doi: 10.4172/mso.1000101
- Geilert, S., Vogl, J., Rosner, M., and Eichert, T. (2019). Boron isotope variability related to boron speciation (change during uptake and transport) in bell pepper plants and SI traceable n(¹¹B)/n(¹⁰B) ratios for plant reference materials. *Rapid Commun. Mass Spectrom.* 33, 1137–1147. doi: 10.1002/rcm.8455
- Gélalbert, A., Pokrovsky, O. S., Viers, J., Schott, J., Boudou, A., and Feurtet-Mazel, A. (2006). Interaction between zinc and freshwater and marine diatom species: surface complexation and Zn isotope fractionation. *Geochim. Cosmochim. Acta* 70, 839–857. doi: 10.1016/j.gca.2005.10.026
- Geldner, N. (2013). Casparian strips. *Curr. Biol.* 23, R1025–R1026. doi: 10.1016/j.cub.2013.08.052
- Guelke, M., and Von Blanckenburg, F. (2007). Fractionation of stable iron isotopes in higher plants. *Environ. Sci. Technol.* 41, 1896–1901. doi: 10.1021/es062288j
- Guelke-Stelling, M., and von Blanckenburg, F. (2012). Fe isotope fractionation caused by translocation of iron during growth of bean and oat as models of strategy I and II plants. *Plant Soil* 352, 217–231. doi: 10.1007/s11104-011-0990-9
- Guinoiseau, D., Galer, S. J. G., and Abouchami, W. (2018). Effect of cadmium sulphide precipitation on the partitioning of Cd isotopes: Implications for the oceanic Cd cycle. *Earth Planet. Sci. Lett.* 498, 300–308. doi: 10.1016/j.epsl.2018.06.039
- Hanikenne, M., Esteves, S. M., Fanara, S., and Rouached, H. (2021). Coordinated homeostasis of essential mineral nutrients: a focus on iron. *J. Exp. Bot.* 72, 2136–2153. doi: 10.1093/jxb/eraa483
- Hansen, T. H., de Bang, T. C., Laursen, K. H., Pedas, P., Husted, S., and Schjoerring, J. K. (2013). “Multielement plant tissue analysis using ICP spectrometry,” in *Plant Mineral Nutrients.* ed. F. J. M. Maathuis (Totowa, NJ: Humana Press), 121–141. doi: 10.1007/978-1-62703-152-3_8
- Hindshaw, R. S., Reynolds, B. C., Wiederhold, J. G., Kiczka, M., Kretzschmar, R., and Bourdon, B. (2013). Calcium isotope fractionation in alpine plants. *Biogeochemistry* 112, 373–388. doi: 10.1007/s10533-012-9732-1
- Hoefs, J. (2018). *Stable Isotope Geochemistry.* Cham: Springer International Publishing.
- Hofmann, A. E., Bourg, I. C., and DePaolo, D. J. (2012). Ion desolvation as a mechanism for kinetic isotope fractionation in aqueous systems. *Proc. Natl. Acad. Sci.* 109, 18689–18694. doi: 10.1073/pnas.1208184109

- Holmden, C., and Bélanger, N. (2010). Ca isotope cycling in a forested ecosystem. *Geochim. Cosmochim. Acta* 74, 995–1015. doi: 10.1016/j.gca.2009.10.020
- Horner, T. J., Lee, R. B. Y., Henderson, G. M., and Rickaby, R. E. M. (2013). Nonspecific uptake and homeostasis drive the oceanic cadmium cycle. *Proc. Natl. Acad. Sci.* 110, 2500–2505. doi: 10.1073/pnas.1213857110
- Houben, D., Sonnet, P., Tricot, G., Mattioli, N., Couder, E., and Opfergelt, S. (2014). Impact of root-induced mobilization of zinc on stable Zn isotope variation in the soil-plant system. *Environ. Sci. Technol.* 48, 7866–7873. doi: 10.1021/es5002874
- Husson, J. M., Higgins, J. A., Maloof, A. C., and Schoene, B. (2015). Ca and Mg isotope constraints on the origin of Earth's deepest $\delta^{13}\text{C}$ excursion. *Geochim. Cosmochim. Acta* 160, 243–266. doi: 10.1016/j.gca.2015.03.012
- Husted, S., Persson, D. P., Laursen, K. H., Hansen, T. H., Pedas, P., Schiller, M., et al. (2011). Review: The role of atomic spectrometry in plant science. *J. Anal. Spectrom.* 26, 52–79. doi: 10.1039/C0JA00058B
- Imseng, M., Wiggenhauser, M., Keller, A., Müller, M., Rehkämper, M., Murphy, K., et al. (2019). Towards an understanding of the Cd isotope fractionation during transfer from the soil to the cereal grain. *Environ. Pollut.* 244, 834–844. doi: 10.1016/j.envpol.2018.09.149
- John, S. G., Geis, R. W., Saito, M. A., and Boyle, E. A. (2007). Zinc isotope fractionation during high-affinity and low-affinity zinc transport by the marine diatom *Thalassiosira oceanica*. *Limnol. Oceanogr.* 52, 2710–2714. doi: 10.4319/lo.2007.52.6.2710
- John, S. G., Rouxel, O. J., Craddock, P. R., Engwall, A. M., and Boyle, E. A. (2008). Zinc stable isotopes in seafloor hydrothermal vent fluids and chimneys. *Earth Planet. Sci. Lett.* 269, 17–28. doi: 10.1016/j.epsl.2007.12.011
- Johnson, C., Beard, B., and Weyer, S. (2020). “Fe isotope fractionation factors,” in *Iron Geochemistry: An Isotopic Perspective Advances in Isotope Geochemistry*. ed. J. Hoefs (Cham: Springer International Publishing), 39–84.
- Jouvin, D., Louvat, P., Juillot, F., Maréchal, C. N., and Benedetti, M. F. (2009). Zinc isotopic fractionation: why organic matters. *Environ. Sci. Technol.* 43, 5747–5754. doi: 10.1021/es803012e
- Jouvin, D., Weiss, D. J., Mason, T. F. M., Bravin, M. N., Louvat, P., Zhao, F., et al. (2012). Stable isotopes of Cu and Zn in higher plants: evidence for Cu reduction at the root surface and two conceptual models for isotopic fractionation processes. *Environ. Sci. Technol.* 46, 2652–2660. doi: 10.1021/es202587m
- Kersten, M., Xiao, T., Kreissig, K., Brett, A., Coles, B. J., and Rehkämper, M. (2014). Tracing anthropogenic thallium in soil using stable isotope compositions. *Environ. Sci. Technol.* 48, 9030–9036. doi: 10.1021/es501968d
- Kiczka, M., Wiederhold, J. G., Kraemer, S. M., Bourdon, B., and Kretzschmar, R. (2010). Iron isotope fractionation during Fe uptake and translocation in alpine plants. *Environ. Sci. Technol.* 44, 6144–6150. doi: 10.1021/es100863b
- Kidder, J. A., Voinot, A., Sullivan, K. V., Chipley, D., Valentino, M., Layton-Matthews, D., et al. (2020). Improved ion-exchange column chromatography for Cu purification from high-Na matrices and isotopic analysis by MC-ICPMS. *J. Anal. At. Spectrom.* 35, 776–783. doi: 10.1039/C9JA00359B
- Kimmig, S. R., Holmden, C., and Bélanger, N. (2018). Biogeochemical cycling of Mg and its isotopes in a sugar maple forest in Québec. *Geochim. Cosmochim. Acta* 230, 60–82. doi: 10.1016/j.gca.2018.03.020
- Kleczkowski, L. A., and Igamberdiev, A. U. (2021). Magnesium Signaling in Plants. *IJMS* 22:1159. doi: 10.3390/ijms22031159
- Köbberich, M., and Vance, D. (2017). Kinetic control on Zn isotope signatures recorded in marine diatoms. *Geochim. Cosmochim. Acta* 210, 97–113. doi: 10.1016/j.gca.2017.04.014
- Komárek, M., Ratič, G., Vaňková, Z., Šípková, A., and Chrástný, V. (2021). *Plant Micronutrient Use Efficiency*. Eds. H. Mohammad Anwar, K. Takehiro, J. B. David, T. Lam-Son Phan, and F. Toru. London: Elsevier.
- Köster, J. R., Bol, R., Leng, M. J., Parker, A. G., Sloane, H. J., and Ma, J. F. (2009). Effects of active silicon uptake by rice on ^{28}Si fractionation in various plant parts. *Rapid Commun. Mass Spectrom.* 23, 2398–2402. doi: 10.1002/rcm.3971
- Langowski, Ł., Růžička, K., Naramoto, S., Kleine-Vehn, J., and Friml, J. (2010). Trafficking to the outer polar domain defines the root-soil interface. *Curr. Biol.* 20, 904–908. doi: 10.1016/j.cub.2010.03.059
- Larner, F., McLean, C. A., Halliday, A. N., Roberts, B. R., Larner, F., McLean, C. A., et al. (2019). Copper isotope compositions of superoxide dismutase and Metallothionein from post-mortem human frontal cortex. *Inorganics* 7:86. doi: 10.3390/inorganics7070086
- Larsen, B., Xu, D., Halkier, B. A., and Nour-Eldin, H. H. (2017). Advances in methods for identification and characterization of plant transporter function. *J. Exp. Bot.* 68, 4045–4056. doi: 10.1093/jxb/erx140
- Laursen, K. H., Mihailova, A., Kelly, S. D., Epov, V. N., Bérail, S., Schjoerring, J. K., et al. (2013). Is it really organic? – multi-isotopic analysis as a tool to discriminate between organic and conventional plants. *Food Chem.* 141, 2812–2820. doi: 10.1016/j.foodchem.2013.05.068
- Li, W., Liu, X.-M., Hu, Y., Teng, F.-Z., Hu, Y.-F., and Chadwick, O. A. (2021). Potassium isotopic fractionation in a humid and an arid soil-plant system in Hawai'i. *Geoderma* 400:115219. doi: 10.1016/j.geoderma.2021.115219
- Li, S. Z., Zhu, X. K., Wu, L. H., and Luo, Y. M. (2016). Cu isotopic compositions in *Elsholtzia splendens*: influence of soil condition and growth period on Cu isotopic fractionation in plant tissue. *Chem. Geol.* 444, 49–58. doi: 10.1016/j.chemgeo.2016.09.036
- Li, S.-Z., Zhu, X.-K., Wu, L.-H., and Luo, Y.-M. (2020). Zinc, iron, and copper isotopic fractionation in *Elsholtzia splendens* Nakai: a study of elemental uptake and (re)translocation mechanisms. *J. Asian Earth Sci.* 192:104227. doi: 10.1016/j.jseaes.2020.104227
- Liu, C., Gao, T., Liu, Y., Liu, J., Li, F., Chen, Z., et al. (2019). Isotopic fingerprints indicate distinct strategies of Fe uptake in rice. *Chem. Geol.* 524, 323–328. doi: 10.1016/j.chemgeo.2019.07.002
- Lobo, L., Pereira, R., and Fernández, B. (2018). Opportunities and challenges of isotopic analysis by laser ablation ICP-MS in biological studies. *Trends Anal. Chem.* 105, 380–390. doi: 10.1016/j.trac.2018.05.020
- Lockless, S. W., Zhou, M., and MacKinnon, R. (2007). Structural and thermodynamic properties of selective ion binding in a K^+ channel. *PLoS Biol.* 5:e121. doi: 10.1371/journal.pbio.0050121
- Ma, J. F. (2015). A cooperative system of silicon transport in plants. *Trends Plant Sci.* 20:8. doi: 10.1016/j.tplants.2015.04.007
- Ma, J. F., and Yamaji, N. (2015). A cooperative system of silicon transport in plants. *Trends Plant Sci.* 20, 435–442. doi: 10.1016/j.tplants.2015.04.007
- MacKinnon, R. (2003). Potassium channels. *FEBS Lett.* 555, 62–65. doi: 10.1016/S0014-5793(03)01104-9
- Maillard, A., Diquélou, S., Billard, V., Lainé, P., Garnica, M., Prudent, M., et al. (2015). Leaf mineral nutrient remobilization during leaf senescence and modulation by nutrient deficiency. *Front. Plant Sci.* 6:317. doi: 10.3389/fpls.2015.00317
- Malinovsky, D., and Kashulin, N. A. (2018). Molybdenum isotope fractionation in plants measured by MC-ICPMS. *Anal. Methods* 10, 131–137. doi: 10.1039/C7AY02316B
- Maréchal, C., and Albarède, F. (2002). Ion-exchange fractionation of copper and zinc isotopes. *Geochim. Cosmochim. Acta* 66, 1499–1509. doi: 10.1016/S0016-7037(01)00815-8
- Maréchal, C. N., Télouk, P., and Albarède, F. (1999). Precise analysis of copper and zinc isotopic compositions by plasma-source mass spectrometry. *Chem. Geol.* 156, 251–273. doi: 10.1016/S0009-2541(98)00191-0
- Maret, W., and Moulis, J.-M. (2013). “The bioinorganic chemistry of cadmium in the context of its toxicity,” in *Cadmium: From Toxicity to Essentiality*. eds. A. Sigel, H. Sigel and R. K. Sigel (Dordrecht: Springer Netherlands), 1–29.
- Marentes, E., Vanderpool, R. A., and Shelp, B. J. (1997). Boron-isotope fractionation in plants. *Can. J. Plant Sci.* 77, 627–629. doi: 10.4141/P97-010
- Marković, T., Manzoor, S., Humphreys-Williams, E., Kirk, G. J. D., Vilar, R., and Weiss, D. J. (2017). Experimental determination of zinc isotope fractionation in complexes with the phyto siderophore 2'-deoxymugeneic acid (DMA) and its structural analogues, and implications for plant uptake mechanisms. *Environ. Sci. Technol.* 51, 98–107. doi: 10.1021/acs.est.6b00566
- Marschner, H., and Marschner, P. (2012). *Marschner's Mineral Nutrition of Higher Plants*. London; Waltham, MA: Academic Press.
- Meier, U. (2001). *Growth Stages of Mono and Dicotyledonous Plants: BBCH-Monograph*. Berlin, Boston: Blackwell Science.
- Mellman, I., and Nelson, W. J. (2008). Coordinated protein sorting, targeting and distribution in polarized cells. *Nat. Rev. Mol. Cell Biol.* 9, 833–845. doi: 10.1038/nrm2525

- Meychik, N., Nikolaeva, Y., and Kushunina, M. (2021). The significance of ion-exchange properties of plant root cell walls for nutrient and water uptake by plants. *Plant Physiol. Biochem.* 166, 140–147. doi: 10.1016/j.plaphy.2021.05.048
- Mir, A. R., Pichtel, J., and Hayat, S. (2021). Copper: uptake, toxicity and tolerance in plants and management of Cu-contaminated soil. *BioMetals* 34, 737–759. doi: 10.1007/s10534-021-00306-z
- Moore, R. E. T., Ullah, I., de Oliveira, V. H., Hammond, S. J., Strekopytov, S., Tibbett, M., et al. (2020). Cadmium isotope fractionation reveals genetic variation in Cd uptake and translocation by *Theobroma cacao* and role of natural resistance-associated macrophage protein 5 and heavy metal ATPase-family transporters. *Hortic. Res.* 7:71. doi: 10.1038/s41438-020-0292-6
- Morgan, L. E., Santiago Ramos, D. P., Davidheiser-Kroll, B., Faithfull, J., Lloyd, N. S., Ellam, R. M., et al. (2018). High-precision $^{41}\text{K}/^{39}\text{K}$ measurements by MC-ICP-MS indicate terrestrial variability of: $\delta^{41}\text{K}$. *J. Anal. At. Spectrom.* 33, 175–186. doi: 10.1039/C7JA00257B
- Moynier, F., and Fujii, T. (2017). Theoretical isotopic fractionation of magnesium between chlorophylls. *Sci. Rep.* 7:6973. doi: 10.1038/s41598-017-07305-6
- Moynier, F., Fujii, T., Wang, K., and Foriel, J. (2013). Ab initio calculations of the Fe(II) and Fe(III) isotopic effects in citrates, nicotianamine, and phyto siderophore, and new Fe isotopic measurements in higher plants. *C. R. Geosci.* 345, 230–240. doi: 10.1016/j.crte.2013.05.003
- Moynier, F., Hu, Y., Wang, K., Zhao, Y., Gérard, Y., Deng, Z., et al. (2021). Potassium isotopic composition of various samples using a dual-path collision cell-capable multiple-collector inductively coupled plasma mass spectrometer, Nu instruments Sapphire. *Chem. Geol.* 571:120144. doi: 10.1016/j.chemgeo.2021.120144
- Murphy, D. T., Allen, C. M., Ghidan, O., Dickson, A., Hu, W., Briggs, E., et al. (2020). Analysing Sr isotopes in low-Sr samples such as single insects with inductively coupled plasma tandem mass spectrometry using N_2O as a reaction gas for in-line Rb separation. *Rapid Commun. Mass Spectrom.* 34:e8604. doi: 10.1002/rcm.8604
- Newton, J. (2016). “Stable Isotopes as Tools in Ecological Research.” Chichester, UK: John Wiley & Sons, Ltd, 1–8.
- Noctio, F. F., Lancilli, C., Dendena, B., Lucchini, G., and Sacchi, G. A. (2011). Cadmium retention in rice roots is influenced by cadmium availability, chelation and translocation: cadmium retention in rice roots. *Plant Cell Environ.* 34, 994–1008. doi: 10.1111/j.1365-3040.2011.02299.x
- Novak, V., Adler, J., Husted, S., Fromberg, A., and Laursen, K. H. (2019). Authenticity testing of organically grown vegetables by stable isotope ratio analysis of oxygen in plant-derived sulphate. *Food Chem.* 291, 59–67. doi: 10.1016/j.foodchem.2019.03.125
- Novak, V., Khatri, P. K., and Laursen, K. H. (2021). The oxygen isotopic signature of soil- and plant-derived sulphate is controlled by fertilizer type and water source. *Plant Cell Environ.* 44, 203–215. doi: 10.1111/pce.13877
- O’Leary, M. H. (1993). Biochemical basis of carbon isotope fractionation. *Stable Isot. Plant Carbon-Water Relat.*, 19–28. doi: 10.1016/B978-0-08-091801-3.50009-X
- O’Leary, M. H., Madhavan, S., and Paneth, P. (1992). Physical and chemical basis of carbon isotope fractionation in plants. *Plant Cell Environ.* 15, 1099–1104. doi: 10.1111/j.1365-3040.1992.tb01660.x
- Opfergelt, S., Burton, K. W., Georg, R. B., West, A. J., Guicharnaud, R. A., Sigfusson, B., et al. (2014). Magnesium retention on the soil exchange complex controlling Mg isotope variations in soils, soil solutions and vegetation in volcanic soils, Iceland. *Geochim. Cosmochim. Acta* 125, 110–130. doi: 10.1016/j.gca.2013.09.036
- Opfergelt, S., Cardinal, D., Henriot, C., André, L., and Delvaux, B. (2006). Silicon isotope fractionation between plant parts in banana: In situ vs. in vitro. *J. Geochem. Explor.* 88, 224–227. doi: 10.1016/j.gexplo.2005.08.044
- Page, B. D., Bullen, T. D., and Mitchell, M. J. (2008). Influences of calcium availability and tree species on Ca isotope fractionation in soil and vegetation. *Biogeochemistry* 88, 1–13. doi: 10.1007/s10533-008-9188-5
- Parker, M. D., and Boron, W. F. (2013). The divergence, actions, roles, and relatives of sodium-coupled bicarbonate transporters. *Physiol. Rev.* 93, 803–959. doi: 10.1152/physrev.00023.2012
- Pilon-Smits, E. A., Quinn, C. F., Tapken, W., Malagoli, M., and Schiavon, M. (2009). Physiological functions of beneficial elements. *Curr. Opin. Plant Biol.* 12, 267–274. doi: 10.1016/j.pbi.2009.04.009
- Pokharel, R., Gerrits, R., Schuessler, J. A., Frings, P. J., Sobotka, R., Gorbushina, A. A., et al. (2018). Magnesium stable isotope fractionation on a cellular level explored by cyanobacteria and black fungi with implications for higher plants. *Environ. Sci. Technol.* 52, 12216–12224. doi: 10.1021/acs.est.8b02238
- Rader, S. T., Maier, R. M., Barton, M. D., and Mazdab, F. K. (2019). Uptake and fractionation of thallium by *Brassica juncea* in a geogenic thallium-amended substrate. *Environ. Sci. Technol.* 53, 2441–2449. doi: 10.1021/acs.est.8b06222
- Rahnemaie, R., Hiemstra, T., and van Riemsdijk, W. H. (2006). Inner- and outer-sphere complexation of ions at the goethite–solution interface. *J. Colloid Interface Sci.* 297, 379–388. doi: 10.1016/j.jcis.2005.11.003
- Ratié, G., Chrastný, V., Guinoiseau, D., Marsac, R., Vaňková, Z., and Komárek, M. (2021). Cadmium isotope fractionation during complexation with humic acid. *Environ. Sci. Technol.* 55, 7430–7444. doi: 10.1021/acs.est.1c00646
- Ratié, G., Quantin, C., Maia De Freitas, A., Echevarria, G., Ponzevera, E., and Garnier, J. (2019). The behavior of nickel isotopes at the biogeochemical interface between ultramafic soils and Ni accumulator species. *J. Geochem. Explor.* 196, 182–191. doi: 10.1016/j.gexplo.2018.10.008
- Rehkämper, M., Schönbächler, M., and Stirling, C. H. (2001). Multiple collector ICP-MS: introduction to instrumentation, measurement techniques and analytical capabilities. *Geostand. Newslett.* 25, 23–40. doi: 10.1111/j.1751-908X.2001.tb00785.x
- Rodushkin, I., Stenberg, A., André, H., Malinovsky, D., and Baxter, D. C. (2004). Isotopic fractionation during diffusion of transition metal ions in solution. *Anal. Chem.* 76, 2148–2151. doi: 10.1021/ac035296g
- Rudge, J. F., Reynolds, B. C., and Bourdon, B. (2009). The double spike toolbox. *Chem. Geol.* 265, 420–431. doi: 10.1016/j.chemgeo.2009.05.010
- Ryan, B. M., Kirby, J. K., Degryse, F., Harris, H., McLaughlin, M. J., and Scheiderich, K. (2013). Copper speciation and isotopic fractionation in plants: uptake and translocation mechanisms. *New Phytol.* 199, 367–378. doi: 10.1111/nph.12276
- Schmitt, A. D., Borrelli, N., Ertlen, D., Gangloff, S., Chabaux, F., and Osterrieth, M. (2018). Stable calcium isotope speciation and calcium oxalate production within beech tree (*Fagus sylvatica* L.) organs. *Biogeochemistry* 137, 197–217. doi: 10.1007/s10533-017-0411-0
- Schmitt, A. D., Cobert, F., Bourgeade, P., Ertlen, D., Labolle, F., Gangloff, S., et al. (2013). Calcium isotope fractionation during plant growth under a limited nutrient supply. *Geochim. Cosmochim. Acta* 110, 70–83. doi: 10.1016/j.gca.2013.02.002
- Schmitt, A. D., Gangloff, S., Labolle, F., Chabaux, F., and Stille, P. (2017). Calcium biogeochemical cycle at the beech tree-soil solution interface from the Strengbach CZO (NE France): insights from stable Ca and radiogenic Sr isotopes. *Geochim. Cosmochim. Acta* 213, 91–109. doi: 10.1016/j.gca.2017.06.039
- Schoenberg, R., and von Blanckenburg, F. (2005). An assessment of the accuracy of stable Fe isotope ratio measurements on samples with organic and inorganic matrices by high-resolution multicollector ICP-MS. *Int. J. Mass Spectrom.* 242, 257–272. doi: 10.1016/j.ijms.2004.11.025
- Schott, J., Mavromatis, V., Fujii, T., Pearce, C. R., and Oelkers, E. H. (2016). The control of carbonate mineral Mg isotope composition by aqueous speciation: theoretical and experimental modeling. *Chem. Geol.* 445, 120–134. doi: 10.1016/j.chemgeo.2016.03.011
- Smolders, E., Versieren, L., Shuofei, D., Mattielli, N., Weiss, D., Petrov, I., et al. (2013). Isotopic fractionation of Zn in tomato plants suggests the role of root exudates on Zn uptake. *Plant Soil* 370, 605–613. doi: 10.1007/s11104-013-1655-7
- Song, W.-Y., Mendoza-Cozatl, D. G., Lee, Y., Schroeder, J. I., Ahn, S.-N., Lee, H.-S., et al. (2014). Phytochelatin-metal(loid) transport into vacuoles shows different substrate preferences in barley and Arabidopsis: Phytochelatin-metal(loid) vacuolar transport in barley. *Plant Cell Environ.* 37, 1192–1201. doi: 10.1111/pce.12227
- Stein, W. D., and Litman, T. (eds.) (2015). “Front-matter,” in *Channels, Carriers, and Pumps. 2nd Edn.* (London: Elsevier), i–iii.
- Sun, L., Wu, L. H., Ding, T. P., and Tian, S. H. (2008). Silicon isotope fractionation in rice plants, an experimental study on rice growth under hydroponic conditions. *Plant Soil* 304, 291–300. doi: 10.1007/s11104-008-9552-1
- Sun, Y., Wu, L. H., and Li, X. Y. (2016a). Experimental determination of silicon isotope fractionation in rice. *PLoS One* 11:e0168970. doi: 10.1371/journal.pone.0168970

- Sun, Y., Wu, L., Li, X., Sun, L., Gao, J., and Ding, T. (2016b). Silicon isotope fractionation in rice and cucumber plants over a life cycle: laboratory studies at different external silicon concentrations. *J. Geophys. Res. Biogeosci.* 121, 2829–2841. doi: 10.1002/2016JG003443
- Taiz, L., Zeiger, E., Møller, I. M., and Murphy, A. (2015). *Plant Physiology and Development*. 6th Edn. Sunderland, Massachusetts.
- Tang, Y.-T., Cloquet, C., Deng, T.-H.-B., Sterckeman, T., Echevarria, G., Yang, W.-J., et al. (2016). Zinc isotope fractionation in the Hyperaccumulator *Noccaea caerulescens* and the nonaccumulating plant *Thlaspi arvense* at low and high Zn supply. *Environ. Sci. Technol.* 50, 8020–8027. doi: 10.1021/acs.est.6b00167
- Tang, R.-J., Luan, M., Wang, C., Lhamo, D., Yang, Y., Zhao, F.-G., et al. (2020). Plant membrane transport research in the post-genomic era. *Plant Commun.* 1:100013. doi: 10.1016/j.xplc.2019.100013
- Tcherkez, G. (2011). Natural $^{15}\text{N}/^{14}\text{N}$ isotope composition in C3 leaves: are enzymatic isotope effects informative for predicting the ^{15}N -abundance in key metabolites? *Funct. Plant Biol.* 38: 1. doi: 10.1071/FP10091
- Tcherkez, G., and Tea, I. (2013). $^{32}\text{S}/^{34}\text{S}$ S isotope fractionation in plant sulphur metabolism. *New Phytol.* 200, 44–53. doi: 10.1111/nph.12314
- Tejada-Jimenez, M., Alejandro, C.-A., Angel, L., Aurora, G., and Emilio, F. (2018). *Plant Micronutrient Use Efficiency*. eds. M. Anwar Hossain, T. Kamiya, D. J. Burritt, L.-S. Phan Tran and T. Fujiwara (London: Elsevier).
- Tipper, E. T., Gaillardet, J., Louvat, P., Capmas, F., and White, A. F. (2010). Mg isotope constraints on soil pore-fluid chemistry: evidence from Santa Cruz, California. *Geochim. Cosmochim. Acta* 74, 3883–3896. doi: 10.1016/j.gca.2010.04.021
- Thor, K. (2019). Calcium—Nutrient and Messenger. *Front. Plant Sci.* 10:440. doi: 10.3389/fpls.2019.00440
- Vaněk, A., Holubík, O., Oborná, V., Mihaljevič, M., Trubač, J., Ettler, V., et al. (2019). Thallium stable isotope fractionation in white mustard: implications for metal transfers and incorporation in plants. *J. Hazard. Mater.* 369, 521–527. doi: 10.1016/j.jhazmat.2019.02.060
- Wang, L., Jin, Y., Weiss, D. J., Schleicher, N. J., Wilcke, W., Wu, L., et al. (2021). Possible application of stable isotope compositions for the identification of metal sources in soil. *J. Hazard. Mater.* 407:124812. doi: 10.1016/j.jhazmat.2020.124812
- Wang, Y., and Wu, W.-H. (2017). Regulation of potassium transport and signaling in plants. *Curr. Opin. Plant Biol.* 39, 123–128. doi: 10.1016/j.cpb.2017.06.006
- Wang, Y., Wu, B., Berns, A. E., Xing, Y., Kuhn, A. J., and Amelung, W. (2020). Magnesium isotope fractionation reflects plant response to magnesium deficiency in magnesium uptake and allocation: a greenhouse study with wheat. *Plant Soil* 455, 93–105. doi: 10.1007/s11104-020-04604-2
- Wei, R., Guo, Q., Yu, G., Kong, J., Li, S., Song, Z., et al. (2018). Stable isotope fractionation during uptake and translocation of cadmium by tolerant *Ricinus communis* and hyperaccumulator *Solanum nigrum* as influenced by EDTA. *Environ. Pollut.* 236, 634–644. doi: 10.1016/j.envpol.2018.01.103
- Weinstein, C., Moynier, F., Wang, K., Paniello, R., Foriel, J., Catalano, J., et al. (2011). Isotopic fractionation of Cu in plants. *Chem. Geol.* 286, 266–271. doi: 10.1016/j.chemgeo.2011.05.010
- Weiss, D., Northover, G., Hanif, M., García-España, E., Vilar, R., Arnold, T., et al. (2021). Isotope fractionation of zinc in the paddy rice soil-water environment and the role of 2-deoxymugineic acid (DMA) as zincophore under Zn limiting conditions. *Chem. Geol.* 577:120271. doi: 10.1016/j.chemgeo.2021.120271
- Wiegand, B. A., Chadwick, O. A., Vitousek, P. M., and Wooden, J. L. (2005). Ca cycling and isotopic fluxes in forested ecosystems in Hawaii. *Geophys. Res. Lett.* 32, 1–4. doi: 10.1029/2005GL022746
- Wiggenhauser, M., Aucour, A.-M., Bureau, S., Campillo, S., Telouk, P., Romani, M., et al. (2021a). Cadmium transfer in contaminated soil-rice systems: insights from solid-state speciation analysis and stable isotope fractionation. *Environ. Pollut.* 269:115934. doi: 10.1016/j.envpol.2020.115934
- Wiggenhauser, M., Aucour, A.-M., Telouk, P., Blommaert, H., and Sarret, G. (2021b). Changes of cadmium storage forms and isotope ratios in Rice During grain filling. *Front. Plant Sci.* 12:645150. doi: 10.3389/fpls.2021.645150
- Wiggenhauser, M., Bigalke, M., Imseng, M., Keller, A., Archer, C., Wilcke, W., et al. (2018). Zinc isotope fractionation during grain filling of wheat and a comparison of zinc and cadmium isotope ratios in identical soil-plant systems. *New Phytol.* 219, 195–205. doi: 10.1111/nph.15146
- Wiggenhauser, M., Bigalke, M., Imseng, M., Müller, M., Keller, A., Murphy, K., et al. (2016). Cadmium isotope fractionation in soil-wheat systems. *Environ. Sci. Technol.* 50, 9223–9231. doi: 10.1021/acs.est.6b01568
- Wrobel, K., Karasiński, J., Tupys, A., Arroyo Negrete, M. A., Halicz, L., Wrobel, K., et al. (2020). Magnesium-isotope fractionation in chlorophyll-a extracted from two plants with different pathways of carbon fixation (C3, C4). *Molecules* 25:1644. doi: 10.3390/molecules25071644
- Wu, B., Amelung, W., Xing, Y., Bol, R., and Berns, A. E. (2019). Iron cycling and isotope fractionation in terrestrial ecosystems. *Earth Sci. Rev.* 190, 323–352. doi: 10.1016/j.earscirev.2018.12.012
- Wu, B., Wang, Y., Berns, A. E., Schweitzer, K., Bauke, S. L., Bol, R., et al. (2021). Iron isotope fractionation in soil and graminaceous crops after 100 years of liming in the long-term agricultural experimental site at Berlin-Dahlem, Germany. *Eur. J. Soil Sci.* 72, 289–299. doi: 10.1111/ejss.12944
- Xie, H. (2008). Activity assay of membrane transport proteins. *Acta Biochim. Biophys. Sin.* 40, 269–277. doi: 10.1111/j.1745-7270.2008.00400.x
- Yamaji, N., and Ma, J. F. (2014). The node, a hub for mineral nutrient distribution in graminaceous plants. *Trends Plant Sci.* 19, 556–563. doi: 10.1016/j.tplants.2014.05.007
- Yan, B.-F., Nguyen, C., Pokrovsky, O. S., Candaudap, F., Coriou, C., Bussièrre, S., et al. (2018). Contribution of remobilization to the loading of cadmium in durum wheat grains: impact of post-anthesis nitrogen supply. *Plant Soil* 424, 591–606. doi: 10.1007/s11104-018-3560-6
- Yan, J., Wang, P., Wang, P., Yang, M., Lian, X., Tang, Z., et al. (2016). A loss-of-function allele of OsHMA3 associated with high cadmium accumulation in shoots and grain of japonica rice cultivars. *Plant Cell Environ.* 39, 1941–1954. doi: 10.1111/pce.12747
- Yoshinari, A., and Takano, J. (2017). Insights into the mechanisms underlying boron homeostasis in plants. *Front. Plant Sci.* 8:1951. doi: 10.3389/fpls.2017.01951
- Zandi, P., Yang, J., Moždžen, K., and Barabasz-Krasny, B. (2020). A review of copper speciation and transformation in plant and soil/wetland systems. *Adv. Agron.* 160, 249–293. doi: 10.1016/bs.agron.2019.11.001
- Zeebe, R. E. (2005). Stable boron isotope fractionation between dissolved $\text{B}(\text{OH})_3$ and $\text{B}(\text{OH})_4^-$. *Geochim. Cosmochim. Acta* 69, 2753–2766. doi: 10.1016/j.gca.2004.12.011
- Zelano, I. O., Cloquet, C., Fraysse, F., Dong, S., Janot, N., Echevarria, G., et al. (2018). The influence of organic complexation on Ni isotopic fractionation and Ni recycling in the upper soil layers. *Chem. Geol.* 483, 47–55. doi: 10.1016/j.chemgeo.2018.02.023
- Zelano, I. O., Cloquet, C., van der Ent, A., Echevarria, G., Gley, R., Landrot, G., et al. (2020). Coupling nickel chemical speciation and isotope ratios to decipher nickel dynamics in the Rinorea cf. bengalensis-soil system in Malaysian Borneo. *Plant Soil* 454, 225–243. doi: 10.1007/s11104-020-04541-0
- Zhang, S. N., Gu, Y., Zhu, Z. L., Hu, S. H., Kopittke, P. M., Zhao, F. J., et al. (2021). Stable isotope fractionation of cadmium in the soil-rice-human continuum. *Sci. Total Environ.* 761:143262. doi: 10.1016/j.scitotenv.2020.143262
- Zhao, Y., Li, Y., Wiggenhauser, M., Yang, J., Sarret, G., Cheng, Q., et al. (2021). Theoretical isotope fractionation of cadmium during complexation with organic ligands. *Chem. Geol.* 571:120178. doi: 10.1016/j.chemgeo.2021.120178
- Zhong, S., Li, X., Li, F., Huang, Y., Liu, T., Yin, H., et al. (2022). Cadmium uptake and transport processes in rice revealed by stable isotope fractionation and Cd-related gene expression. *Sci. Total Environ.* 806:150633. doi: 10.1016/j.scitotenv.2021.150633
- Zhong, S., Li, X., Li, F., Liu, T., Huang, F., Yin, H., et al. (2021). Water management alters cadmium isotope fractionation between shoots and nodes/leaves in a soil-rice system. *Environ. Sci. Technol.* 55, 12902–12913. doi: 10.1021/acs.est.0c04713
- Zhou, F. Y., He, D., Miao, X., Yang, C., Dong, J. H., Zheng, H. T., et al. (2021). Development of an automatic column chromatography separation device for metal isotope analysis based on droplet counting. *Anal. Chem.* 93, 7196–7203. doi: 10.1021/acs.analchem.1c00145
- Zhou, J., Li, Z., Liu, M., Yu, H., Wu, L., Huang, F., et al. (2020). Cadmium isotopic fractionation in the soil-plant system during repeated phytoextraction

with a cadmium hyperaccumulating plant species. *Environ. Sci. Technol.* 54, 13598–13609. doi: 10.1021/acs.est.0c03142

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may

be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2022 Wiggenhauser, Moore, Wang, Bienert, Laursen and Blotevogel. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.