



OPEN ACCESS

EDITED AND REVIEWED BY
Isabel Nogues,
National Research Council (CNR), Italy

*CORRESPONDENCE
Peter Christian Morris
✉ p.c.morris@hw.ac.uk

RECEIVED 22 January 2024
ACCEPTED 30 January 2024
PUBLISHED 07 February 2024

CITATION
Morris PC, Reuveni M and Ortiz R (2024)
Editorial: Impact and mitigation of abiotic
stress in cereals.
Front. Plant Sci. 15:1374631.
doi: 10.3389/fpls.2024.1374631

COPYRIGHT
© 2024 Morris, Reuveni and Ortiz. This is an
open-access article distributed under the terms
of the [Creative Commons Attribution License
\(CC BY\)](#). The use, distribution or reproduction
in other forums is permitted, provided the
original author(s) and the copyright owner(s)
are credited and that the original publication
in this journal is cited, in accordance with
accepted academic practice. No use,
distribution or reproduction is permitted
which does not comply with these terms.

Editorial: Impact and mitigation of abiotic stress in cereals

Peter Christian Morris^{1*}, Moshe Reuveni² and Rodomiro Ortiz³

¹Institute of Life and Earth Sciences, Heriot-Watt University, Edinburgh, United Kingdom, ²Institute of Plant Sciences, Agricultural Research Organization, Volcani Center, Rishon LeZion, Israel, ³Department of Plant Breeding, Swedish University of Agricultural Sciences, Uppsala, Sweden

KEYWORDS

abiotic stress, drought, heat, salinity, cereals

Editorial on the Research Topic

Impact and mitigation of abiotic stress in cereals

Humanity has entered an unprecedented era of climatic change with the past decade showing the fastest rise of global temperature on record and 2023 being the warmest year in 174 years of observation ([Intergovernmental Panel on Climate Change \(IPCC\), 2023](#)). These facts bring into sharp focus the impact of abiotic stresses on food supplies. Human sustenance depends on daily caloric intake mainly by means of the consumption of proteins, fats and carbohydrates from various cellular sources. A single human needs to consume at least 2000 calories in food every day or about 730,000 a year ([Dietary guidelines.gov, 2020](#)). Current global food production (including that destined for animal feed) is about 2,900 calories per person per day ([FAO, 2022](#)). However, the projected increase in world population is to around 10 billion people in the next twenty to thirty years, and feeding the increasing populace is a primary concern for many governments and policymakers in agriculture. Food production for the expected 10 billion population by the year 2050 needs to supply at least 20¹² calories a day. Cereals in particular form a major component of global human diets with nearly half of calories directly derived from the seeds of these cultivated grasses ([Jones, 2023](#)), and even more from the use of cereals as animal feed and in the beverage industries. Abiotic stresses (for example, heat, drought) are exacerbated by climate change and have a negative impact on cereal crop quality and yield. Climate change is currently thought to be a major contributor to reductions in cereal yields worldwide ([Wang et al., 2018](#)). In order to provide the estimated 70 to 100% increase in cereals to feed an expanding world population ([Godfray et al., 2010](#)), it is imperative for plant scientists to better understand all aspects of abiotic stress on cereal crops, from perception and signal transduction through plant strategies for stress avoidance and mitigation to consider genetic resources and agronomic solutions to enhance production under stressful conditions ([Sato et al., 2024](#)).

The major abiotic factors that impact plant growth are heat, drought and salinity, which have distinct but overlapping effects on plant physiology but other abiotic factors (for example, anoxia and cold) are also of importance. In this Research Topic we explore some current advances in our understanding of how cereal crop plants can perceive, and how they respond to abiotic stresses, including both stress avoidance and stress tolerance. Strategies for mitigation through agronomic practice, and also longer-term breeding goals are considered.

The study of abiotic stress in crop plants covers a gamut of approaches, ranging from environmental research through physiological investigations and molecular and genetic aspects of plant responses to stresses. The Research Topic presented here reflect these different aspects of the science. The paper by [Bal et al.](#) analyses the impact of large-scale weather patterns across different environments on the yield of one rice cultivar over a four-year period. The results not only show how the climate (in particular, temperature and rainfall) impacts upon yield, but shows how the local environment at particular growth stages such as seedling establishment or seed filling can affect this crop.

The local crop environment can cause stress to plants through direct anthropogenic factors such as fertiliser regimes, and the paper by [Marti-Jerez et al.](#) has used remote satellite sensing together with ground-based assessments in a six-year longitudinal study to monitor the effect of mineral and organic fertilisers on rice performance in the field. This work shows that under the conditions used, mineral fertilisers gave a slight overall yield advantage, which was related to nitrogen deficiencies at later stages of growth when the organic fertilisers were used; however, organic fertilisers have the advantage of enhancing the available soil phosphorus and potassium levels. The satellite-based monitoring system showed promise for the detection of early signs of nutrient deficiencies which could then be addressed in a targeted manner.

The direct physiological effects of drought on crops have been well investigated by numerous studies, in particular crop water content ([Ievinsh, 2023](#)). The parameters defining water content have been more closely examined in the paper by [Yan et al.](#) who investigated the allocation of water and biomass within maize during periods of abiotic stress. The authors found differential retention of water under drought, which was reduced in roots and stems, and also of biomass which increased in roots and seeds. This work shows that simplistic whole-plant models of drought response are insufficient for a proper understanding of drought stress in crops, and that a more nuanced approach is required for a full understanding of water content fluctuations.

An understanding of the molecular mechanisms underlying stress tolerance is of critical importance when considering long term breeding strategies for stress resilient crops. The paper by [Xing et al.](#) investigates some fundamental mechanisms that regulate primary root growth under salt stress in maize. They find that the microRNA ZmmiR169q is a negative regulator of the transcription

factor ZmNF-YA8. Salt stress reduces ZmmiR169q expression and thus enhances ZmNF-YA8, which together with input from ethylene signalling, leads to the upregulation of ASA1 and ASA2, involved in tryptophan and auxin synthesis. High levels of auxin lead then to inhibition of primary root growth and arrested growth.

Bringing together environmental, physiological, and molecular research on stress responses of crops opens up the possibility of engineering metabolic pathways for stress tolerance. This is explored in the review paper by [Liu et al.](#) Their analysis suggests that selection for or modification of single gene traits in plants is unlikely to yield large improvements in stress tolerance. What may be more fruitful is an approach that targets multiple genes or loci that modify metabolic pathways with multiple outputs such as starch metabolism, phenylpropanoid metabolism or phytohormone signalling.

Taken together, this Research Topic illustrates the range and impact that plant scientists are having on addressing one of the most important problems to face the world in modern time; i.e., how to continue to feed the world's population in a sustainable manner.

Author contributions

PM: Writing original draft, Writing review & editing. MR: Writing review & editing. RO: Writing review & editing.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

References

- FAO. (2022). World Food and Agriculture – Statistical Yearbook 2022. (Rome). Available at: <https://doi.org/10.4060/cc2211en>.
- Godfray, H. C., Beddington, J. R., Crute, I. R., Haddad, L., Lawrence, D., Muir, J. F., et al. (2010). Food security: the challenge of feeding 9 billion people. *Science* 327, 812–818. doi: 10.1126/science.1185383
- Ievinsh, G. (2023). Water content of plant tissues: so simple that almost forgotten. *Plants (Basel)* 12, 1238. doi: 10.3390/plants12061238
- Intergovernmental Panel on Climate Change (IPCC) (2023). *Climate Change 2023: Climate Change 2023: Synthesis Report. A Report of the Intergovernmental Panel on Climate Change in: Contribution of Working Groups I, II and III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*. Eds. Core Writing Team., H. Lee and J. Romero (Geneva, Switzerland: IPCC). doi: 10.59327/IPCC/AR6-9789291691647.001
- Jones, J. M. (2023). "Role of cereals in nutrition and health," in *ICC Handbook of 21st Century Cereal Science and Technology* (Elsevier), 31–43.
- Sato, H., Mizoi, J., Shinozaki, K., and Yamaguchi-Shinozaki, K. (2024). Complex plant responses to drought and heat stress under climate change. *Plant J.* doi: 10.1111/tpj.16612
- U.S. Department of Agriculture and U.S. Department of Health and Human Services. (2022). *Dietary Guidelines for Americans, 2020-2025*. 9th Edition. December 2020. Available at: [DietaryGuidelines.gov](https://www.dietaryguidelines.gov).
- Wang, J., Vanga, S., Saxena, R., Orsat, V., and Raghavan, V. (2018). Effect of Climate change on the yield of cereal crops: A review. *Climate* 6, 41. doi: 10.3390/cli6020041