



# Investigating the Residual Effects of Chronic Cannabis Use and Abstinence on Verbal and Visuospatial Learning

Valentina Lorenzetti<sup>1\*</sup>, Michael Takagi<sup>2</sup>, Yvonne van Dalen<sup>3</sup>, Murat Yücel<sup>4</sup> and Nadia Solowij<sup>5</sup>

<sup>1</sup> Neuroscience of Addiction and Mental Health Program, Faculty of Health Sciences, Healthy Brain and Mind Research Centre, School of Behavioural and Health Sciences, Australian Catholic University, Melbourne, VIC, Australia, <sup>2</sup> Child Neuropsychology Unit, Murdoch Children's Research Institute, Melbourne, VIC, Australia, <sup>3</sup> Faculty of Science, University of Amsterdam, Amsterdam, Netherlands, <sup>4</sup> BrainPark, School of Psychological Sciences and Monash Biomedical Imaging Facility, The Turner Institute for Brain and Mental Health, Monash University, Melbourne, VIC, Australia, <sup>5</sup> School of Psychology and Illawarra Health and Medical Research Institute, University of Wollongong, Wollongong, NSW, Australia

## OPEN ACCESS

### Edited by:

Aviv M. Weinstein,  
Ariel University, Israel

### Reviewed by:

Anna Franceschini,  
Center for Addiction Medicine, Italy  
Delin Sun,  
Duke University, United States

### \*Correspondence:

Valentina Lorenzetti  
valentina.lorenzetti@gmail.com

### Specialty section:

This article was submitted to  
Addictive Disorders,  
a section of the journal  
Frontiers in Psychiatry

**Received:** 03 February 2021

**Accepted:** 10 May 2021

**Published:** 17 June 2021

### Citation:

Lorenzetti V, Takagi M, van Dalen Y,  
Yücel M and Solowij N (2021)  
Investigating the Residual Effects  
of Chronic Cannabis Use and  
Abstinence on Verbal and Visuospatial  
Learning.  
Front. Psychiatry 12:663701.  
doi: 10.3389/fpsy.2021.663701

**Rationale:** Regular cannabis users have been shown to differ from non-using controls in learning performance. It is unclear if these differences are specific to distinct domains of learning (verbal, visuospatial), exacerbate with extent of cannabis exposure and dissipate with sustained abstinence.

**Objective:** This study examines different domains of learning (verbal, visuospatial) in current and abstaining cannabis users, and the role of chronicity of use.

**Methods:** In a cross-sectional design, we examined 127 psychiatrically healthy participants (65 female) with mean aged of 34 years. Of these, 69 individuals were current regular cannabis users (mean 15 years use), 12 were former cannabis users abstinent for ~2.5 yrs (after a mean of 16 years use), and 46 were non-cannabis using controls. Groups were compared on verbal learning performance assessed via the California Verbal Learning Test (CVLT-II) and for visuospatial learning measured with the Brown Location Test (BLT). We explored the association between CVLT/BLT performance and cannabis use levels in current and former users.

**Results:** Current cannabis use compared to non-use was associated with worse performance on select aspects of verbal learning (*Long Delay Cued Recall*) and of visuospatial learning (*Retroactive Interference* and *LD Rotated Recall*). Prolonged abstinence was associated with altered verbal learning but intact visuospatial learning. There were non-significant correlations between distinct cannabis use measures, age and learning in both current and former users.

**Conclusions:** Our findings suggest cannabis use status (current use, former use) affects different domains of learning (verbal and visuospatial) in a distinct fashion. These findings might be accounted for in the design of cognitive interventions aimed to support abstinence in cannabis users.

**Keywords:** cannabis (marijuana), verbal learning and memory, visuospatial learning, abstinence, tetrahydrocannabinol, California Verbal Learning Test—Second Edition, Brown learning test

## INTRODUCTION

Cannabis is the most widely used illicit substance globally (1) and its potency has doubled over the past decade (2, 3). These statistics are concerning as a substantial proportion of cannabis users consume it on a regular basis (4, 5) and a significant minority of people with regular use experience lower school attainment, depression, anxiety, psychosis, impulse-control disorders, suicidal ideation and addiction (6). While there is much to learn about the cognitive correlates of regular cannabis use, a growing body of research has produced increasing knowledge in this area.

One of the core features of regular cannabis use entails alteration of cognitive performance that last beyond acute intoxication, which is thought to reflect chronic residual effects of regular cannabis use (7–12). In particular, altered cognitive function in regular cannabis users affects the domains of learning and memory, which are critical for performing daily tasks, at school and at work (7–13). Therefore, learning and memory alterations in regular cannabis users may underscore lower academic attainment (14, 15) and occupational performance found in cannabis using samples (16, 17).

Poorer learning and memory performance have been documented in regular cannabis users in both verbal and visuospatial domains. Specifically, relative to controls, regular cannabis users have shown lower verbal learning and recall of words (7–12), lower recall and accuracy of visuospatial performance in a checkerboard test (18), and lower retrieval in the virtual Morris water maze task (19). There are some inconsistencies however, as some of the examined samples have not shown verbal/visuospatial learning and memory alterations, or alterations with small to moderate effect sizes (10–12, 18–22). Replication studies in larger samples are required to validate and further examine the association between cannabis use and learning/memory performance, particularly for the visuospatial domain which has been examined by few studies to date.

The role of the extent of cannabis exposure on verbal/visuospatial learning and memory alterations is unclear. Evidence suggests that higher chronicity of use predicts worse verbal learning and memory performance (23–29). However, there is a lack of empirical studies that tested in detail how distinct measures of cannabis exposure affect verbal/visuospatial learning. Such measures include: dosage, duration, age of use onset, hours from last use, potency (9, 10, 12, 30) and cannabinoids such as the main psychoactive compound  $\Delta^9$ -tetrahydrocannabinol (THC) that determines cannabis potency (31–33). The lack of studies on how cannabis use patterns and THC levels affect learning performance in chronic users, creates a knowledge gap to inform users, educators, clinicians and policy makers about which measures and levels of cannabis exposure may be more harmful for verbal/visuospatial learning in chronic cannabis users.

Another issue yet to be elucidated is whether poorer verbal/visuospatial learning performance in regular cannabis users persists beyond prolonged abstinence, and the relevant evidence to date is mixed. Some studies show that learning deficits persist beyond abstinence [i.e., after 1 month (17, 30)].

Other studies found that learning alterations are attenuated (e.g., lower effect size) with longer abstinence (12) [e.g., over 4 to 8 weeks (34)]. Other studies show attenuated learning deficits in cannabis users who abstain for a variety of periods (11): 3 weeks (35), 1 month (36–40), 3 months (41)]; 12 months (35, 42–46). The inconsistency between study findings may be due to methodological confounds, such as long-lasting residual effects of chronic exposure e.g., cumulative lifetime exposure prior to quitting (30).

In sum, lower verbal learning performance has been (largely) consistently identified in chronic, long-term cannabis users. However, visuospatial learning deficits are largely unexplored, as well as the role of cannabis exposure levels and of prolonged abstinence.

The primary aim of this study was to address this evidence gap and to examine whether current cannabis use is associated with selective impairment of either verbal or visuospatial learning and memory. We also aimed to explore the role of extent of cannabis exposure and of prolonged abstinence on verbal/visuospatial learning and memory in chronic cannabis users.

To do this, we recruited 127 people (65 females), consisting of 69 current users and 12 former cannabis users and 46 non-using controls, comprehensively characterized for extent of substance use (alcohol, tobacco and other illicit drug use; cannabis use frequency, quantity, duration, age of onset, time from last cannabis use) and mental health (anxiety, depression and psychotic symptoms). Based on the existing evidence, we hypothesized that worse verbal and visuospatial learning performance would be apparent in cannabis users with more chronic levels of exposure.

## MATERIALS AND METHODS

### Participants

We recruited 127 people aged between 18 and 55 years via advertisements in local newspapers and Internet websites, and screened using a structured telephone interview to determine study eligibility. Participants included 69 current chronic regular cannabis users, 12 former chronic cannabis users and 46 non-using controls (henceforth called “current users,” “former users,” and “controls,” respectively). All groups were age matched.

### Inclusion and Exclusion Criteria

Cannabis users were included if they: (i) used cannabis at least twice a month for >2 years (the vast majority were currently using >3 days a week over many years, with a median of 30 smoking days/month and of 13 years of regular use); (ii) refrained from using substances other than cannabis, alcohol and tobacco in the month prior to assessment. Exclusion criteria for all participants were: (i) neurological disorders or serious head injury; (ii) Intelligence Quotient (IQ) <70; (iii) current regular use of illicit substances other than cannabis (amphetamines, benzodiazepines, cocaine, ecstasy, hallucinogens, inhalants and opiates; median lifetime use was between 0 and 6 occasions for any other drug).

All participants were requested to abstain from cannabis for at least 12 h prior to testing to enable examining cognitive

function in a non-intoxicated state, and provided written informed consent in accordance with local ethics committee guidelines. Ethics approval was given by the Mental Health Research and Ethics Committee (MHREC, I.D. number 459111), the Melbourne Health and North Western Mental Health (Melbourne, Australia), the Wollongong Human Research Ethics Committee (HREC, project number NSA07/03), and the ethics committee of the Murdoch Children's Research Institute (MCRI).

## Procedure

Participants underwent a comprehensive 2.5 h long assessment of mental health, substance use and cognitive function.

### Mental Health

We screened for psychiatric disorders through the Structured Clinical Interview for the Diagnostic and Statistical Manual for Mental Disorders IV-R (47), and assessed global functioning via the Global Assessment of Functioning module of the DSM-IV (48). We examined psychopathology symptoms of anxiety (State and Trait Anxiety Inventory, STAI; (49), depression and psychosis (Community Assessment of Psychic Experiences, CAPE (50).

### Substance Use

We assessed lifetime and past month substance use through semi-structured interviews, including the Substance Use History [Orygen Youth Health Research Centre, Melbourne, Australia (51–53)], a detailed structured assessment interview for cannabis (27), and the Timeline Follow-Back (54). From these interviews, we derived levels of tobacco use (cigarettes per week) and cannabis use (i.e., lifetime and past year cumulative dosages and frequencies of use, duration of use and age of onset).

We converted cannabis dosage to standardized units (i.e., cones, approximately equivalent to  $\sim 0.1$  g) (55). We measured urinary levels of the carboxy metabolite of THC (THC-COOH) via toxicology analysis, and THC accumulated in hair. Alcohol use (standard drinks per month) were quantified from the structured interviews and the Alcohol Use Disorders Identification Test (56).

### Cognitive Function

We assessed *current IQ*, via the Wechsler Abbreviated Scale of Intelligence (WASI) (57) and *premorbid IQ* using the Wechsler Test of Adult Reading (WTAR) (58), respectively.

We measured *verbal learning and memory* via the California Verbal Learning Test, Second Edition (CVLT-II) (59), which was administered according to the manualized instructions. First, participants were asked to recall a list of 16 words presented orally (*List A*) for five consecutive trials (learning *Trials 1 to 5*). Then, participants were instructed to recall a new list of 16 words (*List B*). Subsequently, participants were asked to recall the words from *List A* without any cues to aid memory, and then with cues (*Short Delay Free Recall* and *Short Delay Cued Recall*). After a 20 min interval, the latter procedure was repeated (*Long Delay Free Recall* and *Long Delay Cued Recall*). Finally, participants were asked if they recognized, from among a list of 48 words

including distractors, those words that were previously presented in *List A* (*Recognition Trial*).

*Visuospatial learning* was examined using the Brown Location Test [BLT (60)], a visuospatial analog of the CVLT-type verbal learning tasks. Participants were presented with 12 pages, one at a time, on which 58 identically sized black outlined circles were located. At each presentation, one of the circles was filled with a red dot and the location of the red dot was different on each page, thus forming a "list" of 12 red dot-locations to remember akin to the list of words to be remembered on each trial of a verbal learning task. After each trial (i.e., the serial presentation of the 12 pages), the participant was provided an identical page of circles where none were filled with red, and was asked to place red chips in the locations where the red dots had been presented. This procedure was repeated 5 times (learning *Trials 1 to 5*).

Then, participants were presented with a new series of 12 black dots, which they were also asked to recall as above (*Interference trial*). Subsequently, participants were asked to recall the locations of the red dots from *Trials 1–5* immediately after Interference (*Short Delay Recall*), after 20 min delay (*Long Delay Recall*), and after rotating the recall page 90 degrees. Finally, participants were presented with a page containing the original location of the red dots, and additional distractor red dots. Participants were asked to distinguish known 12 dot locations from 12 distractor dot locations (*Recognition trial*).

## Statistical Analyses

### Descriptive Demographic, Clinical, and Substance Use Data

As the majority of variables were skewed and not transformable, group comparisons for descriptive purposes were performed using chi-square tests for categorical variables (sex); as well as ANCOVAs for normally distributed discrete variables (IQ, STAI) and Kruskal-Wallis tests, followed by *post hoc* Mann-Whitney U tests, for non-normally distributed discrete variables.

### Primary Aim

To examine group differences for CVLT and BLT performance, Quade's method (61) was used for not normally distributed data from the CVLT and the BLT. Quade's method enables running non-parametric tests comparing groups using covariates. With Quade's method, the dependent variable in ANOVA is the unstandardized residual of a linear regression between the ranked (in ascending order) dependent variable (CVLT and BLT scores) and the ranked covariates (IQ for CVLT, IQ, and age for BLT).

Comparisons between CVLT and BLT trials *within* groups were performed using Friedman- and *post-hoc* Wilcoxon signed rank tests.

### Exploratory Correlations

Spearman's correlations were run to investigate how performance on the CVLT and BLT (residualized data after regressing out the effects of IQ on the CVLT data, and that of IQ and age on the BLT data) was associated with (i) THC or THC-COOH in hair and urine, respectively; and (ii) extent of cannabis use [duration, age at use onset, dosage (lifetime cumulative cones), frequency (smoking days/month), and hours since last cannabis use].

For all correlations, we utilized the conservative Bonferroni method to control for multiple tests and therefore readjusted the significance threshold to  $\alpha = 0.0005$ .

### Covariates

We retained IQ as a covariate in all analyses of CVLT and BLT data. We used age as an additional covariate in BLT data analyses, as age was significantly associated with BLT measures in non-users. Sex was used as a within-groups factor in analyses of CVLT but not BLT performance, as it significantly affected the former but not the latter. Alcohol standard drinks/month, tobacco cigarettes/week and sub-diagnostic psychopathology symptoms (i.e., anxiety, depression, positive and negative psychotic symptoms) were not included as covariates because they did not significantly affect CVLT and BLT performance.

### Sensitivity analyses

A series of two sensitivity analyses were performed to confirm the robustness of the effects. First, all analyses were repeated excluding 12 chronic users who used any illicit substances other than cannabis and these confirmed the results from the analyses run with the whole sample. Therefore, we report the results from the whole group analyses.

Second, we reran group comparisons without 7 current users who on the day of testing admitted to having used cannabis for less than the required at least 12 h—abstinence (range 3–10 h, median of 4 h). As these individuals did not show overt signs of acute intoxication, we proceeded with testing.

All analyses were conducted using SPSS, Version 21 (IBM).

## RESULTS

**Table 1** summarizes data on demographic, clinical and substance use measures for chronic users, former users and non-users.

### Socio-Demographic and IQ Data

All groups were matched by age and premorbid IQ. Chronic users and controls had an equal composition of males and females. However, former users had a lower proportion of females to males, relative to both chronic users ( $\chi^2 = 5.82$ ,  $p = 0.016$ , respectively) and non-users ( $\chi^2 = 6.05$ ,  $p = 0.014$ ).

Education years were lower in chronic users than non-users ( $Z = -2.79$ ,  $p = 0.005$ ), and IQ was also lower in chronic users than the other groups (i.e., controls [ $t = -3.71$ ,  $p < 0.001$ ], former users [ $t = -2.42$ ,  $p = 0.018$ ]). Global functioning was lower in both cannabis groups than controls (i.e., chronic users,  $Z = -6.96$ ,  $p < 0.001$ ; and former users,  $Z = -3.80$ ,  $p < 0.001$ ).

### Alcohol and tobacco Use level

Alcohol use (standard drinks/month) was greater in former users compared to all groups (i.e., current users:  $Z = -2.48$ ,  $p = 0.013$ , and controls:  $Z = -2.83$ ,  $p = 0.005$ ). Tobacco use (cigarettes/week) was greater in current users than former users, and lowest in controls ( $Z = -2.13$ ,  $p = 0.034$  and  $Z = -3.24$ ,  $p = 0.001$ , respectively).

### Cannabis Use level

The cannabis groups had similar cannabis use' duration, age of onset and lifetime dosage (see **Table 1**). However, current users smoked more days/week and consumed a greater amount of cannabis in the past year than former users.

Abstinence duration in current users was median of 16 h (range 3–336 h). Seven current cannabis users reported abstaining for 3 to 10 h despite our request to abstain for at least 12 h; and therefore analyses were repeated excluding these very recent users. Abstinence duration in former users was a mean of 2.5 years (median 6 months, range 1 month–19 years); 9 former users had ceased cannabis use within the past 12 months.

### Subclinical Psychopathology Symptoms

Symptom severity for anxiety, depression and psychosis was greater in current users than controls ( $t = 4.11$ ,  $p < 0.001$ ;  $Z = -4.04$ ,  $p = <0.001$ ;  $Z = -2.66$ ,  $p = 0.008$ ;  $Z = -3.41$ ,  $p = 0.001$ , respectively), but did not differ between the other groups ( $p = n.s.$ ).

### Group Differences in CVLT Performance

**Table 1** Overviews group differences in CVLT performance. Group differences in CVLT trials are overviewed below, followed by learning curves and learning trials in each of the three groups (controls, current users, former users).

#### CVLT Trials

Group differences emerged for 7 out of the 18 CVLT variables: *Trials 1, 3, 1–5, B, Short Delay Free Recall, Long Delay Free Recall* and *Long Delay Cued Recall* (see **Table 2**).

#### Current Users vs. Controls

Cannabis users performed worse than controls for CVLT *Trial 1* ( $F = 6.52$ ,  $p = 0.012$ ), and learning from *Sum Trials 1–5* ( $F = 4.54$ ,  $p = 0.035$ ). Additionally, current users recalled less words than controls for CVLT *List B* ( $F = 10.08$ ,  $p = 0.002$ ), *Short Delay Free Recall* ( $F = 4.37$ ,  $p = 0.039$ ) and *Long Delay Free Recall* ( $F = 5.12$ ,  $p = 0.026$ ). These group differences did not survive a sensitivity analyses that we ran after excluding seven current users who consumed cannabis recently <12 h before testing.

After running the sensitivity analysis, the only performance difference that emerged in cannabis users vs. controls was poorer performance by users in *Long Delay Cued Recall* ( $F = 4.35$ ,  $p = 0.015$ ).

#### Former Users vs. Controls

Similarly to current users, former users vs. controls showed lower CVLT performance for *Long Delay Cued Recall* ( $F = 4.42$ ,  $p = 0.040$ ), *Trial 1* ( $F = 6.95$ ,  $p = 0.011$ ), and learning from *Sum Trials 1–5* ( $F = 5.14$ ,  $p = 0.027$ ). Additionally, former users performed worse than controls on *Trial 3* ( $F = 6.04$ ,  $p = 0.017$ ).

### CVLT Learning Curves

As shown in **Figure 1**, CVLT learning curves improved for all groups at every trial from *Trial 1* to *Trial 4* (all  $p < 0.001$ ). Controls improved word recall at every trial from *Trial 1* to *Trial 4* ( $Z$  range from  $-5.71$  to  $-4.54$ ,  $p < 0.001$ ). Similarly to controls, current users improved word recall at every trial from *Trial 1* to



**TABLE 1** | Socio-demographic, cannabis and other substance use, and psychopathology symptoms current and former chronic cannabis users and controls.

	Controls	Cannabis users	Former cannabis users	Z	p	df	
<b>Socio-demographic, IQ, alcohol/tobacco</b>							
Total N [female] <sup>†‡</sup>	46 (26)	69 (37)	12 (2)	$\chi^2$ 6.53	0.038	2	
Age, yrs	31.17 (12.83)	32.68 (11.17)	37.83 (11.03)	3.34	0.188	126	
Education, yrs*	13.97 (1.60)	12.82 (2.25)	13.17 (2.41)	7.90	0.019	122	
Premorbid IQ	106.78 (10.18)	101.40 (13.29)	106.75 (5.01)	5.55	0.062	125	
IQ <sup>†‡</sup>	111.76 (10.91)	104.15 (10.72)	112.25 (11.21)	7.22	0.001	125	
Global functioning <sup>†‡</sup>	85.5 (4.95)	73.7 (9.77)	75.58 (9.64)	50.32	<0.001	126	
Alcohol use (drinks/mo) <sup>†‡</sup>	18.49 (23.89)	24.52 (32.74)	52.48 (32.80)	9.43	0.009	126	
Tobacco (cigarettes/week) <sup>†‡</sup>	4.95 (16.79)	58.95 (53.15)	29.33 (38.16)	56.56	<0.001	126	
<b>Cannabis use</b>							
Frequency, days/mo	Lifetime <sup>†</sup>	NA	23.24 (6.89)	28.83 (4.04)	-3.63	<0.001	80
	Past 12 mo <sup>†</sup>	NA	24.59 (8.62)	3.33 (8.51)	-4.81	<0.001	80
Dosage, cones	Lifetime	NA	69,183 (73,271)	43,036 (36,492)	-1.17	0.241	80
	Past 12 mo <sup>†</sup>	NA	5,070 (4039)	136.3 (375.3)	-5.30	<0.001	80
Duration of regular use, yrs	NA	15.13 (10.00)	15.92 (9.61)	-0.297	0.767	80	
Onset age, yrs	NA	16.96 (3.93)	17.33 (3.75)	-0.510	0.610	80	
Abstinence duration, yrs	NA	NA	2.46 (5.56)	NA	NA	12	
<b>Psychopathology symptom scores</b>							
Trait anxiety, STAI	33.64 (7.39)	41.99 (12.19)	35.33 (13.39)	2.13	0.12	124	
Depression, CAPE*	12.29 (2.69)	15.06 (3.87)	13.58 (2.75)	16.34	<0.001	125	
Positive psychotic, CAPE*	24.56 (3.37)	27.58 (6.21)	25.82 (3.22)	7.46	0.024	121	
Negative psychotic, CAPE*	22.26 (5.11)	26.51 (6.95)	23.42 (13.39)	11.95	0.003	122	

NA, not applicable; <sup>†</sup>Significant difference between cannabis users and controls. <sup>††</sup>Significant difference between former cannabis users and controls. <sup>†††</sup>Significant difference between cannabis users and former cannabis users.  $\chi^2$ , for results from Kruskal-Wallis test; Z for results from Mann-Whitney U test; F values for IQ and anxiety. Current and premorbid IQ measured with the Wechsler Abbreviated Scale of Intelligence and the Wechsler Test of Adult Reading, respectively; Alcohol use measured with the Alcohol Use Disorders Identification Test; Anxiety symptoms measured with the State and Trait Anxiety Inventory (STAI), depression, positive and negative symptoms measured with the subscales of the Community Assessment of Psychic Experiences (CAPE); Global functioning measured with the module of the DSM-IV.

Trial (Z range from -6.83 to -2.58,  $p < 0.01$ ). Former users only significantly improved their word recall from Trial 1 to Trial 2 (Z = -3.08,  $p = 0.002$ ).

## CVLT Delayed Recall

### Controls

Controls showed a significant increase in the **Long Delay Free Recall vs. Short Delay Free Recall** (Z = -3.09,  $p = 0.002$ ) and **Long Delay Cued Recall vs. Short Delay Cued Recall** (Z = -3.25,  $p = 0.001$ ).

Additionally, controls improved significantly only after cues were given, in the **Short Delay Cued Recall** trial (Z = -2.14,  $p = 0.032$ ).

### Current Users

Similarly to controls, current users showed a significant increase in the **Free Long Delay Recall vs. Free Short Delay Recall** (Z = -2.79,  $p < 0.005$ ) and **Cued Long Delay Recall vs. Cued Short Delay Recall** (Z = -2.28,  $p = 0.023$ ). In contrast to controls, current users performed better during **Cued Recall vs. Free Recall** for the **Short Delay** trial (Z = -3.06,  $p = 0.002$ ) and for the **Long Delay** trial (Z = -2.12,  $p = 0.034$ ).

### Former Users

Former users did not show differences in any delayed recall trials.

## Exploratory Correlations Between CVLT Performance, Cannabis Use Measures, and Age

Older age was not associated with any CVLT variables in cannabis users, but was associated with better recall at Trial 5 ( $r_s = 0.465$ ,  $p = 0.001$ ) and more **Intrusions** in controls ( $r_s = 0.291$ ,  $p = 0.045$ ) and with greater **Recognition of False Positives** in former users ( $r_s = 0.602$ ,  $p = 0.029$ ).

### Current Users

Greater cannabis frequency over the lifetime was associated with lower **Retroactive Interference** and greater **Loss After Consolidation**; and greater frequency of cannabis use in the past year was associated with worse recall at Trial 1. More cumulative cannabis dosage in the lifetime was correlated with worse recall at Trial 1 and worse **Retroactive Interference**; and greater past year cumulative dosage was correlated with worse recall at Trial 1. Earlier age of onset was correlated with less **Intrusions**, lower **Retroactive Interference** and greater **Proactive Interference**.


### Former Users

In former users, greater frequency of cannabis use in the past year was associated with greater recall at Trial 1, greater **Recognition of False Positives** and lower **Intrusions**. Greater past year cumulative

**TABLE 2** | CVLT-II performance in *current* users, *former* users, and controls: mean (standard deviation).

CVLT trials	Controls	Current users	Former users	F-value	p	df
<b>Trial 1*</b> <sup>†</sup>	8.30 (2.05)	6.96 (2.29)	6.62 (2.14)	5.09	<b>0.01</b>	127
Trial 2	11.39 (2.33)	10.30 (2.65)	9.92 (2.14)	1.53	0.22	127
<b>Trial 3*</b> <sup>†</sup>	13.07 (1.93)	11.77 (2.66)	11.00 (2.68)	3.67	<b>0.03</b>	127
Trial 4	13.59 (2.04)	12.35 (2.51)	12.15 (2.94)	1.95	0.15	127
Trial 5	13.91 (1.87)	13.00 (2.51)	12.69 (2.69)	0.91	0.41	127
<b>Trials 1–5*</b> <sup>†</sup>	60.26 (7.98)	54.38 (10.92)	52.38 (10.56)	3.85	<b>0.02</b>	127
<b>List B*</b>	7.07 (2.02)	5.59 (2.18)	6.00 (2.16)	5.40	<b>0.01</b>	127
<b>Short delay free recall*</b>	12.98 (2.24)	11.42 (2.87)	11.38 (2.99)	2.74	<b>0.07</b>	127
Short delay cued recall	13.46 (2.00)	11.96 (3.07)	12.00 (2.77)	2.10	0.13	127
<b>Long delay free recall*</b>	13.65 (2.07)	11.91 (3.11)	11.77 (2.98)	3.65	<b>0.03</b>	127
<b>Long delay cued recall*</b> <sup>†</sup>	13.87 (1.89)	12.28 (2.95)	11.77 (3.11)	3.35	<b>0.04</b>	127
Recognition	15.28 (0.958)	14.67 (1.50)	15.38 (0.961)	2.14	0.12	127
Recognition false positives	0.91 (1.75)	1.67 (2.42)	1.92 (2.96)	0.62	0.54	127
Repetitions	4.30 (3.81)	4.49 (4.06)	4.46 (5.24)	0.02	0.98	127
Intrusions	2.07 (2.99)	2.22 (2.92)	3.77 (5.31)	0.10	0.91	127
Proactive interference	1.24 (2.43)	1.36 (2.14)	0.62 (1.50)	1.09	0.34	127
Retroactive interference	0.94 (1.78)	1.58 (1.81)	1.31 (1.75)	1.03	0.36	127
Loss after consolidation	0.26 (1.73)	1.09 (1.88)	0.92 (1.71)	1.50	0.23	127

\*Significant difference between cannabis users and controls. <sup>†</sup>Significant difference between former cannabis users and controls. CVLT-II, California Verbal Learning Test Version 2. Proactive Interference, prior learning interfering with new learning (Trial 1 - List B); Retroactive Interference, later learning interfering with previous learning (Trial 5—Short Delay Free Recall) and Loss after Consolidation, loss of recalled words after delay (Trial 5—Long Delay Free Recall).



**FIGURE 1** | CVLT learning curves across Trials 1–5, in current cannabis users, former cannabis users and controls, and mean performance for List B, Short Delay Free Recall (SD), Long Delay Free Recall (LD), and Recognition Trial (Recogn.).

dosage was correlated with worse recall at *Trial 4*. Earlier age of onset was correlated with better recall at *Trials 1, 2, 3, 4, 1–5*, and *Trial B*.

None of the correlations run between CVLT performance and cannabis use levels in current and former users survived Bonferroni correction for multiple tests (see **Supplementary Tables 1, 2**).

## Group Differences in BLT Performance

### BLT trials

One cannabis user did not complete the BLT, hence for these analyses  $n = 68$  current cannabis users. BLT performance differed between groups for 2 of the 15 variables (see **Table 3**). Current users performed significantly worse than controls on *Retroactive Interference* and on *Long Delay Rotated Recall* ( $F = 5.95, p = 0.016$  and  $F = 1.62, p = 0.014$ ). Current users performed worse than former users on *Short Delay Free Recall* ( $F = 4.28, p = 0.042$ ).

### Sensitivity analyses for BLT trials

After exclusion of seven participants who reported using cannabis within 12 h of the assessment, impaired performance persisted in current users vs. controls, for both *Retroactive Interference* ( $F = 3.29, p = 0.041$ ) and *Long Delay Rotated Recall* ( $F = 3.13, p = 0.048$ ).

### BLT learning curves

BLT learning curves are shown in **Figure 2**. The pattern of results was identical to that of the CVLT: current users, former users and controls showed significant improvement in recall [ $\chi^2_{(4,68)} = 109.0, p < 0.001$ ;  $\chi^2_{(4,12)} = 24.89, p < 0.001$  and  $\chi^2_{(4,46)} = 114.7, p < 0.001$  respectively]. Current users and controls improved at every trial from *Trial 1* to *Trial 4* ( $Z$  range between  $-4.38$  and  $-2.01, p < 0.04$ ; and  $Z$  range between  $-4.88$  and  $-3.52, p < 0.001$ , respectively). Former cannabis users only improved between *Trial 1* and *Trial 2* (see learning curves in **Figure 2**,  $Z = -2.57, p = 0.010$ ).

### BLT Delayed Recall

Current users showed worse BLT performance in *Delayed Recall* only after *Page Rotation* ( $Z = -4.76, p < 0.001$ ). None of the groups showed a difference between *Short Delay Recall* and *Long Delay Recall*.

## Exploratory correlations Between BLT Performance, Age, and Cannabis Use Measures

There was no association between age and BLT performance in neither current users, former users and controls.

### Current users

In current cannabis users, greater lifetime cannabis use frequency was correlated with lower BLT recall at *Trial 1–5 Total*, and greater *Recognition of False Positives*. Greater THC-COOH in urine was associated with lower performance during *Interference* and better performance during the *Proactive Interference* trial.

### Former users

In former cannabis users, greater frequency of cannabis use in the lifetime was associated with greater recall at BLT *Trial 3*, *Trial 4*, *Trial 1–5 Total*, *Long Delay Free Recall*, *Long Delay Rotated Recall* and greater *Recognition Hits*. Later age of cannabis use onset was associated with lower recall at *Trial 3* and *Trial 4*, and lower *Recognition Hits*. Greater abstinence duration was associated with greater *Long Delay Free Recall* and *Long Delay Rotated Recall*.

None of the correlations between BLT performance, cannabis use levels and age in any groups survived Bonferroni correction for multiple tests (**Supplementary Tables 3, 4**).

## DISCUSSION

This study shows that cannabis use status (current vs. former use) and extent of cannabinoid exposure have a differential impact on specific measures of verbal and visuospatial learning. Specifically, select measures of verbal learning in current and former cannabis users were impaired i.e., *Long Delay Cued Recall* of the CVLT. Instead, visuospatial learning was impaired only in current users i.e., *Retroactive Interference* and *Long Delay Rotated Recall* of the BLT. Performance on the CVLT and BLT was not significantly associated with any measures of cannabis exposure.

Poorer CVLT *Long Delay Cued Recall* performance in current cannabis users was robust to sensitivity analyses which excluded 7 people who did not appear intoxicated but admitted having smoked cannabis recently i.e.,  $<12$  h before testing (instead of the required  $>12$  h), and persisted in former users. Therefore, lower *Long Delay Cued Recall* may reflect the residual effects of chronic cannabis use that are long-lasting and detectable well-beyond prolonged abstinence. This interpretation is consistent with findings that verbal learning alterations in cannabis users are apparent after THC metabolites are no longer detected in urine (9), and in cannabis users with chronic exposure reflected by long duration [i.e., 16 yrs (25) and 23 yrs (24)], dependent and almost daily use (62).


However, these notions are not supported by the lack of robust correlations between *Long Delay Cued Recall* and any measure of cannabis exposure including abstinence durations and chronicity of use. Therefore, the specific indices of cannabis exposure driving CVLT alterations in these groups of current and former users are unclear. Of note, verbal learning is ascribed by the function of posterior and frontoparietal cortices (41, 46, 63–65) that are high in cannabinoid receptors and thus might be vulnerable to the long lasting effects of repeated impact of cannabinoid exposure via complex neural mechanisms (66).

Surprisingly, most of the CVLT performance differences between current users and controls (i.e., *Trial 1*, *Sum Trials 1–5*, *List B*, *Short Delay Free Recall*, *Long Delay Free Recall*) were no longer detectable after sensitivity analyses without 7 people who recently smoked cannabis i.e.,  $<12$  h before testing. Therefore, recent cannabis exposure may drive alterations of select components of verbal learning alterations (17, 24, 27, 41, 46, 62). This notion is supported by other study findings that better Ray Auditory Verbal Learning Test performance is associated with longer abstinence duration (67) (*List B*

**TABLE 3** | BLT performance in *current* cannabis users, *former* cannabis users and *controls*: mean (standard deviation).

BLT trials	Controls	Current users	Former users	F	p	df
Trial 1	3.93 (2.03)	3.93 (2.01)	3.54 (1.56)	0.05	0.95	126
Trial 2	5.85 (2.38)	5.24 (2.10)	5.85 (1.95)	0.47	0.62	126
Trial 3	7.00 (2.69)	6.22 (2.44)	6.46 (2.79)	0.54	0.58	126
Trial 4	8.72 (2.66)	7.26 (2.69)	7.85 (3.39)	1.80	0.17	126
Trial 5	8.72 (2.86)	7.78 (2.92)	8.77 (2.92)	0.82	0.45	126
Trials 1–5	34.22 (10.84)	10.43 (9.90)	32.46 (9.17)	0.73	0.49	126
Interference	3.72 (2.04)	3.19 (1.72)	3.23 (1.42)	0.04	0.96	126
<b>Short delay free recall<sup>‡</sup></b>	7.93 (2.93)	6.22 (2.92)	8.15 (3.13)	3.74	<b>0.03</b>	126
Long delay free recall	7.76 (2.81)	6.50 (2.87)	7.23 (3.68)	1.40	0.25	126
<b>Long delay rotated recall*</b>	7.15 (3.18)	5.34 (2.87)	6.69 (3.43)	2.53	<b>0.08</b>	126
Recognition	18.70 (3.41)	16.78 (3.46)	17.69 (4.68)	1.80	0.17	126
Recognition false positives	3.00 (2.29)	3.99 (2.28)	3.00 (2.31)	1.60	0.21	126
Proactive interference	0.22 (2.30)	0.74 (2.35)	0.31 (2.53)	0.19	0.83	126
<b>Retroactive interference*</b>	0.78 (1.53)	1.56 (2.10)	0.62 (1.56)	3.51	<b>0.03</b>	126
Loss after consolidation	0.96 (1.66)	1.28 (1.91)	1.54 (1.51)	1.15	0.32	126

\*Significant difference between cannabis users and controls. ‡Significant difference between cannabis users and former cannabis users. Proactive Interference, prior learning interfering with new learning (Trial 1—List B); Retroactive Interference, later learning interfering with previous learning (Trial 5—Short Delay Free Recall) and Loss after Consolidation, loss of recalled words after delay (Trial 5—Long Delay Free Recall).



**FIGURE 2** | BLT learning curves for Trials 1–5, in current users, former cannabis users and controls, as well as for Interference Trial (Interf.), Short Delay Free Recall (SD), Long Delay Free Recall (LD), Rotated Long Delay Recall (Rotated), and Recognition Trial (Recogn.).

(27), total words recalled (12, 24). However, we found no significant correlation between better CVLT performance and number of hours from last cannabis use or urinary THC

metabolites. Thus, the strength of this finding needs to be verified in future studies that carefully measure abstinence duration (68), as only a few studies of verbal learning to



date report how long people abstained from cannabis before testing (25, 35).

Poorer visuospatial learning performance in cannabis users emerged in select BLT measures (*Retroactive Interference* and *Long Delay Rotated Recall*). These differences were robust and survived sensitivity analyses without a subgroup of recent users. Our findings are consistent with reports that visuospatial learning alterations in cannabis users affect *recall* but not *acquisition* trials (19, 45), but contrast previous meta-analytic findings that failed to find group differences (12).

The discrepancy between our results and those from previous work, might be due to systematic differences between sample characteristics and the tools used to assess visuospatial learning. First, our sample was older (i.e., ~34 years) than the meta-analyzed samples i.e., <26 yrs (12). Interestingly, previous evidence shows that aging affects visuospatial learning (69–71). Therefore, lower visuospatial *recall* may be due to altered aging processes in cannabis users. However, the lack of significant correlations between age and BLT performance in this study does not support this notion and is to be further tested in future work. Second, the meta-analyzed studies to date used measures of visuospatial learning other than the BLT. Such measures (e.g., accuracy and total scores from the Rey-Osterrieth and Bender Visual-Motor Gestalt Test) might not have been sensitive enough to detect alterations specific to *recall* rather than *acquisition* trials (12).

The mechanisms underlying altered BLT in *recall* trials in current users are unclear. One candidate mechanism is impaired executive functioning (45, 72–74). Indeed, recall—but not acquisition—relies on executive function (75). Also, aging significantly affects the integrity of para-hippocampal and cingulate cortices that are concurrently ascribed to visuospatial learning, and to the residual effects of regular cannabis exposure (19).

Interestingly, we found for the first time that both current and former users showed similar *Learning Curves* across the verbal and visuospatial domains, suggesting the employment of common learning strategies across both domains (73). This similarity was apparent, given the similar structure of the BLT and the CVLT (i.e., both tests start with 5 *Learning Trials*, followed by an *Interference* trial and *Delayed Recall* trials). The overlapping *learning curves* between current and former users (see **Figure 1**), suggest that alteration of *learning curves* commences during regular use and does not recover after prolonged abstinence.

There are several limitations to this study. First, we examined a small sample of former users, and our findings require validation in larger samples. Second, our group of current users were *not* matched to controls for level of education, IQ and severity of sub-clinical psychopathology symptoms. Nevertheless, premorbid IQ was matched between groups, and we controlled for current IQ i.e., by using IQ as a covariate in all group comparisons (years of education were not associated with any performance measures and were thus not included as a covariate).

We also minimized potential confounding impact of severe mental health conditions on cognitive performance, by screening for any diagnoses of psychopathology. However, we cannot rule

out that sub-clinical psychopathology symptoms in our sample, which are shown to exacerbate cognitive deficits (15), may have driven our findings on cognitive alterations. Since sub-clinical psychopathology symptoms as measured in this study did not exert a significant effect on performance, they were therefore not included as covariates in the analyses. On the other hand, our sample is representative of cannabis users within the general community, where higher (although not diagnosable) psychopathology symptoms and worse cognitive outcomes have been consistently reported (14, 76–79).

Third, correlational analyses were run in current users and former users including males and females, and not in different sexes separately, therefore precluding a detailed understanding of possible sex differences in the emerging alterations. Our strategy mitigated type-1 errors, as we have run a substantial number of correlational analyses in current and former cannabis users across all cognitive variables examined. Fourth, the number of abstinent users ( $n = 12$ ) was small for statistical analyses, and findings pertaining to this group require replication in larger samples. Further, the cross-sectional study design and the lack of information on the sample's resilience levels, prevented to determine if abstinence was the reason for better performance compared to current usage, or the consequence of a latent factor such as resilience that also leads to better performance.

Our findings might be accounted for in the design of cognitive interventions aimed to support abstinence in cannabis users. For example, if a clinical practitioner knew that a patient uses cannabis regularly, and that regular cannabis use is associated with impaired *Long Delay Cued Recall*, the practitioner may implement strategies to ensure that their client recalls information/instructions critical for engaging with the treatment (e.g., repeating or asking the client to repeat the instruction/information, sharing written instructions). Also, knowing that regular cannabis users have worse performance on *Retroactive Interference* and *LD Rotate Recall*, could indicate to a practitioner that visuospatial information relevant for the treatment (e.g., the location of appointment/testing/treatment sites) may not be retained when learning occurs; and may prompt the practitioner to implement strategies to boost recall of visuospatial information relevant for the patient to attend treatment (e.g., sharing a map or sending a text reminder with information on appointment/testing/treatment sites).

In conclusion, our findings suggest that current cannabis use is associated with chronic residual effects on select aspects of verbal learning (*Long Delay Cued Recall*) and of visuospatial learning (*Retroactive Interference* and *Long Delay Rotated Recall*). Prolonged abstinence in former users was associated with altered verbal learning but intact visuospatial learning, suggesting that abstinence has a different impact on distinct domains of learning. Our findings warrant the conduct of future work that systematically tracks how learning performance in chronic cannabis users is affected during intoxication to map acute-on-chronic effects, residual effects beyond intoxication, and those that remain with prolonged abstinence, to track how the residual-on-chronic effects dissipate over time. Additional studies with careful assessment of cannabis use indices and exposure to THC and other cannabinoids, should examine how these findings

extend to cannabis user groups who experience worse mental health outcomes.

## DATA AVAILABILITY STATEMENT

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation.

## ETHICS STATEMENT

Ethics approval was given by the Mental Health Research and Ethics Committee (MHREC, I.D. number 459111), the Melbourne Health and North Western Mental Health (Melbourne, Australia), the Wollongong Human Research Ethics Committee (HREC, project number NSA07/03), and the ethics committee of the Murdoch Children's Research Institute (MCRI). The patients/participants provided their written informed consent to participate in this study.

## AUTHOR CONTRIBUTIONS

MT and VL drove data collection. VL drafted and revised the manuscript writing and statistical analyses and interpretation. YD contributed to the drafting and the running of the statistical analyses. MY and NS drove the study design. All authors

revised the manuscript and contributed to the interpretation of the findings.

## FUNDING

This study was funded by the National Medical & Health Research Council (NHMRC ID: 459111). MY has received funding from Monash University, the National Health and Medical Research Council (NHMRC; including Fellowship #APP1117188), the Australian Research Council (ARC), Australian Defence Science and Technology (DST), and the Department of Industry, Innovation and Science (DIIS). He has also received philanthropic donations from the David Winston Turner Endowment Fund, the Wilson Foundation, as well as payment from law firms in relation to court and/or expert witness reports. NS was supported by an Australian Research Council Future Fellowship (FT110100752). VL was supported by The Australian Catholic University through a competitive scheme.

## SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fpsy.2021.663701/full#supplementary-material>

## REFERENCES

- UNODC. *World Drug Report 2020. World Drug Report Series 2020*. United Nations Office on Drugs and Crime (2020).
- Chandra S, Radwan MM, Majumdar CG, Church JC, Freeman TP, ElSohly MA. New trends in cannabis potency in USA and Europe during the last decade (2008–2017). *Eur Arch Psychiatry Clin Neurosci*. (2019) 269:5–15. doi: 10.1007/s00406-019-00983-5
- Freeman TP, Groshkova T, Cunningham A, Sedefov R, Griffiths P, Lynskey MT. Increasing potency and price of cannabis in Europe, 2006–16. *Addiction*. (2019) 114:1015–23. doi: 10.1111/add.14525
- Budney AJ, Sofis MJ, Borodovsky JT. An update on cannabis use disorder with comment on the impact of policy related to therapeutic and recreational cannabis use. *Eur Arch Psychiatry Clin Neurosci*. (2019) 269:73–86. doi: 10.1007/s00406-018-0976-1
- Lopez-Quintero C, de los Cobos JP, Hasin DS, Okuda M, Wang A, Grant BF, Blanco C. Probability and predictors of transition from first use to dependence on nicotine, alcohol, cannabis, and cocaine: results of the National Epidemiologic Survey on Alcohol and Related Conditions (NESARC). *Drug Alcohol Depend*. (2011) 115:120–30. doi: 10.1016/j.drugalcdep.2010.11.004
- Lorenzetti V, Hoch E, Hall W. Adolescent cannabis use, cognition, brain health and educational outcomes: a review of the evidence. *Eur Neuropsychopharmacol*. (2020) 26:169–80. doi: 10.1016/j.euroneuro.2020.03.012
- Broyd SJ, van Hell HH, Beale C, Yücel M, Solowij N. Acute and chronic effects of cannabinoids on human cognition—a systematic review. *Biol Psychiatry*. (2016) 79:557–67. doi: 10.1016/j.biopsych.2015.12.002
- Bogaty SER, Lee RSC, Hickie IB, Hermens DF. Meta-analysis of neurocognition in young psychosis patients with current cannabis use. *J Psychiatr Res*. (2018) 99:22–32. doi: 10.1016/j.jpsychires.2018.01.010
- Duperrouzel JC, Granja K, Pacheco-Colón I, Gonzalez R. Adverse effects of cannabis use on neurocognitive functioning: a systematic review of meta-analytic studies. *J Dual Diagn*. (2020) 16:43–57. doi: 10.1080/15504263.2019.1626030
- Figueiredo PR, Tolomeo S, Steele JD, Baldacchino A. Neurocognitive consequences of chronic cannabis use: a systematic review and meta-analysis. *Neurosci Biobehav Rev*. (2020) 108:358–69. doi: 10.1016/j.neubiorev.2019.10.014
- Lovell ME, Akhurst J, Padgett C, Garry MI, Matthews A. Cognitive outcomes associated with long-term, regular, recreational cannabis use in adults: a meta-analysis. *Exp Clin Psychopharmacol*. (2020) 28:471–94. doi: 10.1037/pha0000326
- Scott JC, Wolf DH, Calkins ME, Bach EC, Weidner J, Ruparel K, et al. Cognitive functioning of adolescent and young adult cannabis users in the Philadelphia Neurodevelopmental Cohort. *Psychol Addict Behav*. (2017) 31:423–34. doi: 10.1037/adb0000268
- Solowij N, Battisti R. The chronic effects of cannabis on memory in humans: a review. *Curr Drug Abusers Rev*. (2008) 1:81–98. doi: 10.2174/1874473710801010081
- Volkow ND, Baler RD, Compton WM, Weiss SR. Adverse health effects of marijuana use. *N Engl J Med*. (2014) 370:2219–27. doi: 10.1056/NEJMra1402309
- Farmer RE, Kosty DB, Seeley JR, Duncan SC, Lynskey MT, Rohde P, et al. Natural course of cannabis use disorders. *Psychol Med*. (2015) 45:63–72. doi: 10.1017/S003329171400107X
- Ellickson PL, Martino SC, Collins RL. Marijuana use from adolescence to young adulthood: multiple developmental trajectories and their associated outcomes. *Health Psychol*. (2004) 23:299–307. doi: 10.1037/0278-6133.23.3.299
- Meier MH, Caspi A, Ambler A, Harrington H, Houts R, Keefe RS, et al. Persistent cannabis users show neuropsychological decline from childhood to midlife. *Proc Natl Acad Sci USA*. (2012) 109:E2657–64. doi: 10.1073/pnas.1206820109
- Pope HG, Jacobs A, Mialet J-P, Yurgelun-Todd D, Gruber S. Evidence for a sex-specific residual effect of cannabis on visuospatial memory. *Psychother Psychosom*. (1997) 66:179–84. doi: 10.1159/000289132
- Sneider JT, Gruber SA, Rogowska J, Silveri MM, Yurgelun-Todd DA. A preliminary study of functional brain activation among marijuana

- users during performance of a virtual water maze task. *J Addict.* (2013) 2013:461029. doi: 10.1155/2013/461029
20. Chang L, Yakupov R, Cloak C, Ernst T. Marijuana use is associated with a reorganized visual-attention network and cerebellar hypoactivation. *Brain.* (2006) 129:1096–12. doi: 10.1093/brain/awl064
  21. Ashtari M, Avants B, Cyckowski L, Cervellione KL, Roofeh D, Cook P, et al. Medial temporal structures and memory functions in adolescents with heavy cannabis use. *J Psychiatr Res.* (2011) 45:1055–66. doi: 10.1016/j.jpsychires.2011.01.004
  22. Gruber SA, Sagar KA, Dahlgren MK, Racine M, Lukas SE. Age of onset of marijuana use and executive function. *Psychol Addict Behav.* (2012) 26:496–506. doi: 10.1037/a0026269
  23. Solowij N, Stephens R, Roffman RA, Babor T. Does marijuana use cause long-term cognitive deficits? *JAMA.* (2002) 287:2653–4. doi: 10.1001/jama.287.9.1123
  24. Solowij N, Stephens RS, Roffman RA, Babor T, Kadden R, Miller M, et al. Cognitive functioning of long-term heavy cannabis users seeking treatment. *JAMA.* (2002) 287:1123–31.
  25. Messinis L, Kyprianidou A, Malefaki S, Papathanasopoulos P. Neuropsychological deficits in long-term frequent cannabis users. *Neurology.* (2006) 66:737–9. doi: 10.1212/01.wnl.0000201279.83203.c6
  26. Becker B, Wagner D, Gouzoulis-Mayfrank E, Spuentrup E, Daumann J. Altered parahippocampal functioning in cannabis users is related to the frequency of use. *Psychopharmacology.* (2010) 209:361–74. doi: 10.1007/s00213-010-1805-z
  27. Solowij N, Jones KA, Rozman ME, Davis SM, Ciarrochi J, Heaven PCL, et al. Verbal learning and memory in adolescent cannabis users, alcohol users and non-users. *Psychopharmacology.* (2011) 216:131–44. doi: 10.1007/s00213-011-2203-x
  28. Cuttler C, McLaughlin RJ, Graf P. Mechanisms underlying the link between cannabis use and prospective memory. *PLoS ONE.* (2012) 7:e36820. doi: 10.1371/journal.pone.0036820
  29. Thames AD, Arbid N, Sayegh P. Cannabis use and neurocognitive functioning in a non-clinical sample of users. *Addict Behav.* (2014) 39:994–9. doi: 10.1016/j.addbeh.2014.01.019
  30. Bolla KI, Brown K, Eldreth D, Tate K, Cadet JL. Dose-related neurocognitive effects of marijuana use. *Neurology.* (2002) 59:1337–43. doi: 10.1212/01.WNL.0000031422.66442.49
  31. Gaoni Y, Mechoulam R. The isolation and structure of delta-1-tetrahydrocannabinol and other neutral cannabinoids from hashish. *J Am Chem Soc.* (1971) 93:217–24. doi: 10.1021/ja00730a036
  32. Landfield PW, Caldwelllader LB, Vinsant S. Quantitative changes in hippocampal structure following long-term exposure to  $\Delta^9$ -tetrahydrocannabinol: possible mediation by glucocorticoid systems. *Brain Res.* (1988) 443:47–62. doi: 10.1016/0006-8993(88)91597-1
  33. Di Forti M, Marconi A, Carra E, Fraiteta S, Trotta A, Bonomo M, et al. Proportion of patients in south London with first-episode psychosis attributable to use of high potency cannabis: a case-control study. *Lancet Psychiatry.* (2015) 2:233–8. doi: 10.1016/S2215-0366(14)00117-5
  34. Roten A, Baker NL, Gray KM. Cognitive performance in a placebo-controlled pharmacotherapy trial for youth with marijuana dependence. *Addict Behav.* (2015) 45:119–23. doi: 10.1016/j.addbeh.2015.01.013
  35. Hanson KL, Winward JL, Schweinsburg AD, Medina KL, Brown SA, Tapert SF. Longitudinal study of cognition among adolescent marijuana users over three weeks of abstinence. *Addict Behav.* (2010) 35:970–6. doi: 10.1016/j.addbeh.2010.06.012
  36. Pope HG Jr, Gruber AJ, Hudson JL, Huestis MA, Yurgelun-Todd D. Neuropsychological performance in long-term cannabis users. *Arch Gen Psychiatry.* (2001) 58:909–15. doi: 10.1001/archpsyc.58.10.909
  37. Medina KL, Nagel BJ, Park A, McQueeney T, Tapert SF. Depressive symptoms in adolescents: associations with white matter volume and marijuana use. *J Child Psychol Psychiatry.* (2007) 48:592–600. doi: 10.1111/j.1469-7610.2007.01728.x
  38. Medina KL, Schweinsburg AD, Cohen-Zion M, Nagel BJ, Tapert SF. Effects of alcohol and combined marijuana and alcohol use hippocampal volume and asymmetry. *Neurotoxicol Teratol.* (2007) 29:141–52. doi: 10.1016/j.ntt.2006.10.010
  39. Schoeler T, Bhattacharyya S. The effect of cannabis use on memory function: an update. *Subst Abuse Rehabil.* (2013) 4:11. doi: 10.2147/SAR.S25869
  40. Winward JL, Hanson KL, Tapert SF, Brown SA. Heavy alcohol use, marijuana use, and concomitant use by adolescents are associated with unique and shared cognitive decrements. *J Int Neuropsychol Soc.* (2014) 20:784–95. doi: 10.1017/S1355617714000666
  41. Fried PA, Watkinson B, Gray R. Neurocognitive consequences of marijuana—a comparison with pre-drug performance. *Neurotoxicol Teratol.* (2005) 27:231–9. doi: 10.1016/j.ntt.2004.11.003
  42. Lyons MJ, Bar JL, Panizzon MS, Toomey R, Eisen S, Xian H, et al. Neuropsychological consequences of regular marijuana use: a twin study. *Psychol Med.* (2004) 34:1239–50. doi: 10.1017/S0033291704002260
  43. Solowij N. Do cognitive impairments recover following cessation of cannabis use? *Life Sci.* (1995) 56:2119–26. doi: 10.1016/0024-3205(95)00197-E
  44. Medina KL, Hanson KL, Schweinsburg AD, Cohen-Zion M, Nagel BJ, Tapert SF. Neuropsychological functioning in adolescent marijuana users: subtle deficits detectable after a month of abstinence. *J Int Neuropsychol Soc.* (2007) 13:807–20. doi: 10.1017/S1355617707071032
  45. McHale S, Hunt N. Executive function deficits in short-term abstinent cannabis users. *Hum Psychopharmacol Clin Exp.* (2008) 23:409–15. doi: 10.1002/hup.941
  46. Tait RJ, Mackinnon A, Christensen H. Cannabis use and cognitive function: eight year trajectory in a young adult cohort. *Addict Postprint.* (2011) 106:2195–203. doi: 10.1111/j.1360-0443.2011.03574.x
  47. American Psychiatric Association. *DSM-V Diagnostic and Statistical Manual of Mental Disorders*, 4th ed (2013).
  48. American Psychiatric Association. *DSM-IV Diagnostic and Statistical Manual of Mental Disorders*, 4th ed (2006).
  49. Spielberger CD. *State-Trait Anxiety Inventory*. John Wiley & Sons, Inc. (2010).
  50. Konings M, Bak M, Hanssen M, Van Os J, Krabbendam L. Validity and reliability of the CAPE: a self-report instrument for the measurement of psychotic experiences in the general population. *Acta Psychiatr Scand.* (2006) 114:55–61. doi: 10.1111/j.1600-0447.2005.00741.x
  51. Yücel M, Zalesky A, Takagi MJ, Bora E, Fornito A, Ditchfield M, et al. White-matter abnormalities in adolescents with long-term inhalant and cannabis use: a diffusion magnetic resonance imaging study. *J Psychiatry Neurosci.* (2010) 35:409–12. doi: 10.1503/jpn.090177
  52. Takagi M, Lubman DI, Cotton S, Fornito A, Baliz Y, Tucker A, et al. Executive control among adolescent inhalant and cannabis users. *Drug Alcohol Rev.* (2011) 30:629–37. doi: 10.1111/j.1465-3362.2010.00256.x
  53. Takagi M, Yücel M, Cotton SM, Baliz Y, Tucker A, Elkins K, et al. Verbal memory, learning, and executive functioning among adolescent inhalant and cannabis users. *J Stud Alcohol Drugs.* (2011) 72:96–105. doi: 10.15288/jsad.2011.72.96
  54. Sobell LC, Sobell MB. *Timeline Follow-Back: A Technique for Assessing Self-Reported Alcohol Consumption*. Totowa, NJ: Humana Press (1992).
  55. Lorenzetti V. *The association between cannabis use and human brain structure*. [Ph.D. thesis]. Melbourne, VIC: The University of Melbourne (2012).
  56. Bohn MJ, Babor TF, Kranzler HR. The alcohol use disorders identification test (AUDIT): validation of a screening instrument for use in medical settings. *J Stud Alcohol.* (1995) 56:423–32. doi: 10.15288/jsa.1995.56.423
  57. Wechsler D. *Wechsler Adult Intelligence Scale*, 3rd ed (1997).
  58. Wechsler D. *Manual for the Wechsler Test of Adult Reading (WTAR)*. San Antonio, TX: The Psychological Corporation (2001).
  59. Delis DC, Kramer JH, Kaplan E, Ober B. *The California Verbal Learning Test*. San Antonio, TX (1987).
  60. Brown FC, Roth RM, Saykin AJ, Beverly-Gibson G. A new measure of visual location learning and memory: development and psychometric properties for the Brown Location Test (BLT). *Clin Neuropsychol.* (2007) 21:811–25. doi: 10.1080/13854040600878777
  61. Quade D. Rank analysis of covariance. *J Am Stat Assoc.* (1967) 62:1187–200. doi: 10.1080/01621459.1967.10500925
  62. Roebke, PV, Vadhan NP, Brooks DP, Levin FR. Verbal learning in marijuana users seeking treatment: a comparison between depressed and non-depressed samples. *Am J Drug Alcohol Abuse.* (2014) 40:274–9. doi: 10.3109/00952990.2013.875551

63. Pope HG Jr, Gruber AJ, Yurgelun-Todd D. Residual neuropsychologic effects of cannabis. *Curr Psychiatry Rep.* (2001) 3:507–12. doi: 10.1007/s11920-001-0045-7
64. Ravizza SM, Delgado MR, Chein JM, Becker JT, Fiez JA. Functional dissociations within the inferior parietal cortex in verbal working memory. *NeuroImage.* (2004) 22:562–73. doi: 10.1016/j.neuroimage.2004.01.039
65. Jacobsen LK, Pugh KR, Constable RT, Westerveld M, Mencl WE. Functional correlates of verbal memory deficits emerging during nicotine withdrawal in abstinent adolescent cannabis users. *Biol Psychiatry.* (2007) 61:31–40. doi: 10.1016/j.biopsych.2006.02.014
66. Jacobus J, Bava S, Cohen-Zion M, Mahmood O, Tapert S. Functional consequences of marijuana use in adolescents. *Pharmacol Biochem Behav.* (2009) 92:559–65. doi: 10.1016/j.pbb.2009.04.001
67. Rey A. L'examen psychologique dans les cas d'encéphalopathie traumatique (Les problems). *Arch Psychol.* (1941).
68. Harvey MA, Sellman JD, Porter RJ, Frampton CM. The relationship between non-acute adolescent cannabis use and cognition. *Drug Alcohol Rev.* (2007) 26:309–19. doi: 10.1080/09595230701247772
69. Bopp KL, Verhaeghen P. Age-related differences in control processes in verbal and visuospatial working memory: storage, transformation, supervision, and coordination. *J Gerontol Ser B Psychol Sci Soc Sci.* (2007) 62:P239–46. doi: 10.1093/geronb/62.5.P239
70. Fiore E, Borella E, Mammarella IC, De Beni R. Age differences in verbal and visuo-spatial working memory updating: evidence from analysis of serial position curves. *Memory.* (2012) 20:14–27. doi: 10.1080/09658211.2011.628320
71. Davis HP, Small SA, Stern Y, Mayeux R, Feldstein SN, Keller FR. Acquisition, recall, and forgetting of verbal information in long-term memory by young, middle-aged, and elderly individuals. *Cortex.* (2003) 39:1063–91. doi: 10.1016/S0010-9452(08)70878-5
72. Whitlow CT, Liguori A, Livengood LB, Hart SL, Mussat-Whitlow BJ, Lamborn CM, et al. Long-term heavy marijuana users make costly decisions on a gambling task. *Drug Alcohol Depend.* (2004) 76:107–11. doi: 10.1016/j.drugalcdep.2004.04.009
73. Crean RD, Tapert SF, Minassian A, MacDonald K, Crane NA, Mason BJ. Effects of chronic, heavy cannabis use on executive functions. *J Addict Med.* (2011) 5:9. doi: 10.1097/ADM.0b013e31820cdd57
74. Montgomery C, Seddon A, Fisk J, Murphy PN, Jansari A. Cannabis-related deficits in real-world memory. *Hum Psychopharmacol Clin Exp.* (2012) 27:217–25. doi: 10.1002/hup.1273
75. Lundqvist T. Cognitive consequences of cannabis use: comparison with abuse of stimulants and heroin with regard to attention, memory and executive functions. *Pharmacol Biochem Behav.* (2005) 81:319–30. doi: 10.1016/j.pbb.2005.02.017
76. Thomas H. A community survey of adverse effects of cannabis use. *Drug Alcohol Depend.* (1996) 42:201–7. doi: 10.1016/S0376-8716(96)01277-X
77. Troisi A, Pasini A, Saracco M, Spalletta G. Psychiatric symptoms in male cannabis users not using other illicit drugs. *Addiction.* (1998) 93:487–92. doi: 10.1046/j.1360-0443.1998.9344874.x
78. Reilly D, Didcott P, Swift W, Hall W. Long-term cannabis use: characteristics of users in an Australian rural area. *Addiction.* (1998) 93:837–46. doi: 10.1046/j.1360-0443.1998.9368375.x
79. Patton GC, Coffey C, Carlin JB, Degenhardt L, Lynskey M, Hall W. Cannabis use and mental health in young people: cohort study. *BMJ.* (2002) 325:1195. doi: 10.1136/bmj.325.7374.1195

**Conflict of Interest:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2021 Lorenzetti, Takagi, van Dalen, Yücel and Solowij. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.