



OPEN ACCESS

EDITED BY

Linda J. Larson-Prior,
University of Arkansas for Medical
Sciences, United States

REVIEWED BY

Barbara Galland,
University of Otago, New Zealand
Nanloh Samuel Jimam,
University of Jos, Nigeria

*CORRESPONDENCE

Yanhui Chen
yanhui_0655@163.com
Xiaoxia Lin
linxia1230@sina.com

SPECIALTY SECTION

This article was submitted to
Sleep Disorders,
a section of the journal
Frontiers in Psychiatry

RECEIVED 06 July 2022

ACCEPTED 20 October 2022

PUBLISHED 10 November 2022

CITATION

Chen X, Xu P, Chen Y, Chen S, Yao Y
and Lin X (2022) Validation of the
sleep disturbance scale for children
(SDSC) in infants and toddlers from
mainland China.
Front. Psychiatry 13:987304.
doi: 10.3389/fpsy.2022.987304

COPYRIGHT

© 2022 Chen, Xu, Chen, Chen, Yao
and Lin. This is an open-access article
distributed under the terms of the
[Creative Commons Attribution License
\(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or
reproduction in other forums is
permitted, provided the original
author(s) and the copyright owner(s)
are credited and that the original
publication in this journal is cited, in
accordance with accepted academic
practice. No use, distribution or
reproduction is permitted which does
not comply with these terms.

Validation of the sleep disturbance scale for children (SDSC) in infants and toddlers from mainland China

Xianrui Chen¹, Ping Xu², Yanhui Chen^{3*}, Shan Chen³,
Yonghua Yao⁴ and Xiaoxia Lin^{3*}

¹Department of Pediatric Rehabilitation, Xiamen Rehabilitation Hospital, Xiamen, China, ²Fujian Family Planning Association, Fuzhou, China, ³Department of Pediatrics, Fujian Medical University Union Hospital, Fuzhou, China, ⁴Pediatric Key Laboratory of Xiamen, Department of Pediatrics, The First Affiliated Hospital of Xiamen University, School of Medicine, Institute of Pediatrics, Xiamen University, Xiamen, China

The sleep disturbance scale for children (SDSC) has been validated in the population of Chinese children (aged 5–16 years) and has good psychometric properties. However, valid assessment tools for sleep problems in the population of infants and young children from mainland China are relatively scarce, and rates of screening and management for pediatric sleep problems are low.

Objective: This study is to evaluate the reliability and validity of SDSC among infants (aged 6–36 months), and to provide a reference for expanding the application of the SDSC for Chinese infants (SDSC-I).

Materials and methods: From April to November 2021, parents of infants from Fuzhou, Quanzhou, Longyan, Sanming, and Nanping cities in Fujian Province, China completed the SDSC-I. Several factor analyses were performed to evaluate the reliability and validity of the scale.

Results: Of note, 432 out of 469 samples were valid. After item selections and exploratory factor analyses, the SDSC-I concluded six dimensions (disorders of initiating sleep, disorders of maintaining sleep, sleep hyperhidrosis, sleep breathing disorders, parasomnias, and non-restorative sleep and excessive somnolence) with 23 items. The Cronbach's α coefficient of the scale was 0.863, and those for the six dimensions were within 0.576–0.835. The values of parameters for content validity of the scale were: IR = 0.87, I-CVI > 0.78, Kappa value > 0.74, S-CVI/UA = 0.87, S-CVI/Ave = 0.98. Principal component analysis revealed that the Kaiser-Meyer-Olkin (KMO) value was 0.84, and the factor loading of items ranged from 0.328 to 0.849, with six factors of eigenvalue more than one, which could explain 58.274% of the total variance. The confirmatory factor analysis results showed that χ^2/DF was 3.556, root-mean-square error of approximation (RMSEA) was 0.077, comparative fit index (CFI) was 0.809, and standardized RMR (SRMR) was 0.070.

Conclusion: Our study provides evidence that the SDSC-I is reliable and valid, and it is effective for the screening and management of sleep disturbances among infants (aged 6–36 months). Compared with other questionnaires such as the Brief Infant Sleep Questionnaire (BISQ), it is worthy of popularization and application in pediatric primary care.

KEYWORDS

sleep, Chinese version, sleep disturbance scale for children, infants, toddlers

Introduction

Sleep in children is an essential and dynamic process that affects numerous aspects of health and development. It has numerous functions, including growth, development, immune strengthening, memory, learning, regulation of behavior, and emotion. Several important changes in development during the first years of life lead to expected patterns of sleep and wakefulness in adults. Sleep problems are relatively common but they often can be challenging to recognize (1–3). A meta-analysis showed that the prevalence of sleep problems for children in mainland China was 37.6% (95% CI: 34.3–40.9%), with 33.3% in infants, 38.9% in preschoolers, and 43.7% in school-age children (4). Infancy and early childhood are critical periods for the development of sleep and the formation of healthy sleep patterns and habits in children. Few pediatricians in mainland China have received formal training in sleep medicine during medical school or residency. The practitioners have been shown to inadequately address sleep issues during routine well-child healthcare and to underdiagnose sleep problems. As a result, pediatricians are not able to effectively and confidently screen patients for sleep problems and provide sleep-related counseling. Therefore, early screening for sleep problems in infants and toddlers can help to correctly offer the diagnosis and the treatment of sleep disorders.

Since Sen et al. (5) conducted a comprehensive review of the overall deficiencies of pediatric questionnaires in sleep in 2011, many existing tools have been further developed and

evaluated by sleep researchers worldwide. There are 144 “tool”-studies (70 tools) published aiming at investigating sleep in primarily 6–18 years old per parental report (5). In 1996, Bruni et al. (6) developed a 26-item Likert-type rating scale (Sleep Disturbance Scale for Children: SDSC) used and validated in an Italian population of 1,304 children. The SDSC appears to be a useful tool for assessing the sleep disturbances of school-age children in both clinical and non-clinical populations. In 2013, they used it to assess the psychometric properties of the SDSC in a population of healthy preschool children (3–6 years old) in Italian. This further demonstrated that the SDSC had good reliability and validity for the application in studies of sleep in young children (7). In the same year, Huang et al. (8) translated the SDSC and determined its reliability and validity in 3,525 children aged 5–16 years. The results showed that its psychometric properties were also satisfactory as original, and indicated that it could be used as an epidemiological screening tool for parent-reported sleep disorders in Chinese school-age children. In 2019, Lecuelle et al. (9) first validated the SDSC for French preschoolers (aged from 6 months to 4 years), and the convergent validity and the internal reliability of a 22-item SDSC for young children (SDSC-Y) were acceptable. Subsequently, Romeo et al. (10) applied a 22-item version of the SDSC to a convenience sample of 193 Italian infants and toddlers (aged 6–36 months), and a 19-item questionnaire was eventually established for the evaluation of sleep problems.

It is believed that the factors affecting children's sleep are inconsistent, and there is no clear definition of sleep disorders in children. The concepts of sleep disorders and sleep problems are also vague (4, 8). Despite that the studies on children's sleep in mainland China have gradually increased over the past 20 years, valid assessment tools for pediatric sleep problems are few, and rates of screening and management are low. The SDSC was a parent-report scale for screening sleep disturbances within the preceding 6 months, but it now has been validated and adapted in sleep studies in different languages such as Australian (11), Persian (12), Indonesian (13), Flemish (14), Brazilian Portuguese (15), Portuguese (16), Finnish (17), Malay (18), and Turkish (19). In 2017, the “Guideline for Sleep Hygiene among Children Aged 0–5 Years” was published in mainland China (20), recommending the “Brief Infant Sleep

Abbreviations: SDSC, sleep disturbance scale for children; SDSC-I, SDSC for the Chinese infants; EFA, exploratory factor analysis; CFA, confirmatory factorial analyses; SDSC-Y, a French version of the SDSC adapted for preschool (6-month to 4-year-old) children; BISQ, Brief Infant Sleep Questionnaire; CSHQ, Children's Sleep Habits Questionnaire; IR, inter-rater reliability; I-CVI, item-content validity index; S-CVI, scale level-content validity index; CFI, comparative fit index; RMSEA, root-mean-square error of approximation; SRMR, standardized RMR; DIS, disorders of initiating sleep; DMS, disorders of maintaining sleep; SHY, sleep hyperhidrosis; SBD, sleep breathing disorders; PARA, parasomnias; NRSES, non-restorative sleep and excessive somnolence; ICSD-3, the third edition of the international classification of sleep disorders; KMO, Kaiser-Meyer-Olkin; SPSS, Statistical Product and Service Solutions.

TABLE 1 Characteristics of the sample ($n = 432$).

| Characteristic | | <i>n</i> | % |
|-------------------------|-------------------------------|----------|-------|
| Region | Rural | 226 | 52.30 |
| | Urban | 206 | 47.70 |
| Household income | <10,000 yuan | 37 | 8.60 |
| | 10,000–5,000 yuan | 120 | 27.80 |
| | 50,000–100,000 yuan | 160 | 37.00 |
| | >100,000 yuan | 115 | 26.60 |
| Mother's education | Primary school and lower | 90 | 20.80 |
| | Junior high school | 92 | 21.30 |
| | High school or junior college | 246 | 56.90 |
| | College or university | 4 | 0.90 |
| Mother's work | Unemployed | 125 | 28.90 |
| | Full-time | 270 | 62.50 |
| | Part-time | 37 | 8.60 |
| Housing area per capita | <10 m ² | 34 | 7.90 |
| | 10–30 m ² | 231 | 53.50 |
| | >30 m ² | 167 | 38.70 |

Questionnaire (BISQ) (21), and the “Children’s Sleep Habits Questionnaire (CSHQ)” (22) as screening tests for children’s sleep. Nevertheless, BISQ was mainly used to assess children’s sleep, such as sleep ecology, infant daytime sleep, nocturnal sleep patterns, bedtime, hours of nighttime, and the number of night wakings (23).

Therefore, this present study was primarily to assess the psychometric properties of the SDSC in a Chinese population of infants and toddlers (6–36 months), which had the potential to assist pediatricians in early screening and diagnosis of sleep disturbances in young children at community hospitals.

Materials and methods

Sampling

For from April to November 2021, several community hospitals in Fuzhou, Quanzhou, Longyan, Sanming, and Nanping cities in Fujian Province were selected for sampling using a convenience sampling method. Ethical approval was obtained by the Scientific Research Ethics Committee of Fujian Medical University Union Hospital (approval date and decision number: 2021KY131). And informed consent was acquired from all the parents of the infants who participated in our study.

Participants

Parents of healthy infants and toddlers aged 6 months to 3 years were selected to exclude children with psycho-behavioral developmental abnormalities, rheumatological and

TABLE 2 Item analysis for the Chinese version of SDSC-I ($n = 432$).

| Item | High group | Low group | <i>T</i> | <i>P</i> |
|------|-------------|-------------|----------|----------|
| D1 | 1.61 ± 0.63 | 1.27 ± 0.48 | 4.85 | <0.001 |
| D2 | 3.30 ± 1.09 | 1.95 ± 0.77 | 11.56 | <0.001 |
| D3 | 2.99 ± 1.17 | 1.47 ± 0.64 | 13.14 | <0.001 |
| D4 | 3.12 ± 1.15 | 1.35 ± 0.53 | 16.04 | <0.001 |
| D5 | 4.76 ± 0.59 | 3.28 ± 1.58 | 9.98 | <0.001 |
| D6 | 4.36 ± 1.12 | 2.10 ± 1.47 | 13.93 | <0.001 |
| M7 | 2.80 ± 1.30 | 1.45 ± 0.64 | 10.73 | <0.001 |
| M8 | 2.05 ± 1.04 | 1.09 ± 0.28 | 10.35 | <0.001 |
| M9 | 4.26 ± 0.88 | 2.26 ± 1.24 | 14.95 | <0.001 |
| H10 | 3.03 ± 1.28 | 1.29 ± 0.58 | 14.27 | <0.001 |
| H11 | 2.67 ± 1.28 | 1.19 ± 0.45 | 12.56 | <0.001 |
| B12 | 1.39 ± 0.75 | 1.00 ± 0.00 | 6.04 | <0.001 |
| B13 | 1.15 ± 0.50 | 1.00 ± 0.00 | 3.47 | 0.001 |
| B14 | 1.97 ± 1.02 | 1.20 ± 0.42 | 7.99 | <0.001 |
| P15 | 1.99 ± 1.01 | 1.21 ± 0.41 | 8.26 | <0.001 |
| P16 | 2.38 ± 0.94 | 1.30 ± 0.49 | 11.65 | <0.001 |
| P17 | 1.86 ± 0.91 | 1.23 ± 0.46 | 7.02 | <0.001 |
| P18 | 1.48 ± 0.78 | 1.10 ± 0.33 | 5.23 | <0.001 |
| N19 | 1.85 ± 1.13 | 1.16 ± 0.56 | 6.25 | <0.001 |
| N20 | 1.70 ± 0.86 | 1.04 ± 0.23 | 8.56 | <0.001 |
| N21 | 1.52 ± 0.84 | 1.02 ± 0.15 | 6.60 | <0.001 |
| N22 | 1.68 ± 0.86 | 1.09 ± 0.32 | 7.38 | <0.001 |
| N23 | 1.55 ± 0.78 | 1.21 ± 0.51 | 4.24 | <0.001 |

immune diseases, oncological diseases, or the use of medications known to cause sleepiness (e.g., anti-epileptics, antihistamines, and benzodiazepines). Children with developmental, physical, or mental disabilities, or receiving medications such as antihistaminic drugs, antiepileptic drugs, glucocorticoids, and melatonin, were excluded as well, as these factors may affect sleep architecture including sleep cycles of rapid eye movement (REM) and non-rapid eye movement (NREM) sleep. The participants were approached face to face and all the responses were collected digitally.

The SDSC for the Chinese infants based on the sleep disturbance scale for children

The SDSC adapted in the present study was authorized by Oliviero Bruni. The translation and the adaptation of the scale were performed independently by six senior experts, including a pediatric neurologist, a healthcare physician, a pediatric psychiatrist, a pediatric psychiatrist, an English language specialist, and a nurse. The entries were revised, screened, and evaluated for validation. All specialists independently provided a clarity score between one and four for each item to evaluate the scale’s content validity. Drawing on previous

TABLE 3 Content validity analysis of each item for the Chinese version of SDSC-I ($n = 432$).

| Item | Expert rating (score) | | | | | | I-CVI | Kappa |
|------|-----------------------|---|---|---|---|---|-------|-------|
| | A | B | C | D | E | F | | |
| D1 | 4 | 3 | 4 | 4 | 4 | 4 | 1.00 | 1.00 |
| D2 | 4 | 4 | 4 | 4 | 4 | 4 | 1.00 | 1.00 |
| D3 | 3 | 3 | 4 | 3 | 3 | 4 | 1.00 | 1.00 |
| D4 | 4 | 4 | 3 | 3 | 3 | 4 | 1.00 | 1.00 |
| D5 | 3 | 3 | 4 | 2 | 4 | 4 | 0.83 | 0.81 |
| D6 | 3 | 2 | 4 | 3 | 4 | 3 | 0.83 | 0.81 |
| M7 | 4 | 3 | 4 | 3 | 4 | 3 | 1.00 | 1.00 |
| M8 | 4 | 4 | 4 | 3 | 4 | 3 | 1.00 | 1.00 |
| M9 | 4 | 4 | 4 | 4 | 4 | 3 | 1.00 | 1.00 |
| H10 | 4 | 4 | 4 | 4 | 3 | 3 | 1.00 | 1.00 |
| H11 | 4 | 4 | 4 | 3 | 3 | 4 | 1.00 | 1.00 |
| B12 | 3 | 3 | 4 | 4 | 3 | 3 | 1.00 | 1.00 |
| B13 | 3 | 3 | 4 | 4 | 4 | 3 | 1.00 | 1.00 |
| B14 | 4 | 4 | 4 | 3 | 4 | 3 | 1.00 | 1.00 |
| P15 | 4 | 4 | 4 | 2 | 4 | 4 | 0.83 | 0.81 |
| P16 | 4 | 4 | 4 | 3 | 4 | 4 | 1.00 | 1.00 |
| P17 | 4 | 4 | 4 | 3 | 4 | 4 | 1.00 | 1.00 |
| P18 | 4 | 3 | 4 | 4 | 3 | 4 | 1.00 | 1.00 |
| N19 | 4 | 4 | 4 | 3 | 4 | 4 | 1.00 | 1.00 |
| N20 | 4 | 4 | 4 | 3 | 4 | 4 | 1.00 | 1.00 |
| N21 | 3 | 3 | 4 | 3 | 4 | 3 | 1.00 | 1.00 |
| N22 | 4 | 4 | 4 | 3 | 3 | 4 | 1.00 | 1.00 |
| N23 | 4 | 4 | 4 | 3 | 4 | 4 | 1.00 | 1.00 |

sleep questionnaires, particularly the recent validation of the Italian SDSC (a 19-item version) and the French SDSC-Y (a 22-item version), we finally removed two items (sleepwalking and hypnagogic hallucinations). Meanwhile, the two items “sleep terrors” and “nightmares” were combined into one (sleep terrors or nightmares). And an item (Need of parental accompaniment for sleep) was added, resulting in a 23-item questionnaire. This questionnaire investigates the occurrence of sleep disturbance during the preceding 6 months and each item is scored on a 5-point Likert scale. A sociodemographic data form was filled out by the parents, which included the date of birth, gender, developmental history, and medical history (i.e., past or present diseases, therapies, etc.) of the children. All the respondents were the mothers of the children.

Statistical analysis

The normality of the data was examined by using the Shapiro–Wilks test. Arithmetic means \pm standard deviations of item scores were reported as the previous study reported (10). Item analysis was used to evaluate the appropriateness of the items. The subjects were ranked in descending order

of their total scores. The content validity of the scale was evaluated using inter-rater reliability (IR), item-content validity index (I-CVI), scale-content validity index (S-CVI), universal agreement S-CVI (S-CVI/UA), average S-CVI (S-CVI/Ave), and Kappa value (24–26). The reliability analysis of the scale was evaluated by Cronbach’s alpha, inter-item correlation, and internal correlation consistency. The T-score table was prepared similarly to that in the original study by converting the mean scores of the scale into T-scores using the following formula: $T\text{-score} = 50 + (\text{value} - \text{mean}) / \text{standard deviation} * 10$. A child with a T-score of more than 70 was considered to have a significant sleep disorder (14). Confirmatory factorial analyses (CFA) were performed by a structural equation model. Comparative fit index (CFI) value > 0.80 was considered a moderate fit (27). The values of root-mean-square error of approximation (RMSEA) and standardized RMR (SRMR) < 0.08 were accept (28). The data analysis was performed using the Statistical Product and Service Solutions (SPSS) version 21.0 and Mplus version 8.3 software, and $p < 0.05$ was considered to be statistically significant.

Results

Participants

The ages of the 432 children ranging from 6 months to 3 years, and the average was 2 years and 1 month \pm 7 months. The mean age for 242 boys (56.02% of total) was 2 years with a standard deviation of 6 months, and the mean age for 190 girls (43.98% of total) was 2 years with a standard deviation of 7 months. A total of 188 of the participants (43.5%) were infants younger than 24 months. They were from a non-deprived population relative to the norm because the average household income of our population was more than 4,000 yuan/year. Other characteristics of the sample are listed in Table 1.

Item analysis

Based on the total scores, the critical ratio method was used to divide 27% into the “high” group ($n = 132$, ≥ 48 points, numbered 2) and the “low” group ($n = 129$, ≤ 37 points, numbered 1), and t -tests were conducted to determine the differences between the two groups on each entry score. The results showed statistically significant differences between the high and the low subgroups ($p < 0.05$, see Table 2), indicating a strong differentiation in each dimension.

Content validity

The results showed as follows: the inter-rater reliability of the SDSC-I = 0.87, I-CVI > 0.78 , Kappa values > 0.74 ,

TABLE 4 Item-total correlation analysis and exploratory factor analysis of the SDSC-I ($n = 432$).

| | Item-total correlation | Alpha if deleted | Variance explained (%) | Factor loading |
|--|------------------------|------------------|------------------------|----------------|
| Factor 1: Disorders of initiating sleep (DIS) | | | 11.747 | |
| D1. Sleep duration | 0.155 | 0.839 | | 0.463 |
| D2. Sleep latency | 0.423 | 0.830 | | 0.328 |
| D3. Going to bed reluctantly | 0.511 | 0.826 | | 0.679 |
| D4. Difficulty in falling asleep | 0.610 | 0.821 | | 0.721 |
| D5. Need of parental accompaniment for sleep | 0.351 | 0.835 | | 0.849 |
| D6. Anxiety when falling asleep alone | 0.412 | 0.835 | | 0.817 |
| Factor 2: Disorders of Maintaining Sleep (DMS) | | | 11.129 | |
| M7. Night awakenings more than twice | 0.431 | 0.830 | | 0.586 |
| M8. Difficulty in falling asleep after waking | 0.518 | 0.827 | | 0.660 |
| M9. Nocturnal hyperkinesia | 0.473 | 0.829 | | 0.419 |
| Factor 3: Sleep Hyperhidrosis (SHY) | | | 10.994 | |
| H10. Falling asleep sweating | 0.473 | 0.828 | | 0.837 |
| H11. Night sweating | 0.497 | 0.827 | | 0.824 |
| Factor 4: Sleep Breathing Disorders (SBD) | | | 9.061 | |
| B12. Breathing problems | 0.344 | 0.834 | | 0.765 |
| B13. Sleep apnea | 0.229 | 0.837 | | 0.806 |
| B14. Snoring | 0.418 | 0.831 | | 0.603 |
| Factor 5: Parasomnias (PARA) | | | 8.112 | |
| P15. Startles or jerks while falling asleep | 0.456 | 0.830 | | 0.668 |
| P16. Sleep terrors or nightmares | 0.551 | 0.826 | | 0.735 |
| P17. Sleep talking | 0.368 | 0.833 | | 0.545 |
| P18. Sleep bruxism | 0.257 | 0.836 | | 0.421 |
| Factor 6: Non-restorative sleep and excessive somnolence (NRSES) | | | 7.232 | |
| N19. Unusually difficult to wake up in the morning | 0.282 | 0.836 | | 0.638 |
| N20. Feeling tired with non-restorative sleep | 0.445 | 0.831 | | 0.787 |
| N21. Sleep paralysis | 0.387 | 0.833 | | 0.773 |
| N22. Daytime somnolence | 0.381 | 0.833 | | 0.664 |
| N23. Sleep attacks | 0.276 | 0.845 | | 0.543 |

S-CVI/UA = 0.87, and S-CVI/Ave = 0.98. All values met the requirements. The content validity of the scale was desirable, presented in [Table 3](#).

Reliability analysis

The Cronbach's alpha value for the scores of 23 items was 0.838, indicating good reliability of the scale. The Cronbach's alpha value for all dimensions ranged from 0.576 to 0.835. Except for Item 1 and Item 24, item-total correlations were greater than 0.3. After eliminating Item 1 or Item 24, the alpha was unchanged (see [Table 4](#)).

A Shapiro–Wilks test of the total scale and subscale scores demonstrated a non-normal distribution. Non-parametric Spearman correlations confirmed these results. All correlations between the two subscales (DIS, DMS, SHY, SBD, PARA, and NRSES) of the SDSC-I were smaller than 0.40, except for the correlation between DIS and DMS (0.525). Between the total

scores and the subscales, a strong correlation was observed ranging from 0.501 (SBD) to 0.820 (DIS).

Construct validity

This scale was suitable for exploratory factor analysis (EFA) using geomin rotation, with a Kaiser–Meyer–Olkin (KMO) value of 0.82 and Bartlett's test of sphericity of χ^2 value of 3,088.36 ($p < 0.001$). Based on the eigenvalues, a six-factor solution explained 58.27% of the total variance. The factor loads of all the items varied between 0.328 (Item 2) and 0.849 (Item 5). The 23-item scale, the factor loadings, and the six factors were depicted in [Figure 1](#). In the model described in CFA, values were determined as follows: $\chi^2/DF = 3.556$, CFI = 0.809, SRMR = 0.070, and the RMSEA = 0.077. The factor loadings of the 23 items were from 0.22 to 0.87 (see [Figure 1](#)). The factor structure of the Chinese SDSC in infants was consistent with previous studies and appeared to be compatible.

Scale score distribution

The minimum score of the scale was 23, and the minimum scores for the subscales of DIS, DMS, SHY, SBD, PARA, and NRSES were 6, 3, 2, 3, 4, and 5, respectively. The distribution of SDSC scores showed the maximum scores of the total scale and subscales were 78, 27, 15, 10, 11, 15, and 20, respectively. The single factor scores showed the percentages of the sample with minimum and maximum values ranging from 0.20 to 0.20% for DIS, 0.70 to 0.70% for DMS, 6.5 to 0.20% for SHY, 37.5 to 3.5% for SBD, 58.3 to 0.20% for PARA, 22.2 to 0.70% for NRSES, and 48.8 to 0.20% for T-score. All the total scores demonstrated left-tailed distribution (Wilk's $W = 0.973$, $p = 0.000$) (see Figure 2). Differences in total scores by age and gender were not found.


Other measurements

As previously done in similar studies (10), most of the infants (62.0%) slept more than 9 h per night and none slept less than 5 h. A total of 59.7% of the young children fell asleep 30 min after going to bed. The highest frequency of parent-reported sleeping disturbance was Item 5 (77.3%), and the lowest ones were Item 1 (0.2%) and Item 13 (0.2%) (see Table 5). The analysis of the participant's compliance and the completion time of the scale showed that most of the parents chose to fill out the questionnaire issued in this study and completed it within 10–15 min, and the recovery rate of the questionnaire was over 90%.

Discussion

The main objective of the present study was to assess sleep disturbance in Chinese infants and toddlers using the SDSC. The validity and reliability were investigated by a modified version of SDSC-I. The 23-item SDSC-I consisted of six factors (DIS, DMS, SHY, SBD, PARA, and NRSES) and was highly reliable and compatible.

Our study demonstrated that there was a strong correlation between each dimension of the SDSC-I scale and the total score, with all the correlations more than 0.5. It indicated that the factors (DIS, DMS, SHY, SBD, PARA, and NRSES) were consistent with the clinical concept, as previously reported (9, 10). Except for Item 1, the correlation coefficient values were all greater than 0.5 between each item and the corresponding dimension. Therefore, our study showed that the dimensions in which the entries were located were reasonable and suggested that the scale had good construct validity. According to the exploratory factor analyses, six factors with eigenvalues greater than one were extracted, which was basically consistent with



the findings from the Italian and French scholars (9, 10). And all six areas of sleep disorders based on clinical concepts and previous studies were helpful for clinicians to identify the areas that require more in-depth investigations (10). All of the factor loadings of items ranged from 0.328 to 0.849, which was similar to the results (from 0.26 to 0.91) in the previous French study and slightly lower than the results (from 0.50 to 0.87) in the Italian study. Six-factor model in the present study matched the theoretical constructs of the six dimensions designed by the SDSC-I, indicating that the scale had good structural validity.

In the present study, it was the first time that the SDSC was applied to infants and toddlers within 3 years old in mainland China. Sleep problems are prevalent in Chinese preschool and school-age children, including difficulties in falling asleep, enuresis, night waking, teeth grinding, sleep breathing disorder, night terrors, open mouth breathing, sleep talking, nightmares, night sweats, and daytime sleepiness (29). The existing sleep assessment tools for children are mainly applicable to preschool or school-age children, and most of them are limited to the




FIGURE 2
Distribution of the SDSC-I total score.

assessment of one specific sleep problem but cannot accurately and comprehensively reflect the sleep problems of infants in China (30). The clinical evidence demonstrated that pediatric sleep disturbances overlapped with each other (10). The SDSC not only evaluates sleep duration and sleep behavior habits but also assesses sleep disorder-related problems, which is more beneficial for the early screening of sleep disorders in clinical practice. The SDSC is currently used in China to only evaluate school-age children aged from 5 to 16 years (8), but it has been used to assess preschool-age children or infants in other countries (9, 10). The validity and reliability of the Chinese version of SDSC for children in mainland China first conducted by Huang et al. (8) revealed a good internal consistency (Cronbach's $\alpha = 0.81$). Our study showed that the SDSC-I scale had an acceptable and stable internal consistency with a Cronbach's α coefficient of 0.847, slightly higher than the result of previous research from Chinese scholars (8), but it was similar to the French population and the Italian population (9, 10). The internal consistency of all the dimensions was observed across infants and school-age children in China (8). Therefore, this study indicated that the SDSC-I had good reliability for evaluating sleep disturbance in Chinese infants.

The current international classification of sleep disorders is based on the "The Third Edition of the International Classification of Sleep Disorders (ICSD-3)" released in 2014, identifying seven categories (31) including insomnia disorders, sleep-related breathing disorders, central disorders of hypersomnolence, circadian rhythm sleep-wake disorders, parasomnias, sleep-related movement disorders, and other sleep disorders. Insomnia in children is usually reported by parents or other caregivers and is mainly characterized by going to bed reluctantly, frequent awakenings during the night, and difficulty falling asleep alone. In addition to nocturnal symptoms, daytime symptoms of functional impairment such as fatigue, irritability, hyperactivity and impulsivity, and daytime sleepiness may also result from night sleep difficulties (32). The 23-item version of the SDSC in our study was also an adapted version of the original SDSC as Lecuelle et al. (9) and Romeo et al. reported (10). In the SDSC-I, we removed three items regarding sleepwalking and hypnagogic hallucinations. Bruxism, on the other hand, was retained in our study due to the results of research on sleep problems in children in mainland China (4, 29, 30). Because, even though bruxism is reported to be poorly prevalent in French and Italian infants and toddlers, it is common among infants and young children in mainland China (29). Furthermore, in French, rhythmic movement

TABLE 5 Frequencies (%) of parent-reported sleeping disturbances in infants ($n = 432$).

| Item | Never | Occasionally 1–2x/month | Sometimes 1–2x/week | Often 3–5x/week | Always |
|------|--------|-------------------------|---------------------|-----------------|--------|
| D3 | 32.200 | 37.300 | 16.900 | 8.600 | 5.100 |
| D4 | 33.600 | 37.300 | 15.300 | 9.300 | 4.600 |
| D5 | 6.900 | 8.800 | 6.900 | 14.800 | 62.500 |
| D6 | 23.400 | 14.800 | 8.300 | 12.700 | 40.700 |
| M7 | 37.300 | 38.900 | 9.700 | 9.700 | 4.400 |
| M8 | 64.100 | 25.000 | 7.900 | 2.300 | 0.700 |
| M9 | 11.100 | 22.200 | 12.300 | 30.600 | 23.800 |
| H10 | 42.400 | 26.900 | 13.900 | 9.000 | 7.900 |
| H11 | 50.200 | 27.800 | 10.600 | 7.400 | 3.900 |
| B12 | 88.200 | 7.600 | 2.800 | 0.900 | 0.500 |
| B13 | 95.600 | 3.000 | 1.200 | 0.200 | 0.000 |
| B14 | 62.000 | 28.000 | 6.700 | 2.300 | 0.900 |
| P15 | 55.800 | 35.600 | 4.900 | 3.000 | 0.700 |
| P16 | 40.500 | 44.400 | 10.600 | 4.200 | 0.200 |
| P17 | 58.100 | 32.900 | 6.900 | 2.100 | 0.000 |
| P18 | 78.900 | 16.400 | 3.500 | 1.200 | 0.000 |
| N19 | 73.600 | 14.400 | 7.200 | 3.000 | 1.900 |
| N20 | 78.200 | 15.700 | 4.600 | 1.400 | 0.000 |
| N21 | 85.200 | 10.900 | 3.000 | 0.500 | 0.500 |
| N22 | 75.900 | 18.800 | 4.200 | 0.900 | 0.200 |
| N23 | 74.100 | 19.200 | 5.600 | 1.200 | 0.000 |

disorders and hypnic jerks are frequently misinterpreted by parents as that the child is agitated or turning several times in bed when unable to fall asleep (9), but parents, as well as researchers of sleep problems in mainland China, are more concerned about children's frequency of awakenings, restless sleep, leg movements, bruxism, hyperhidrosis, snoring, night awakening, apnea, choking or gasping, mouth breathing, sleep talking, nightmare, enuresis, trouble falling asleep, or other problems during sleep (4). For instance, Chinese parents of young children pay particularly close attention to bruxism as they often associate bruxism with ascariasis or other diseases. Therefore, bruxism was not removed after consultation with specialists and parents. Despite that the national and cultural differences led to slight inconsistency in the application of the SDSC, the scale still illustrated good applicability.

Compared with the previous study by Romeo et al. (10), 58.5% of infants in our group slept 9–11 h, which was higher than the Italian population (46.6%). A sleep latency shorter than 30 min to fall asleep was about 59.7% of the Chinese infants, rather than 54.5% of the Italian infants. The incidence of bedtime resistance and difficulty in falling asleep among Chinese infants were lower than those in Italian infants (13.6 vs. 20.9% and 13.2 vs. 16.6%, respectively). Chinese infants and toddlers were awakened less frequently during the night (13.6 vs. 20.9%) and presented nocturnal hyperkinesia (13.2 vs. 16.6%). Snoring was found similar in the Chinese and Italian

populations (5.7 vs. 4.8%). As for parasomnias, a prevalence of 3.1% of sleep talking was reported in our research while 3.5% in the Italian research and frequent twitching or jerking of legs were reported in 54.4% of Chinese infants, significantly higher than in Italian (24.0%).

The practice of parent and child sharing a sleeping surface, or “bed-sharing,” is one of the most controversial topics in parenting research. The lay literature has popularized and polarized this debate, offering claims of dangers and benefits, both physical and psychological, associated with bed-sharing (33). The bed-sharing practice is very prevalent at any age, which ranges from 69.9 to 78.3%. Most infants fall asleep while feeding or being rocked/held before age 12 months. By the age of 35 months, 62.4% of the children fall asleep in bed near their parents. The most common reasons for bed-sharing are breastfeeding/feeding and convenience. Parental involvement when falling asleep is significantly related to frequent night awakenings and difficulty falling asleep. Bed-sharing and parental involvement were very common among Chinese children aged <years (34). This may be due to the fact that Chinese parents prefer to share a bed with their children for sleep, as studies have shown that Chinese children usually need to be held or patted to sleep by their parents (35). The percentages of sweating while falling asleep and during sleep were respectively 16.9% (vs. 7.7% in Italian) and 11.3% (vs. 8.4% in Italian). It may be related to the high parental concern about night sweating or

excessive warmth of children in mainland China. Interestingly, parents of Chinese infants often associate excessive sweating with vitamin D deficiency or physical weakness (a symptom defined in traditional Chinese medicine). The percentage of T-score greater than 70 was higher than that in the Italian population (4.6 vs. 3.6%) and similar to that of Chinese school-age children in the previous study (4.1%). Our study showed that 10.9% of infants and toddlers were within the critical range of 61–70 points, while 12% in the Italian study were within that critical value. The influencing factors relevant to sleep disorders or sleep problems in children vary widely from study to study and may be closely related to sociocultural, climatic environment, and individual or family factors (29).

Several limitations of our study need to be acknowledged as follows: (a) the sensitivity and specificity of the scale could not be calculated in our study because of lacking a patient group with sleep disorders. Due to methodological issues, the rates of sleep disorders determined in this study cannot be used as epidemiological data (19); (b) the parents may lack awareness and attention to sleep problems because of social, familial, cultural, and environmental factors. Therefore, data collected from parental reports had a potential recall bias (8); and (c) our study also has some statistical validation limitations such as lacking test–retest fidelity, comparisons of different age groups, and multi-group modeling CFA (9). Nevertheless, the potential sleep-related problems can be evaluated by pediatricians by easily administrating, scoring, and interpreting the SDSC-I, which would be conducive to addressing the prevention of sleep disorders (8).

Conclusion

Our study provides evidence that SDSC-I is a reliable and valid version as qualified as in other languages, and it can assess the sleep problems of infants in a comprehensive and detailed manner. Therefore, this scale is useful for screening and detection of sleep disturbances in pediatric primary care and is worthy of popularization and application. This study is the first step to applying the SDSC to Chinese infants, and further research is needed to provide a more in-depth assessment of sleep disorders like polysomnography or actigraphy. Recommendations for improving sleep services in pediatric primary care are also needed in the areas of research, practice, and education.

Data availability statement

The original contributions presented in this study are included in the article/supplementary material, further inquiries can be directed to the corresponding authors.

Ethics statement

The studies involving human participants were reviewed and approved by the Fujian Medical University Union Hospital, Scientific Research Ethics Committee (approval date and decision number: 2021KY131). Written informed consent to participate in this study was provided by the participants' legal guardian/next of kin.

Author contributions

XC and XL participated in the design of the study and data collection and performed the statistical analysis. XL, XC, YC, YY, PX, and SC conceived of the study and participated in its design as well as coordination and helped to write and revise the manuscript. All authors read and approved the final version of the manuscript.

Funding

This work was supported by the Fujian Medical University Union Hospital Key Clinical Specialty Construction Project and the Joint Funds for the Innovation of Science and Technology, Fujian Province (No. 2018Y9056), China.

Acknowledgments

We thank all our colleagues working in the Department of Epidemiology and Health Statistics, School of Public Health of Central South University.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

References

1. Matricciani L, Paquet C, Galland B, Short M, Olds T. Children's sleep and health: a meta-review. *Sleep Med Rev.* (2019) 46:136–50. doi: 10.1016/j.smrv.2019.04.011
2. Carter JC, Wrede JE. Overview of sleep and sleep disorders in infancy and childhood. *Pediatr Ann.* (2017) 46:e133–8. doi: 10.3928/19382359-20170316-02
3. Rana M, Riffo Allende C, Mesa Latorre T, Rosso Astorga K, Torres AR. Sleep in children: physiology and update of a literature review. *Medicina.* (2019) 79:25–8.
4. Chen X, Ke ZL, Chen Y, Lin X. The prevalence of sleep problems among children in mainland China: a meta-analysis and systemic-analysis. *Sleep Med.* (2021) 83:248–55. doi: 10.1016/j.sleep.2021.04.014
5. Sen T, Spruyt K. Pediatric sleep tools: an updated literature review. *Front Psychiatry.* (2020) 11:317. doi: 10.3389/fpsy.2020.00317
6. Bruni O, Ottaviano S, Guidetti V, Romoli M, Innocenzi M, Cortesi F, et al. The sleep disturbance scale for children (SDSC). Construction and validation of an instrument to evaluate sleep disturbances in childhood and adolescence. *J Sleep Res.* (1996) 5:251–61. doi: 10.1111/j.1365-2869.1996.00251.x
7. Romeo DM, Bruni O, Brogna C, Ferri R, Galluccio C, De Clemente V, et al. Application of the Sleep Disturbance Scale for Children (SDSC) in preschool age. *Eur J Paediatr Neurol.* (2013) 17:374–82. doi: 10.1016/j.ejpn.2012.12.009
8. Huang MM, Qian Z, Wang J, Vaughn MG, Lee YL, Dong GH. Validation of the Sleep Disturbance Scale for Children and prevalence of parent-reported sleep disorder symptoms in Chinese children. *Sleep Med.* (2014) 15:923–8. doi: 10.1016/j.sleep.2014.03.023
9. Lecuelle F, Gustin MP, Leslie W, Mindell JA, Franco P, Putois B. French validation of the sleep disturbance scale for children (SDSC) in infants (aged 6 months to 4 years). *Sleep Med.* (2020) 67:56–65. doi: 10.1016/j.sleep.2019.09.014
10. Romeo DM, Cordaro G, Macchione E, Venezia I, Brogna C, Mercuri E, et al. Application of the sleep disturbance scale for children (SDSC) in infants and toddlers (6–36 months). *Sleep Med.* (2021) 81:62–8. doi: 10.1016/j.sleep.2021.02.001
11. Marriner AM, Pestell C, Bayliss DM, McCann M, Bucks RS. Confirmatory factor analysis of the sleep disturbance scale for children (SDSC) in a clinical sample of children and adolescents. *J Sleep Res.* (2017) 26:587–94. doi: 10.1111/jsr.12526
12. Saffari M, Gholamrezaei A, Saneian H, Attari A, Bruni O. Linguistic validation of the sleep disturbance scale for children (SDSC) in Iranian children with Persian language. *Sleep Med.* (2014) 15:998–1001. doi: 10.1016/j.sleep.2014.03.021
13. Herwanto H, Lestari H, Warouw SM, Salendu PM. Sleep disturbance scale for children as a diagnostic tool for sleep disorders in adolescents. *Paediatr Indones.* (2018) 8:133–7. doi: 10.14238/pi58.3.2018.133-7
14. Spruyt K, Cluydts R, Verleye GB. Pediatric sleep disorders: exploratory modulation of their 434 relationships. *Sleep.* (2004) 27:495–501. doi: 10.1093/sleep/27.3.495
15. Sousa VD, Rojjanasrirat W. Translation, adaptation and validation of instruments or scales for use in cross-cultural health care research: a clear and user-friendly guideline. *J Eval Clin Pract.* (2011) 17:268–74. doi: 10.1111/j.1365-2753.2010.01434.x
16. Ferreira VR, Carvalho LBC, Ruotolo F, de Moraes JE, Prado LBF, Prado GF. Sleep disturbance scale for children: translation, cultural adaptation, and validation. *Sleep Med.* (2009) 10:457–63. doi: 10.1016/j.sleep.2008.03.018
17. Simola P, Niskakangas M, Liukkonen K, Virkkula P, Pitkäranta A, Kirjavainen T, et al. Sleep problems and daytime tiredness in Finnish preschool-aged children—a community survey. *Child Care Health Dev.* (2010) 36:805–11. doi: 10.1111/j.1365-2214.2010.01112.x
18. Ong LC, Yang WW, Wong SW, alSiddiq F, Khu YS. Sleep habits and disturbances in Malaysian children with epilepsy. *J Paediatr Child Health.* (2010) 46:80–4. doi: 10.1111/j.1440-1754.2009.01642.x
19. Ađca S, Görker I, Turan FN, Öztürk L. Validity and reliability of the Turkish version of Sleep Disturbance Scale for Children. *Sleep Med.* (2021) 84:56–62. doi: 10.1016/j.sleep.2021.05.016
20. National Health and Family Planning Commission of the People's Republic of China. *Guideline for sleep hygiene among children aged 0 to 5 years:WS/T 579-2017[S].* (2017).
21. Sadeh A. A brief screening questionnaire for infant sleep problems: validation and findings for an Internet sample. *Pediatrics.* (2004) 113:e570–7. doi: 10.1542/peds.113.6.e
22. Owens JA, Spirito A, McGuinn M. The children's sleep habits questionnaire (CSHQ): psychometric properties of a survey instrument for school-aged children. *Sleep.* (2000) 23:1043–51.
23. Mindell JA, Gould RA, Tikotzy L, Leichman ES, Walters RM. Norm-referenced scoring system for the brief infant sleep questionnaire - revised (BISQ-R). *Sleep Med.* (2019) 63:106–14. doi: 10.1016/j.sleep.2019.05.010
24. Beaton DE, Bombardier C, Guillemin F, Ferraz MB. Guidelines for the process of cross-cultural adaptation of self-report measures. *Spine.* (2000) 25:3186–91. doi: 10.1097/00007632-200012150-00014
25. Lynn MR. Determination and quantification of content validity. *Nurs Res.* (1986) 35:382.
26. Polit DF, Beck C T. The content validity index: are you sure you know what's being reported? Critique and recommendations. *Res Nurs Health.* (2006) 29:489–97. doi: 10.1002/nur.20147
27. Hair J, Black W, Anderson R, Babin B. *Multivariate Data Analysis: A Global Perspective.* 7th ed. Upper Saddle River, NJ: Pearson (2010).
28. Brown TA. *Confirmatory factor Analysis for Applied Research.* New York, NY: Guilford Press (2006).
29. Chen YY, Zhou N. Research progress on sleep problems in preschool children in China. *Chin J Sch Health.* (2020) 41:1433–7. doi: 10.16835/j.cnki.1000-9817.2020.09.044
30. Feng WW, Pan XP, Wang HS, Zhang Y, Jin X, Huang XN, et al. Development and evaluation of the Chinese infants sleep assessment scales. *Chin J Child Health Care.* (2021) 29:468–72. doi: 10.11852/zgetbjzz2020-1376
31. Sateia MJ. International classification of sleep disorders-third edition: highlights and modifications. *Chest.* (2014) 146:1387–94. doi: 10.1378/chest.14-0970
32. American Academy of Sleep Medicine. *International Classification of Sleep Disorders.* 3rd ed. Darien, IL: American Academy of Sleep Medicine (2014).
33. Mileva-Seitz VR, Bakermans-Kranenburg MJ, Battaini C, Luijk MP. Parent-child bed-sharing: the good, the bad, and the burden of evidence. *Sleep Med Rev.* (2017) 32:4–27. doi: 10.1016/j.smrv.2016.03.003
34. Feng WW, Zhang Y, Wang HS, Pan XP, Jin X, Xu T, et al. Understanding the choice of sleep arrangements and soothing methods and their associations with sleep problems among children under 3 years old: a Chinese population-based study. *Biomed Environ Sci.* (2022) 35:225–33. doi: 10.3967/bes2022.031
35. Feng WW, Zhang Y, Wang HS, Pan XP, Huang XN, Xu T, et al. Development and evaluation of the Chinese toddler sleep assessment scale. *Chin J Child Health Care.* (2021) 29:1295–9. doi: 10.11852/zgetbjzz2021-0272