



OPEN ACCESS

EDITED BY

Susanne Fischer,
University of Zurich, Switzerland

REVIEWED BY

Badari Birur,
University of Alabama at Birmingham,
United States

*CORRESPONDENCE

Nanouk Bakker
✉ nanoukbakker@heelzorg.nl
Eline M. Vissia
✉ elinevissia@gmail.com

[†]These authors share first authorship

RECEIVED 15 September 2023

ACCEPTED 14 February 2024

PUBLISHED 28 February 2024

CITATION

Bakker N, Vissia EM, van den Dungen M,
Kraaij C, Tjink D and Boon S (2024)
Commentary: Schema therapy for
Dissociative Identity Disorder: a case report.
Front. Psychiatry 15:1292432.
doi: 10.3389/fpsy.2024.1292432

COPYRIGHT

© 2024 Bakker, Vissia, van den Dungen, Kraaij,
Tjink and Boon. This is an open-access article
distributed under the terms of the [Creative
Commons Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The
use, distribution or reproduction in other
forums is permitted, provided the original
author(s) and the copyright owner(s) are
credited and that the original publication in
this journal is cited, in accordance with
accepted academic practice. No use,
distribution or reproduction is permitted
which does not comply with these terms.

RETRACTED: Commentary: Schema therapy for Dissociative Identity Disorder: a case report

Nanouk Bakker^{1*†}, Eline M. Vissia^{1*†}, Maaïke van den Dungen²,
Christel Kraaij³, Desiree Tjink⁴ and Suzette Boon²

¹Heelzorg, Centre for Psychotrauma, Zwolle, Netherlands, ²Private Practitioner, Utrecht, Netherlands,
³Trauma and Transition, Doorn, Netherlands, ⁴GGz Centraal, Centre for Psychotherapy TRTC Transit,
Ermelo, Netherlands

KEYWORDS

dissociative identity disorder (DID), trauma, schematherapy, phase oriented trauma
treatment, dissociation

A Commentary on

Schema therapy for Dissociative Identity Disorder: a case report

By Bachrach N, Rijkeboer MM, Arntz A and Huntjens RJC (2023) *Front. Psychiatry* 14:1151872.
doi: 10.3389/fpsy.2023.1151872

Introduction

We sincerely appreciate new research on the treatment of dissociative identity disorder (DID), which adds important information on a relatively understudied diagnosis. More specifically, we value the new avenues Bachrach et al. (1) explore and describe the course of Schema Therapy (ST) in DID in more detail. We have some concerns and comments to raise, many of which relate to issues that have been discussed before (2, 3). These are combined with clinical consensus on how to treat individuals with DID.

Subsections

First, we would like to highlight that even though ST is presented as a new treatment for individuals with DID, there are many overlapping interventions with the guidelines for phase-oriented trauma treatment (POTT) (4). As described in this case report, ST interventions cover many stabilization techniques frequently described and used within POTT treatment, which are therefore not unique to ST. This also contradicts Bachrach et al.'s (1) notion that ST pays little attention to stabilization in the first phase of treatment. Furthermore, ST's biweekly sessions may involve a more intensive approach than POTT's standard weekly sessions (3). The total length of both treatments is, therefore, probably comparable. Bachrach et al. (1) also mistakenly present POTT as a solely practice-based psychodynamic psychotherapy approach. POTT is an eclectic treatment that incorporates various approaches such as psychodynamic psychotherapy, cognitive behavioural therapy,

systemic therapy, mindfulness, imagery rescripting, sensorimotor psychotherapy, and EMDR (5, 6). Both models focus on trauma and its effects when treating Dissociative Identity Disorder (DID), including reducing avoidance during the stabilisation phase. As for Bachrach et al.'s (1) statement that research results indicate that POTT effects are small or absent regarding the core symptoms, we disagree. The review by Brand et al. (7) suggests that dissociative disorder (DD) treatment is associated with improvement in symptoms of dissociation. In addition, in their TOP DD study (Treatment of Patients with Dissociative Disorders; 8), patients showed reduced manifestations of dissociated self-states, including subjective self-division and hearing voices of self-states. The last imprecise assumption is that the randomized controlled trial by Bækkelund et al. (9) would be exemplary for regular POTT; the Bækkelund study only covers a 20-session skill-based group treatment.

Secondly, we would like to stress our concerns regarding the intervention described by Bachrach et al. (1) in summoning a punitive part to leave. In DID treatment, close attention is given to the function of aggressive and critical dissociative parts while pacing and validating their needs (10, 11). This basic assumption holds for ST and POTT; therefore, rejecting the punitive part seems inappropriate.

In our regard, this is a gravely invalidating intervention, dismissing the patient's natural ability to cope with highly threatening circumstances. When appropriately acknowledged, these punitive parts become the patient's strongest allies in their healing process and in finding the strength to cope with daily life issues (e.g. being assertive). Removing punitive parts, if that is possible at all, would undermine the patients' development to establish a healthy balance between dependent and autonomous functioning. Furthermore, the expert clinical consensus is that critical or aggressive parts will not respond to being sent away by disappearing altogether in authentic DID. It is more likely that the working alliance between the therapist and the patient and between the patient and the punitive part becomes harmed and for the punitive part to reappear as time passes.

Moreover, the particular structure of the inner world of the patient described in this paper seems highly atypical for genuine DID. This makes us question the validity of the DID diagnosis given to this patient. In general, a dissociative system consists of more parts carrying trauma material, mostly child parts that are stuck in the past, portraying bodily behaviours matching the stress reactions (fight, flight, freeze and collapse), than parts that are highly avoidant of trauma content and emotions, which enables them to keep performing tasks in daily life. In the case described by Bachrach et al., (1) significantly more adult avoidant parts are reported than child parts. Diagnosing DID is a rather complex endeavour which needs sufficient clinical experience to differentiate between genuine DID and factitious cases of DID and cases where identity confusion is due to borderline personality disorder (BPD) (2, 12). The differential diagnosis of these conditions can be challenging and is crucial to ensure the reliability of DID diagnoses (4, 13).

Discussion

We aim to highlight the intricacies of treating Dissociative Identity Disorder (DID) and essential considerations. Some final questions about the ST study can therefore be added. The study is introduced as a case report to illustrate the application of an adapted form of ST for DID. In this respect, we would appreciate more detailed information on how "ST for DID is personalized to each patient" and what adaptations they made to ST for DID patients, a suggestion previously done by Brand et al. (2), in response to the original rationale and study protocol (14). Furthermore, we would appreciate a more comprehensive outline of how trauma-focused treatments are administered for sexual abuse cases and the level of expertise of the diagnosing clinician to ensure accuracy in this report. The presentation of baseline data in the absence of post-treatment measures, together with mentioning a "strong" reduction in dissociative symptoms, leaves us with the suggestion that these symptoms reflect an improvement in DID. It remains unclear, however, whether these reduced dissociative symptoms reflect core DID symptoms, as improvements may result from reduced comorbid BPD- and posttraumatic stress disorder symptoms, or the termination of cannabis usage since the use of this substance is associated with derealization and depersonalization (15). We are interested in whether the described patient's modi were integrated after the treatment or whether she merely learned to cope with them, at least for now. A longer follow-up is crucial as DID can have a relapsing and remitting course across the lifespan (2, 4). Overall, it is essential to continue research on DID treatment, and while we raised concerns as the above, our goal is to consolidate DID treatment and expand knowledge on how these severely traumatized individuals have the best chances to heal.

Author contributions

NB: Writing – original draft, Writing – review & editing. EV: Writing – original draft, Writing – review & editing. MvdD: Writing – review & editing. CK: Writing – review & editing. DT: Writing – review & editing. SB: Writing – review & editing.

Funding

The author(s) declare that no financial support was received for the research, authorship, and/or publication of this article.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated

organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

References

1. Bachrach N, Rijkeboer MM, Arntz A, Huntjens RJC. Schema therapy for Dissociative Identity Disorder: a case report. *Front Psychiatry*. (2023) 14:1151872. doi: 10.3389/fpsy.2023.1151872
2. Brand BL, Loewenstein RJ, Schielke HJ, van der Hart O, Nijenhuis ERS, Schlumpf YR, et al. Cautions and concerns about Huntjens et al.'s Schema Therapy for Dissociative Identity Disorder. *Eur J Psychotraumatol*. (2019) 10(1):1631698. doi: 10.1080/20008198.2019.1631698
3. Nijenhuis ERS, van der Hart O, Schlumpf YR, Vissia EM, Reinders AATS. Considerations regarding treatment efficiency, dissociative parts and dissociative amnesia for Huntjens et al.'s Schema Therapy for Dissociative Identity Disorder. *Eur J Psychotraumatol*. (2019) 10(1):1687081. doi: 10.1080/20008198.2019.1687081
4. International Society for the Study of Trauma and Dissociation. Chu, J.A., Dell, P.F., van der Hart, O., Cardena, E., Barach, P.M., Somer, E., Loewenstein, R.J., Brand, B., et al. Guidelines for treating dissociative identity disorder in adults, 3rd revision. *J Trauma Dissociation*. (2011) 12:115–8. doi: 10.1080/15299732.2011.537247
5. Van der Hart O, Groenendijk M, Gonzalez A, Mosquera D, Solomon R. Dissociation of the personality and EMDR therapy in complex trauma-related disorders: Applications in the stabilization phase. *J EMDR Pract Res*. (2013) 7:81–94. doi: 10.1891/1933-3196.7.2.81
6. Steele K, Boon S, van der Hart O. *Treating trauma-related dissociation: A practical, integrative approach*. New York City: W Norton & Co (2017).
7. Brand BL, Classen CC, McNary SW, Zaveri P. A review of dissociative disorders treatment studies. *J Nerv Ment Dis*. (2009) 197(9):646–54. doi: 10.1097/NMD.0b013e3181b3afaa
8. Brand B, Loewenstein RJ. Does phasic trauma treatment make patients with dissociative identity disorder treatment more dissociative? *J Trauma Dissociation*. (2014) 15:52–65. doi: 10.1080/15299732.2013.828150
9. Bækkelund H, Ulvenes P, Boon-Langelaan S, Arnevik EA. Group treatment for complex dissociative disorders: a randomized clinical trial. *BMC Psychiatry*. (2022) 22:1–13. doi: 10.1186/s12888-022-03970-8
10. Van der Hart O, Nijenhuis ERS, Steele K. *The haunted self: Structural dissociation and the treatment of chronic traumatization*. New York: Norton (2006).
11. Nijenhuis ERS. *The trinity of trauma: Ignorance, fragility, and control* Vol. III. . Göttingen: Vandenhoeck & Ruprecht (2017).
12. Draijer N, Boon S. The imitation of dissociative identity disorder: Patients at risk, therapists at risk. *J Psychiatry Law*. (1999) 27:423–58. doi: 10.1177/009318539902700304
13. Boon S. *Assessing Trauma-Related Dissociation*. New York: Norton Publishers (2023).
14. Huntjens RJC, Rijkeboer MM, Arntz A. Schema therapy for Dissociative Identity Disorder (DID): further explanation about the rationale and study protocol. *Eur J Psychotraumatol*. (2019) 10(1):1571377. doi: 10.1080/20008198.2019.1684629
15. Madden SP, Einhorn PM. Cannabis-induced depersonalization-derealization disorder. *Am J Psychiatry*. (2018) 175:3–6. doi: 10.1176/appi.ajp-rj.2018.130202

RETRACTED