



## OPEN ACCESS

## EDITED BY

Athanasia Pataka,  
Aristotle University of Thessaloniki, Greece

## REVIEWED BY

Kimberlyn Maravet Baig-Ward,  
University of Texas Southwestern Medical  
Center, United States  
Antonio Santagostini,  
University of Parma, Italy

## \*CORRESPONDENCE

Liwei Cui

✉ mscui0552@163.com

RECEIVED 22 March 2024

ACCEPTED 23 September 2024

PUBLISHED 09 October 2024

## CITATION

Zhang Y, Liu H, Feng C, Yang Y and Cui L  
(2024) Prevalence of sleep disturbances in  
endometriosis patients: a systematic  
review and meta-analysis.  
*Front. Psychiatry* 15:1405320.  
doi: 10.3389/fpsy.2024.1405320

## COPYRIGHT

© 2024 Zhang, Liu, Feng, Yang and Cui. This is  
an open-access article distributed under the  
terms of the [Creative Commons Attribution  
License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or  
reproduction in other forums is permitted,  
provided the original author(s) and the  
copyright owner(s) are credited and that the  
original publication in this journal is cited, in  
accordance with accepted academic  
practice. No use, distribution or reproduction  
is permitted which does not comply with  
these terms.

# Prevalence of sleep disturbances in endometriosis patients: a systematic review and meta-analysis

Yujie Zhang<sup>1</sup>, Hui Liu<sup>2</sup>, Chaochen Feng<sup>2</sup>, Yadi Yang<sup>2</sup>  
and Liwei Cui<sup>2\*</sup>

<sup>1</sup>School of Humanities and Management, Zhejiang Chinese Medical University, Hangzhou, Zhejiang, China, <sup>2</sup>Department of Quality Management, Jining N0.1 People's Hospital, Jining, Shandong, China

**Objective:** This study systematically analyzes the prevalence of sleep disturbance in patients with endometriosis.

**Methods:** The PubMed, Web of Science, Embase, Wanfang, China National Knowledge Internet Database (CNKI), China Science and Technology Journal Database were searched from their establishment to January 2024, using the search terms endometriosis and sleep disturbance to collect relevant literature on the prevalence of sleep disturbance in patients with endometriosis. Two researchers independently screened the literature, extracted data, and evaluated the risk of bias. The prevalence of sleep disorders in patients with endometriosis was systematically analyzed using Stata17.0 software.

**Results:** Sixteen studies with 2573 participants were included. The prevalence of sleep disturbance in patients with endometriosis was 70.8% (95% confidence interval: 60.7%~80.9%). The said prevalence was higher in China than in Iran and the European countries (78.2 vs. 57.6 vs. 64.4,  $Q=9.27$ ,  $P=0.010$ ) and increased significantly since 2018 (79.0 vs. 61.3,  $Q=3.97$ ,  $P=0.046$ ). This prevalence was significantly higher in the cohort study than that in cross-sectional and case-control studies (84.0 vs. 74.0 vs. 59.5,  $Q=7.16$ ,  $P=0.028$ ).

**Conclusion:** The prevalence of sleep disturbance is high in patients with endometriosis, particularly in China and its prevalence has increased significantly in recent years. Appropriate interventions are recommended to effectively prevent or minimize sleep disturbances in patients with endometriosis.

## KEYWORDS

sleep disturbance, endometriosis, prevalence, meta-analysis, systematic review

## 1 Introduction

Endometriosis is a common gynecological disorder affecting women, where active endometrial cells are planted in a location other than the endometrium (1–3). The prevalence of endometriosis is high (4–8). The disease affects approximately 10–15% of the female population of reproductive age, or 176 million women worldwide (9, 10). Sleep disorders constitute major health problems worldwide, causing impairments in initiating and maintaining sleep, as well as abnormal sleep events that interfere with an individual's normal daily functioning and mood while awake (11). The most common sleep disorders include insomnia, obstructive sleep apnea, restless leg syndrome, and circadian rhythm disorders (12). Endometriosis can lead to a variety of painful symptoms such as dysmenorrhea, painful intercourse, painful defecation or urination, and chronic pelvic pain (13). The various pain symptoms in patients with endometriosis affect the quality of their sleep, making them prone to sleep disorders. Studies have shown that the higher the pain score, the lower the patient's sleep quality score for endometriosis as a progressive disease (14). Nunes' study also showed that pain in endometriosis patients has a negative impact on sleep (15). One study showed that chronic pelvic pain worsened subjective sleep quality by more than three times, increased sleep disturbances by nearly six times, and decreased sleep duration by almost seven times (16). In addition, painful bladder syndrome increased sleep disturbances by almost five times (16).

Other factors that cause endometriosis patients to be prone to sleep disorders are endocrine changes and psychological states. There is a higher prevalence of depression and symptoms of anxiety in patients with endometriosis, and these psychological conditions are strongly associated with sleep disturbances (17). A case-control study showed higher levels of depression in patients with endometriosis compared to controls (18). Roomaney's findings showed that 71% of patients with endometriosis reported moderate to severe depressive symptoms (19). Maulitz, for his part, hypothesized that at least one-third of patients with endometriosis suffer from mental disorders (primarily depression or anxiety) (20). Current evidence suggests that women with endometriosis have more psychological disorders that are strongly associated with sleep disorders (15, 21). For example, a quantitative study showed that a 1-point increase in the PHQ-9 (worsening depression) increased the primary outcome (poorer sleep quality) by 1.62 points (21).

In addition, changes in ectopic endothelial tissue during the menstrual cycle may lead to elevated levels of inflammatory mediators and prostaglandins in the patient's body (22), which may in turn lead to altered sleep patterns and decreased sleep quality. Further, sleep disorders in endometriosis patients have been associated with a variety of adverse health outcomes that can be detrimental to the individual. For example, studies have shown that endometriosis patients who experience sleep disorders exhibit more fatigue (23). Maulitz and Mundo-Lrses's (20, 24) study showed that sleep disorders adversely affect the quality of life of patients with endometriosis. Therefore, it is crucial to identify severity of sleep disorders in patients with endometriosis and provide timely and effective intervention.

In recent times, researchers have focused on sleep disorders in patients with endometriosis. And most of the studies have shown that the prevalence of sleep disorders in endometriosis patients is high and more attention needs to be given to these patients. For example, the studies of Souza (16), Davie (25), and Goksu (26) showed that the prevalence of sleep disorders in endometriosis patients was 87.14%, 80.00%, and 90.48%, respectively. However, owing to the differences in the type of study, survey area, and survey instruments used in different studies, the results differ. As shown in a prospective cross-sectional questionnaire study (27), the prevalence of sleep disorders in patients with endometriosis was 42.58%.

Therefore, the prevalence of sleep disorders in patients with endometriosis has not yet been systematically determined. This study aims to assess the prevalence of sleep disorders quantitatively and accurately in patients with endometriosis through a single-rate meta-analysis and to clarify the current status of sleep disorders in these patients. We also conducted stratified analyses of the incidence of sleep disorders based on geographic region, year of publication, study type, sample size, and survey instrument to clarify the factors affecting the incidence of sleep disorders. This will provide a reference for effective prevention and intervention of sleep disorders in patients with endometriosis.

## 2 Methods

### 2.1 Protocol

The literature search was conducted following the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines (28), and the research protocol was registered in PROSPERO (CRD42023463967).

### 2.2 Search strategy

Two researchers independently searched the PubMed, Web of Science, Embase, Wanfang, China National Knowledge Internet Database (CNKI), China Science and Technology Journal Database to identify relevant studies on the incidence of sleep disturbance in patients with endometriosis. The timeframe for this search was from database construction to January 2024. The search terms primarily included endometriosis, endometrioma, adenomyosis, sleep disturbance, sleep quality, insomnia, sleep problem, sleep disorder, and sleep symptom. References in the included studies were manually searched. We determined whether relevant articles met the inclusion criteria and could be included in this study.

### 2.3 Study selection

Studies that met the following criteria were included: (1) Observational studies; (2) the study population comprised of patients with endometriosis; and (3) the outcome indicator was the prevalence of sleep disturbance or any type of sleep disorder,

including insomnia, obstructive sleep apnea, and restless leg syndrome. Studies for which the full text was not available was excluded. Additionally, studies with duplicate publications or similar full-text data were excluded. If the same data appeared in more than one study, the studies with complete data and the largest sample size were included in the meta-analysis.

## 2.4 Data extraction and quality assessment

Two researchers independently screened the literature, extracted information from those who met the inclusion criteria, and performed crosschecking. In cases of disagreement, a third party negotiated the judgment. Relevant literature was initially screened by reading the title and abstract and then further screened for final inclusion by reading the full text.

The information excerpts mainly included: (1) Literature information: authors, publication year, survey time, country, and assessment tools. We placed no restriction on the instruments used to assess sleep disorders in this study, accepting well-established generic scales, self-developed questionnaires or entries. As long as the article provided the total number of people with endometriosis and the number of people presenting with sleep disorders, it was eligible for inclusion in this study. (2) Participant information: age, gender, and sample size. (3) Outcome indicators: prevalence of sleep disturbance. If there was no specific incidence rate in the literature, the incidence rate was considered to be calculated thus: incidence rate = number of people presenting with sleep disorders/total number of people in the sample  $\times$  100%. (4) Quality evaluation information: the Newcastle-Ottawa Scale (NOS) was used to evaluate the quality of cohort and case-control studies, which was scored out of 9, and a score of  $\geq 7$  was considered high-quality literature (29). The risk of bias evaluation criteria developed by the Joanna Briggs Institute (JBI) was used to evaluate cross-sectional studies and consisted of ten entries (30). The entries were scored according to their degree of compliance: 0, non-compliance; 1, mention but no detailed description; and 2, detailed and comprehensive description. Generally, scores greater than 70% of the total score are considered high quality.

## 2.5 Statistical analysis

Statistical analysis was conducted using Stata 17.0 software. The data type of this study's outcome indicators was dichotomous information, and Odds ratio (OR) was chosen as the effect indicator. If the heterogeneity among the included studies was low ( $I^2 < 50\%$ ,  $P > 0.1$ ), a fixed-effects model was selected for analysis. In cases of high heterogeneity among the included studies ( $I^2 > 50\%$ ,  $P < 0.1$ ), a random-effects model was selected for the analysis. Sensitivity and subgroup analyses were used to analyze the sources of heterogeneity. Factors for subgroup analyses mainly included country or region (China, Iran, and Europe), survey time ( $\leq 2018$  and  $> 2018$ ), type of research (Cross-sectional study, Case-control study, and Cohort study), sample size ( $\leq 100$  and  $> 100$ ),

assessment tools (Pittsburgh Sleep Quality Index and Others), and pelvic pain (Yes and No). Funnel plots, Begg's test, and Egger's test were used to analyze publication bias. The significance level was set at  $P < 0.05$  (two-tailed).

## 3 Results

### 3.1 Selection of studies and basic characteristics

A total of 679 relevant studies were obtained through the search. After a layer-by-layer screening, 16 studies were finally included (Figure 1). The included studies were 3 Chinese and 13 English studies. A total of 2573 patients with endometriosis were included.


The investigation was conducted from January 2010 to August 2021. Two studies were conducted in China, Iran, Italy, Turkey. Other countries of study included Australia, Thailand, Spain, Switzerland, Germany, Austria, Brazil, the United Kingdom and the United States. The 16 included studies comprised 9 cross-sectional, 5 case-control, and 2 cohort studies. Eleven papers chose the Pittsburgh Sleep Quality Index (PSQI) as the assessment tool while five chose other assessment tools. The prevalence of sleep disorders in these studies ranged from 42.58% to 90.48% (Table 1).

### 3.2 Quality assessment

The quality scores of the cross-sectional studies ranged from 13–15, and were mostly of medium to high quality (Table 2). However, the scores were relatively low in terms of the choice of sampling method, reliability and validity of the data collection instruments, and verification of data authenticity. No studies chose a random sampling method or reported on the reliability and validity of the data collection instrument. Only one study used more complete measures to verify the authenticity of information. Case-control and cohort studies had quality scores of 7–8, rendering them as high-quality studies (Table 3). However, all studies had deficiencies in the completeness of follow-up or response rates.

### 3.3 Prevalence of sleep disturbances in patients with endometriosis

The results of the random-effects model showed that the prevalence of sleep disturbance in patients with endometriosis was 70.8% (Figure 2). The 95% confidence interval (CI) was 60.7%–80.9%. After the sequential removal of each study, the prevalence of sleep disturbance in such patients fluctuated between 69.5% and 72.8%, with the combined value being at most 2.0 percentage points higher than before the exclusion, suggesting that the results of this study are more stable and reliable (Figure 3).



### 3.4 Publication bias

The funnel plot results showed symmetry in the distribution of the graphs on both sides (Figure 4). Egger's and Begg's tests showed *p*-values of 0.115 and 0.822, respectively. This method indicated a low probability of publication bias.

### 3.5 Subgroup analysis

Subgroups were analyzed according to the country or region, time of investigation, study type, sample size, assessment tools, and pelvic pain. Subgroup analyses showed that the prevalence of sleep disturbance in Chinese patients with endometriosis (78.2%, 95% CI: 67.8%~88.6%) was higher than that in Iran (57.6%, 95% CI: 49.4%~65.8%) and the European countries (64.4%, 95% CI: 49.4%~79.5%) (Table 4). This prevalence was higher in patients with endometriosis from 2018 to 2021 (79.0%, 95% CI: 69.5%~88.6%) than pre-2018 (61.3%, 95% CI: 46.8%~75.9%) (Table 4). Furthermore, the prevalence was significantly higher in the cohort study (84.0%, 95% CI: 70.7%~97.4%) than that in cross-sectional (74.0%, 95% CI: 61.4%~86.6%) and case-control studies (59.5%, 95% CI: 47.1%~71.8%) (Table 4). Additionally, the prevalence was not significantly correlated with the sample size, assessment tool, or presence of pelvic pain ( $P>0.05$ ) (Table 4).

## 4 Discussion

The results of this study show that the prevalence of sleep disorders in patients with endometriosis was 70.8% (95% CI: 60.7%~80.9%). Multiple meta-analyses (11, 40–46) showed that patients

with cancer (60.7%), HIV infection (58%), ankylosing spondylitis (53%), traumatic brain injury (50%), stroke (41.85%), irritable bowel syndrome (37.6%), and dementia (26%) had relatively lower sleep disorders. Only patients with chronic non-cancer conditions had a higher prevalence of sleep disorders (76.3%) than those with endometriosis. The prevalence of sleep disorders in women with health issues, such as breast cancer survivors (47) and women with recurrent pregnancy loss (48), was 62% and 31.2%, respectively. This finding indicates that the incidence of sleep disorders is high in patients with endometriosis.

The development of sleep disorders in patients with endometriosis is a multifactorial process involving disease symptoms, adverse outcomes, physiological effects, and lifestyle changes. Dysmenorrhea, chronic pelvic pain, menstrual abnormalities, and painful intercourse are the primary symptoms of endometriosis (49), with secondary dysmenorrhea being the most common symptom and often worsening progressively. Endometriosis of the uterorectal sulcus and the vaginorectal septum may also cause painful intercourse. Research has shown a significant correlation between chronic pain and sleep disorders (50). Pain can make it difficult for patients to find a comfortable sleeping position and may cause frequent awakenings, thus interfering with normal sleep patterns.

Second, endometriosis is associated with infertility: Infertility rates range from 30% to 50% in patients with endometriosis (51). This is mainly because endometriosis can cause adhesions around the fallopian tubes, affecting oocytes or ovulation due to ovarian pathology (52). Previous studies have shown that endometriosis is strongly associated with the risk of developing chronic diseases, such as coronary artery disease (53), type 2 diabetes (54), stroke (55), ankylosing spondylitis (56), and rheumatoid arthritis (57). Li (53) and Xue (57) showed that patients with endometriosis had higher rates of coronary heart disease and rheumatoid arthritis than those

TABLE 1 Characteristics of the studies included in the meta-analysis.

Authors (Publication year)	Survey time	Country	Type of research	Age (years) (Mean eaSD)	Sample size (n)	Number of Incidence (n)	Incidence rate (%)	Assessment tools	Quality score
Halici(2023) (14)	2020-2021	Turkey	Cohort study	36.82 6.8.77	56	43	76.79	PSQI	8
Souza (a) (2023) (16)	2020.1-2021.2	Brazil	Cross-sectional study	32-41	140	122	87.14	PSQI	15
Facchin(2021) (23)	2019.7-2020.3	Italy	Case control study	34.11 4.1.34	123	66	53.66	PSQI	8
Mundo-L2020.3al01 (24)	2019.1-7	Spain	Cross-sectional study	36.7 6.7.2	230	187	81.30	PSQI	15
Davie (2020) (25)	—	Australia	Case control study	35.0 5.00.35	30	24	80.00	PSQI	7
Goksu(2021) (26)	2019-2020	Turkey	Cohort study	33.8 3.8.6	42	38	90.48	PSQI	7
Tempest(2021) (27)	2017.10-2018.1	United Kingdom	Cross-sectional study	—	465	198	42.58	IRLSSG+BSGE	15
Cai(2022) (31)	2020.2-2021.8	China	Cross-sectional study	40.25 0.24.12	50	42	84.00	PSQI	13
Wang(2017) (32)	2016.3-2017.3	China	Cross-sectional study	21-47	120	88	73.33	Self-designed	15
Han(2019) (33)	2017.1-2018.12	Thailand	Cross-sectional study	16-40	200	177	88.50	Self-designed	14
Tanha(2014) (34)	2013.3-9	Iran	Cross-sectional study	31.4 1.4.7	61	33	54.10	PSQI	13
DiVasta(2018) (35)	2012.11-2016.3	America	Cross-sectional study	—	117	75	64.10	Self-designed	14
Souza (b) (36)	—	—	Cross-sectional study	37 7ud	161	144	89.44	PSQI	15
Ramin-Wright(2018) (37)	2010-2016	Switzerland, Germany, Austria	Case control study	37.9 7.9.2	555	238	42.88	Self-designed	8
Youseflu(2020) (38)	2016.5-2017.2	Iran	Case control study	31.00 1.0.63	78	47	60.26	PSQI	8
Maggiore(2015) (39)	2012.5-2013.12	Italy	Case control study	32.9 2.9.3	145	94	64.83	PSQI	8




FIGURE 2 Forest plot of the prevalence of sleep disturbances based on the random-effects model.

without endometriosis. There is also a link between endometriosis and ovarian (58) and endometrial cancer (59). Patients with endometriosis have a significantly higher risk of endometrial cancer than those in the control group, and they also demonstrate an association with poor gestational and perinatal outcomes (60–62).

Endometriosis not only causes physical pain to the patient but may also result in adverse outcomes such as infertility and miscarriage. Patients with endometriosis are also at risk for a variety of diseases, especially endometrial cancer, ovarian cancer, and other malignant tumors. A variety of factors cause physical pain





FIGURE 3 Sensitivity analysis.





and mental stress in patients with endometriosis, making their sleep quality generally poorer, and the incidence of sleep disorders is generally higher than that in other disease populations.

Third, endometriosis may affect a patient’s endocrine system. Studies have found that patients with endometriosis have higher levels of prostaglandins, which may interfere with the balance of sleep-regulating hormones (22). In addition, endometriosis may cause other physiological changes, such as chronic fatigue and altered immune system function, which may affect sleep. The relationship between chronic fatigue syndrome and sleep disorders has been demonstrated in several studies, and chronic conditions such as endometriosis may exacerbate this fatigue (63).

Fourth, due to the physical discomfort caused by the disease, people with endometriosis may be less physically active, which may lead to decreased physical functioning and sleep quality. Studies have reported a positive association between physical activity and sleep quality, and reduced physical activity may impair sleep quality (64). Finally, medications used to treat endometriosis, such as

nonsteroidal anti-inflammatory drugs (NSAIDs) and hormonal drugs, may have an impact on sleep. Studies have found that certain medications such as NSAIDs may affect the structure and continuity of sleep (65).

As there are multiple reasons for sleep disorders in individuals with endometriosis, the issue requires urgent attention. The study findings suggest that timely and effective interventions are necessary to reduce the incidence of sleep disorders in patients with endometriosis. Physical therapy may be used in addition to medication, and transcutaneous electrical nerve stimulation (TENS) and massage therapy can also relieve chronic pain and help improve sleep efficiency (66, 67). In addition, healthcare professionals can provide information about good sleep habits, such as maintaining a regular sleep schedule and avoiding the use of electronic devices close to bedtime, which can help improve sleep quality. Additionally, patients with endometriosis can engage in regular moderate-intensity exercise (64), and patients experiencing mood disorders can seek emotional support and help through

TABLE 2 Results of bias risk assessment of included studies (Cross-sectional study).

Authors	①	②	③	④	⑤	⑥	⑦	⑧	⑨	⑩	JBI score
Souza (a) (2023) (16)	2	0	2	2	0	1	2	2	2	2	15
Mundo-lsectional1 (24)	2	0	2	2	1	0	2	2	2	2	15
Tempest(2021) (27)	2	0	2	2	0	1	2	2	2	2	15
Cai(2022) (31)	2	0	2	1	0	0	2	2	2	2	13
Wang(2017) (32)	2	0	2	2	0	2	1	2	2	2	15
Han(2019) (33)	2	0	2	2	0	1	1	2	2	2	14
Tanha(2014) (34)	2	0	2	2	1	0	0	2	2	2	13
DiVasta(2018) (35)	2	0	1	2	0	1	2	2	2	2	14
Souza (b) (36)	2	0	2	1	1	1	2	2	2	2	15

TABLE 3 Results of bias risk assessment of included studies (Case control study and Cohort study).

Authors	Selection of population				Comparability	Evaluation of exposure or outcome			NOS score
Halici(2023) (14)	1	1	1	1	1	1	1	0	7
Facchin(2021) (23)	1	1	1	1	2	1	1	0	8
Davie(2020) (25)	1	1	1	1	2	1	1	0	8
Goksu(2021) (26)	1	1	1	1	1	1	1	0	7
Ramin-Wright(2018) (37)	1	1	1	1	2	1	1	0	8
Youseflu(2020) (38)	1	1	1	1	2	1	1	0	8
Maggiore(2015) (39)	1	1	1	1	2	1	1	0	8

meditation and deep breathing exercises. They can also obtain counseling or join support groups to relieve stress and anxiety and improve their sleep.

More notably, the results showed that the prevalence of sleep disorders in patients with endometriosis since 2018 reached 79.0, which was much higher than the pre-2018 prevalence. With the gradual improvement in living standards, people’s healthy life expectancy and disability-free life expectancy have also increased, and their concern for health continues to grow (68, 69). Once a disease occurs, especially endometriosis, which has a greater impact on the individual, it is more devastating. People are more psychologically stressed and more likely to experience adverse conditions such as sleep disorders. It was also found that the prevalence of sleep disorders in patients with endometriosis in the cohort study was significantly higher than in other types of studies. The main reason for this finding may be the small sample size (98 patients) of the cohort study. The results of small-sample studies are

usually unstable and may lead to biased prevalence of sleep disorders.

This study found a higher significantly prevalence of sleep disorders in Chinese patients with endometriosis than in other countries and regions. This may be related to the traditional Chinese stigma associated with infertility. In China, traditional beliefs emphasize the importance of having children (70). Infertility may thus bring stigma upon the individual and his/her family, causing greater psychological stress and making sleep disorders more likely to occur (71). Chinese patients may also be more inclined to tolerate symptoms without seeking medical help, potentially leading to severe disease progression and exacerbation of associated sleep disorders (72). At the same time, China’s health system has a relatively limited capacity and lacks multidisciplinary teams with the broad range of skills and equipment needed to diagnose endometriosis early and effectively treat it. At the same time, primary healthcare providers have limited knowledge of

TABLE 4 Prevalence rate of effort-reward imbalance by demographic characteristics.

Subgroups	Categories	No. of studies	Sample size	Heterogeneity		Incidence rate (%)	95%CI	Q	P
				I <sup>2</sup> (%)	P				
Country	China	2	170	62.1	0.105	78.2	(67.8, 88.6)	9.27	0.010
	Iran	2	139	0	0.466	57.6	(49.4, 65.8)		
	Europe	7	1493	97.6	<0.001	64.4	(49.4, 79.5)		
Survey time	≤imee	6	498	97.6	<0.001	61.3	(46.8, 75.9)	3.97	0.046
	>2018	8	950	89.4	<0.001	79.0	(69.5, 88.6)		
Type of research	Cross-sectional study	9	1544	97.5	<0.001	74.0	(61.4, 86.6)	7.16	0.028
	Case control study	5	931	91.3	<0.001	59.5	(47.1, 71.8)		
	Cohort study	2	98	72.1	0.058	84.0	(70.7, 97.4)		
Sample size	≤size	6	317	84.7	<0.001	74.5	(62.9, 86.1)	0.40	0.527
	>100	10	2256	98.2	<0.001	68.8	(55.6, 82.1)		
Scale	PSQI	11	1116	90.6	<0.001	75.1	(67.4, 82.9)	1.33	0.248
	other	5	1457	98.6	<0.001	62.2	(41.7, 82.7)		
Pelvic pain	Yes	2	141	82.1	0.018	82.0	(64.8, 99.2)	0.85	0.356
	No	2	77	91.3	0.001	63.8	(29.3, 98.3)		



endometriosis and play a limited role in its management. This may have contributed to the relatively high prevalence of sleep disorders in Chinese patients with endometriosis.

Other countries, especially in Europe, have a lower incidence of sleep disorders in endometriosis patients than in China, which may reflect differences in healthcare systems, cultural differences, and other such factors. Healthcare systems in developed European countries may pay more attention to individualized treatment and health education, and education and awareness of sleep health may be higher (73). This would increase patients' awareness and management of the disease and help the public to identify and manage factors that may contribute to sleep disorders, potentially reducing the incidence of sleep disorders. Studies have shown that cultural differences influence patients' attitudes and behaviors toward the disease and that Western cultures strongly encourage personal expression and seeking psychological support (74). This could aid patients with endometriosis in managing the psychological stress caused by the disease and potentially reduce sleep disorders.

This study has several limitations. First, individuals with endometriosis may be hesitant to share personal information due to concerns about reproductive health and privacy implications, making it challenging to collect information on such patients. In addition, recruiting a sufficient number of subjects for high-quality research is fraught with challenges due to the heterogeneity of the disease and the difficulty of diagnosis (75). Therefore, the number of studies that could be included was relatively small, and the studies themselves had small sample sizes. In the subgroup analysis, the number of studies included in each subgroup was limited. Second, the included studies used different sleep disorder questionnaires to determine the prevalence of sleep disorders. Moreover, reports on sleep disorders are self-assessed, and there is a certain degree of subjectivity in the evaluation of sleep quality. This may result in the lower precision and accuracy of prevalence estimates. Third, we only included publications in Chinese and English. This implies that important local studies published in journals of other languages may have been overlooked. This may have led to bias in our findings.

## 5 Conclusion

Although our study had some limitations, we systematically evaluated the prevalence of sleep disorders in patients with endometriosis using a meta-analysis. The incidence of sleep disorders in patients with endometriosis was relatively high. Its incidence has gradually increased over time. In addition, there were some differences in the incidence between the different regions. The incidence of sleep disorders in Chinese patients with endometriosis is significantly higher than those of patients from other regions. These results suggest a possible link between endometriosis and sleep disorders and indicate

that further research is needed to better understand this correlation. For patients with endometriosis, various forms of physical therapy, physical exercise, and psychological counseling can be used to avoid or alleviate sleep disorders and improve sleep quality. Due to limitations in the number and quality of included studies, the above conclusions have yet to be confirmed by more high quality studies. Future studies should prioritize conducting large, multicenter prospective studies and ensure rigorous measurement of the reliability and validity of data collection tools. In addition, attention to quality control issues during data collection is crucial, including verification of data, bias reduction, and measurement accuracy.

## Author contributions

YZ: Writing – review & editing, Writing – original draft, Software, Project administration, Data curation, Conceptualization. HL: Writing – review & editing, Data curation. CF: Writing – review & editing, Data curation. YY: Writing – review & editing, Data curation. LC: Writing – review & editing, Writing – original draft, Supervision, Software, Project administration, Data curation, Conceptualization.

## Funding

The author(s) declare that no financial support was received for the research, authorship, and/or publication of this article.

## Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

## Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

## Supplementary material

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fpsy.2024.1405320/full#supplementary-material>

## References

- Shafir AL, Farland LV, Shah DK, Harris HR, Kvaskoff M, Zondervan K, et al. Risk for and consequences of endometriosis: A critical epidemiologic review. *Best Pract Res Clin Obstet Gynaecol.* (2018) 51:1–15. doi: 10.1016/j.bpobgyn.2018.06.001
- Giudice LC, Kao LC. Endometriosis. *Lancet.* (2004) 364:1789–99. doi: 10.1016/S0140-6736(04)17403-5
- Vercellini P, Viganò P, Somigliana E, Fedele L. Endometriosis: pathogenesis and treatment. *Nat Rev Endocrinol.* (2014) 10:261–75. doi: 10.1038/nrendo.2013.255
- Kim H, Lee M, Hwang H, Chung YJ, Cho HH, Yoon H, et al. The estimated prevalence and incidence of endometriosis with the Korean National Health Insurance Service-national sample cohort (NHIS-NSC): A national population-based study. *J Epidemiol.* (2021) 31:593–600. doi: 10.2188/jea.JE20200002
- Le Moal J, Gorla S, Chesneau J, Fauconnier A, Kvaskoff M, De Crouy-Chanel P, et al. Increasing incidence and spatial hotspots of hospitalized endometriosis in France from 2011 to 2017. *Sci Rep.* (2022) 12:6966. doi: 10.1038/s41598-022-11017-x
- Ávalos MA, Barranco CE, Martos GR, Mendoza Ladrón de Guevara N, Mazheika M. Epidemiology of endometriosis in Spain and its autonomous communities: A large, nationwide study. *Int J Environ Res Public Health.* (2021) 18:7861. doi: 10.3390/ijerph18157861
- Rowlands JJ, Abbott JA, Montgomery GW, Hockey R, Rogers P, Mishra GD. Prevalence and incidence of endometriosis in Australian women: A data linkage cohort study. *BJOG.* (2021) 128:657–65. doi: 10.1111/1471-0528.16447
- Illum LRH, Forman A, Melgaard A, Hansen KE, Hansen SN, Nyegaard M, et al. Temporal and regional differences in the incidence of hospital-diagnosed endometriosis: A Danish population-based study. *Acta Obstet Gynecol Scand.* (2022) 101:737–46. doi: 10.1111/aogs.14364
- Zondervan KT, Becker CM, Koga K, Missmer SA, Taylor RN, Viganò P. Endometriosis. *Nat Rev Dis Primers.* (2018) 4:9. doi: 10.1038/s41572-018-0008-5
- Adamson GD, Kennedy S, Hummelshoj L. Creating solutions in endometriosis: Global collaboration through the world endometriosis Research Foundation. *J Endometriosis.* (2010) 2:3–6. doi: 10.1177/228402651000200102
- Sun Y, Laksono I, Selvanathan J, Saripella A, Nagappa M, Pham C, et al. Prevalence of sleep disturbances in patients with chronic non-cancer pain: A systematic review and meta-analysis. *Sleep Med Rev.* (2021) 57:101467. doi: 10.1016/j.smrv.2021.101467
- Sumbodo CD, Tyson K, Mooney S, Lamont J, McMahan M, Holdsworth-Carson SJ. The relationship between sleep disturbances and endometriosis: A systematic review. *Eur J Obstet Gynecol Reprod Biol.* (2024) 293:1–8. doi: 10.1016/j.ejogrb.2023.12.010
- Zondervan KT, Becker CM, Missmer SA. Endometriosis. *N Engl J Med.* (2020) 382:1244–56. doi: 10.1056/NEJMra1810764
- Halici BNA, Aktöz F, Kabakci M, Kiran G, Özcan P. Analysis of preoperative and postoperative quality of life, sexual function, and sleep in patients with endometriosis: a prospective cohort study. *Arch Gynecol Obstet.* (2023) 307:113–20. doi: 10.1007/s00404-022-06562-9
- Nunes FR, Ferreira JM, Bahamondes L. Pain threshold and sleep quality in women with endometriosis. *Eur J Pain.* (2015) 19:15–20. doi: 10.1002/ejp.514
- de Souza RJ, Villela NR, Broglio LCS, Oliveira MAP. Impact of chronic pelvic pain and painful bladder syndrome on the Pittsburgh Sleep Quality Index on women with deep endometriosis: a cross-sectional study. *Int Urogynecol J.* (2023) 34:2487–93. doi: 10.1007/s00192-023-05560-y
- Arion K, Orr NL, Noga H, Allaire C, Williams C, Bedaiwy MA, et al. A quantitative analysis of sleep quality in women with endometriosis. *J Womens Health (Larchmt).* (2020) 29:1209–15. doi: 10.1089/jwh.2019.8008
- Gambadauro P, Carli V, Hadlaczky G. Depressive symptoms among women with endometriosis: a systematic review and meta-analysis. *Am J Obstet Gynecol.* (2019) 220:230–41. doi: 10.1016/j.ajog.2018.11.123
- Roomaney R, Mitchell H. Psychosocial correlates of symptoms of depression among patients with endometriosis in the United Kingdom. *Women Health.* (2022) 62:764–74. doi: 10.1080/03630242.2022.2144985
- Maulitz L, Stickeler E, Stickel S, Habel U, Tchaikovski SN, Chechko N. Endometriosis, psychiatric comorbidities and neuroimaging: Estimating the odds of an endometriosis brain. *Front Neuroendocrinol.* (2022) 65:100988. doi: 10.1016/j.yfrne.2022.100988
- Casalechi M, Vieira-Lopes M, Quessada MP, Araújo TC, Reis FM. Endometriosis and related pelvic pain: association with stress, anxiety and depressive symptoms. *Minerva Obstet Gynecol.* (2021) 73:283–9. doi: 10.23736/S2724-606X.21.04704-3
- Sacco K, Portelli M, Pollacco J, Schembri-Wismayer P, Calleja-Agius J. The role of prostaglandin E2 in endometriosis. *Gynecol Endocrinol.* (2012) 28:134–8. doi: 10.3109/09513590.2011.588753
- Facchin F, Buggio L, Roncella E, Somigliana E, Ottolini F, Drudi D, et al. Sleep disturbances, fatigue and psychological health in women with endometriosis: a matched pair case-control study. *Reprod BioMed Online.* (2021) 43:1027–34. doi: 10.1016/j.rbmo.2021.08.011
- Mundo-López A, Ocón-Hernández O, San-Sebastián AP, Galiano-Castillo N, Rodríguez-Pérez O, Arroyo-Luque MS, et al. Contribution of chronic fatigue to psychosocial status and quality of life in Spanish women diagnosed with endometriosis. *Int J Environ Res Public Health.* (2020) 17:3831. doi: 10.3390/ijerph17113831
- Davie S, Hamilton Y, Webb L, Amoako AA. Sleep quality and endometriosis: A group comparison study. *J Endometr Pelvic Pain Disord.* (2020) 12:94–100. doi: 10.1177/2284026520909979
- Goksu M, Kadirogullari P, Seckin KD. Evaluation of depression and sleep disorders in the preoperative and postoperative period in stage 4 endometriosis patients. *Eur J Obstet Gynecol Reprod Biol.* (2021) 264:254–8. doi: 10.1016/j.ejogrb.2021.07.037
- Tempest N, Boyers M, Carter A, Lane S, Hapangama DK. Premenopausal women with a diagnosis of endometriosis have a significantly higher prevalence of a diagnosis or symptoms suggestive of restless leg syndrome: A prospective cross-sectional questionnaire study. *Front Endocrinol (Lausanne).* (2021) 12:599306. doi: 10.3389/fendo.2021.599306
- Moher D, Shamseer L, Clarke M, Ghersi D, Liberati A, Petticrew M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Syst Rev.* (2015) 4:1. doi: 10.1186/2046-4053-4-1
- Wells GA, Shea B, O'Connell D, Peterson J, Welch V, Losos M, et al. The Newcastle-Ottawa Scale (NOS) for assessing the quality of non-randomised studies in meta-analyses(2019). Available online at: [http://www.ohri.ca/programs/clinical\\_epidemiology/oxford.asp](http://www.ohri.ca/programs/clinical_epidemiology/oxford.asp) (Accessed January 11, 2024).
- Joanna Briggs Institute. Checklist for analytical cross sectional studies, critical appraisal tools(2017). Available online at: <https://jbi.global/critical-appraisal-tools> (Accessed January 24, 2024).
- Cai H. Effect of dysmenorrhea induced by endometriosis on depression and sleep disorders. *World J Sleep Med.* (2022) 9:449–52. doi: 10.3969/j.issn.2095-7130.2022.03.026
- Wang Z. *Research of endometriosis through syndromes with TCM, body constitutions and disease related factors.* Chengdu, China: Chengdu University of Traditional Chinese Medicine (2017).
- Han R. *Research on Thailand dysmenorrhea of endometriosis through syndromes with TCM, body constitution and other disease related factors.* Heilongjiang, China: Heilongjiang University of Chinese Medicine (2020). doi: 10.27127/d.cnki.ghlzu.2019.000445
- Davari-Tanha F, Askari F, Akrami M, Mohseni M, Ghajarzadeh M. Sleep quality in women with endometriosis. *Acad J Surg.* (2014) 1:57–9.
- DiVasta AD, Vitonis AF, Laufer MR, Missmer SA. Spectrum of symptoms in women diagnosed with endometriosis during adolescence vs adulthood. *Am J Obstet Gynecol.* (2018) 218:324.e1–324.e11. doi: 10.1016/j.ajog.2017.12.007
- de Souza RJ, Broglio LCS, Carrerette FB, Villela NR, Oliveira MAP. Challenges in measuring sleep quality among women with endometriosis: A comparison of two questionnaires. *Sleep Med.* (2024) 114:250–4. doi: 10.1016/j.sleep.2024.01.008
- Ramin-Wright A, Schwartz ASK, Geraedts K, Rauchfuss M, Wölfler MM, Haeblerlin F, et al. Fatigue - a symptom in endometriosis. *Hum Reprod.* (2018) 33:1459–65. doi: 10.1093/humrep/dey115
- Youseflu S, Jahanian Sadatmahalleh S, Roshanzadeh G, Mottaghi A, Kazemnejad A, Moini A. Effects of endometriosis on sleep quality of women: does life style factor make a difference? *BMC Womens Health.* (2020) 20:168. doi: 10.1186/s12905-020-01036-z
- Leone Roberti Maggiore U, Bizzarri N, Scala C, Tafi E, Siesto G, Alessandri F, et al. Symptomatic endometriosis of the posterior cul-de-sac is associated with impaired sleep quality, excessive daytime sleepiness and insomnia: a case-control study. *Eur J Obstet Gynecol Reprod Biol.* (2017) 209:39–43. doi: 10.1016/j.ejogrb.2015.11.026
- Al Maqbali M, Al Sinani M, Alsayed A, Gleason AM. Prevalence of sleep disturbance in patients with cancer: A systematic review and meta-analysis. *Clin Nurs Res.* (2022) 31:1107–23. doi: 10.1177/10547738221092146
- Wu J, Wu H, Lu C, Guo L, Li P. Self-reported sleep disturbances in HIV-infected people: A meta-analysis of prevalence and moderators. *Sleep Med.* (2015) 16:901–7. doi: 10.1016/j.sleep.2015.03.027
- Salari N, Sadeghi N, Hosseini-Far A, Hasheminezhad R, Khaeia H, Shohaimi S, et al. Prevalence of sleep disturbance in patients with ankylosing spondylitis: A systematic review and meta-analysis. *Adv Rheumatol.* (2023) 63:33. doi: 10.1186/s42358-023-00315-1
- Mathias JL, Alvaro PK. Prevalence of sleep disturbances, disorders, and problems following traumatic brain injury: A meta-analysis. *Sleep Med.* (2012) 13:898–905. doi: 10.1016/j.sleep.2012.04.006
- Xiaolin Gu MM. Risk factors of sleep disorder after stroke: A meta-analysis. *Top Stroke Rehabil.* (2017) 24:34–40. doi: 10.1080/10749357.2016.1188474
- Wang B, Duan R, Duan L. Prevalence of sleep disorder in irritable bowel syndrome: A systematic review with meta-analysis. *Saudi J Gastroenterol.* (2018) 24:141–50. doi: 10.4103/sjg.SJG\_603\_17

46. Zhang J, Ye J, Tao X, Lu W, Chen X, Liu C. Sleep disturbances, sleep quality, and cardiovascular risk factors in women with polycystic ovary syndrome: Systematic review and meta-analysis. *Front Endocrinol (Lausanne)*. (2022) 13:971604. doi: 10.3389/fendo.2022.971604
47. Cheng WH, Teo RH, Cheng LJ, Lau Y, Lau ST. Global prevalence of sleep disturbances among breast cancer survivors: A systematic review with meta-analysis. *Sleep Health*. (2023) 9:704–16. doi: 10.1016/j.sleh.2023.04.004
48. Wang TT, Liu YL, Hou Y, Li JP, Qiao C. The risk factors of progestational anxiety, depression, and sleep disturbance in women with recurrent pregnancy loss: A cross-sectional study in China. *Front Psychol*. (2023) 14:1116331. doi: 10.3389/fpsyg.2023.1116331
49. Kurman RJ. Blaustein's pathology of the female genital tract. *Springer Berlin Heidelberg*. (2019). doi: 10.1007/978-3-319-46334-6
50. Leuenberger J, Kohl Schwartz AS, Geraedts K, Haeberlin F, Eberhard M, von Orellie S, et al. Living with endometriosis: Comorbid pain disorders, characteristics of pain and relevance for daily life. *Eur J Pain*. (2022) 26:1021–38. doi: 10.1002/ejp.1926
51. Bulletti C, Coccia ME, Battistoni S, Borini A. Endometriosis and infertility. *J Assist Reprod Genet*. (2010) 27:441–7. doi: 10.1007/s10815-010-9436-1
52. Kanellopoulos D, Karagianni D, Pergialiotis V, Nikiteas N, Lazaris AC, Iliopoulos D. Endometriosis and subfertility: A literature review. *Maedica (Bucur)*. (2022) 17:458–63. doi: 10.26574/maedica.2022.17.2.458
53. Li PC, Yang YC, Wang JH, Lin SZ, Ding DC. Endometriosis is associated with an increased risk of coronary artery disease in Asian women. *J Clin Med*. (2021) 10:4173. doi: 10.3390/jcm10184173
54. Farland LV, Degnan WJ, Harris HR, Tobias DK, Missmer SA. A prospective study of endometriosis and risk of type 2 diabetes. *Diabetologia*. (2021) 64:552–60. doi: 10.1007/s00125-020-05347-6
55. Zheng M, Zheng S. Endometriosis increases the risk of stroke: A Mendelian randomization study. *Stroke*. (2023) 54:e30–3. doi: 10.1161/STROKEAHA.122.041163
56. Yin Z, Low HY, Chen BS, Huang KS, Zhang Y, Wang YH, et al. Risk of ankylosing spondylitis in patients with endometriosis: A population-based retrospective cohort study. *Front Immunol*. (2022) 13:877942. doi: 10.3389/fimmu.2022.877942
57. Xue YH, You LT, Ting HF, Chen YW, Sheng ZY, Xie YD, et al. Increased risk of rheumatoid arthritis among patients with endometriosis: A nationwide population-based cohort study. *Rheumatol (Oxf Engl)*. (2021) 60:3326–33. doi: 10.1093/rheumatology/keaa784
58. Rueda-Martínez A, Garitazelaia A, Cilleros-Portet A, Marí S, Arauzo R, de Miguel J, et al. Genetic contribution of endometriosis to the risk of developing hormone-related cancers. *Int J Mol Sci*. (2021) 22:6083. doi: 10.3390/ijms22116083
59. Kim H, Kim HJ, Ahn HS. Does endometriosis increase the risks of endometrial hyperplasia and endometrial cancer? *Gynecol Oncol*. (2023) 169:147–53. doi: 10.1016/j.ygyno.2022.06.021
60. Breintoft K, Pinnerup R, Henriksen TB, Rytter D, Ulbjerg N, Forman A, et al. Endometriosis and risk of adverse pregnancy outcome: A systematic review and meta-analysis. *J Clin Med*. (2021) 10:667. doi: 10.3390/jcm10040667
61. Lafleur N, Wei SQ, Bilodeau-Bertrand M, Auger N. Association of endometriosis and severe maternal morbidity. *Obstet Gynecol*. (2022) 140:1008–16. doi: 10.1097/AOG.0000000000004969
62. Namazi M, Behboodi Moghadam Z, Zareyan A, Jafarabadi M. Impact of endometriosis on reproductive health: An integrative review. *J Obstet Gynaecol*. (2021) 41:1183–91. doi: 10.1080/01443615.2020.1862772
63. Álvarez-Salvago F, Lara-Ramos A, Cantarero-Villanueva I, Mazheika M, Mundo-Laeva A, Galiano-Castillo N, et al. Chronic fatigue, physical impairments and quality of life in women with endometriosis: A case-control study. *Int J Environ Res Public Health*. (2020) 17:3610. doi: 10.3390/ijerph17103610
64. Passos NF, Freitas PD, Carvalho-Pinto RM, Cukier A, Carvalho CRF. Increased physical activity reduces sleep disturbances in asthma: A randomized controlled trial. *Respirology*. (2023) 28:20–8. doi: 10.1111/resp.14359
65. Murphy PJ, Badia P, Myers BL, Boecker MR, Wright KP Jr. Nonsteroidal anti-inflammatory drugs affect normal sleep patterns in humans. *Physiol Behav*. (1994) 55:1063–6. doi: 10.1016/0031-9384(94)90388-3
66. Field T. Massage therapy research review. *Complement Ther Clin Pract*. (2014) 20:224–9. doi: 10.1016/j.ctcp.2014.07.002
67. Johnson MI, Paley CA, Jones G, Mulvey MR, Wittkopf PG. Efficacy and safety of transcutaneous electrical nerve stimulation (TENS) for acute and chronic pain in adults: a systematic review and meta-analysis of 381 studies (the meta-TENS study). *BMJ Open*. (2022) 12:e051073. doi: 10.1136/bmjopen-2021-051073
68. Yu HT, Xia T, Wang CF, Fang B, Cai RZ, Chen L, et al. Analysis on the difference between life expectancy and healthy life expectancy in Shanghai. *Zhonghua Liu Xing Bing Xue Za Zhi*. (2021) 42:1018–23. doi: 10.3760/cma.j.cn112338-20200417-00591
69. Crimmins EM, Zhang Y, Saito Y. Trends over 4 decades in disability-free life expectancy in the United States. *Am J Public Health*. (2016) 106:1287–93. doi: 10.2105/AJPH.2016.303120
70. Yu X, Liang J. Social norms and fertility intentions: Evidence from China. *Front Psychol*. (2022) 13:947134. doi: 10.3389/fpsyg.2022.947134
71. Yao H, Chan CHY, Chan CLW. Childbearing importance: A qualitative study of women with infertility in China. *Res Nurs Health*. (2018) 41:69–77. doi: 10.1002/nur.21846
72. Ma J, Luo Y, Yang S, Liu X, Peng Y, Wang H, et al. Patient delay and related influencing factors in Chinese women under 35 years diagnosed with cervical cancer: A cross-sectional study. *Asia Pac J Oncol Nurs*. (2022) 10:100165. doi: 10.1016/j.apjon.2022.100165
73. Ferrie JE, Shipley MJ, Cappuccio FP, Brunner E, Miller MA, Kumari M, et al. A prospective study of change in sleep duration: associations with mortality in the Whitehall II cohort. *Sleep*. (2007) 30:1659–66. doi: 10.1093/sleep/30.12.1659
74. Sue S, Zane N, Nagayama Hall GC, Berger LK. The case for cultural competency in psychotherapeutic interventions. *Annu Rev Psychol*. (2009) 60:525–48. doi: 10.1146/annurev.psych.60.110707.163651
75. Parasar P, Ozcan P, Terry KL. Endometriosis: epidemiology, diagnosis and clinical management. *Curr Obstet Gynecol Rep*. (2017) 6:347|e. doi: 10.1007/s13669-017-0187-1