



OPEN ACCESS

APPROVED BY
Frontiers Editorial Office,
Frontiers Media SA, Switzerland

*CORRESPONDENCE

Yuan-Jian Yang

✉ yuanjimyang@yeah.net

Chun-Nuan Chen

✉ chenchunnuan1983@aliyun.com

†These authors have contributed
equally to this work and share
first authorship

RECEIVED 29 March 2024

ACCEPTED 02 April 2024

PUBLISHED 12 April 2024

CITATION

Ye S-Y, Zhao Y, Liu Z-B, Luo C-P,
Xiong J-W, Zhan J-Q, Li Y-H, Wei B,
Chen C-N and Yang Y-J (2024)

Corrigendum: Lower serum insulin-like
growth factor 2 level in patients with bipolar
disorder is associated with the severity of
manic symptoms during manic episodes.

Front. Psychiatry 15:1408899.

doi: 10.3389/fpsy.2024.1408899

COPYRIGHT

© 2024 Ye, Zhao, Liu, Luo, Xiong, Zhan, Li, Wei,
Chen and Yang. This is an open-access article
distributed under the terms of the [Creative
Commons Attribution License \(CC BY\)](#). The
use, distribution or reproduction in other
forums is permitted, provided the original
author(s) and the copyright owner(s) are
credited and that the original publication in
this journal is cited, in accordance with
accepted academic practice. No use,
distribution or reproduction is permitted
which does not comply with these terms.

Corrigendum: Lower serum insulin-like growth factor 2 level in patients with bipolar disorder is associated with the severity of manic symptoms during manic episodes

Shi-Yi Ye^{1,2†}, Ying Zhao^{3†}, Zhao-Bo Liu^{4†}, Cui-Pin Luo¹,
Jian-Wen Xiong^{1,5}, Jin-Qiong Zhan^{1,5}, Yi-Heng Li^{1,5}, Bo Wei^{1,5},
Chun-Nuan Chen^{6*} and Yuan-Jian Yang^{1,5*}

¹Department of Psychiatry and Biological Psychiatry Laboratory, Jiangxi Mental Hospital & Affiliated Mental Hospital, Jiangxi Medical College, Nanchang University, Nanchang, Jiangxi, China, ²The 3rd Clinical Medical College, Jiangxi Medical College, Nanchang University, Nanchang, Jiangxi, China, ³Department of Pharmacy, Union Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, China, ⁴Department of Psychiatry, Third People's Hospital of Ji'an City, Ji'an, China, ⁵Nanchang City Key Laboratory of Biological Psychiatry, Jiangxi Provincial Clinical Research Center on Mental Disorders, Jiangxi Mental Hospital, Nanchang, Jiangxi, China, ⁶Department of Neurology, The Second Clinical Medical College, The Second Affiliated Hospital, Fujian Medical University, Quanzhou, China

KEYWORDS

bipolar disorder, insulin-like growth factor-2, manic symptoms, severity, serum

A Corrigendum on

Lower serum insulin-like growth factor 2 level in patients with bipolar disorder is associated with the severity of manic symptoms during manic episodes

by Ye S-Y, Zhao Y, Liu Z-B, Luo C-P, Xiong J-W, Zhan J-Q, Li Y-H, Wei B, Chen C-N and Yang Y-J (2024). *Front. Psychiatry* 15:1354999. doi: 10.3389/fpsy.2024.1354999

In the published article, an author name was incorrectly written as Yin Zhao. The correct spelling is Ying Zhao.

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.