



Short-term meditation increases blood flow in anterior cingulate cortex and insula

Yi-Yuan Tang^{1,2}*, Qilin Lu³, Hongbo Feng^{3,4}, Rongxiang Tang⁵ and Michael I. Posner²

¹ Department of Psychological Sciences, Texas Tech University, Lubbock, TX, USA

² Department of Psychology, University of Oregon, Eugene, OR, USA

³ Institute of Neuroinformatics and Lab for Body and Mind, Dalian University of Technology, Dalian, China

⁴ First Affiliated Hospital of Dalian Medical University, Dalian, China

⁵ Department of Psychology, The University of Texas at Austin, Austin, TX, USA

Edited by:

Barbara Tomasino, University of Udine, Italy

Reviewed by:

Zoran Josipovic, New York University, USA

Joseph John Loizzo, Weill Cornell Medical College, USA

*Correspondence:

Yi-Yuan Tang, Department of Psychological Sciences, Texas Tech University, Lubbock, TX 79409, USA
e-mail: yiyuan.tang@ttu.edu

Asymmetry in frontal electrical activity has been reported to be associated with positive mood. One form of mindfulness meditation, integrative body-mind training (IBMT) improves positive mood and neuroplasticity. The purpose of this study is to determine whether short-term IBMT improves mood and induces frontal asymmetry. This study showed that 5-days (30-min per day) IBMT significantly enhanced cerebral blood flow (CBF) in subgenual/adjacent ventral anterior cingulate cortex (ACC), medial prefrontal cortex and insula. The results showed that both IBMT and relaxation training increased left laterality of CBF, but only IBMT improved CBF in left ACC and insula, critical brain areas in self-regulation.

Keywords: integrative body-mind training, cerebral blood flow, positive mood, frontal asymmetry, anterior cingulate cortex

INTRODUCTION

Mindfulness meditation has been shown to produce positive effects on psychological wellbeing (Hölzel et al., 2011). As an important benefit of meditation practice, changes in self-reported positive mood or emotion are often observed (Kabat-Zinn et al., 1992; Tang et al., 2007). In particular, one form of meditation, the integrative body-mind training (IBMT) that originates from traditional Chinese medicine, improves attention, self-regulation, and mood after only few hours of training in comparison with relaxation training in a random assignment design (Tang et al., 2007; Ding et al., 2014).

Since late 1970s, frontal EEG asymmetry has had widespread use in measuring individual differences in emotional state. In general, it is believed a left-sided frontal activation indicates a positive emotion, although the evidence is not always consistent with this view (Allen and Kline, 2004; Allen et al., 2004; Cacioppo, 2004; Travis and Arenander, 2004, 2006).

A number of EEG studies have examined the relationship between meditation and frontal asymmetry (Davidson et al., 2003; Moyer et al., 2011). For example, 8-weeks mindfulness-based stress reduction, in comparison to a wait-list control, increased left-sided lateralization of alpha power and decreased negative affect (Davidson et al., 2003). Another study suggested 5-weeks of meditation shifts EEG asymmetry toward a pattern associated with positive emotion compared to a waiting-list control although there was no significant difference between the two groups (Moyer et al., 2011).

However, previous studies have used EEG with low spatial resolution and have not involved an active control group. To our knowledge, only one single photon emission computed tomography (SPECT) imaging study reported a marginal significant left frontal asymmetry of cerebral blood flow (CBF) but this study involved a sample of 12 long-term meditators compared to normal

controls (Newberg et al., 2010) and was not a randomized test with an active control. Thus, we set out to apply brain imaging to investigate the CBF asymmetry induced by short-term training in a relatively large sample of 40 undergraduates with a random assignment to IBMT or relaxation training groups. We hypothesize that IBMT could improve left frontal CBF at resting state, which may underlie the promotion of positive emotion.

MATERIALS AND METHODS

PARTICIPANTS

Forty right-handed Chinese undergraduates at Dalian University of Technology without any training experience and a history of psychiatric or neurological conditions were recruited. They were randomly assigned to IBMT or relaxation group (20:20 each group, age = 22.75 ± 2.02), a brief self-report mood scale, similar to short form of Positive and Negative Affect Schedule (PANAS) was used to measure positive (PA) and negative affect (NA; Moyer et al., 2011; Ding et al., 2014). A written informed consent was obtained and local IRB approved the study.

TRAINING

Participants received 30-min of IBMT or relaxation training from Monday through Friday, with a total of 2.5 h training. IBMT involves body relaxation, mental imagery, and mindfulness training, guided by an IBMT coach and compact disk. Cooperation between the body and the mind is emphasized in facilitating and achieving a meditative state. The trainees concentrated on achieving a balanced state of body and mind. The method stresses no effort to control thoughts, but instead a state of restful alertness that allows a high degree of awareness of body, mind, and external instructions (Tang et al., 2007, 2010; Tang, 2011). Relaxation training involves the relaxing of different muscle groups

over the face, head, shoulders, arms, legs, chest, back, and abdomen, etc., guided by a tutor and compact disk. With eyes closed and in a sequential pattern, one is forced to concentrate on the sensation of relaxation such as the feelings of warmth and heaviness. This progressive training helps the participant achieve physical and mental relaxation and calmness (Tang et al., 2007, 2010).

IMAGE ACQUISITION AND ANALYSIS

Single photon emission computed tomography neuroimaging data were acquired on a double head detector of GE HAWK EYE SPECT system (Millenium VG, GE Healthcare) from all participants before and after a 5-days IBMT or relaxation training. Participants were instructed to stay in a quiet dark room with eyes closed and ears unoccluded for 10 min, and then they were injected with 25 mCi of ^{99m}Tc -ECD. Approximately 20 min following the injection, participants were scanned by SPECT system for 30 min (Tang et al., 2009; Newberg et al., 2010). Two different statistical methods were used to estimate CBF before and after training.

Statistical parametric mapping (SPM¹) was applied to evaluate whole brain CBF differences between these two groups. All images were normalized to the SPECT template in SPM, and then 8 mm Gaussian kernel was used to smooth these images. Two groups' data were compared using two sample *t*-test with ANCOVA for removing global effect. The significance threshold was set at $p < 0.001$ (Tang et al., 2009; Newberg et al., 2010). The location and peaks of activation was identified by *xjview*².

Frontal asymmetry analysis was performed by SPSS (SPSS Inc., Chicago, IL, USA). CBF of the prefrontal regions, including inferior, middle, and superior frontal gyrus were extracted. All brain regions were identified by AAL atlas (Tzourio-Mazoyer et al., 2002). The laterality indexes were calculated by the following formula (Newberg et al., 2010):

$$\text{Laterality Index} = \frac{(\text{Right} - \text{Left})}{0.5 \times (\text{Right} + \text{Left})} \times 100$$

RESULTS

Consistent with our previous studies (Tang et al., 2007; Ding et al., 2014), no significant difference in mood was detected before training between two groups ($p > 0.05$). After training, the IBMT group showed significantly better scores in positive and negative affect in comparison with the RT group; IBMT group (not RT) also showed significant post vs. pre difference in PN and NA (all $p < 0.05$). These results suggested that short-term IBMT induced higher positive mood and lower negative mood states than relaxation training. However there was no significant time by group interaction.

Imaging results showed that two groups did not differ in frontal CBF asymmetry before training ($p > 0.05$). However, the IBMT group (not relaxation group) had a significant CBF increase in subgenual/adjacent ventral anterior cingulate cortex

Table 1 | Statistical parametric mapping (SPM) activation results in integrative body-mind training (IBMT) group (after vs. before).

Brodmann area	<i>p</i> value
Left hemisphere	
BA 32	0.043
Insula	0.026
BA 10	0.001
Right hemisphere	
BA 25	0.005
BA 32	0.021

(ACC; BA 25, BA 32), medial prefrontal cortex (BA 10) and insula after training (all $p < 0.05$). The group \times session interaction was significant for BA 25 [$F(1,38) = 4.652$; $p < 0.05$], BA 10 [$F(1,38) = 10.652$; $p < 0.01$], and marginally significant for insula [$F(1,38) = 3.73$; $p = 0.06$], respectively. SPM activation results of Brodmann areas in IBMT group were shown in Table 1 (after vs. before).

There were no significant differences in before vs. after laterality scores in frontal areas between the two groups. Instead, both groups showed significant differences favoring the left hemisphere in tIFG (triangular part of inferior frontal gyrus, $p = 0.033$) and oSFG (orbital part of superior frontal gyrus, $p = 0.045$) as shown in Figure 1. The IBMT group revealed larger laterality indexes that implied greater left frontal CBF lateralization, compared to relaxation group after training, but the group \times session interaction was not significant.

DISCUSSION

Consistent with our previous studies, short-term IBMT improves CBF in the midfrontal lobe and insula compared to the relaxation training (Tang et al., 2007, 2009). This result is in line with the neural correlates of mindfulness meditation (Cahn and Polich,




FIGURE 1 | Statistical results of laterality indexes in frontal regions.

Left: significant frontal laterality differences in integrative body-mind training (IBMT) and relaxation groups. Right: axial section of two frontal regions ($z = -1$). tIFG, triangular part of inferior frontal gyrus; oSFG, orbital part of superior frontal gyrus. * $p < 0.05$.

¹<http://www.fil.ion.ucl.ac.uk/spm/>

²<http://www.alivelearn.net/xjview8/>

2006; Hölzel et al., 2011; Tang and Posner, 2014). The specific brain areas showing greater CBF following IBMT training suggests that IBMT works, in part, by improving self-regulation (Posner and Rothbart, 2006; Tang et al., 2007; Hölzel et al., 2011; Tang, 2011; Tang and Posner, 2014).

Our study, using the frontal asymmetry analysis, indicated only a few hours of training increases left-sided CBF asymmetry in both groups but IBMT's increase is greater than that of relaxation training.

In previous studies meditation training promoted positive emotions and degraded negative affect (Kabat-Zinn et al., 1992; Tang et al., 2007; Ding et al., 2014). Studies have found 8-weeks mindfulness meditation produced EEG frontal asymmetry and reduced negative mood state (Davidson et al., 2003). In our previous study of 80 Chinese undergraduates, a few hours of IBMT improved positive moods using a 65-item Profile of Mood States (POMS; Tang et al., 2007, 2009; Newberg et al., 2010; Tang, 2011). The current study did not repeat the POMS measures, instead, a brief self-report positive and negative mood scale was used before and after training (Moyer et al., 2011; Ding et al., 2014). After training a significant between-group differences were observed and IBMT group performed better in mood states than relaxation group. However, we did not find a significant difference in the group by time of testing interaction in mood. It is likely that the brief measure of mood was not as sensitive as the POMS. These findings could also be due to power considerations and sensitivity of imaging techniques (Gatzke-Kopp et al., 2001), but with 20 participants per group our study has a larger number than most imaging studies.

Although studies have shown mindfulness meditation increased left-sided anterior activation, a pattern previously associated with positive affect (Davidson et al., 2003; Moyer et al., 2011) comparing to a wait-list control group, alternation of EEG frontal asymmetry was not detected during transcendental meditation, but the participants' emotion states were improved (Travis and Arenander, 2004, 2006). Given the low spatial resolution of EEG, it remains to be elusive whether the brain regions involved in these studies are the same. In our study both meditation and relaxation resulted in increased left lateralization, but only IBMT changed mood and brain activity in ACC and insula associated with self-control, consistent with our series of research (Tang et al., 2007, 2009, 2010). Since we did not detect the significant correlation between frontal asymmetry and mood using EEG, future studies will be needed to demonstrate the mechanisms of changes in frontal asymmetry and mood state using ERP/fMRI fusion approach.

Our study shows that few hours of IBMT increases resting CBF in specific brain areas often shown to be involved in attention and self-regulation (Posner and Rothbart, 2006; Tang et al., 2009, 2010; Hölzel et al., 2011; Tang, 2011). However, both IBMT and relaxation training increase frontal asymmetry significantly, although IBMT has a somewhat larger increase. The IBMT group has better overall mood state following training, but it is not clear if the brain differences found in this study are the cause of the changes in affect and mood.

ACKNOWLEDGMENTS

We thank the staff of the Institute of Neuroinformatics for data collection. This work was supported by the Office of Naval Research.

REFERENCES

- Allen, J. J., Coan, J. A., and Nazarian, M. (2004). Issues and assumptions on the road from raw signals to metrics of frontal EEG asymmetry in emotion. *Biol. Psychol.* 67, 183–218. doi: 10.1016/j.biopsycho.2004.03.007
- Allen, J. J., and Kline, J. P. (2004). Frontal EEG asymmetry, emotion, and psychopathology: the first, and the next 25 years. *Biol. Psychol.* 67, 1–5. doi: 10.1016/j.biopsycho.2004.03.001
- Cacioppo, J. T. (2004). Feelings and emotions: roles for electrophysiological markers. *Biol. Psychol.* 67, 235–243. doi: 10.1016/j.biopsycho.2004.03.009
- Cahn, B. R., and Polich, J. (2006). Meditation states and traits: EEG, ERP, and neuroimaging studies. *Psychol. Bull.* 132, 180–211. doi: 10.1037/0033-2909.132.2.180
- Davidson, R. J., Kabat-Zinn, J., Schumacher, J., Rosenkranz, M., Muller, D., Santorelli, S. F., et al. (2003). Alterations in brain and immune function produced by mindfulness meditation. *Psychosom. Med.* 65, 564–570. doi: 10.1097/01.PSY.0000077505.67574.E3
- Ding, X., Tang, Y. Y., Tang, R., and Posner, M. I. (2014). Improving creativity performance by short-term meditation. *Behav. Brain Funct.* 10:9. doi: 10.1186/1744-9081-10-9
- Gatzke-Kopp, L. M., Raine, A., Buchsbaum, M., and LaCasse, L. (2001). Temporal lobe deficits in murderers: EEG findings undetected by PET. *J. Neuropsychiatry Clin. Neurosci.* 13, 486–491.
- Hölzel, B. K., Lazar, S. W., Gard, T., Schuman-Olivier, Z., Vago, D. R., and Ott, U. (2011). How does mindfulness meditation work? Proposing mechanisms of action from a conceptual and neural perspective. *Perspect. Psychol. Sci.* 6, 537–559. doi: 10.1177/1745691611419671
- Kabat-Zinn, J., Massion, A. O., Kristeller, J., Peterson, L. G., Fletcher, K. E., Pbert, L., et al. (1992). Effectiveness of a meditation-based stress reduction program in the treatment of anxiety disorders. *Am. J. Psychiatry* 149, 936–943. doi: 10.1176/ajp.149.7.936
- Moyer, C. A., Donnelly, M. P. W., Anderson, J. C., Valek, K. C., Huckaby, S. J., Wiederholt, D. A., et al. (2011). Frontal electroencephalographic asymmetry associated with positive emotion is produced by very brief meditation training. *Psychol. Sci.* 22, 1277–1279. doi: 10.1177/0956797611418985
- Newberg, A. B., Wintering, N., Waldman, M. R., Amen, D., Khalsa, D. S., and Alavi, A. (2010). Cerebral blood flow differences between long-term meditators and non-meditators. *Conscious. Cogn.* 19, 899–905. doi: 10.1016/j.concog.2010.05.003
- Posner, M. I., and Rothbart, M. K. (2006). *Educating the Human Brain*. Washington, DC: APA Press.
- Tang, Y. Y. (2011). Mechanism of integrative body-mind training. *Neurosci. Bull.* 27, 383–388. doi: 10.1007/s12264-011-1141-2
- Tang, Y. Y., Lu, Q., Geng, X., Stein, E. A., Yang, Y., and Posner, M. I. (2010). Short-term meditation induces white matter changes in the anterior cingulate. *Proc. Natl. Acad. Sci. U.S.A.* 107, 15649–15652. doi: 10.1073/pnas.1011043107
- Tang, Y. Y., Ma, Y., Fan, Y., Feng, H., Wang, J., Feng, S., et al. (2009). Central and autonomic nervous system interaction is altered by short term meditation. *Proc. Natl. Acad. Sci. U.S.A.* 106, 8865–8870. doi: 10.1073/pnas.0904031106
- Tang, Y. Y., Ma, Y., Wang, J., Fan, Y., Feng, S., Lu, Q., et al. (2007). Short term meditation training improves attention and self regulation. *Proc. Natl. Acad. Sci. U.S.A.* 104, 17152–17156. doi: 10.1073/pnas.0707678104
- Tang, Y. Y., and Posner, M. I. (2014). Training brain networks and states. *Trends Cogn. Sci.* 18, 345–350. doi: 10.1016/j.tics.2014.04.002
- Travis, F., and Arenander, A. (2004). EEG asymmetry and mindfulness meditation. *Psychosom. Med.* 66, 147–147. doi: 10.1097/00006842-200401000-00020
- Travis, F., and Arenander, A. (2006). Cross-sectional and longitudinal study of effects of transcendental meditation practice on interhemispheric frontal asymmetry and frontal coherence. *Int. J. Neurosci.* 116, 1519–1538. doi: 10.1080/00207450600575482

Tzourio-Mazoyer, N., Landeau, B., Papathanassiou, D., Crivello, F., Etard, O., Delcroix, N., et al. (2002). Automated anatomical labeling of activations in SPM using a macroscopic anatomical parcellation of the MNI MRI single-subject brain. *Neuroimage* 15, 273–289. doi: 10.1006/nimg.2001.0978

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 28 October 2014; accepted: 11 February 2015; published online: 26 February 2015.

Citation: Tang Y-Y, Lu Q, Feng H, Tang R and Posner MI (2015) Short-term meditation increases blood flow in anterior cingulate cortex and insula. *Front. Psychol.* 6:212. doi: 10.3389/fpsyg.2015.00212

This article was submitted to *Cognition*, a section of the journal *Frontiers in Psychology*. Copyright © 2015 Tang, Lu, Feng, Tang and Posner. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.