



Psychometric Properties of the Chinese Version of Anxiety Sensitivity Index-3 in Women Diagnosed With Breast Cancer

Yan Han^{1,2†}, Jiang Zhu^{1,2†}, Lingyan Li³, Huan Zhou¹, Shichen Li¹, Jinqiang Zhang¹, Jie Fan¹, Yanjie Yang⁴, Xingwei Luo^{1*} and Xiongzhao Zhu^{1,2*}

¹ Medical Psychological Center, Second Xiangya Hospital, Central South University, Changsha, China, ² Medical Psychological Institute of Central South University, Changsha, China, ³ School of Nursing, Nanchang University, Nanchang, China, ⁴ School of Public Health, Harbin Medical University, Harbin, China

OPEN ACCESS

Edited by:

Laura Badenes-Ribera,
University of Valencia, Spain

Reviewed by:

Javier Sánchez-Rosas,
National University of Córdoba,
Argentina
Cesar Merino-Soto,
University of San Martín de Porres,
Peru

*Correspondence:

Xingwei Luo
luoxingwei6@csu.edu.cn
Xiongzhao Zhu
xiongzhaozhu@csu.edu.cn

[†]These authors have contributed
equally to this work and share first
authorship

Specialty section:

This article was submitted to
Quantitative Psychology
and Measurement,
a section of the journal
Frontiers in Psychology

Received: 09 October 2019

Accepted: 06 January 2020

Published: 07 February 2020

Citation:

Han Y, Zhu J, Li L, Zhou H, Li S,
Zhang J, Fan J, Yang Y, Luo X and
Zhu X (2020) Psychometric Properties
of the Chinese Version of Anxiety
Sensitivity Index-3 in Women
Diagnosed With Breast Cancer.
Front. Psychol. 11:12.
doi: 10.3389/fpsyg.2020.00012

Background: Anxiety sensitivity (AS) is a trait-like predisposing factor for the prevalence of anxiety in patients diagnosed with breast cancer. The Anxiety Sensitivity Index-3 (ASI-3) has been widely used in both general and clinical samples for measuring AS. However, the data about its psychometric properties in women with breast cancer are deficient. In addition, there is no evidence proving the measure equivalence of ASI-3 across sociodemographic variables in the specific sample. Thus, the present study examined the psychometric properties and conducted measure equivalence testing of ASI-3 in Chinese women diagnosed with breast cancer.

Methods: This study included 815 Chinese women diagnosed with breast cancer. Single group confirmatory factor analysis (CFA) was administrated to detect the factorial validity of ASI-3, and multigroup CFAs were conducted to test the measurement equivalence of ASI-3 across various sociodemographic variables. The reliability of ASI-3 was tested by Cronbach's alpha coefficient, Guttman split-half coefficient, McDonald's omega coefficient, and test-retest coefficient. The standardized factor loadings, construct reliability, and the average variance extracted of factors were used to measure the construct validity of ASI-3, and the partial correlations were conducted to examine the criterion-related validity of ASI-3.

Results: The ASI-3 had satisfactory reliability and validity in Chinese women diagnosed with breast cancer; three-factor model fitted the data well in CFA and reached partial strict invariances across age, education, and residence groups.

Conclusion: This study explored the psychometric properties and measurement invariance across sociodemographic variables of ASI-3 in Chinese women diagnosed with breast cancer. Our results not only proved that the ASI-3 is an appropriate instrument for measuring AS but also deepened the understanding of ASI-3 in Chinese women with malignancy.

Keywords: anxiety sensitivity, Anxiety Sensitivity Index-3, breast cancer, psychometric properties, measurement invariance

INTRODUCTION

Nowadays, breast cancer is the most common cancer and becomes a leading cause of cancer-related death in Chinese women (Fan et al., 2014). People often experience varying levels of anxiety after the diagnosis of breast cancer, which can bring about harmful effects on their mental health (Burgess et al., 2005). Many studies have proved that anxiety severity was correlated with the levels of anxiety sensitivity (AS) and AS is a trait-like predisposing factor of anxiety (Olatunji and Wolitzky-Taylor, 2009; Knapp et al., 2016; Mohammadkhani et al., 2016). AS refers to individuals' fear of anxiety-related sensory arousal from the beliefs that the sensation have adverse effects on their body, cognition, and social evaluation (Taylor et al., 2007). Individuals may experience more severe anxiety symptoms by misinterpreting their physical sensations as danger signals (Olatunji and Wolitzky-Taylor, 2009). For example, there is evidence suggesting that women with breast cancer showed more health anxiety, which is strongly associated with the cognitive AS (Jones et al., 2014). Ren et al. (2018) found that neuroticism and AS can positively predict insomnia in women with breast cancer. In general, AS is a specific vulnerability trait for the prevalence of anxiety in patients diagnosed with breast cancer.

The most common instrument for measuring AS is the Anxiety Sensitivity Index-3 (ASI-3), which was established by Taylor et al. (2007) and was proved to have adequate reliability, validity, as well as stable three-factor structure in the United States and Canadian non-clinical samples. At present, ASI-3 has been widely used in different countries. For instance, Ghisi et al. (2016) verified that the ASI-3 was a reliable and valid scale to evaluate AS in the Italian community sample; Kemper et al. (2012) demonstrated that the ASI-3 had a high degree of internal consistency and construct validity in German clinical samples. In addition to these, ASI-3 has been evidenced to have excellent psychometric properties among other countries such as Korean, China, Turkish, and so on (Mantar et al., 2010; Lim and Kim, 2012; Cai et al., 2018). Besides, ASI-3 was also widely applied in various clinical populations such as psychotic disorders, migraine, and cranial meningiomas (Rifkin et al., 2015; Farris et al., 2019; Wagner A. et al., 2019). In terms of factor structure, most previous studies have supported that ASI-3 has a three-factor structure, named physical (six items), social (six items), and cognitive concerns (six items), respectively (Taylor et al., 2007; Kemper et al., 2012; Lim and Kim, 2012; Cai et al., 2018). The physical concerns are positively correlated with body vigilance, social concerns are positively correlated with fear of negative evaluation, and cognitive concerns are most strongly correlated with depression symptoms (Kemper et al., 2012). All in all, the ASI-3 is a satisfactory instrument to evaluate AS in various samples.

Anxiety sensitivity is a trait-like cognitive vulnerability for elevating anxiety levels in patients diagnosed with breast cancer (Olatunji and Wolitzky-Taylor, 2009). However, studies about the application of the ASI-3 in patients with breast cancer are deficient. To explore that whether ASI-3 is also an appropriate tool for measuring AS in patients diagnosed with breast cancer, one aim of the present study was to examine the factor

structure, reliability, and validity of ASI-3 in a sample of Chinese women diagnosed with breast cancer. Beyond these, according to previous research, patients with breast cancer in different demographic groups were found to score differently on anxiety-related scales. For example, in Hassan et al.'s (2015) study, the younger age group had a higher risk of anxiety, Wagner J. F. et al. (2019) found patients with young age and low education level performed higher anxiety level. Long-term residence was found to have a significant effect on anxiety in a Chinese sample with breast cancer (Li et al., 2016). More work is needed to explore whether there are differences in vulnerability factors such as AS across various sociodemographic groups in patients with breast cancer. People have studied the invariance of measurement properties of ASI-3 across sex, sexual minority status, age, and race/ethnicity in university students (Jardin et al., 2018). To our knowledge, there is no study proving the measurement invariance of ASI-3 across different demographic groups in patients with breast cancer. It is meaningless to conduct cross-groups comparison of mean differences if the measurement invariance is not established (Schmitt and Kuljanin, 2008). Hence, in current study, we also examined the invariance of factor structure and different performance on ASI-3 across different demographic variables in Chinese women diagnosed with breast cancer.

MATERIALS AND METHODS

Participants

We recruited 1,131 female patients diagnosed with breast cancer at two hospitals in Changsha from March 2011 to March 2016. According to the inclusion criteria, e.g., (1) 20–75 years old, (2) been diagnosed with breast cancer pathologically, (3) are able to read Chinese, (4) were voluntary and completed written informed consent, and (5) no history of major psychiatric disorders or substance abuse, there are 986 qualified candidates. Subsequently, we removed 171 subjects whose questionnaires were incomplete. Finally, this study included 815 patients.

Measures

Anxiety Sensitivity Index-3

It is an 18-item self-report tool for assessing anxiety-related physical (items 3, 4, 7, 8, 12, and 15), cognitive (items 2, 5, 10, 14, 16, and 18), or social (items 1, 6, 9, 11, 13, and 17) concerns. It is a 5-point Likert scale. Each item is scored from 0 to 4, with 0 meaning "I agree very little" and 4 indicating "I agree very much." Total score is the sum of the score on all items, ranging from 0 to 72. The higher scores on the scale present more severe AS level (Cai et al., 2018).

Hamilton Rating Scale for Anxiety

Hamilton et al. developed this questionnaire to assess the anxiety level, which includes 14 items. Half of the items are aimed to test psychic anxiety, and the remaining items measure somatic anxiety. The values on each item range from 0 to 4 (0 = mild anxiety, 2 = moderate anxiety, 3 = severe anxiety, and 4 = very severe or grossly disabling anxiety), and the total scores range

from 0 to 56 (Garalejić et al., 2010). The Hamilton Rating Scale for Anxiety (HAMA) is a reliable and valid instrument for measuring anxiety symptoms and widely used in China (Wang C. et al., 2011; Zimmerman et al., 2017).

Center for Epidemiologic Studies Depression Scale

The Chinese version Center for Epidemiologic Studies Depression Scale (CESD) is a reliable and effective instrument to assess individuals' depressive symptoms, including 20 items. Each item is rated from 0 (rarely or none of the time) to 3 (most or all of the time); total scores range from 0 to 60. Higher CESD scores indicated more severe depressive symptoms. In addition, the CESD has a three-factor structure, which includes positive affect, interpersonal problems, as well as depressive mood and somatic symptoms (Zhang et al., 2012).

Statistical Analysis

SPSS version 22.0 and Mplus version 7.0 were used in this study. Single-group confirmatory factor analysis (CFA) was administrated to verify the three-factor model of ASI-3. Owing to that fact the chi-square (χ^2) test is sensitive to the sample size, the goodness-of-fit was assessed by the comparative fit index (CFI), Tucker–Lewis index (TLI), root-mean-square error of approximation (RMSEA), and standardized root mean square residual (SRMR). CFI and TLI > 0.90, RMSEA and SRMR < 0.08 indicate an acceptable model fit (Sun, 2005).

Multigroup CFAs were conducted to test the measurement equivalence of ASI-3 across age (younger and older), levels of education (lower and higher), and places of residence (rural and urban). We assessed configural, metric, scalar, and strict invariances across groups in succession, which demonstrated that the composition of latent variables, factor loading of each item, intercepts of the observed variables, and latent variable variation are equal between groups, respectively (Schmitt and Kuljanin, 2008). Given that the χ^2 change test is also sensitive to the sample size, the results mainly consider the value of the difference of CFI (Δ CFI). According to Cheung and Rensvold (2002) the equivalent model is considered acceptable when Δ CFI < 0.010. As all items obtained significant skewness and kurtosis values ($P < 0.001$) in the Kolmogorov–Smirnov normality test, which means that the data did not fit the normal distribution, so the robust maximum likelihood with mean and variance adjustments estimator was used in all CFAs (Muthén and Muthén, 2012).

The Cronbach's alpha coefficient, Guttman split-half coefficient, and McDonald's omega coefficient were used to test the internal consistency of the ASI-3 (Guttman, 1945; Cronbach, 1951; McDonald, 1985). To understand the test–retest reliability of the ASI-3, we randomly recruited 24 patients to complete the ASI-3 again in 2 weeks later, which can test the stability of the ASI-3 across time. Moreover, Mann–Whitney *U*-tests were used to compare the mean differences of ASI-3 scores between age, education, and residence groups.

In terms of construct validity, we tested the criterion-related validity, convergent validity, and discriminant validity of the ASI-3. Partial correlations were conducted to assess the criterion-related validity of ASI-3 against the HAMA and CESD when controlling the effect of age. In addition, the Harman's

single factor test was performed to examine the effects caused by common method variance. The hypothesis is that if the covariation could largely be explained by method variance, a single factor (method factor) should exist in the variables of these measures (Harris and Mossholder, 1996). According to Hair et al. (2010), the convergent validity could be measured by standardized factor loadings, construct reliability (CR), and average variance extracted (AVE) of factors. If the standardized factor loadings of ASI-3 are >0.5 and statistically significant, the value of CR and AVE of each factor is higher than 0.7 and 0.5, respectively, the ASI-3 has satisfied convergent validity. Discriminant validity was examined by comparing the AVE with the square of the latent correlations among factors. When the AVE of the ASI-3 is above the square of the latent correlations, the ASI-3 has an acceptable discriminant validity (Hair et al., 2010).

RESULTS

Descriptive Statistics

A sample of 815 female patients diagnosed with breast cancer from two hospitals in Changsha participated in this study. These participants aged 26–71 years (mean age = 47.66 years, SD = 8.35). The specific information is summarized in **Tables 1, 2**.

Confirmatory Factor Analysis

Confirmatory factor analysis was administrated to examine the three-factor model of ASI-3 in this study. The fit indices were as follows: $\chi^2 = 474.881$, $df = 132$, CFI = 0.930, TLI = 0.919, RMSEA = 0.056 (90% confidence interval = 0.051–0.062), and SRMR = 0.046. All indices meet the fitting requirements, demonstrating that the three-factor model had adequate factorial validity in current sample. The standardized factor loadings

TABLE 1 | Demographic and disease-related information of the patients.

Item	N (%)
Years of age	
≤50	555 (68.1%)
>50	260 (31.9%)
Marital status	
Married	765 (93.8%)
Single	3 (0.4%)
Divorced	29 (3.6%)
Widowed	18 (2.2%)
Educational level	
≤6 years	148 (18.2%)
>6 years	667 (81.8%)
Places of residence	
Urban	409 (50.2%)
Rural	406 (49.8%)
Period	
Preoperative	235 (28.8%)
Postoperative	550 (67.5%)
Convalescent	30 (3.7%)

obtained in CFA and the latent correlations among factors are presented in **Table 3**.

Measurement Invariance of ASI-3 Three-Factor Model Across Demographic Groups

The fit indices of CFA in demographic subgroups are presented in **Table 4**. We found that the fitting indices did not totally meet the fitting conditions in older group (>50), lower education group (≤6 years), and urban group. According to the modified

index (MI), we selected the largest value of MI in turn to free the parameters. In the older group (>50), there were correlated measurement errors between items 2 and 5 [Cov(y₂, y₅)] as well as items 11 and 13 [Cov(y₁₁, y₁₃)], the error correlation coefficient was 0.595 (*p* < 0.01) for Cov(y₂, y₅) and 0.489 (*p* < 0.01) for Cov(y₁₁, y₁₃); with regard to lower education group (≤6 years), the measurement error between items 10 and 14 was correlated [Cov(y₁₀, y₁₄)], and the correlation coefficient was 0.611 (*p* < 0.01). Concerning the urban group, we correlated the measurement error of item 2 with item 5, and the error correlation coefficient was 0.510 (*p* < 0.01). The three-factor

TABLE 2 | The descriptive statistics and the reliability parameters of the instruments used in this study.

Instruments	<i>M</i>	<i>SD</i>	Skewness	Kurtosis	Cronbach's alpha coefficients (90% CI)	McDonald's omega coefficient (90% CI)
ASI-3 (<i>N</i> = 815)	13.659	11.766	1.075	0.779		
Physical concerns	3.993	4.488	0.990	0.082	0.936 (0.930–0.942)	0.902 (0.890–0.914)
Social concerns	5.371	5.221	0.960	0.378	0.924 (0.917–0.930)	0.885 (0.871–0.899)
Cognitive concerns	4.296	4.474	1.228	1.253	0.935 (0.929–0.941)	0.898 (0.885–0.911)
HAMA (<i>N</i> = 604)	10.320	5.271	0.170	−0.227	0.774 (0.751–0.796)	0.733 (0.715–0.752)
CESD (<i>N</i> = 428)	19.923	10.099	−0.051	−1.096	0.683 (0.646–0.719)	0.783 (0.766–0.799)

ASI-3, Anxiety Sensitivity Index-3; HAMA, Hamilton Anxiety Scale; CESD, Center for Epidemiologic Studies Depression Scale.

TABLE 3 | Item loadings of three-factor structure and latent correlations among factors in CFA.

		Standardized factor loadings		
		Factor I	Factor II	Factor III
Physical concerns				
Item 3	It scares me when my heart beats rapidly.	0.844		
Item 4	When my stomach is upset, I worry that I might be seriously ill.	0.844		
Item 7	When my chest feels tight, I get scared that I won't be able to breathe properly.	0.885		
Item 8	When I feel pain in my chest, I worry that I'm going to have a heart attack.	0.903		
Item 12	When I notice my heart skipping a beat, I worry seriously wrong with me.	0.775		
Item 15	When my throat feels tight, I worry that I could choke to death.	0.816		
Social concerns				
Item 1	It is important for me not to appear nervous.		0.740	
Item 6	When I tremble in the presence of others, I fear what people might think of me.		0.816	
Item 9	I worry that other people will notice my anxiety.		0.917	
Item 11	It scares me when I blush in front of people.		0.838	
Item 13	When I begin to sweat in a social situation, I fear people will think negatively of me.		0.870	
Item 17	I think it would be horrible for me to faint in public.		0.750	
Cognitive concerns				
Item 2	When I cannot keep my mind on a task, I worry that I might be going crazy.			0.851
Item 5	It scares me when I am unable to keep my mind on a task.			0.883
Item 10	When I feel "spacey" or spaced out, I worry that I may be mentally ill.			0.830
Item 14	When my thoughts seem to speed up, I worry that I might be going crazy.			0.834
Item 16	When I have trouble thinking clearly, I worry there is something wrong with me.			0.846
Item 18	When my mind goes blank, I worry there is something terribly wrong with me.			0.800
		Latent correlations (90% CI)		
Physical concerns with social concerns		0.495 (0.440–0.549)		
Physical concerns with cognitive concerns		0.531 (0.482–0.580)		
Social concerns with cognitive concerns		0.658 (0.616–0.700)		

All of the standardized factor loadings are statistically significant.

TABLE 4 | Goodness-of-fit indices of confirmatory factor analysis (CFA) in demographic subgroups.

Item		χ^2	df	CFI	TLI	RMSEA (90% CI)	SRMR
Years of age	≤50	393.432	132	0.917	0.903	0.060 (0.053–0.067)	0.048
	>50	261.747	132	0.897	0.880	0.061 (0.051–0.072)	0.066
Modification	>50*§	223.255	130	0.926	0.912	0.053 (0.041–0.064)	0.062
Educational level	≤6 years	210.817	132	0.897	0.881	0.064 (0.047–0.079)	0.048
	>6 years	413.639	132	0.920	0.908	0.057 (0.050–0.063)	0.051
Modification	≤6 years#	197.065	131	0.914	0.900	0.058 (0.041–0.075)	0.047
Places of residence	Urban	332.046	132	0.903	0.887	0.061 (0.053–0.069)	0.057
	Rural	326.636	132	0.924	0.912	0.057(0.049–0.065)	0.056
Modification	Urban*	304.219	131	0.916	0.901	0.057 (0.049–0.065)	0.053

*Items 2 and 5 correlated. §Items 11 and 13 correlated. #Items 10 and 14 correlated. CFI, comparative fit index; TLI, Tucker–Lewis index; RMSEA, root-mean-square error of approximation; SRMR, standardized root mean square residual.

model of ASI-3 had adequate fit among all subgroups after freed above parameters which are also summarized in **Table 4**. Actually, any parameter to be released must be highly consistent with the theory (MacCallum et al., 1992). We found that the description of item 2 (When I cannot keep my mind on a task, I worry that I might be going crazy) and item 5 (It scares me when I am unable to keep my mind on a task) are similar; meanwhile, they both load on cognitive concerns dimension; as for item 11 (It scares me when I blush in front of people) and item 13 (When I begin to sweat in a social situation, I fear people will think negatively of me), both of them express individual's worry about somatic symptoms in front of others and belong to the same factor; item 10 (When I feel “spacey” or spaced out, I worry that I may be mentally ill) and item 14 (When my thoughts seem to speed up, I worry that I might be going crazy) express the concerns about the instability of cognitive state and belong to cognitive concerns.

According to Byrne et al. (1989), baseline models are not required to be identical across groups in invariance testing; therefore, we established the multigroup model based on different baseline models in measurement invariance testing.

First, we examined the measurement invariance across different age groups (≤50 years old vs. >50 years old). In the multigroup model, there were $Cov(y_2, y_5)$ and $Cov(y_{11}, y_{13})$ in the older group (>50), and there is no modification in the younger group (≤50). We test the configural, metric, scalar, and strict invariance between two groups. The ΔCFI between the configural invariance model and metric invariance model was 0.001, ΔCFI between the metric invariance model and scalar invariance model was 0.003, and ΔCFI between the scalar invariance model and strict invariance model was 0.002. All of them were below than 0.01.

Second, we evaluated the equivalence of the factor structure between two groups with lower (≤6 years) and higher (>6 years) educational level, respectively. In the multigroup model, there was just a $Cov(y_{10}, y_{14})$ in the lower education group (≤6 years). All of the ΔCFI s were 0.002 when compared the constrained model with the unconstrained model.

Finally, we tested invariance between urban and rural groups. In the multigroup model, there was a $Cov(y_2, y_5)$ only in the urban group. We found that all invariance models fit the data well. The differences in CFI between the unconstrained model and the constrained model were 0.004 (configural

invariance vs. metric invariance), 0.003 (metric invariance vs. scalar invariance), and 0.004 (scalar invariance vs. strict invariance). **Table 5** summarized the goodness-of-fit statistics of measurement equivalence analysis.

In summary, given the modifications in the multigroup models, there were partial strict invariances across age, education, and residence groups.

Reliability

In our study, ASI-3 was proved to have a three-factor structure. Therefore, we examined the reliability coefficient for each factor of the ASI-3. The statistics of Cronbach's alpha coefficients (0.924–0.936) and McDonald's omega coefficients (0.885–0.902) are summarized in **Table 2**, and the Guttman split-half coefficient of the physical concerns, social concerns, and cognitive concerns was 0.938, 0.906, and 0.918, respectively, suggesting that the three factors of ASI-3 had satisfactory reliability. The test–retest coefficient for the physical concerns, social concerns, and cognitive concerns was 0.775, 0.596, and 0.730, respectively ($P < 0.01$), which revealed a moderate correlation between test and retest results (Lee et al., 2018). Moreover, we summarized the reliability parameters of other instruments used in our study in **Table 2**.

Mann–Whitney U-Tests

The specific information is shown in **Table 6**. We found that there was just a significant difference in cognitive concerns between age groups ($P < 0.01$). Lower education group had significantly higher scores on ASI-3 compared with higher education group except for cognitive concerns dimension ($P < 0.01$). The ASI-3 scores in rural group were significantly higher than urban group ($P < 0.05$). However, the effect sizes of all significant differences were lower than 0.3 (Fritz et al., 2012).

Construct Validity

Harman's single factor test revealed that the one-factor structure did not fit the data well, and the fit indices were as follows: $\chi^2 = 11,715.484$, $df = 1,274$, $CFI = 0.296$, $TLI = 0.268$, $RMSEA = 0.139$ (90% confidence interval = 0.137–0.141), and $SRMR = 0.151$. Therefore, the common method variance did not have a serious effect on our following results.

TABLE 5 | Measurement invariance testing of the Anxiety Sensitivity Index-3 (ASI-3).

Item	χ^2	df	CFI	TLI	RMSEA (90% CI)	Δ CFI
Years of age (≤ 50 vs. > 50)						
Configural invariance	591.776	262	0.919	0.906	0.058 (0.050–0.062)	
Metric invariance	612.192	277	0.918	0.909	0.054 (0.049–0.060)	0.001
Scalar invariance	639.807	292	0.915	0.911	0.054 (0.048–0.060)	0.003
Strict invariance	648.743	310	0.917	0.918	0.052(0.046–0.057)	0.002
Educational level (≤ 6 vs. > 6 years)						
Configural invariance	577.834	263	0.920	0.906	0.054 (0.048–0.060)	
Metric invariance	597.364	278	0.918	0.910	0.053 (0.047–0.059)	0.002
Scalar invariance	622.582	293	0.916	0.912	0.053 (0.047–0.058)	0.002
Strict invariance	633.086	311	0.918	0.919	0.050(0.045–0.056)	0.002
Places of residence (urban vs. rural)						
Configural invariance	598.601	263	0.922	0.909	0.056 (0.050–0.062)	
Metric invariance	631.396	278	0.918	0.910	0.056 (0.050–0.062)	0.004
Scalar invariance	660.025	293	0.915	0.911	0.055 (0.050–0.061)	0.003
Strict invariance	694.632	311	0.911	0.912	0.055(0.050–0.061)	0.004

CFI, comparative fit index; TLI, Tucker–Lewis index; RMSEA, root-mean-square error of approximation; Δ CFI, change in comparative fit index.

TABLE 6 | Comparison of ASI-3 scores across groups.

Variables			Mean difference	P	r
Years of age	≤ 50	> 50			
	ASI-3	13.85 \pm 11.81	13.26 \pm 11.67	0.59	0.378
	Physical concerns	3.91 \pm 4.21	4.17 \pm 4.63	0.26	0.622
	Cognitive concerns	4.57 \pm 4.56	3.71 \pm 4.24	0.86	0.004
	Social concerns	5.37 \pm 5.10	5.38 \pm 5.47	0.01	0.622
Educational level	≤ 6 years	> 6 years			
	ASI-3	17.10 \pm 13.84	12.90 \pm 11.12	4.20	0.002
	Physical concerns	5.39 \pm 4.76	3.68 \pm 4.37	1.71	0.000
	Cognitive concerns	5.05 \pm 5.18	4.13 \pm 4.29	0.92	0.102
	Social concerns	6.66 \pm 5.89	5.09 \pm 5.02	1.57	0.008
Places of residence	Urban	Rural			
	ASI-3	12.55 \pm 11.04	14.78 \pm 12.37	2.23	0.013
	Physical concerns	3.70 \pm 4.46	4.29 \pm 4.50	0.59	0.018
	Cognitive concerns	3.95 \pm 4.25	4.64 \pm 4.67	0.69	0.027
	Social concerns	4.89 \pm 4.96	5.85 \pm 5.43	0.96	0.011

ASI-3, Anxiety Sensitivity Index-3.

Partial correlations were conducted to test the criterion-related validity of ASI-3 when controlling for the effect of age and shown in **Table 7**. We found that the scores of the three factors were highly and positively interrelated with total score ($r = 0.777$ – 0.867 , $P < 0.01$) and moderately correlated with each other ($r = 0.488$ – 0.581 , $P < 0.01$). The significant correlation between ASI-3 total score and HAMA ($r = 0.505$, $P < 0.01$) revealed that the ASI-3 had an acceptable empirical validation. Furthermore, the cognitive concerns dimension had greater correlations with the total score ($r = 0.394$, $P < 0.01$) and two factors (positive affect, $r = -0.242$, $P < 0.01$; depressive mood and somatic symptoms, $r = 0.436$, $P < 0.01$) of CESD comparing to other factors and total scale of ASI-3, while the social concerns had a greatest correlation with the interpersonal problems dimension of CESD ($r = 0.222$, $P < 0.01$). In addition, the standardized factor loadings for all items in ASI-3 ranged from 0.740 to 0.917 and statistically significant, which are presented in **Table 3**. The value of AVE of each factor ranged from 0.678 to 0.715, and the

value of CR ranged from 0.926 to 0.938, which are summarized in **Table 8**. The standardized factor loadings, AVE of factors, and CR were united to suggest that the ASI-3 had strong convergent validity. In addition, the values of AVE of three factors were higher than the square of the latent correlations among factors (ranged from 0.245 to 0.433), indicating a satisfied discriminant validity of ASI-3.

DISCUSSION

Anxiety Sensitivity Index-3 is a widely used instrument for measuring AS. This study is aimed at exploring the psychometric properties of ASI-3 in Chinese women with breast cancer. Many previous studies have proven that the ASI-3 had a three-factor structure when applied in different cultural backgrounds and populations (Taylor et al., 2007; Mantar et al., 2010; Lim and Kim, 2012; Ghisi et al., 2016). To verify the factor validity of ASI-3, we

TABLE 7 | Partial correlations among ASI-3 and other measures.

	ASI-3	Physical concerns	Cognitive concerns	Social concerns
Physical concerns	0.777**			
Cognitive concerns	0.835**	0.488**		
Social concerns	0.867**	0.513**	0.581**	
HAMA	0.505**	0.294**	0.492**	0.450**
CESD	0.318**	0.130**	0.394**	0.252**
CESD – positive affect	–0.189**	–0.054	–0.242**	–0.159**
CESD – interpersonal problems	0.187**	0.117*	0.110*	0.222**
CESD – depressive mood and somatic symptoms	0.338**	0.129**	0.436**	0.256**

The effects of age were included in the analysis. ASI-3, Anxiety Sensitivity Index-3; HAMA, Hamilton Anxiety Scale; CESD, Center for Epidemiologic Studies Depression Scale. * $P < 0.05$. ** $P < 0.01$.

TABLE 8 | The value of construct reliability (CR) and the average variance extracted (AVE) for each factor in ASI-3.

Factor	CR	AVE
Physical concerns	0.938	0.715
Social concerns	0.926	0.678
Cognitive concerns	0.935	0.707

CR, construct reliability; AVE, average variance extracted.

conducted CFA in current sample. The satisfactory fitting indices illustrated that the three-factor model was practicable in Chinese women with breast cancer. The standardized factor loadings and the latent correlations among factors testified that the three factors are not only relative independent but also correlated with each other.

Measurement equivalence is the premise for comparing the scores on ASI-3 across different groups (Schmitt and Kuljanin, 2008); therefore, we conducted multigroup CFA to test the measurement invariance before compared the scores on ASI-3 across various sociodemographic groups. We found that the baseline models were different across groups when analyzing the three-factor structure separately for each group. Specifically, there were $Cov(y_2, y_5)$ and $Cov(y_{11}, y_{13})$ in the older group (> 50), $Cov(y_{10}, y_{14})$ in the lower education group (≤ 6 years), and $Cov(y_2, y_5)$ in the urban group. All of the modifications were based on the MI applied by Mplus and sufficient theoretical meaning because the three pairs of correlated items have similar directionality and load on the same factor in our study. Particularly, for the older group and lower education group, it might be difficult for them to distinguish the related items due to their relatively lower faculty of understanding. Moreover, the sample size could affect the analysis of CFA and model modifications (MacCallum et al., 1992). In terms of the modification in the urban group, generally, urban residents are more engaged in mental activities, while rural residents are more engaged in manual labor; thus, when people cannot keep focus on their work, urban residents are more likely to ascribe the reason to cognitive problems, while rural residents are more likely to

attribute the reason to tired body. It may be the reason why we correlated measurement errors between items 2 and 5 in urban group but not in rural group. All in all, baseline models are not required to be identical across groups (Byrne et al., 1989; Wang J. et al., 2011). Therefore, we carried out measurement equivalence testing based on these results and found the model of three-factor structure reached partial strict invariances across demographic groups. In addition, Cronbach's alpha coefficients, McDonald's omega coefficients, and Guttman split-half coefficients of the three factors of ASI-3 all indicated that the ASI-3 was a reliable instrument to test the three aspects – physical, social, and cognitive concerns of AS in Chinese women with breast cancer. The test–retest coefficient revealed that the three factors of ASI-3 remained relatively stable over a 2-week period.

Subsequently, the results of Mann–Whitney U tests revealed that the score on cognitive concerns was significantly higher in younger group than in older group. Previous studies have shown that breast cancer patients with younger age showed higher levels of anxiety (Jones et al., 2014; Hassan et al., 2015; Wagner J. F. et al., 2019). AS is a trait-like cognitive vulnerability of anxiety; therefore, it is understandable to get a higher score of the ASI-3 for younger patients. However, the differences were not observed in physical and social concerns. In addition, the ASI-3 scores were significantly higher in the lower education group than in the higher education group, except for cognitive concerns dimension, and the ASI-3 scores were significantly higher in the rural group compared with the urban group. Actually, for patients with breast cancer, long-term residence and education level have an impact on their cognitive range and ability (Li et al., 2016). Relatively speaking, patients in rural and with lower education might have more fragile and sensitive perception, which may be the reason for the higher scores of the ASI-3 in the rural group and in the lower education group. Notably, the effect sizes of these significant differences were small, which indicated that the results might be overvalued because of the large sample size (Fritz et al., 2012; Tomczak and Tomczak, 2014).

The moderate correlation between ASI-3 and the measures of anxiety indicated that the ASI-3 had an adequate empirical validation. In the meantime, the cognitive concerns were positively and moderately related with the depressive mood and somatic symptoms as well as the total score of CESD, which manifested that the cognitive concerns were uniquely relevant to depressive symptoms; it is consistent with the findings of previous research (Olthuis et al., 2014). Moreover, the social concerns performed a greatest correlation with the interpersonal problems, which was accordant with our hypothesis. Moreover, either the standardized factor loadings or the value of AVE and CR were well above the threshold suggested by Hair et al. (2010), indicating that the variances were more explained by each factor and all of the items of each factor were consistent for measuring the same latent construct. Meanwhile, the AVE of any factor was higher than the square of the latent correlations among the factors, which indicated that the three factors could extract more variance than the sharing among factors. To sum up, the ASI-3 had satisfied convergent and discriminant validity in Chinese women diagnosed with breast cancer (Hair et al., 2010).

There were several limitations in this study. First, there are other contents about criterion-related validity of ASI-3 that were not explored; thus, more researches are needed to complement relevant information. Second, we just roughly divided the age and educational level into two groups; the contents about the measurement equivalence across more age and education groups need further supplement. Third, we only collected 24 patients in the test–retest reliability analysis; the number of cases is relatively small, so the cross-time stability of the factor structure needs to be further confirmed by future studies. Finally, the social desirability problems might have an effect on our results, and more researches are needed to explore the degree of the effects.

In summary, we explored the factor structure and psychometric properties of the Chinese version ASI-3 in female patients with breast cancer. In spite of the limitations mentioned above, we have expanded the use of ASI-3 in Chinese women diagnosed with breast cancer. Our findings not only demonstrated that the ASI-3 had adequate reliability, validity, and a stable three-factor structure but also initially approved its strict sociodemographic measurement equivalence.

DATA AVAILABILITY STATEMENT

The datasets generated for this study are available on request to the corresponding author.

REFERENCES

- Burgess, C., Cornelius, V., Love, S., Graham, J., Richards, M., and Ramirez, A. (2005). Depression and anxiety in women with early breast cancer: five year observational cohort study. *BMJ* 330:702. doi: 10.1136/bmj.38343.670868.D3
- Byrne, B. M., Shavelson, R. J., and Muthén, B. (1989). Testing for the equivalence of factor covariance and mean structures: the issue of partial measurement invariance. *Psychol. Bull.* 105, 456–466. doi: 10.1037/0033-2909.105.3.456
- Cai, W., Dong, W., Pan, Y., Wei, C., Zhang, S., Tian, B., et al. (2018). Reliability, validation and norms of the Chinese version of Anxiety Sensitivity Index 3 in a sample of military personnel. *PLoS One* 13:e0201778. doi: 10.1371/journal.pone.0201778
- Cheung, G. W., and Rensvold, R. B. (2002). Evaluating goodness-of-fit indexes for testing measurement invariance. *Struct. Equ. Modeling* 9, 233–255. doi: 10.1207/S15328007SEM0902_5
- Cronbach, L. J. (1951). Coefficient alpha and the internal structure of tests. *Psychometrika* 16, 297–334. doi: 10.1007/bf02310555
- Fan, L., Strasser-Weippl, K., Li, J. J., St Louis, J., Finkelstein, D. M., Yu, K. D., et al. (2014). Breast cancer in China. *Lancet Oncol.* 15, e279–e289. doi: 10.1016/S1470-2045(13)70567-9
- Farris, S. G., Thomas, J. G., Abrantes, A. M., Lipton, R. B., Burr, E. K., Godley, F. A., et al. (2019). Anxiety sensitivity and intentional avoidance of physical activity in women with probable migraine. *Cephalalgia* 39, 1465–1469. doi: 10.1177/0333102419861712
- Fritz, C. O., Morris, P. E., and Richler, J. J. (2012). Effect size estimates: current use, calculations, and interpretation. *J. Exp. Psychol. Gen.* 141, 2–18. doi: 10.1037/a0024338
- Garajčić, E., Bojović-Jović, D., Damjanović, A., Arsić, B., Pantić, I., Turjačanin-Pantelić, D., et al. (2010). Hamilton anxiety scale (HAMA) in infertile women with endometriosis and its correlation with magnesium levels in peritoneal fluid. *Psychiatr. Danub.* 22, 64–67.
- Ghisi, M., Bottesi, G., Altoè, G., Razzetti, E., Melli, G., and Sica, C. (2016). Factor structure and psychometric properties of the anxiety sensitivity index-3 in an Italian community sample. *Front. Psychol.* 7:160. doi: 10.3389/fpsyg.2016.00160

ETHICS STATEMENT

The studies involving human participants were reviewed and approved by the Ethics Committee of the Second Xiangya Hospital, Central South University. The patients/participants provided their written informed consent to participate in this study.

AUTHOR CONTRIBUTIONS

YH, LL, XL, and XZ conceived and designed the study. JZ collected the data and revised the manuscript. YH, HZ, and SL organized and supervised the data collection and inputting. YH drafted the manuscript and organized and supervised the data analysis. JF, JZ, and YY provided the critical comments on various drafts of the manuscript. All authors read and approved the final manuscript.

FUNDING

This work was supported by the National Natural Science Foundation of China (Grant No. 81671341).

- Guttman, L. (1945). A basis for analyzing test-retest reliability. *Psychometrika* 10, 255–282. doi: 10.1007/bf02288892
- Hair, J. F., Black, W. C., Babin, B. J., and Anderson, R. E. (2010). *Multivariate Data Analysis: A Global Perspective*. Upper Saddle River, NJ: Pearson.
- Harris, S. G., and Mossholder, K. W. (1996). The affective implications of perceived congruence with culture dimensions during organizational transformation. *J. Manag.* 22, 527–547. doi: 10.1177/014920639602200401
- Hassan, M. R., Shah, S. A., Ghazi, H. F., Mujar, N. M. M., Samsuri, M. F., and Baharom, N. (2015). Anxiety and depression among breast cancer patients in an urban setting in Malaysia. *Asian Pac. J. Cancer Prev.* 16, 4031–4035. doi: 10.7314/APJCP.2015.16.9.4031
- Jardin, C., Paulus, D. J., Garey, L., Kauffman, B., Bakhshaei, J., Manning, K., et al. (2018). Towards a greater understanding of anxiety sensitivity across groups: the construct validity of the anxiety sensitivity index-3. *Psychiatry Res.* 268, 72–81. doi: 10.1016/j.psychres.2018.07.007
- Jones, S. L., Hadjistavropoulos, H. D., and Gullickson, K. (2014). Understanding health anxiety following breast cancer diagnosis. *Psychol. Health Med.* 19, 525–535. doi: 10.1080/13548506.2013.845300
- Kemper, C. J., Lutz, J., Bähr, T., Rüdell, H., and Hock, M. (2012). Construct validity of the anxiety sensitivity index-3 in clinical samples. *Assessment* 19, 89–100. doi: 10.1177/1073191111429389
- Knapp, A. A., Blumenthal, H., Mischel, E. R., Badour, C. L., and Leen-Feldner, E. W. (2016). Anxiety sensitivity and its factors in relation to generalized anxiety disorder among adolescents. *J. Abnorm. Child Psychol.* 44, 233–244. doi: 10.1007/s10802-015-9991-0
- Lee, J., Yim, M. H., and Kim, J. Y. (2018). Test-retest reliability of the questionnaire in the Sasang constitutional analysis tool (SCAT). *Integr. Med. Res.* 7, 136–140. doi: 10.1016/j.imr.2018.02.001
- Li, D., Cui, Y., Wang, S., Chen, X., Wang, L., and Shao, S. (2016). Study on emotional disorders and health-related quality of life in patients with breast cancer during postoperative chemotherapy. *Chin. J. Cancer Prev. Treat.* 23, 979–982. doi: 10.16073/j.cnki.cjcp.2016.15.002
- Lim, Y. J., and Kim, J. H. (2012). Korean anxiety sensitivity index-3: its factor structure, reliability, and validity in non-clinical samples. *Psychiatry Investig.* 9, 45–53. doi: 10.4306/pi.2012.9.1.45

- MacCallum, R. C., Roznowski, M., and Necowitz, L. B. (1992). Model modifications in covariance structure analysis: the problem of capitalization on chance. *Psychol. Bull.* 111, 490–504. doi: 10.1037/0033-2909.111.3.490
- Mantar, A., Yemez, B., and Alkin, T. (2010). The validity and reliability of the Turkish version of the anxiety sensitivity index-3. *Turk Psikiyatri Derg.* 21, 225–234.
- McDonald, R. (1985). *Factor Analysis and Related Methods*. Hillsdale, NJ: Erlbaum.
- Mohammadkhani, P., Pourshahbaz, A., Kami, M., and Mazidi, M. (2016). Anxiety sensitivity dimensions and generalized anxiety severity: the mediating role of experiential avoidance and repetitive negative thinking. *Iran. J. Psychiatry* 11, 140–146.
- Muthén, L. K., and Muthén, B. O. (2012). *MPlus: Statistical Analysis With Latent Variables—User's Guide*. Los Angeles, CA: Muthén & Muthén.
- Olatunji, B. O., and Wolitzky-Taylor, K. B. (2009). Anxiety sensitivity and the anxiety disorders: a meta-analytic review and synthesis. *Psychol. Bull.* 135, 974–999. doi: 10.1037/a0017428
- Olthuis, J. V., Watt, M. C., and Stewart, S. H. (2014). Anxiety Sensitivity Index (ASI-3) subscales predict unique variance in anxiety and depressive symptoms. *J. Anxiety Disord.* 28, 115–124. doi: 10.1016/j.janxdis.2013.04.009
- Ren, Y., Li, S., Zhou, S., Wang, Y., Li, L., and Zhang, J. (2018). Optimism outweighs neuroticism and anxiety sensitivity to predict insomnia symptoms in women after surgery for breast cancer. *Support. Care Cancer* 27, 2903–2909. doi: 10.1007/s00520-018-4610-6
- Rifkin, L. S., Beard, C., Hsu, K. J., Garner, L., and Björgvinsson, T. (2015). Psychometric properties of the anxiety sensitivity index-3 in an acute and heterogeneous treatment sample. *J. Anxiety Disord.* 36, 99–102. doi: 10.1016/j.janxdis.2015.09.010
- Schmitt, N., and Kuljanin, G. (2008). Measurement invariance: Review of practice and implications. *Hum. Resour. Manag. Rev.* 18, 210–222. doi: 10.1016/j.hrmr.2008.03.003
- Sun, J. (2005). Assessing goodness of fit in confirmatory factor analysis. *Meas. Eval. Couns. Dev.* 37, 240–256. doi: 10.1080/07481756.2005.11909764
- Taylor, S., Zvolensky, M. J., Cox, B. J., Deacon, B., Heimberg, R. G., Ledley, D. R., et al. (2007). Robust dimensions of anxiety sensitivity: development and initial validation of the anxiety sensitivity index-3. *Psychol. Assess.* 19, 176–188. doi: 10.1037/1040-3590.19.2.176
- Tomczak, M., and Tomczak, E. W. A. (2014). The need to report effect size estimates revisited. an overview of some recommended measures of effect size. *Trends Sport Sci.* 1, 19–25.
- Wagner, A., Shiban, Y., Lange, N., Joerger, A. K., Hoffmann, U., and Meyer, B. (2019). The relevant psychological burden of having a benign brain tumor: a prospective study of patients undergoing surgical treatment of cranial meningiomas. *J. Neurosurg.* 11, 1–8. doi: 10.3171/2018.8.JNS181343
- Wagner, J. F., Lüdders, D., Hoellen, F., Rody, A., and Banz-Jansen, C. (2019). Treatment anxiety in breast cancer patients. *Arch. Gynecol. Obstet.* 299, 1365–1371.
- Wang, C., Chu, Y., Zhang, Y., Zhang, N., Zhang, J., and Yang, H. (2011). Study on factor structure of hamilton rating scale for anxiety. *J. Clin. Psychiatry* 21, 299–301.
- Wang, J., Wang, X., and Jiang, B. (2011). *Structural Equation Model: Methods and Applications (in Chinese)*. Beijing: Higher education press.
- Zhang, J., Sun, W., Kong, Y., and Wang, C. (2012). Reliability and validity of the center for epidemiological studies depression scale in 2 special adult samples from rural China. *Compr. Psychiatry* 53, 1243–1251. doi: 10.1016/j.comppsy.2012.03.015
- Zimmerman, M., Martin, J., Clark, H., McGonigal, P., Harris, L., and Holst, C. G. (2017). Measuring anxiety in depressed patients: a comparison of the hamilton anxiety rating scale and the DSM-5 anxious distress specifier interview. *J. Psychiatr. Res.* 93, 59–63. doi: 10.1016/j.jpsychires.2017.05.014

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Han, Zhu, Li, Zhou, Li, Zhang, Fan, Yang, Luo and Zhu. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.