



OPEN ACCESS

EDITED BY

Ruoyu Wang,
University of Edinburgh,
United Kingdom

REVIEWED BY

Vera Ling Hui Phung,
National Institute for Environmental
Studies (NIES), Japan
Shazia Rehman,
Pak-Austria Fachhochschule Institute
of Applied Sciences and
Technology, Pakistan

*CORRESPONDENCE

Nguyen Thi Trang Nhung
ntrn2@huph.edu.vn;
nhungnguyenbio@gmail.com

SPECIALTY SECTION

This article was submitted to
Environmental Health and Exposome,
a section of the journal
Frontiers in Public Health

RECEIVED 28 September 2022

ACCEPTED 03 November 2022

PUBLISHED 18 November 2022

CITATION

Nhung NTT, Duc VT, Ngoc VD,
Dien TM, Hoang LT, Ha TTT, Khue PM,
Truong NX, Thanh NTN, Jegasothy E,
Marks GB and Morgan G (2022)
Mortality benefits of reduction fine
particulate matter in Vietnam, 2019.
Front. Public Health 10:1056370.
doi: 10.3389/fpubh.2022.1056370

COPYRIGHT

© 2022 Nhung, Duc, Ngoc, Dien,
Hoang, Ha, Khue, Truong, Thanh,
Jegasothy, Marks and Morgan. This is
an open-access article distributed
under the terms of the [Creative
Commons Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/).
The use, distribution or reproduction
in other forums is permitted, provided
the original author(s) and the copyright
owner(s) are credited and that the
original publication in this journal is
cited, in accordance with accepted
academic practice. No use, distribution
or reproduction is permitted which
does not comply with these terms.

Mortality benefits of reduction fine particulate matter in Vietnam, 2019

Nguyen Thi Trang Nhung^{1,2*}, Vu Tri Duc², Vo Duc Ngoc²,
Tran Minh Dien², Le Tu Hoang¹, Tran Thi Thuy Ha³,
Pham Minh Khue³, Ngo Xuan Truong⁴,
Nguyen Thi Nhat Thanh⁴, Edward Jegasothy⁵, Guy B. Marks^{5,6}
and Geoffrey Morgan^{5,6}

¹Department of Biostatistics, Faculty of Fundamental Sciences, Hanoi University of Public Health, Hanoi, Vietnam, ²Vietnam National Children's Hospital, The Training and Research Institute for Child Health, Hanoi, Vietnam, ³Faculty of Public Health, Haiphong University of Medicine and Pharmacy, Haiphong, Vietnam, ⁴University of Engineering and Technology, Vietnam National University, Hanoi, Vietnam, ⁵Faculty of Medicine and Health, Sydney School of Public Health, University Centre for Rural Health, University of Sydney, Sydney, NSW, Australia, ⁶Centre for Air Pollution, Energy and Health Research, University of New South Wales, Sydney, NSW, Australia

Introduction and objectives: Studies assessing the health benefits of air pollution reduction in Vietnam are scarce. This study quantified the annual mortality burden due to PM_{2.5} pollution in Vietnam above the World Health Organization recommendation for community health (AQG: 5 µg/m³) and the proposed National Technical Regulation on Ambient Air Quality (*proposed QCVN*: 15 µg/m³).

Methodology: This study applied a health impact assessment methodology with the hazard risk function for non-communicable diseases (NCDs) and lower respiratory infections (LRIs) in the Global Exposure Mortality Model (GEMM) to calculate attributable deaths, Years of Life lost, and Loss of Life expectancy at birth due to air pollution in the Vietnamese population above 25 years of age in 11 provinces. We obtained annual average PM_{2.5} concentrations for Vietnam in 2019 at a 3x3 km grid modeled using Mixed Linear regression and multi-data sources. Population and baseline mortality data were obtained from administrative data system in Vietnam. We reported the findings at both the provincial and smaller district levels.

Results: Annual PM_{2.5} concentrations in all studied provinces exceeded both the AQG and the *proposed QCVN*. The maximum annual number of attributable deaths in the studied provinces if they had complied with WHO air quality guidelines was in Ha Noi City, with 5,090 (95%CI: 4,253–5,888) attributable deaths. At the district level, the highest annual rate of attributable deaths if the WHO recommendation for community health had been met was 104.6 (95%CI: 87.0–121.5) attributable deaths per 100,000 population in Ly Nhan (Ha Nam province).

Conclusion: A much larger number of premature deaths in Vietnam could potentially be avoided by lowering the recommended air quality standard. These results highlight the need for effective clean air action plans by local authorities to reduce air pollution and improve community health.

KEYWORDS

air pollution, PM_{2.5}, Vietnam, mortality burden, health benefits

Introduction

Fine particulate matter (denoted as $PM_{2.5}$) concentrations have declined globally for the last decade. For example, the annual mean $PM_{2.5}$ in the United States was reduced by approximately 41% from 2000 to 2020 (1). In Europe, the decrease in $PM_{2.5}$ concentrations across countries was 29% from 2005 to 2019 (2). In China, a decline in annual $PM_{2.5}$ concentration was observed, ranging from 22.2 to 56.6% during 2015–2020 (3). These improvements in outdoor air quality throughout the world are the result of affordable clean air action. For example, the “Air Pollution Prevention and Control Action Plan,” first implemented in China in 2013, contributed to 15% reduction in $PM_{2.5}$ in the Pearl River Delta (4). In other words, evidence quantifying the health benefits of reducing exposure to $PM_{2.5}$ is urgently needed to support further clean air policy development and implementation, particularly in high-exposure countries such as Vietnam.

Though air quality in Vietnam has improved in recent years, it is still poor by international comparisons. Annual mean $PM_{2.5}$ concentrations in Vietnamese provinces ranged from 9–41 $\mu\text{g}/\text{m}^3$ in 2019 to 8–36 $\mu\text{g}/\text{m}^3$ in 2020 (5). However, the annual $PM_{2.5}$ concentration in 10 out of 63 provinces in Vietnam in 2020 exceeded the National Technical Regulation on Ambient Air Quality- QCVN 05:2013 (set at 25 $\mu\text{g}/\text{m}^3$), and no provinces reached the new World Health Organization’s air quality guidelines (AQG, set at 5 $\mu\text{g}/\text{m}^3$) (5, 6). Provinces in the Red River Delta such as Hung Yen (32.7 $\mu\text{g}/\text{m}^3$) and Bac Ninh (33.0 $\mu\text{g}/\text{m}^3$) experienced the poorest air quality, rather than the major cities such as Ha Noi and Ho Chi Minh City (5).

Exposure to the high concentration of $PM_{2.5}$ can cause many diseases including cardiovascular, respiratory diseases, diabetes, lung cancer (7), and contribute to premature deaths (8–10). The evidence also indicates no threshold of air pollutant concentration for community health. The effect of $PM_{2.5}$ remained even where the exposure level is low such as in the European Union region and Australia (7, 11, 12). Therefore, in September 2021, the World Health Organization (WHO) updated its 2015 recommendation in which it set the annual mean concentration of $PM_{2.5}$ at 5 $\mu\text{g}/\text{m}^3$ (6). Though this guideline is not a regulation, it can drive the air quality policy worldwide.

In recent years, the Vietnamese government has attempted to address poor air quality through a series of actions. In 2016, the Vietnamese Prime Minister issued the National Action Plan on air quality management by 2020 including a vision for 2025 (Decision No 985a/QĐ-TTg) which targeted improved air quality management by controlling emissions and increasing ambient air quality monitoring (13). Directive No 03/CT-TTg on enhancing air quality management was signed by the Prime Minister on 18 January 2021 (14). This latter document also called for the assessment of the adverse health impacts of air

pollution on community health (14). Decision No 1973/QĐ-TTg states that Vietnam will take actions to improve air quality management *via* controlling emissions and monitoring ambient air pollutants and also conduct research to provide information *via* early warning systems that predict air quality in each province (15). These documents emphasize the importance of assessing the health impacts of air pollution to evaluate the implemented policy and associated interventions in local areas (13–15). The Official Dispatch No 3051/BTNMT-TCMT on the technical guidelines of building air management plan at the provincial level recommends the use of AirQ+ software (a software developed by the World Health Organization–WHO) to assess the health impacts of air pollution (16). Ha Noi, in particular, issued Directive No 15/CT-UBND on entirely replacing and eliminating the use of the beehive coal fuel to reduce the adverse impacts of air pollution in the city (17). Also, in 2021, the Ministry of Natural Resources and Environment issued a draft of the National Technical Regulation on Ambient Air Quality (called *proposed QCVN* in this study) (18) in order to establish a policy to improve ambient air quality in Vietnam. This study quantifies the mortality impact of air pollution in Vietnam in response to the call for such policies by the Vietnamese government.

Evidence of the health impacts of ambient air pollution in Vietnam is scarce, and mainly in Ha Noi and Ho Chi Minh City. Two studies showed an association between air pollutants, including $PM_{2.5}$, and hospital admissions amongst Hanoi children (19, 20). Another study demonstrated the relationship between ambient air pollutants and hospital admissions for cardiovascular and respiratory diseases in adults in Hanoi, Quang Ninh, and Phu Tho (21) and a study in Ho Chi Minh City showed the association between hospitalization due to acute lower respiratory infection and $PM_{2.5}$ (22). The first Vietnamese health impact assessment study of exposure to $PM_{2.5}$ was conducted in Ha Noi in 2017 (23). The findings from this study showed the life expectancy at birth of the Hanoi population due to exposure to $PM_{2.5}$ had decreased by 1.8 years in 2017 (23). Research in Ho Chi Minh City also found that 1,136 premature deaths were attributed to $PM_{2.5}$ in 2017 (24). Such findings highlight the large burden of diseases due to $PM_{2.5}$ exposure in Vietnam.

This study aimed to quantify the health burden due to ambient $PM_{2.5}$ concentrations above the World Health Organization guideline and the proposed National Technical Regulation on Ambient Air Quality in several provinces of Vietnam, including Bac Ninh, Hung Yen, Hai Duong, Ha Noi, Thai Binh, Hai Phong, Ninh Binh, Ha Nam, Ho Chi Minh, Quang Ninh, and Dien Bien. The burden of mortality is expressed as attributable deaths (AD) (number and rate per 100,000 population), Years of life lost (YLL) (number and rate per 100,000 population), and Loss of Life Expectancy (LLE) (in years) for these provinces in Vietnam. Moreover, we also

calculated the attributable deaths (rate per 100,000 population) using the AirQ+ software to compare our results with the estimate according to the technical guidelines of the Official Dispatch No 3051/BTNMT-TCMT of the Ministry of Natural Resources and Environmental (16). This study not only provides scientific data on the health impact of air pollution in Vietnam but also responds to the Vietnamese government's call for evidence to establish air quality control that will improve health.

Materials and methods

Study areas

We conducted a health impact assessment of ambient PM_{2.5} concentrations in 11 Vietnamese provinces, including 10 provinces in the northern region (Ha Noi, Bac Ninh, Hung Yen, Ha Nam, Hai Duong, Thai Binh, Hai Phong, Ninh Binh, Quang Ninh, Dien Bien) and Ho Chi Minh City in the south of Vietnam. The total 2019 population of Vietnam was 96,208,984 people and our study region included an exposed population of 18,342,836 (19%) inhabitants (25) covering the most polluted provinces (5) and also the two megacities of Hanoi (the capital) and Ho Chi Minh City.

PM_{2.5} exposure

We assessed exposure using a previously published study (23). Daily PM_{2.5} maps for Vietnam at 3x3 km resolution were estimated by Mixed Effect Models developed on a 9-year dataset (2012-2020) of PM_{2.5} measured at ground stations, combined with satellite Aerosol Optical Depth (AOD), meteorological conditions (i.e., humidity, planetary boundary layer height) and land use (i.e., normalized difference vegetation index (NDVI), road density) (26). The detailed method, data source, and model validation for estimating daily PM_{2.5} maps is described in our previous study (23). The daily mean PM_{2.5} maps were validated by comparing them with ground-based PM_{2.5} measurements, with a Pearson coefficient (r) of 0.87 and a Root Mean Square Error (RMSE) of 11.76 $\mu\text{g}/\text{m}^3$. The daily maps were then aggregated into monthly maps and annual maps. The annual average PM_{2.5} map in 2019 had high agreement with ground PM_{2.5} measurements ($r = 0.84$, $\text{RMSE} = 5.07 \mu\text{g}/\text{m}^3$). The annual PM_{2.5} mean concentration data in 2019 was calculated for each district of the selected provinces for the health impact assessment process.

Mortality data and exposure population

Death cases including age, address (district and province), gender, and specific causes of death were obtained from the

Vietnam A6 registration. A6 document is a death registration at the commune health, the lowest level of the health sector in Vietnam—and data on death cases of permanent inhabitants was collected by community health staff. This document covers approximately 89% of the total deaths in the commune (27). The report also confirmed that data on death due to injury is robust (27) and so we were able to exclude injury-related deaths from our analysis.

The number of inhabitants in each district by age group and gender were extracted from the Population and Household census in 2019 by the General Statistics Office of Vietnam (hereafter called GSO) (25, 28). This census obtained information on the population including fertility, gender, migration, urbanization, and population aging in Vietnam.

Health impact assessment estimation

We applied the Global Exposure Mortality Model (GEMM) risk function to estimate the attributable deaths due to ambient PM_{2.5} pollution (29) for each province. Burnett et al. developed GEMM by pooling hazard ratios from 41 cohorts (29), including the Chinese cohort where the ambient PM_{2.5} concentration and emission sources are more similar to Vietnam compared to European and North American studies.

In each province, we estimated attributable deaths (AD), years of life lost (YLL) and loss of life expectancy (LLE) due to air pollution for populations above 25 years of age in each province separately. The AD for each 5-year age subgroup in a district is estimated from applying the hazard ratio to specific age groups and PM_{2.5} concentration using the GEMM risk function (29), the death rate for each age group in a district, and the population at that age group for the corresponding district. We applied the estimate for non-communicable diseases (NCDs) and lower respiratory infections (LRIs) in GEMM (denoted as GEMM NCD+LRI). We set the counterfactual concentration as defined by 1) the WHO air quality guidelines (AQG: 5 $\mu\text{g}/\text{m}^3$) and 2) the proposed QCVN (15 $\mu\text{g}/\text{m}^3$). YLL is calculated by multiplying the attributable deaths in each age group by the life expectancy in the corresponding age group. This life expectancy is calculated by applying the life-table method with the baseline death rate per 5-year age group in 2019. Thirdly, the LLE in days was calculated by subtracting the life expectancy without the contribution of PM_{2.5} to the PM_{2.5} attributed life expectancy. For each province and district, LLE was expressed in years by dividing the LLE in days by 365.25. The rates of deaths and YLL for each district per 100,000 inhabitants were calculated by dividing the number of the corresponding indicators by the population size. We present the findings at the provincial and district levels.

The other analysis was conducted in R version 4.0.3 (R Core Team Vienna, R Foundation for Statistical Computing, Vienna,

TABLE 1 Annual means of the PM_{2.5} concentration (in $\mu\text{g}/\text{m}^3$), population and death rates per 1,000 population above age 25 years in 11 study region provinces, Vietnam, 2019.

Province's name	PM _{2.5} annual mean concentration ($\mu\text{g}/\text{m}^3$)	Population above age 25	Death rates above aged 25+ years (per 1,000 population)
Ha Noi	40.8	4,889,379	4.6
Bac Ninh	37.8	812,711	5.4
Hung Yen	35.2	789,408	4.1
Ha Nam	31.5	543,823	7.9
Hai Duong	31.2	1,222,568	4.3
Thai Binh	27.1	1,237,051	8.3
Hai Phong	25.8	1,311,556	7.4
Ninh Binh	23.5	622,278	7.8
Ho Chi Minh	20.9	5,795,749	4.8
Quang Ninh	18.9	825,247	5.8
Dien Bien	15.8	293,066	6.3

Austria) with the iomlifetR package (30), apart from the AirQ+ comparison calculations.

Comparison analysis

For comparison, we also estimated the attributable deaths (rate per 100,000 population) by province using the AirQ+ software and selecting the long-term impact assessment option for total natural causes. By default this option utilized the log-linear model with the relative risks retrieved from the literature and meta-analysis of Hoek et al. (31). Since this model only applies to adults 30 years of age or older, we only calculated attributable deaths for this population *via* AirQ+.

Results

Population and mortality data for people above 25 years old in 11 provinces in 2019 are provided in Table 1. Ha Noi and Ho Chi Minh were the two most populous areas, with the population over 25 years old reaching approximately 5 to 6 million inhabitants. Crude death rates by province ranged from 4.1 deaths per 1,000 population in Hung Yen to 8.3 deaths per 1,000 population in Thai Binh.

The annual PM_{2.5} concentration ranged from 15.8 $\mu\text{g}/\text{m}^3$ in Dien Bien to 40.8 $\mu\text{g}/\text{m}^3$ in Ha Noi (Table 1, Figure 1). All annual mean concentrations in each of the 11 provinces exceeded the WHO air quality guidelines and the proposed QCVN. Notably, Bac Ninh, Hung Yen and Ha Nam, adjacent to Hanoi, experienced highly polluted levels. The annual mean of PM_{2.5} concentration in Ha Noi was 2.72 times higher than the proposed QCVN (Table 1, Figure 1).

At the district level, the highest PM_{2.5} concentration was observed in all districts of Bac Ninh, Hai Duong, Hung Yen, Ha Nam province, and the central districts of the capital Ha Noi (Figure 1).


Compliance with the WHO air pollution guideline could potentially avoid 5,090 (95%CI: 4,253–5,888) annual premature deaths in Hanoi from exposure to PM_{2.5} (Table 2) and 4,076 (95% CI: 3,377–4,754) premature deaths in Ho Chi Minh City. The highest PM_{2.5} attributable death rates were Ha Nam, Thai Binh, and Hai Phong, with 95.1 (95% CI: 79.1–110.4), 93.4 (95% CI: 77.6–108.7), and 81.9 (95% CI: 68.0–95.3) deaths per 100,000 population, respectively.

Our study also observed great variability in attributable deaths at district levels (Supplementary Table S1). The attributable death rates range from 22.3 deaths per 100,000 population in Muong Nhe district (Dien Bien) to 104.6 deaths in Ly Nhan (Ha Nam province). In Ha Noi, the capital of Vietnam, districts with the greatest attributable death rate are located in the central districts (Hoan Kiem, Ba Dinh, Hai Ba Trung, Dong Da, and Tay Ho) of the province.

Figure 2 illustrates the number and the rate (per 100,000 population) of attribute deaths due to PM_{2.5} concentrations above the proposed QCVN. The rate of attributable deaths due to annual ambient PM_{2.5} in the selected provinces above the proposed QCVN was 52.3 deaths per 100,000 population in Ha Noi and 66.6 per 100,000 population in Thai Binh.

Discussion

In this study, we estimated 37.0 to 95.1 deaths per 100,000 population are attributable to PM_{2.5} pollution in 2019 above the WHO air quality guidelines for annual average PM_{2.5} in the selected provinces. These numbers were generally higher



than the PM_{2.5} attributable mortality rate in Vietnam (44.8 per 100,000 population) according to the calculation of the Global Burden of Disease (GBD) study (32). The discrepancy will be influenced by the difference in resolution of the PM_{2.5} concentration maps we used of 3 x 3 km that is smaller (higher

spatial resolution) compared to the air pollution maps used in the GBD. Nonetheless, both these analyses present a high rate of attributable mortality due to exposure to PM_{2.5} in Vietnam and highlight the urgency to effectively implement the National Action Plan for air quality management in Vietnam.


TABLE 2 Avoidable deaths, years of life lost and loss of life expectancy (expressed in number and rate per 100,000 population) and confidence interval (95% CI) when compliance with The World Health Organization’s Air Quality Guidelines in Vietnam study region provinces, 2019.

Province’s name	Avoidable deaths		Years of life lost		Loss of life expectancy (years) (95%CI)
	Number (95%CI)	Rate per 100,000 population (95%CI)	Number (95%CI)	Rate per 100,000 population (95%CI)	
Ha Noi	5,090 (4,253–5,888)	63.2 (52.8–73.1)	152,828.6 (124,123.8–181,963.7)	1,897.6 (1,541.2–2,259.4)	4.9 (3.9–5.9)
Bac Ninh	946 (789–1,096)	69.1 (57.7–80.0)	23,779.8 (19,353.7–28,252.8)	1,737.2 (1,413.9–2,064.0)	3.8 (3.1–4.6)
Hung Yen	658 (548–763)	52.5 (43.7–60.9)	23,940.0 (19,392.6–28,573.7)	1,911.0 (1,548.0–2,280.9)	5.7 (4.6–6.8)
Ha Nam	811 (675–942)	95.1 (79.1–110.4)	19,520.0 (15,881.1–23,196.9)	2,288.9 (1,862.2–2,720.1)	3.4 (2.8–4.1)
Hai Duong	968 (805–1,124)	51.1 (42.5–59.4)	32,989.8 (26,753.7–39,327.5)	1,743.4 (1,413.9–2,078.3)	4.8 (3.9–5.8)
Thai Binh	1,738 (1,443–2,022)	93.4 (77.6–108.7)	39,772.4 (32,368.7–47,244.1)	2,137.8 (1,739.8–2,539.4)	2.9 (2.4–3.5)
Hai Phong	1,661 (1,380–1,933)	81.9 (68.0–95.3)	38,360.9 (31,294.0–45,461.7)	1,891.7 (1,543.2–2,241.8)	2.8 (2.2–3.3)
Ninh Binh	758 (628–883)	77.1 (64.0–89.8)	18,422.6 (15,029.6–21,830.2)	1,875.1 (1,529.8–2,221.9)	2.7 (2.2–3.2)
Ho Chi Minh	4,076 (3,377–4,754)	45.3 (37.5–52.9)	88,710.3 (72,525.0–104,897.0)	986.4 (806.5–1,166.4)	1.9 (1.6–2.3)
Quang Ninh	687 (569–801)	52.0 (43.1–60.7)	18,436.3 (15,042.3–21,844.0)	1,396.3 (1,139.3–1,654.4)	2.5 (2.1–3.0)
Dien Bien	222 (183–259)	37.0 (30.6–43.3)	6,437.5 (5,258.7–7,617.9)	1,075.0 (878.1–1,272.1)	2.2 (1.8–2.6)

To address this issue, the Prime Minister of Vietnam stated 6 objectives in Decision No 1973/QĐ-TTg in 2021, including: 1) Complete the structure, policies, and constitutions regarding air quality management; 2) Prevent and reduce emission; 3) Complete the financial structure and diversify the investment for air quality management; 4) International cooperation and scientific research regarding technology and air quality management; 5) Monitor and evaluate the compliance of air quality management; 6) Communicate, educate, and improve the capacity and acknowledgment regarding air quality management (15). Implementing these objectives is an interdisciplinary task and requires the involvement of the entire Vietnamese administrative structures (the details of the assignments were described in Decision No 1973/QĐ-TTg and the Law 72/2020/QH14 on Environmental Protection) (15, 33). Our study findings demonstrate the health benefits which Vietnam can obtain from effectively implementing air pollution

control measures. Parallel with the above strategies, the Vietnam government has planned to reduce QCVN from 25 µg/m³ to 15 µg/m³. However, the proposed QCVN and a time frame for achieving this standard have not yet been officially issued.

The Official Dispatch No. 3051/BTNMT-TCMT suggests using the AirQ+ software developed by the WHO (16). In this study, the death rates for provinces estimated by AirQ+ were higher than the findings using GEMM, though the exposure population is marginally smaller (Table 3). The likely reason for these differences is that the population attributable fraction estimated by GEMM compared to that estimated using the Hoek et al. (31) risk function at lower PM_{2.5} concentrations, but the opposite situation occurs when the PM_{2.5} concentration is high (34). Therefore, using the log-linear model in AiQ+ for Vietnam might overestimate the actual burden of mortality due to the high ambient PM_{2.5} concentrations.



The attributable mortality rates in the central northern provinces (such as Thai Binh or Ha Nam and Hai Duong) were high and this is likely due to the industrial areas located in these provinces (35). A greater number of thermal power stations, cement manufacturer factories, and factories with older

technology are located in the northern region, especially in Hai Duong and Thai Binh (35). Moreover, Bac Ninh includes many villages with traditional industries that lack pollution control systems (35). Our study indicates that the substantial health burden due to air pollution in the Bac Ninh province could be

reduced by implementing effective interventions to manage the air quality in these polluted areas.

The National Action Plan on air quality management for 2021–2025 in Vietnam is in its initial stages and authorities should be aware of the range of air pollutants that need to be reduced to decrease the health burden (15). In this study, we primarily focus on the mortality burden due to a reduction in PM_{2.5}. A study in China found an inverse association between PM_{2.5} and ground-level Ozone (O₃) concentration (36). According to the authors reducing the Nitrogen Oxide (NO_x) and Volatile Organic Compounds (VOCs) components in PM_{2.5} could potentially increase the photochemical formation of O₃, thus resulting in the higher concentration of O₃. Ground-level ozone has been found to have an impact on hospital admissions in Vietnam (37). Therefore, future interventions should consider the proportion of air pollutants in the mixture and their interrelation before implementing interventions to reduce any specific pollutant.

In this study, we applied the GEMM risk function to estimate the attributable deaths due to ambient PM_{2.5}. This risk function has been used in numerous studies to assess not only the health impact of air pollution, but also the economic health benefits of an intervention. For example, Haikun Wang (2020) used GEMM to estimate the health benefits of control programs relating to on-road transportation in China during the period of 1998 to 2015 (38). Another study used GEMM to assess the health and economic impact of China’s air pollution shift from 2013 to 2018, as well as to suggest implications for air pollution control policies during this time period (36). This approach is more suitable for countries and regions with severe pollution. The estimation is similar to the GBD approach. In addition, the Environmental Benefits Mapping and Analysis Program—Community Edition (BenMAP-CE) by the United States Environmental Protection Agency integrated this approach into the software; therefore so it can be easily applied (39). We recommend that Vietnam use this approach to estimate both the health and economic benefits of interventions to reduce air pollution.

Our study has several limitations. While we were unable to collect data from all the areas in Vietnam we obtained data for the most polluted provinces, including both the north and south of Vietnam. The mortality data used in our analysis is estimated to include around 89% of all deaths based on a study in Quang Ninh and Thai Nguyen provinces (27). Although we performed the quality assurance process on the mortality data (details of the procedure were described elsewhere (27)) the attributable mortality due to air pollution we estimated is likely to be an underestimate due to the < 100% covered of the mortality data we used. The air quality data we used was modeled using data from a limited number of fixed site air pollution monitoring stations in Vietnam and this could introduce error in our estimated PM_{2.5} exposures. Our study applied a hazard ratio obtained from the analysis of 41 cohorts, including China (29).

TABLE 3 Attributable deaths (rate per 100,000) between the calculation by Log–linear model in AirQ+ and the calculation by GEMM for each province.

Province’s name	Attributable deaths (Rate per 100,000 population) by Log–linear model in AirQ + (95%CI)	Attributable deaths (Rate per 100,000 population) by GEMM(95%CI)
Ha Noi	103.7	63.2
Bac Ninh	118.8	69.1
Hung Yen	79.3	52.5
Ha Nam	132.8	95.1
Hai Duong	70.5	51.1
Thai Binh	115.2	93.4
Hai Phong	103.7	81.9
Ninh Binh	95.4	77.1
Ho Chi Minh	53	45.3
Quang Ninh	59.7	52.5
Dien Bien	46.6	37.0

While this hazard ratio is estimated from a number of large cohort studies there may be some differences when generalizing to different populations.

Conclusion

Our study found that there is a substantial burden of mortality due to air pollution in Vietnam above the air quality guidelines by the WHO and the proposed Vietnam QCVN. These findings highlight the health benefits of reducing air pollution in Vietnam including actions by local provincial and district authorities to implement policies to improve air quality.

Data availability statement

The original contributions presented in the study are included in the article/[Supplementary material](#), further inquiries can be directed to the corresponding author.

Ethics statement

The studies involving human participants were reviewed and approved by the Ethics Committee of the Hanoi University of Public Health (Ref. No. 020-265/DD-YTCC). Written informed consent for participation was not required for this study in accordance with the National Legislation and the Institutional Requirements.

Author contributions

NTTN, GBM, and GM conceived, designed the study, and acquired data. NTNT and NXT conducted statistical analysis for PM_{2.5} concentration. NTTN, VTD, VDN, and LTH conducted statistical analyses for health impact assessment. NTTN and VTD drafted the manuscript. All authors provide statistical support, interpretation, professional support, and made several critical revisions to the manuscript. All authors read and approved the final manuscript.

Funding

This study received funding from Vietnam National Foundation for Science and Technology Development (NAFOSTED) under grant number 105.08–2019.331. The funder was not involved in the study design, collection, analysis, interpretation of data, the writing of this article, or the decision to submit it for publication.

Acknowledgments

This study also acknowledge the contributions of staffs from Centers of Diseases Control of Bac Ninh, Hung Yen, Hai Duong, Ha Noi, Thai Binh, Hai Phong, Ninh Binh, Ha Nam, Ho Chi Minh, Quang Ninh, and Dien Bien for providing mortality data

References

- USEPA. *Our Nation's Air: Status and Trends Through 2020*. Available online at: <https://gispub.epa.gov/air/trendsreport/2021/> (accessed February 24, 2022).
- European Environment Agency. *Sources and Emissions of Air Pollutants in Europe*. (2021). Available online at: <https://www.eea.europa.eu/publications/air-quality-in-europe-2021/sources-and-emissions-of-air> (accessed May 6, 2022).
- Li Y, Zhang Z, Xing Y. Long-term change analysis of PM_{2.5} and Ozone pollution in China's most polluted region during 2015–2020. *Atmosphere*. (2022) 13:104. doi: 10.3390/atmos13010104
- The State Council of the People's Republic of China. *Air Pollution Prevention and Control Action Plan*. (2013). Available online at: http://www.gov.cn/zwqk/2013-09/12/content_2486773.htm (accessed July 7, 2022).
- VNU-UET, Live&Learn & USAID. *State of PM_{2.5} in Vietnam during 2019-2020 based on multi-source data. Prepared by University of Engineering and Technology under Vietnam National University in Hanoi (VNU-UET), Center of Live and Learn for Environment and Community (Live&Learn) and the United States Agency for International Development (USAID)*. (2021). Available online at: <https://khisachtroixanh.com/tai-lieu/nghien-cuu-hien-trang-bui-pm2-5-o-viet-nam-giai-doan-2019-2020-su-dung-du-lieu-da-nguon/> (accessed February 7, 2022).
- World Health Organization. *WHO Global Air Quality Guidelines: Particulate Matter (PM_{2.5} and PM₁₀), Ozone, Nitrogen Dioxide, Sulfur Dioxide and Carbon Monoxide*. Geneva: World Health Organization (2021).
- Raaschou-Nielsen O, Andersen ZJ, Beelen R, Samoli E, Stafoggia M, Weinmayr G, et al. Air pollution and lung cancer incidence in 17 European cohorts: prospective analyses from the European Study of Cohorts for Air Pollution

and giving support during data cleaning, as well as the staff from the Northern Center Monitoring Station for providing the air quality data.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Supplementary material

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fpubh.2022.1056370/full#supplementary-material>

- Effects (ESCAPE). *Lancet Oncol*. (2013) 14:813–22. doi: 10.1016/S1470-2045(13)0279-1
- Bourbeau J, Doiron D, Biswas S, Smith BM, Benedetti A, Brook JR, et al. Ambient air pollution and dysanapsis: associations with lung function and chronic obstructive pulmonary disease in the canadian cohort obstructive lung disease study. *Am J Respir Crit Care Med*. (2022) 206:44–55. doi: 10.1164/rccm.202106-1439OC
 - Wolf K, Hoffmann B, Andersen ZJ, Atkinson RW, Bauwelinck M, Bellander T, et al. Long-term exposure to low-level ambient air pollution and incidence of stroke and coronary heart disease: a pooled analysis of six European cohorts within the ELAPSE project. *Lancet Planet Health*. (2021) 5:e620–32. doi: 10.1016/S2542-5196(21)00195-9
 - Li Y, Xu L, Shan Z, Teng W, Han C. Association between air pollution and type 2 diabetes: an updated review of the literature. *Ther Adv Endocrinol Metab*. (2019) 10:2042018819897046. doi: 10.1177/2042018819897046
 - Andersen ZJ, Gehring U, De Matteis S, Melen E, Vicedo-Cabrera AM, Katsouyanni K, et al. Clean air for healthy lungs—an urgent call to action: European Respiratory Society position on the launch of the WHO 2021 Air Quality Guidelines. *Eur Respir J*. (2021) 58:2021. doi: 10.1183/13993003.02447-2021
 - Hanigan IC, Broome RA, Chaston TB, Cope M, Dennekamp M, Heyworth JS, et al. Avoidable mortality attributable to anthropogenic fine particulate matter (PM_{2.5}) in Australia. *Int J Environ Res Public Health*. (2021) 18:254. doi: 10.3390/ijerph18010254
 - The Prime Minister. *Decision No 985a/QĐ-TTg on Approving the National Action Plan on Air Quality Management by 2020 With the Vision to 2025*. (2016). Available online at: <https://chinhphu.vn/default.aspx?pageid=27160&docid=184944> (accessed April 5, 2022).

14. The Prime Minister. *Directive No 03/CT-TTg on Enhancing the Air Quality Management*. (2021). Available online at: <https://vanban.chinhphu.vn/default.aspx?pageid=27160&docid=188138> (accessed April 5, 2022).
15. The Prime Minister. *Decision No 1973/QĐ-TTg on Approving the National Action Plan on Air Quality Management During 2021-2025*. (2021). Available online at: <https://vanban.chinhphu.vn/default.aspx?pageid=27160&docid=204535> (accessed April 5, 2022).
16. Ministry of Natural Resources and Environment. *Official Dispatch No 3051/BTNMT-TCMT*. (2021). Available online at: <https://www.monre.gov.vn/VanBan/Pages/ChiTietVanBanChiDao.aspx?piD=2615> (accessed May 31, 2022).
17. Ha Noi City People's Committee. *Directive No 15/CT-UBND: Replace and Eliminate the Use of the Beehive Coal Fuel in Daily Life, and Business Services to Reduce the Adverse Impacts on the Environment in the City*. (2019). Available online at: <https://www.thudo.gov.vn/newsdetail.aspx?NewsID=6e2e099f-0d1f-274b-aa4a-4f4ac2f3069b&CateID=366e5f3c-7d01-9545-9034-49d13bb569c2> (accessed April 10, 2022).
18. Ministry of Natural Resources and Environment. *National Technical Regulation on Ambient Air Quality*. (2021). Available online at: <https://monre.gov.vn/VanBan/Pages/ChiTietVanBanDuThao.aspx?piD=236> (accessed March 31, 2022).
19. Nhung NTT, Schindler C, Dien TM, Probst-Hensch N, Perez L, Künzli N. Acute effects of ambient air pollution on lower respiratory infections in Hanoi children: an eight-year time series study. *Environ Int*. (2018) 110:139–48. doi: 10.1016/j.envint.2017.10.024
20. Luong LMT, Phung D, Sly PD, Morawska L, Thai PK. The association between particulate air pollution and respiratory admissions among young children in Hanoi, Vietnam. *Sci Total Environ*. (2017) 578:249–55. doi: 10.1016/j.scitotenv.2016.08.012
21. Nhung NTT, Schindler C, Chau NQ, Hanh PT, Dien TM, Thanh NTN, et al. Exposure to air pollution and risk of hospitalization for cardiovascular diseases amongst Vietnamese adults: case-crossover study. *Sci Total Environ*. (2020) 703:134637. doi: 10.1016/j.scitotenv.2019.134637
22. Luong LTM, Dang TN, Huong NTT, Phung D, Tran LK, Thai PK. Particulate air pollution in Ho Chi Minh city and risk of hospital admission for Acute Lower Respiratory Infection (ALRI) among young children. *Environ Pollut*. (2020) 257:113424. doi: 10.1016/j.envpol.2019.113424
23. Nhung NT, Jegasothy E, Ngan NT, Truong NX, Thanh NT, Marks GB, et al. Mortality burden due to exposure to outdoor fine particulate matter in Hanoi, Vietnam: health impact assessment. *Int J Public Health*. (2022) 67:1604331. doi: 10.3389/ijph.2022.1604331
24. Vu HNK, Ha QP, Nguyen DH, Nguyen TTT, Nguyen TT, Nguyen TTH, et al. Poor air quality and its association with mortality in Ho Chi Minh City: case study. *Atmosphere*. (2020) 11:750. doi: 10.3390/atmos11070750
25. General Statistics Office of Vietnam. *Area, Population and Population Density by Province*. Gen Stat Off Vietnam. Available online at: <https://www.gso.gov.vn/px-web-2/> (accessed February 7, 2022).
26. Thanh TNN, Truong XN, Ha VP, Hieu DTP, Anh TNN, Luan NV, et al. *LASER Vietnam Air Pollution Monitoring Project PM2.5 Data (2019–2021)*. LASER PULSE Research for Development Program and VNU—University of Engineering and Technology. (2022).
27. Hong TT, Phuong Hoa N, Walker SM, Hill PS, Rao C. Completeness and reliability of mortality data in Viet Nam: implications for the national routine health management information system. *PLoS ONE*. (2018) 13:e0190755. doi: 10.1371/journal.pone.0190755
28. General Statistics Office of Vietnam. *Conference Announced Results of the in-Depth Analysis of the 2019 Vietnam Population and Housing*. Gen Stat Off Vietnam. (2020). Available online at: <https://www.gso.gov.vn/en/events/2021/01/conference-a-nounced-results-of-the-in-depth-analysis-of-the-2019-vietnam-population-and-housing-census/> (accessed April 4, 2022).
29. Burnett R, Chen H, Szyszkowicz M, Fann N, Hubbell B, Pope CA, et al. Global estimates of mortality associated with long-term exposure to outdoor fine particulate matter. *Proc Natl Acad Sci*. (2018) 115:9592–7. doi: 10.1073/pnas.1803222115
30. Broome R, Horsley J, Hanigan I. *iomlifetR. R Package Version 0.2*. (2022). Available online at: <https://github.com/richardbroome2002/iomlifetR> (accessed July 7, 2022).
31. Hoek G, Krishnan RM, Beelen R, Peters A, Ostro B, Brunekreef B, et al. Long-term air pollution exposure and cardio-respiratory mortality: a review. *Environ Health*. (2013) 12:1–16. doi: 10.1186/1476-069X-12-43
32. Institute for Health Metrics and Evaluation. *GBD Results Tool | GHDx*. Available online at: <http://ghdx.healthdata.org/gbd-results-tool> (accessed January 17, 2022).
33. Vietnamese Congress. *Law 72/2020/QH14 on Environmental Protection*. (2020). Available online at: <https://vanban.chinhphu.vn/?pageid=27160&docid=202613&classid=1&typegroupid=3> (accessed April 5, 2022).
34. Burnett R, Cohen A. Relative risk functions for estimating excess mortality attributable to outdoor PM2.5 air pollution: evolution and state-of-the-art. *Atmosphere*. (2020) 11:589. doi: 10.3390/atmos11060589
35. Vietnam Environment Administration (VEA). *National Environment Report 2016-2020*. (2021). Available online at: <http://vea.gov.vn/bao-cao-moi-truong-quoc-gia1105-7662> (accessed April 5, 2022).
36. Liu J, Yin H, Tang X, Zhu T, Zhang Q, Liu Z, et al. Transition in air pollution, disease burden and health cost in China: a comparative study of long-term and short-term exposure. *Environ Pollut*. (2021) 277:116770. doi: 10.1016/j.envpol.2021.116770
37. Luong LM, Phung D, Dang TN, Sly PD, Morawska L, Thai PK. Seasonal association between ambient ozone and hospital admission for respiratory diseases in Hanoi, Vietnam. *PLoS ONE*. (2018) 13:e0203751. doi: 10.1371/journal.pone.0203751
38. Wang H, He X, Liang X, Choma EF, Liu Y, Shan L, et al. Health benefits of on-road transportation pollution control programs in China. *Proc Natl Acad Sci*. (2020) 117:25370–7. doi: 10.1073/pnas.1921271117
39. Hassan Bhat T, Jiawen G, Farzaneh H. *Air Pollution Health Risk Assessment (AP-HRA), Principles and Applications*. *Int J Environ Res Public Health*. (2021) 18:1935. doi: 10.3390/ijerph18041935