



OPEN ACCESS

EDITED BY

Zhen-Yu Zhang,
KU Leuven, Belgium

REVIEWED BY

Malgorzata Wojcik,
Jagiellonian University Medical
College, Poland
Andrea Lukács,
University of Miskolc, Hungary

*CORRESPONDENCE

Mario Leone
✉ mario.leone@uqac.ca

SPECIALTY SECTION

This article was submitted to
Children and Health,
a section of the journal
Frontiers in Public Health

RECEIVED 28 September 2022

ACCEPTED 05 December 2022

PUBLISHED 05 January 2023

CITATION

Leone M, Levesque P,
Bourget-Gaudreault S, Lemoyne J,
Kalinova E, Comtois AS, Bui HT,
Léger L, Frémont P and Allisse M (2023)
Secular trends of cardiorespiratory
fitness in children and adolescents
over a 35-year period: Chronicle of a
predicted foretold.
Front. Public Health 10:1056484.
doi: 10.3389/fpubh.2022.1056484

COPYRIGHT

© 2023 Leone, Levesque,
Bourget-Gaudreault, Lemoyne,
Kalinova, Comtois, Bui, Léger, Frémont
and Allisse. This is an open-access
article distributed under the terms of
the [Creative Commons Attribution
License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution
or reproduction in other forums is
permitted, provided the original
author(s) and the copyright owner(s)
are credited and that the original
publication in this journal is cited, in
accordance with accepted academic
practice. No use, distribution or
reproduction is permitted which does
not comply with these terms.

Secular trends of cardiorespiratory fitness in children and adolescents over a 35-year period: Chronicle of a predicted foretold

Mario Leone^{1,2*}, Patrick Levesque¹,
Sabrina Bourget-Gaudreault¹, Jean Lemoyne³,
Emilia Kalinova⁴, Alain Steve Comtois⁴, Hung Tien Bui²,
Luc Léger⁵, Pierre Frémont⁶ and Maxime Allisse⁷

¹Department of Health Sciences, Université du Québec à Chicoutimi, Chicoutimi, QC, Canada,

²Department of Medicine and Health Sciences, Université de Sherbrooke, Sherbrooke, QC, Canada,

³Department of Physical Activity Sciences, Université du Québec à Trois-Rivières, Trois-Rivières, QC,

Canada, ⁴Department of Physical Activity Sciences, Université du Québec à Montréal, Montreal, QC,

Canada, ⁵School of Kinesiology, Université de Montréal, Montréal, QC, Canada, ⁶Department of

Kinesiology, Laval University, Quebec, QC, Canada, ⁷Department of Kinanthropology, Université de
Sherbrooke, Sherbrooke, QC, Canada

Background: In the context of concerns regarding the cardiorespiratory fitness (CRF) of youth populations, the aims of this study were: (1) to update reference values for the $VO_2\max$ for school-aged Canadians and (2) to document secular trends in CRF after a 35-year interval.

Methods: Between September 2014 and April 2017, the CRF of 3725 students (53.2% boys; 6.0 to 17.9 yrs) was determined using the 20-m shuttle run test. The sample was collected in 36 different schools from six cities of Québec (Canada).

Results: Median values of $VO_2\max$ decreased with age in both sexes ($p \leq 0.05$). By the age of 10, more than 20% of boys showed $VO_2\max$ values below the recommended value ($42 \text{ ml}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$). At the age of 17, that proportion reached 56.8%. A similar proportion of 12 yrs girls (20%) were under the recommended minimal value ($37 \text{ ml}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$) and that value reached 69.9% at the age of 17. Compared to 1982, the $VO_2\max$ at age 17 has declined by 18% for boys and 12% for girls. The situation is worse in terms of functional capacity (number of stages completed) with an overall decrease of more than 30%.

Conclusion: This study demonstrates that, compared to data obtained using the same methodology 35 years ago, the CRF and functional capacity of children and adolescents has declined to levels that should raise concerns from a public health perspective. Thus, the development of strategies to promote a physically active lifestyle in youth is more relevant than ever.

KEYWORDS

normative reference values, $VO_2\max$, functional capacity, secular trends, youth, cardiorespiratory fitness

Introduction

According to Health Canada (2016) (1) the prevalence of obesity in youth (5–17 years) has more than tripled over the last 30 years. One of the most common explanations is related to the marked decline in physical activity levels during childhood and adolescence (2, 3). In fact, some recent studies have also shown the huge impact of a physically active lifestyle on the prevention and management of multiple chronic health problems such as cardiometabolic risk factors, several cancers, mental health problems and more (4–6). Data from the Public Health Agency of Canada report (2016) (7) indicate that the vast majority of young Canadians fail to meet recommended levels of physical activity due to increased sedentary behaviors. In fact, nearly 91% of children and youth aged 5–17 do not reach the Canadian Physical Activity Guidelines (8) recommendation of 60 min of moderate to vigorous physical activity daily.

Cardiorespiratory fitness

CRF is such a key determinant of health (4, 9) that it has been proposed as a vital sign that should be monitored in clinical practice (10). In childhood and adolescence specifically, poor CRF is a major precursor to the development of short-term and later-life cardiometabolic risk factors and chronic diseases (10–13). According to the WHO and based on several studies, $VO_2\text{max}$ which represents the maximal capacity of the organism to consume oxygen during maximum physical exertion, has long been considered the leading indicator of CRF (14–17). Although some authors have questioned the usefulness and relevance of field tests for the evaluation of the CRF (18, 19) there is a strong consensus in favor of the use of this type of test, particularly for population surveillance (20–22). In fact, over the last 4 decades, the most commonly used test to assess aerobic fitness in school is the 20-m shuttle run test (23). The popularity of this field test relies on the fact that it is easy to manage, requires little equipment and time, is inexpensive and can be administered to several individuals simultaneously. In 2019, Statistics Canada released a set of normative percentile reference values including CRF (24). For practical reasons, the aerobic test chosen was the Modified Canadian Aerobic Fitness Test (mCAFT), a submaximal step test used to estimate an individual's CRF. Due to the very different nature of the procedures, the two tests cannot be interchanged for population surveillance purposes since the estimated $VO_2\text{max}$ values will be different.

Thus, the first objective of this study was to provide an update of the reference values for $VO_2\text{max}$ for the Canadian youth population (aged 6–17). The second objective was document the suspected secular trends in youth CRF by comparing the data collected in 1982 by Léger et al. (23) with the results of the present study.

Methods

Design

This study is a descriptive comparative research with a cross-sectional design based on a large sample of children and adolescents from Québec (Canada).

Participants

Between September 2014 and April 2017, a total of 3,725 students (boys = 1,983; girls = 1,742) were recruited for this study. The age varied between 6.0 and 17.9 years, which covers elementary and high school education in Canada. The participants were recruited in 36 different schools (elementary school = 24 and high school = 12) from six cities in the province of Québec (Montréal, Québec city, Saguenay, Trois-Rivières, Laval and Sherbrooke). The data was collected in the gymnasium of each school during physical education classes. Parents and students were informed of our presence and could indicate their refusal to participate in the project (a consent form was signed by the school authorities). The Institutional Ethical Committee Board (University of Québec in Chicoutimi) approved the project (no: 602-225-01).

Selection of the school boards, schools and classrooms

A three-stage sampling approach was used for the selection of a representative number of school boards, schools and classrooms. Each school received an invitation letter in order to take part in the project. Following the sending of approximately 1,200 invitations to school principals, over 300 schools expressed their interest to participate in this project. Particular attention was also paid to the equitable representation of the various socioeconomic status in our sample through a socioeconomic school rating from the Québec government (Ministère de l'Éducation et de l'Enseignement Supérieur, 2017).

All schools and classrooms were randomly selected by lot. Apart from very rare exceptions, all students of the same class were assessed, thus eliminating a selection bias. If a chosen school would withdraw, a new draw was then carried out. All participants were free from illnesses, disabilities, or injuries that could have been aggravated by physical activity. The sample size required for conducting this study was determined from a Cohen's d power analysis in order to detect small effects ($d < 0.1$) with a $1-\beta = 0.95$ for $\alpha = 0.05$ using G Power software version 3.1.9.4. Thus, 1,564 youths per sex were required for a total of 3,128 participants.

TABLE 1 Anthropometric and cardiorespiratory profiles of boys and girls aged 6–17 years old.

Age	Body mass (kg)		Height (cm)		BMI (kg·m ⁻²)		VO ₂ max (ml·kg ⁻¹ ·min ⁻¹)		Stages (number)	
	Mean±SD	CI (95%)	Mean±SD	CI (95%)	Mean±SD	CI (95%)	Mean±SD	CI (95%)	Mean±SD	CI (95%)
Boys										
6.0–6.9 yrs	24.0 ± 5.1	22.7–25.2	121.3 ± 4.9	120.1–122.5	16.1 ± 3.2	15.3–16.8	48.6 ± 2.5	47.7–48.9	2.1 ± 1.0	1.9–2.4
7.0–7.9 yrs	25.2 ± 4.4	24.4–26.0	125.5 ± 5.5	124.5–126.4	15.9 ± 2.2	15.5–16.3	47.9 ± 2.6	47.5–48.4	2.6 ± 1.3	2.4–2.8
8.0–8.9 yrs	28.9 ± 5.6	28.1–29.8	132.1 ± 6.7	131.1–133.1	16.5 ± 2.9	16.1–16.9	47.5 ± 3.9	46.9–48.1	3.2 ± 1.7	3.0–3.5
9.0–9.9 yrs	31.4 ± 5.7	30.6–32.2	136.8 ± 6.9	135.7–137.8	16.8 ± 2.9	16.4–17.3	46.7 ± 4.5	46.0–47.3	3.6 ± 1.9	3.3–3.8
10.0–10.9 yrs	35.3 ± 7.6	34.3–36.4	142.1 ± 6.6	141.2–143.1	17.4 ± 3.2	16.9–17.8	45.9 ± 4.7	45.2–46.6	3.9 ± 2.0	3.6–4.2
11.0–11.9 yrs	40.1 ± 8.0	38.8–41.3	148.1 ± 7.2	146.9–149.2	18.3 ± 3.4	17.7–18.8	43.9 ± 4.5	43.2–44.6	3.8 ± 1.8	3.5–4.1
12.0–12.9 yrs	47.5 ± 12.2	45.8–49.1	153.6 ± 8.5	152.5–154.8	20.0 ± 4.4	19.4–20.6	44.8 ± 5.2	44.1–45.5	5.0 ± 2.0	4.7–5.2
13.0–13.9 yrs	54.4 ± 12.7	52.7–56.0	161.0 ± 8.8	159.9–162.2	20.9 ± 4.1	20.3–21.4	43.8 ± 4.9	43.1–44.4	5.3 ± 1.9	5.0–5.5
14.0–14.9 yrs	57.9 ± 10.4	56.3–59.5	166.4 ± 7.8	165.2–167.6	21.1 ± 4.1	20.5–21.7	44.6 ± 6.7	43.6–45.6	6.1 ± 2.5	5.8–6.5
15.0–15.9 yrs	63.5 ± 12.2	61.9–65.2	170.5 ± 7.0	169.6–171.5	21.9 ± 4.2	21.3–22.5	43.0 ± 6.9	42.1–43.9	6.1 ± 2.5	5.8–6.5
16.0–16.9 yrs	66.6 ± 11.8	64.7–68.4	172.8 ± 7.6	171.6–174.0	22.5 ± 5.0	21.7–23.3	42.8 ± 7.3	41.7–44.0	6.6 ± 2.5	6.2–7.0
17.0–17.9 yrs	71.0 ± 14.9	68.0–74.5	173.5 ± 7.9	171.8–175.1	23.6 ± 4.8	22.6–24.6	40.3 ± 7.0	39.4–41.8	6.5 ± 2.4	6.0–7.0
Girls										
6.0–6.9 yrs	22.5 ± 3.7	21.6–23.5	120.1 ± 5.1	118.8–121.4	15.6 ± 1.9	15.1–16.0	47.5 ± 2.0	47.5–48.0	1.7 ± 0.8	1.5–1.9
7.0–7.9 yrs	25.0 ± 4.7	24.1–25.8	125.4 ± 5.5	124.4–126.3	15.8 ± 2.2	15.4–16.2	47.1 ± 2.5	46.6–47.5	2.3 ± 1.0	2.1–2.5
8.0–8.9 yrs	28.9 ± 7.7	27.6–30.2	130.8 ± 6.4	129.7–131.8	16.8 ± 3.8	16.2–17.4	46.3 ± 3.1	45.8–46.8	2.6 ± 1.3	2.6–2.8
9.0–9.9 yrs	31.0 ± 6.1	30.1–31.9	136.2 ± 6.6	135.2–137.1	16.7 ± 2.6	16.3–17.0	44.5 ± 2.6	44.1–44.9	2.6 ± 1.1	2.5–2.8
10.0–10.9 yrs	37.7 ± 8.9	36.3–39.1	144.1 ± 8.3	142.8–145.4	18.1 ± 3.4	17.5–18.6	43.8 ± 3.7	43.3–44.4	3.1 ± 1.5	2.9–3.3
11.0–11.9 yrs	42.3 ± 9.3	40.8–43.9	149.5 ± 7.7	148.3–150.8	18.8 ± 3.6	18.2–19.4	42.5 ± 3.3	42.0–43.1	3.3 ± 1.3	3.1–3.5
12.0–12.9 yrs	49.4 ± 11.8	47.8–50.9	154.7 ± 6.7	153.8–155.5	20.5 ± 4.4	20.0–21.1	41.6 ± 4.4	41.1–42.2	3.7 ± 1.7	3.5–4.0
13.0–13.9 yrs	52.1 ± 11.2	50.6–53.6	157.2 ± 6.9	156.3–158.2	21.1 ± 4.4	20.5–21.7	39.5 ± 4.6	38.9–40.1	3.7 ± 1.8	3.4–3.9
14.0–14.9 yrs	57.6 ± 13.2	55.5–59.8	157.9 ± 6.4	156.9–159.0	23.1 ± 5.1	22.2–23.9	37.3 ± 4.3	36.6–38.0	3.5 ± 1.6	3.2–3.7
15.0–15.9 yrs	57.4 ± 9.2	55.7–59.1	161.4 ± 7.6	160.0–162.8	22.0 ± 3.6	21.3–22.6	38.4 ± 5.6	37.4–39.4	4.6 ± 2.0	4.2–4.9
16.0–16.9 yrs	59.7 ± 10.8	57.9–61.6	162.3 ± 9.0	160.8–163.9	22.7 ± 4.4	22.0–23.5	35.2 ± 5.1	34.3–36.1	4.2 ± 1.9	3.9–4.5
17.0–17.9 yrs	60.9 ± 13.0	57.8–64.0	162.6 ± 8.8	160.4–164.7	22.7 ± 4.2	21.7–23.7	33.9 ± 5.5	32.6–35.2	4.0 ± 1.8	3.6–4.5

SD, Standard deviation; CI, Confidence interval; BMI, Body mass index.

Anthropometric measures

Anthropometric variables were collected using procedures proposed by Lohman et al. (25). Body mass (BM) was noted to the nearest 0.1 kg using a Detecto scale (Missouri, USA). Body height (BH) was assessed using a Lafayette stadiometer (Louisiana, USA) at the nearest 0.1 cm. Body mass index (BMI) was also calculated. BMI (typical vs. overweight and obese youths) was classified according to age and sex as suggested by Cole et al. (26).

Cardiorespiratory fitness

CRF was determined in accordance with the 20-m shuttle run test described and validated by Léger et al. (23). Briefly, the test took place in a standard size gymnasium of at least 25 m. The entire classroom (generally around twenty students) took up position on the starting line. Whenever a participant could no longer follow the required running speed, he or she was stopped and the number of the last completed stage was recorded. At the end of the test, the following information was then extracted or estimated for each student: the number of the final stage, the associated running speed ($\text{km}\cdot\text{h}^{-1}$) and the estimated VO_2max value ($\text{ml}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$).

Statistical analysis

All descriptive values are reported as mean \pm standard deviation (SD). Confidence intervals (CI) were set at the 95% level. Cohen's effect sizes were calculated for various intergroup comparisons. The Shapiro-Wilk test for normality was compiled for each variable. When normality was violated, a Box-Cox transformation (27) was conducted using the following equation:

$$BC = (\text{VAR}^L - -1) \cdot L^{-1} \text{ when } L \neq 0$$

$$BC = \text{Log}(\text{VAR}) \text{ when } L = 0$$

Where, BC, Box-Cox transformation; VAR, variable; L, lambda

The Box-Cox power exponential method, which smoothed the curves by cubic splines, has been used to create the curves.

Outliers were identified using the method proposed by Hoaglin et al. (28, 29). The equation reads as follows:

$$[(Q75 - Q25) \cdot g] - Q25 \text{ for the lowest value}$$

$$[(Q75 - Q25) \cdot g] + Q75 \text{ for the highest value}$$

Where $Q75 = 3^{\text{rd}}$ quartile; $Q25 = 4^{\text{th}}$ quartile; $g = 2.2$

Percentiles values were computed using the LMS method, (30) which read as follows:

$$P = M \cdot [1 + \text{LSZ}]^{1/L}$$

Where, P = percentile; M = median; L = Lambda; S = coefficient of variation; S = Z-score for the desire percentile.

In order to be able to assess changes that have occurred between 1982 and 2017, the data from the present study were compared with the study carried out by Léger et al. (23) using an unpaired *T*-Test. Statistical analysis was produced by the IBM-SPSS software version 24.

Results

Anthropometric (BM, BH and BMI) and cardiorespiratory (number of stages completed and VO_2max) characteristics as a function of age and sex are shown in Table 1. From the age of 10, girls are heavier and taller than boys until about the age of 13 years, which is consistent with puberty in girls. The cardiorespiratory profile presents a different picture where boys already have higher values for all age groups. This difference is particularly marked for the functional component of the test, which is reflected by the number of stages completed in the 20-m shuttle run test.

Percentile curves of VO_2max and functional capacity, for boys and girls aged 6 to 17, are presented in Figures 1A, B. Between age 6 and 17, the median values for VO_2max declines by about 14% for boys and 27% for girls. Also, this trend appears to be strongly affected by the percentile (VO_2max) reached in early childhood. For example, for the 25th percentile value, a decline of 21% for boys and 33% for girls is observed between the age of 6 and 17. In Figures 1C, D, the percentile curves of the number of stages during the 20-m shuttle run test provide useful information regarding the functional aspect of the cardiorespiratory capacity. Thus, individuals in the upper percentiles tend to considerably improve the number of stages completed throughout the physical growth period compared to individuals in the lower percentiles.

Tables 2, 3 provide the standardized values for VO_2max and functional capacity (stages) respectively. All parameters included in the LMS method are reported for each year of chronological age (6–17 years) for both sexes. Additionally, values for the 3rd, 10th, 25th, 50th, 75th, 90th, and 97th percentiles are also shown.

The impact of BMI on VO_2max was also examined (Figures 2A, B). As shown in Table 4, boys in the overweight/obese zone have VO_2max values markedly lower than individuals with typical BMI across all age groups and this difference increases between the age of 14 and 17. In girls, a similar but less important difference is observed between overweight/obese individuals and those with typical BMI follows a slightly shifted curve which increases with age in favor of the former. For the functional aspect of the 20-m shuttle run test, a very large discrepancy is also observed




FIGURE 1
Age-specific smoothed percentile curves for VO₂max and the number of stages completed estimated using the 20-m shuttle run test for boys (A, C) and girls respectively (B, D).

when comparing individuals from the two BMI categories (Figures 2C, D).

Secular trend for VO₂max (Figures 3A, B) and the number of stages completed in the 20-m shuttle run test (Figures 3C, D) over a 35-year interval are illustrated by comparing data from the present study with data collected in 1982 using the same methodology (23). Over 35 years, median VO₂max decreased by 7.6% for boys and 8.3% for girls at the age of 6 and this difference increases to nearly 18% in boys and 12.2% in girls by the age of 17 (Table 5). Compared to 1982, a significant decrease is also

observed for all age groups for the number of stages completed for boys and girls.

Given that the difference in body mass between 1982 and 2017 could influence the VO₂max observed between the two periods, BM normalization was carried out. The discrepancies observed between the 2 periods are not attenuated by BM standardization as shown in Table 6. Similarly, BM normalization did not affect the secular trends observed between 1982 and 2017 regarding the number of stages completed.

TABLE 2 Percentile references values for VO₂max according to age and gender in Québec children and adolescents (N = 3725).

	N	L	M	S	Percentiles						
					3th	10th	25th	50th	75th	90th	97th
Boys											
6.0–6.9 yrs	66	−1.55	48.3	0.052	44.1	45.3	46.7	48.3	50.1	51.8	53.7
7.0–7.9 yrs	123	−0.45	48.0	0.057	43.2	44.7	46.2	48.0	49.9	51.7	53.6
8.0–8.9 yrs	168	−1.70	47.5	0.083	41.4	43.1	45.0	47.5	50.4	53.4	56.9
9.0–9.9 yrs	178	−1.83	47.0	0.097	40.2	42.1	44.3	47.0	50.5	54.3	58.8
10.0–10.9 yrs	192	0.01	46.2	0.104	38.0	40.4	43.1	46.2	49.6	52.8	56.2
11.0–11.9 yrs	158	−0.70	43.9	0.104	36.5	38.6	41.0	43.9	47.2	50.5	54.2
12.0–12.9 yrs	213	0.43	44.6	0.115	35.5	38.3	41.2	44.6	48.1	51.4	54.8
13.0–13.9 yrs	232	1.06	43.8	0.110	34.7	37.6	39.8	43.8	47.0	50.0	52.8
14.0–14.9 yrs	180	0.54	44.8	0.150	32.0	36.6	40.4	44.8	49.4	53.8	58.3
15.0–15.9 yrs	216	1.09	43.1	0.160	30.0	34.2	38.4	43.1	47.7	51.8	55.9
16.0–16.9 yrs	162	0.90	42.8	0.171	29.3	33.5	37.9	42.8	47.8	52.3	56.8
17.0–17.9 yrs	95	0.90	41.6	0.181	27.8	32.1	36.6	41.6	46.7	51.3	55.9
Girls											
6.0–6.9 yrs	62	−1.00	47.5	0.042	44.0	45.1	46.2	47.5	48.9	50.2	51.6
7.0–7.9 yrs	119	−2.00	46.9	0.051	43.0	44.1	45.4	46.9	48.6	50.3	52.2
8.0–8.9 yrs	139	−1.90	46.4	0.067	41.5	42.9	44.4	46.4	48.6	51.0	53.6
9.0–9.9 yrs	180	−2.64	44.5	0.059	40.4	41.5	42.9	44.5	46.4	48.4	50.8
10.0–10.9 yrs	164	−1.05	43.7	0.089	37.7	39.4	41.3	43.7	46.4	49.0	52.0
11.0–11.9 yrs	143	−1.08	42.3	0.084	36.9	38.5	40.2	42.3	44.7	47.0	49.6
12.0–12.9 yrs	240	−0.96	41.9	0.105	35.0	36.9	39.1	41.9	45.1	48.4	52.2
13.0–13.9 yrs	218	−1.60	39.3	0.119	32.5	34.3	36.5	39.3	42.8	46.8	51.8
14.0–14.9 yrs	146	−0.70	37.1	0.115	30.3	32.2	34.4	37.1	40.2	43.4	47.0
15.0–15.9 yrs	123	0.30	37.9	0.145	28.4	31.2	34.3	37.9	41.8	45.5	49.4
16.0–16.9 yrs	135	−0.40	36.6	0.167	27.2	29.8	32.8	36.6	41.1	45.8	51.3
17.0–17.9 yrs	73	−0.70	34.4	0.162	26.1	28.4	31.0	34.4	38.1	43.0	48.4

N, Number of participants; L, lambda; M, median; S, coefficient of variation ($SD \cdot mean^{-1}$).

TABLE 3 Percentile reference values for the number of stages completed during the 20-m shuttle run test according to age and gender in Québec children and adolescents (N = 3725).

	N	L	M	S	Percentiles						
					3th	10th	25th	50th	75th	90th	97th
Boys											
6.0–6.9 yrs	66	0.25	2.22	0.519	0.75	1.00	1.50	2.25	3.00	4.00	5.25
7.0–7.9 yrs	124	0.34	2.71	0.487	1.00	1.25	2.00	2.75	3.75	4.75	6.00
8.0–8.9 yrs	168	0.23	3.35	0.514	1.00	1.50	2.25	3.25	4.75	6.25	8.00
9.0–9.9 yrs	178	0.34	3.85	0.531	1.25	1.75	2.75	3.75	5.50	7.00	9.00
10.0–10.9 yrs	192	0.63	4.19	0.501	1.00	1.75	2.75	4.25	5.75	7.25	8.50
11.0–11.9 yrs	158	0.63	3.67	0.480	1.00	1.75	2.50	3.75	5.00	6.25	7.50
12.0–12.9 yrs	213	0.69	5.13	0.405	1.75	2.75	3.75	5.00	6.50	8.00	9.50
13.0–13.9 yrs	232	1.00	5.50	0.351	1.75	3.00	4.25	5.50	6.75	8.00	9.25
14.0–14.9 yrs	180	0.66	6.18	0.397	2.25	3.25	4.50	6.25	8.00	9.50	11.25
15.0–15.9 yrs	217	1.00	6.00	0.400	1.50	3.00	4.50	6.00	7.50	9.00	10.50
16.0–16.9 yrs	162	0.86	6.57	0.378	2.25	3.50	5.00	6.50	8.25	9.75	11.50
17.0–17.9 yrs	95	0.81	6.59	0.372	2.25	3.50	5.00	6.50	8.25	9.75	11.50
Girls											
6.0–6.9 yrs	62	−0.60	1.73	0.432	0.75	1.00	1.25	1.75	2.50	3.50	5.25
7.0–7.9 yrs	120	0.01	2.25	0.462	1.00	1.25	1.50	2.25	3.00	4.00	5.25
8.0–8.9 yrs	139	0.12	2.84	0.492	1.00	1.50	2.00	2.75	4.00	5.25	6.75
9.0–9.9 yrs	180	0.26	2.67	0.440	1.00	1.50	2.00	2.75	3.50	4.50	5.75
10.0–10.9 yrs	165	0.34	3.20	0.498	1.00	1.50	2.25	3.25	4.50	5.75	7.25
11.0–11.9 yrs	143	0.38	3.16	0.389	1.25	1.75	2.25	3.25	4.00	5.00	6.00
12.0–12.9 yrs	240	0.48	3.74	0.449	1.25	2.00	2.75	3.75	5.00	6.25	7.50
13.0–13.9 yrs	218	0.31	3.81	0.484	1.25	2.00	2.75	3.75	5.25	6.75	8.50
14.0–14.9 yrs	146	0.62	3.15	0.465	1.00	1.50	2.25	3.25	4.25	5.25	6.50
15.0–15.9 yrs	123	0.63	4.16	0.429	1.25	2.00	3.00	4.25	5.50	6.75	8.00
16.0–16.9 yrs	133	0.37	4.30	0.472	1.50	2.25	3.00	4.25	5.75	7.50	9.25
17.0–17.9 yrs	73	0.61	4.18	0.460	1.25	2.00	3.00	4.25	5.50	7.00	8.50

N, Number of participants; L, lambda; M, median; S, coefficient of variation ($SD \cdot mean^{-1}$); The number of stages has been rounded to the nearest quarter. The speed at the first stage starts at $8.5 \text{ km} \cdot \text{h}^{-1}$ and then increase by $0.5 \text{ km} \cdot \text{h}^{-1}$ every minute.




FIGURE 2 Modeling of VO_{2max} curves and the number of stages completed in youth with typical BMI or with overweight/obese profile in boys (A, C) and girls (B, D). The thick horizontal line represents the critical cutoff health zone.

Discussion

This study provides recent reference values for the maximal aerobic 20-m shuttle run test in children and adolescents of the province of Québec (Canada). It also provides unique opportunity to directly compare recent CRF data with reference values initially documented for this test in the same geographic area and age group in 1982 (31).

CRF reference values

According to different studies and regardless of age, it is estimated that a minimum VO_{2max} value of approximately 42 ml·kg⁻¹·min⁻¹ in boys and 37 ml·kg⁻¹·min⁻¹ in girls is required to minimize the risk of developing severe health problems (32, 33). Considering these CRF thresholds, the reference values documented in this study raise a powerful red

TABLE 4 Comparison of VO₂max and the number of stages completed in children and adolescents with typical or overweight/obese BMI profile.

	Typical BMI		Overweight/Obese		Δ %		P-values		Cohen's d	
	Boys	Girls	Boys	Girls	Boys	Girls	Boys	Girls	Boys	Girls
6.0–6.9 yrs										
VO ₂ max	48.3 ± 2.6	47.3 ± 1.9	48.5 ± 2.6	48.3 ± 2.4	−0.4	−2.0	0.856	0.193	0.08	0.51
Stages	2.2 ± 1.2	1.7 ± 0.8	2.0 ± 1.0	1.5 ± 0.6	9.1	11.8	0.668	0.542	0.17	0.26
7.0–7.9 yrs										
VO ₂ max	48.1 ± 2.8	47.3 ± 2.5	47.1 ± 1.9	45.7 ± 1.8	2.1	3.4	0.159	0.007	0.37	0.67
Stages	2.7 ± 1.3	2.3 ± 1.1	2.1 ± 1.0	1.6 ± 0.6	22.2	30.4	0.104	0.000	0.47	0.68
8.0–8.9 yrs										
VO ₂ max	47.9 ± 3.9	46.7 ± 3.4	44.9 ± 3.3	44.6 ± 3.1	6.3	4.5	0.000	0.001	0.78	0.63
Stages	3.4 ± 1.7	2.9 ± 1.4	2.2 ± 1.2	1.8 ± 0.9	24.1	37.9	0.000	0.000	0.73	0.84
9.0–9.9 yrs										
VO ₂ max	47.1 ± 4.5	44.7 ± 2.6	44.3 ± 4.0	43.4 ± 2.7	2.3	2.9	0.003	0.013	0.63	0.50
Stages	3.8 ± 1.9	2.7 ± 1.1	2.5 ± 1.6	2.2 ± 1.0	34.2	18.5	0.001	0.014	0.70	0.46
10.0–10.9 yrs										
VO ₂ max	46.6 ± 4.6	44.2 ± 3.8	41.9 ± 3.4	42.7 ± 3.3	10.1	3.4	0.000	0.031	1.06	0.41
Stages	4.2 ± 2.0	3.3 ± 1.6	2.4 ± 1.3	2.5 ± 1.3	42.9	24.2	0.000	0.007	0.94	0.52
11.0–11.9 yrs										
VO ₂ max	44.6 ± 4.5	43.1 ± 3.1	40.8 ± 3.6	40.1 ± 3.2	8.5	7.0	0.000	0.000	0.87	0.96
Stages	4.1 ± 1.9	3.5 ± 1.3	2.6 ± 1.3	2.4 ± 1.0	36.6	31.4	0.000	0.000	0.83	0.88
12.0–12.9 yrs										
VO ₂ max	45.8 ± 5.0	41.6 ± 4.1	42.2 ± 4.7	41.8 ± 5.1	7.9	−0.5	0.000	0.714	0.73	0.05
Stages	5.3 ± 1.9	4.2 ± 1.7	4.0 ± 1.9	2.8 ± 1.3	24.5	33.3	0.000	0.000	0.68	0.88
13.0–13.9 yrs										
VO ₂ max	44.7 ± 4.6	40.6 ± 4.5	41.4 ± 4.7	36.6 ± 4.1	8.1	9.9	0.000	0.000	0.71	0.91
Stages	5.6 ± 1.7	4.0 ± 1.8	4.4 ± 1.8	2.7 ± 1.5	21.4	32.5	0.000	0.000	0.69	0.75
14.0–14.9 yrs										
VO ₂ max	46.0 ± 6.2	38.5 ± 4.2	39.8 ± 6.6	35.3 ± 3.4	13.5	8.3	0.000	0.000	0.99	0.81
Stages	6.7 ± 2.3	3.9 ± 1.6	4.4 ± 2.4	2.7 ± 1.3	34.2	30.8	0.000	0.000	0.99	0.80

(Continued)

TABLE 4 (Continued)

	Typical BMI		Overweight/Obese		Δ %		P-values		Cohen's d	
	Boys	Girls	Boys	Girls	Boys	Girls	Boys	Girls	Boys	Girls
15.0–15.9 yrs										
VO ₂ max	44.1 ± 6.7	39.6 ± 5.3	39.6 ± 6.8	34.2 ± 4.7	10.2	13.6	0.000	0.000	0.67	1.05
Stages	6.5 ± 2.4	5.0 ± 1.9	4.9 ± 2.4	3.1 ± 1.6	24.6	38.0	0.000	0.000	0.67	1.04
16.0–16.9 yrs										
VO ₂ max	44.5 ± 6.8	36.4 ± 6.1	38.0 ± 5.9	34.3 ± 5.6	14.6	5.8	0.000	0.109	0.99	0.35
Stages	7.2 ± 2.4	4.5 ± 2.1	5.0 ± 2.0	3.6 ± 1.7	31.4	20.0	0.000	0.040	0.96	0.45
17.0 yrs +										
VO ₂ max	42.9 ± 6.3	34.8 ± 4.8	35.4 ± 6.6	30.5 ± 6.3	17.5	12.4	0.000	0.004	1.18	0.83
Stages	7.2 ± 2.1	4.3 ± 1.7	4.7 ± 2.1	3.0 ± 2.0	34.7	30.2	0.000	0.010	1.19	0.73

Yrs, years; P-values in bold = significant at ≤0.05; Δ% = percentage of difference; Cohen's d = 0.20 marginal; 0.50 moderate; 0.80 large; Cohen d in bold = large effect (≥0.80).

flag by showing that, by the age of 17 in boys and 15 in girls, the median VO₂max value is below the minimal CRF level associated with favorable health outcomes later in life. These results are consistent with the reference values reported by Tomkinson et al. (6) in a meta-analysis of 177 studies, most of which were published between 2000 and 2015.

Functional capacity is also affected, demonstrating that today's youth have a reduced ability to sustain moderate/intense effort. In fact, this decrease begins 1 year earlier than the decline in VO₂max. This finding is certainly as worrying as the decrease in VO₂max.

The results also indicate that the higher the CRF reached at a young age, the greater the chances that it will be maintained during the growth period. Assuming that this tendency persists later in life through adulthood, these results further support the notion that childhood CRF can contribute to prevent the development of cardiometabolic risk factors and diseases later in life (14).


Effect of overweight/obesity on the CRF

Although obesity is recognized as a major cause of morbidity, (1, 32) a recent meta-analysis indicated that it is not an independent factor of premature mortality (14). In the present study, data stratification for BMI (typical vs. overweight/obese) shows that, in the later group, a higher proportion evolves toward a VO₂max value below the minimal CRF level associated with favorable health outcomes. In boys with the overweight/obese profile, it is noted that the critical median cutoff value of VO₂max is crossed as soon as the age of 11, which never happens for the typical BMI group.

The number of stages completed as a function of age is also heavily impaired in the overweight/obese BMI group. These results further support the notion that lower CRF and functional capacity are factors that likely contribute to unfavorable cardiometabolic outcomes in youth with BMI that correspond to the definition of overweight and obesity. However, other factors also need to be taken into account since in girls with a normal BMI median VO₂max will eventually fall below the recommended cutoff by the age of 16. Reduced physical activity combined with increased passive activities has been suggested as a likely factor that explains this situation (32, 34, 35).

Secular trends in CRF

The great heterogeneity of CRF assessment procedures makes comparisons between studies complicated. In 1982, Léger and colleagues developed the 20-m shuttle run test and developed reference values for the CRF of youth living in Quebec (23). International reference values for the 20-m shuttle run



test were recently developed by combining data from studies published up to 2015 (6). However, these more recent reference values do not allow comparison for specific population over time. The present study has the advantage of using the same test, administered under the same conditions, in the same cities and in the same school boards as 35 years earlier by Léger et al. (23).

This methodology resulted in a unique opportunity to objectively appreciate the evolution of CRF in youth between 1982 and 2017. The results confirmed an important decrease of CRF (estimated VO_2max) and functional capacity (number

of stages completed) in the study population. This difference tends to accentuate with age, with a VO_2max decrease reaching nearly -18% for males and -12% for females at the age of 17. The functional impact of this situation is even more important in terms of the number of stages completed with an overall decrease of more than -30% . Furthermore, in 1982, all age groups of both sexes displayed VO_2max values above the minimal recommended threshold associated with positive health outcomes. In the present study population, this is no longer the case from the age of 16 for boys and 15 for girls.

TABLE 5 Comparison of VO₂max and the number of stages completed in children and adolescents between 1982 and 2017.

	Léger et al. (23)		Current study		Δ %		P-values		Cohen's d	
	Boys	Girls	Boys	Girls	Boys	Girls	Boys	Girls	Boys	Girls
6.0–6.9 yrs										
VO ₂ max	52.4 ± 2.8	51.8 ± 2.3	48.4 ± 2.5	47.5 ± 2.0	−7.6	−8.3	0.000	0.000	1.43	1.96
Stages	3.6 ± 1.4	3.4 ± 1.1	2.1 ± 1.1	1.7 ± 0.7	−41.7	−50.0	0.000	0.000	1.15	1.74
7.0–7.9 yrs										
VO ₂ max	51.2 ± 3.3	50.3 ± 2.6	48.0 ± 2.7	47.0 ± 2.4	−6.3	−6.6	0.000	0.000	1.02	1.3
Stages	3.9 ± 1.6	3.5 ± 1.2	2.6 ± 1.3	2.2 ± 1.0	−33.3	−37.1	0.000	0.000	0.86	1.13
8.0–8.9 yrs										
VO ₂ max	51.7 ± 3.9	49.8 ± 3.4	47.5 ± 3.9	46.3 ± 3.1	−8.1	−7.0	0.000	0.000	1.11	1.06
Stages	4.9 ± 1.8	4.1 ± 1.5	3.2 ± 1.7	2.7 ± 1.1	−34.7	−36.6	0.000	0.000	0.96	1.04
9.0–9.9 yrs										
VO ₂ max	51.5 ± 4.4	49.2 ± 3.2	46.7 ± 4.5	44.5 ± 2.6	−9.3	−9.6	0.000	0.000	1.10	1.57
Stages	5.5 ± 1.9	4.5 ± 1.4	3.6 ± 1.9	2.7 ± 1.1	−41.8	−40.0	0.000	0.000	1.26	1.38
10.0–10.9 yrs										
VO ₂ max	51.6 ± 4.2	46.8 ± 2.8	45.9 ± 4.8	43.8 ± 3.7	−11.2	−6.2	0.000	0.000	1.33	0.92
Stages	6.2 ± 1.8	4.9 ± 1.4	3.9 ± 2.0	3.1 ± 1.6	−37.1	−36.7	0.000	0.000	1.23	1.26
11.0–11.9 yrs										
VO ₂ max	51.1 ± 4.5	47.5 ± 4.0	43.9 ± 4.6	42.6 ± 3.3	−14.1	−10.1	0.000	0.000	1.60	1.24
Stages	6.7 ± 1.8	5.2 ± 1.6	3.8 ± 1.8	3.3 ± 1.3	−43.3	−36.5	0.000	0.000	1.61	1.25
12.0–12.9 yrs										
VO ₂ max	51.9 ± 5.2	46.7 ± 4.2	44.8 ± 5.2	41.6 ± 4.4	−13.7	−10.9	0.000	0.000	1.37	1.19
Stages	7.2 ± 2.0	5.5 ± 1.6	5.0 ± 2.0	3.7 ± 1.7	−30.6	−32.7	0.000	0.000	1.10	1.09
13.0–13.9 yrs										
VO ₂ max	50.1 ± 5.2	44.4 ± 4.8	43.7 ± 4.8	39.5 ± 4.7	−12.6	−11.0	0.000	0.000	1.23	1.04
Stages	7.4 ± 2.0	5.3 ± 1.8	5.2 ± 1.8	3.7 ± 1.8	−29.7	−30.2	0.000	0.000	1.15	0.89
14.0–14.9 yrs										
VO ₂ max	50.1 ± 5.2	41.7 ± 4.7	44.6 ± 6.7	37.3 ± 4.3	−11.0	−10.6	0.000	0.000	0.94	0.97
Stages	8.0 ± 1.9	4.8 ± 1.8	6.2 ± 2.5	3.5 ± 1.6	−23.8	−27.1	0.000	0.000	0.88	0.75

(Continued)

TABLE 5 (Continued)

	Léger et al. (23)		Current study				Δ %		P-values		Cohen's d	
	Boys	Girls	Boys	Girls	Boys	Girls	Boys	Girls	Boys	Girls	Boys	Girls
	15.0–15.9 yrs											
VO ₂ max	50.2 ± 6.1	41.2 ± 5.1	43.1 ± 6.9	38.4 ± 5.6	-14.3	-6.8	0.000	0.000	1.12	0.55		
Stages	8.5 ± 2.2	5.2 ± 1.8	6.1 ± 2.5	4.6 ± 2.0	-28.2	-11.5	0.000	0.000	1.03	0.32		
16.0–16.9 yrs												
VO ₂ max	50.0 ± 5.8	39.5 ± 5.0	42.8 ± 7.3	35.9 ± 6.0	-15.2	-9.1	0.000	0.000	1.12	0.68		
Stages	8.9 ± 2.0	5.2 ± 1.7	6.6 ± 2.5	4.3 ± 2.0	-29.2	-17.3	0.000	0.000	1.06	0.67		
17.0 yrs +												
VO ₂ max	49.7 ± 5.9	38.6 ± 5.2	40.9 ± 7.4	33.9 ± 5.5	-17.7	-12.2	0.000	0.000	1.38	0.89		
Stages	9.3 ± 2.0	5.5 ± 1.8	6.5 ± 2.4	4.1 ± 1.9	-30.1	-25.5	0.000	0.000	1.27	0.76		

Yrs, years; P-values in bold = significant at ≤0.05; Δ% = percentage of difference (the negative sign indicates the decrease in percentage since 1982; Cohen's d = 0.20 marginal; 0.50 moderate; 0.80 large; Cohen d in bold = large effect (≥0.80).

While some authors deny the fact that CRF has decreased over the last decades (18, 19) our results clearly show an alarming decline, both in relative (ml·kg⁻¹·min⁻¹) and functional values (number of stages). It has been suggested that the decrease in VO₂max when expressed in relative values is predictable given the increase in body mass in youth over the past decades. However, data from the present study indicates that, after normalization for BM, significant differences remain for VO₂max and for the number of stages completed in all age groups for both sexes. It may be noted that even when the body mass of Léger's cohort age groups was heavier, the relative and functional values remain markedly higher in their favor. Under these circumstances, it is reasonable to assume that body mass alone is insufficient to explain these differences.

It is assumed that, in addition to the body mass gain observed in recent decades in children and adolescents, increased time spent on sedentary activities is the factor that probably best explains the decrease in CRF (14, 21). Back in the 1980s when the 20-m shuttle run test was developed, computers and video games were in their infancy. With the development of the Internet and social networks youth became less physical active with an increase of time spent on sedentary activities (36, 37). As of 2016, combined data from 146 countries indicates that over 80% of adolescents do not meet the recommended levels of physical activity (38). Even more recently, the substantial reduction in physical activity due to containment measures related to COVID-19 is expected to further accelerate this decline (39).

Reference values vs. standard values

In this paper, we use two distinct concepts that deserve to be explained. Based on the Centers for Disease Control and Prevention (DCC) in 2002, (40) reference values reflect the current situation without regard to its impact on health (what is). This information should not be interpreted as an objective to be achieved. This seems quite obvious as the VO₂max values as well as the number of stages completed (i.e., functional capacity) have considerably decreased over the last decades. Thus, the role of the reference values is to make possible to measure the actual changes, and perhaps those that may occur in the future. They also allow comparison of current values with those from other studies.

On the other hand, the standard values represent what is minimally desirable in order to protect against certain potential health problems (what should be). In the case of VO₂max, the minimum threshold should be around 42 ml in boys and 37 ml in girls. In order to stay above these thresholds, the reference values indicate that youths should follow at least around the 65th percentile curve throughout the growth period.

TABLE 6 Comparison of VO₂max and the number of stages completed between Current study and Léger et al. (23) with adjustment for BM.

Age	Boys											
	Current study				Léger et al. (23)				Statistics			
	N	Body mass	VO ₂ max	Stages	N	Body mass	P	N	VO ₂ max	P	Stages	P
6	56	22.0 ± 2.8	48.3 ± 2.6	2.2 ± 1.1	89	23.4 ± 3.0	0.006	121	52.4 ± 2.8	0	3.6 ± 1.4	0
7	108	24.0 ± 3.2	48.1 ± 2.8	2.7 ± 1.3	221	25.2 ± 4.3	0.011	297	51.2 ± 3.3	0	3.9 ± 1.6	0
8	144	27.0 ± 3.7	47.9 ± 3.9	3.4 ± 1.6	211	28.0 ± 4.5	0.028	303	51.7 ± 3.9	0	4.9 ± 1.8	0
9	148	29.6 ± 4.4	47.1 ± 4.5	3.8 ± 1.9	200	31.8 ± 5.7	0	322	51.5 ± 4.4	0	5.5 ± 1.9	0
10	163	32.8 ± 5.1	46.6 ± 4.6	4.2 ± 1.9	253	34.6 ± 5.9	0.002	404	51.6 ± 4.2	0	6.2 ± 1.8	0
11	129	36.9 ± 5.5	44.6 ± 4.5	4.1 ± 1.8	247	38.8 ± 7.8	0.014	386	51.1 ± 4.5	0	6.7 ± 1.8	0
12	153	41.6 ± 7.4	45.8 ± 5.0	5.3 ± 1.9	206	42.7 ± 7.9	0.181	341	51.9 ± 5.2	0	7.2 ± 2.0	0
13	158	48.0 ± 8.3	44.7 ± 4.5	5.6 ± 1.7	233	47.8 ± 8.6	0.819	325	50.1 ± 5.2	0	7.4 ± 2.0	0
14	131	53.4 ± 7.0	46.0 ± 6.2	6.7 ± 2.2	237	53.4 ± 9.8	1	289	50.1 ± 5.2	0	8.0 ± 1.9	0
15	156	58.0 ± 7.7	44.1 ± 6.7	6.5 ± 2.4	254	58.3 ± 9.7	0.743	333	50.2 ± 6.1	0	8.5 ± 2.2	0
16	117	60.6 ± 8.0	44.5 ± 6.8	7.2 ± 2.3	245	62.6 ± 9.2	0.045	336	50.0 ± 5.8	0	8.9 ± 2.0	0
17	67	63.5 ± 6.6	42.9 ± 6.3	7.2 ± 2.1	161	64.5 ± 8.9	0.408	212	49.7 ± 5.9	0	9.3 ± 2.0	0
Girls												
6	51	21.8 ± 2.7	47.3 ± 1.9	1.7 ± 0.8	81	22.8 ± 2.9	0.05	112	51.8 ± 2.3	0	3.4 ± 1.1	0
7	97	23.3 ± 3.0	47.3 ± 2.5	2.3 ± 1.0	227	24.4 ± 3.6	0.009	299	50.3 ± 2.6	0	3.5 ± 1.2	0
8	111	26.2 ± 3.6	46.7 ± 3.1	2.9 ± 1.2	231	28.0 ± 5.2	0.001	308	49.8 ± 3.4	0	4.1 ± 1.5	0
9	152	28.9 ± 3.9	44.7 ± 2.6	2.7 ± 1.1	196	31.4 ± 5.6	0	322	49.2 ± 3.2	0	4.5 ± 1.4	0
10	128	34.2 ± 6.0	44.2 ± 3.8	3.3 ± 1.5	214	34.6 ± 7.0	0.59	335	46.8 ± 2.8	0	4.9 ± 1.5	0
11	118	39.5 ± 7.2	43.1 ± 3.1	3.5 ± 1.2	258	39.2 ± 8.5	0.74	382	47.5 ± 4.0	0	5.2 ± 1.6	0
12	165	43.4 ± 6.3	41.6 ± 4.1	4.2 ± 1.6	204	45.1 ± 9.0	0.041	292	46.7 ± 4.2	0	5.5 ± 1.6	0
13	161	47.2 ± 7.7	40.6 ± 4.5	4.0 ± 1.7	224	49.2 ± 9.0	0.023	298	44.4 ± 4.8	0	5.3 ± 1.8	0
14	93	49.7 ± 5.7	38.5 ± 4.2	3.9 ± 1.6	211	50.4 ± 7.3	0.412	260	41.7 ± 4.7	0	4.8 ± 1.8	0
15	89	53.5 ± 6.8	39.6 ± 5.3	5.0 ± 1.8	189	53.6 ± 7.1	0.912	260	41.2 ± 5.1	0.012	5.2 ± 1.8	0.366
16	102	56.0 ± 7.3	36.6 ± 5.9	4.5 ± 2.0	236	54.2 ± 7.8	0.048	332	39.5 ± 5.0	0	5.2 ± 1.7	0.001
17	54	56.0 ± 8.4	34.8 ± 4.8	4.4 ± 1.6	133	54.4 ± 7.4	0.199	155	38.6 ± 5.2	0	5.5 ± 1.8	0

N, number of participants; Body mass = kg; VO₂max = ml·kg⁻¹·min⁻¹; Stages = number completed; P-values in bold = significant at ≤ 0.05.

Strengths and limitations

The large sample size ($n > 3700$) allows a valid representation of youths living in Québec (Canada). The test used to estimate the $VO_2\text{max}$ is internationally accepted as valid and reliable. The procedure used was repeated under the same conditions: same cities, same test and same school boards as the original 1982 study, which allows to assess the secular trends with a reduced number of biases. However, some limitations should also be noted. The cross-sectional nature of the data restricts inferences. $VO_2\text{max}$ values were estimated instead of measured directly, which affected the accuracy. Finally, although some towns were in suburban areas, cities in rural zones were not represented.

Conclusion

While providing updated reference values for the 20-m shuttle run test, this study provides direct comparative evidence of an alarming decrease of CRF and functional capacity in a population of children and adolescents since the 1980s. This further highlights the threat of an epidemic of cardiometabolic pathologies in the near future. Thus, development of population surveillance tools and public health strategies to promote a physically active lifestyle is more important than ever.

Data availability statement

The raw data supporting the conclusions of this article will be made available from the corresponding author, without undue reservation.

Ethics statement

The studies involving human participants were reviewed and approved by the Université du Québec à Chicoutimi (CER). Written informed consent to participate in this study was provided by the participants' legal guardian/next of kin.

Author contributions

ML was involved in the design and concept of the study, data collection, data analysis, drafted the initial, and final version

of the manuscript. MA and EK co-ordinated, supervised data collection, involved in the study design, drafted the initial manuscript, reviewed, and revised the manuscript. AC, HB, and JL were involved in the study design, initial analyses, data collection, drafted the initial manuscript, reviewed, and revised the manuscript. LL and PF were involved in the data analysis, reviewed, and drafted the manuscript for important intellectual content. PL co-ordinated, supervised data collection, involved in initial data analysis, drafted the initial manuscript, reviewed, and revised the manuscript. SB-G was involved in the data collection, the initial data analysis, drafted the initial manuscript, reviewed, and revised the manuscript. All authors approved the final manuscript as submitted and agree to be accountable for all aspects of the work.

Funding

This study was financially supported by the Center for Interdisciplinary Research on Quality and Healthy Lifestyles – UQAC: Grant no. UBR 324029.

Acknowledgments

The authors wish to acknowledge the principals, physical education teachers, school boards, and students who agreed to participate in this study. Special thanks to the interns who also helped with the data collection.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

References

1. Health Canada (Government of Canada). *Childhood Obesity*. 2016. Available online at: <https://www.canada.ca/en/public-health/services/childhood-obesity/childhoodobesity.html> (accessed September 15, 2020).

2. Moraes Ferrari GL, Bracco MM, Matsudo VK, Fisberg M. Cardiorespiratory fitness and nutritional status of school children: 30-year evolution. *J Pediatr*. (2013) 89:366–73. doi: 10.1016/j.jpeds.2012.12.006

3. Hallal PC, Andersen LB, Bull FC, Guthold R, Haskell W, Ekelund U. Global physical activity levels: surveillance progress, pitfalls, and prospects. *Lancet*. (2012) 380:247–57. doi: 10.1016/S0140-6736(12)60646-1
4. Ozemek C, Laddu DR, Lavie CJ, Claeys H, Kaminski LA, Ross R, et al. An update on the role of cardiorespiratory fitness, Structured Exercise and lifestyle physical activity in preventing cardiovascular disease and health risk. *Prog Cardiovasc Dis*. (2018) 61:484–90. doi: 10.1016/j.pcad.2018.11.005
5. Sun C, Magnussen CG, Ponsosby AL, Schmidt MD, Carlin JB, Huynh Q, et al. The contribution of childhood cardiorespiratory fitness and adiposity to in?ammation in young adults. *Obesity*. (2014) 22:2598–605. doi: 10.1002/oby.20871
6. Tomkinson GR, Lang JJ, Tremblay MS, Dale M, LeBlanc AG, Belanger K, et al. International normative 20 m shuttle run values from 1,142,026 children and youth representing 50 countries. *Br J Sports Med*. (2017) 51:1545–54. doi: 10.1136/bjsports-2016-095987
7. Public Agency of Canada. *How Healthy are Canadians? A Trend Analysis of the Health of Canadians From a Healthy Living and Chronic Disease Perspective*. (2016). Available online at: <http://www.publications.gc.ca/site/eng/9.825964/publication.html> (accessed September 15, 2020).
8. Canadian Society of Exercise Physiology. *Canadian 24-Hour Movement Guidelines for Children and Youth: An Integration of Physical Activity, Sedentary Behaviour, and Sleep*. (2020). Available online at: https://csepguidelines.ca/wpcontent/themes/csep2017/pdf/Canadian24HourMovementGuidelines2016_2.pdf (accessed September 15, 2020).
9. Sui X, Sarzynski MA, Lee DC, Kokkinos PF. Impact of changes in cardiorespiratory fitness on hypertension, dyslipidemia and survival: an overview of the epidemiological evidence. *Prog Cardiovasc Dis*. (2017) 60:56–66. doi: 10.1016/j.pcad.2017.02.006
10. Höglström G, Nordström A, Nordström P. High aerobic fitness in late adolescence is associated with a reduced risk of myocardial infarction later in life: a nationwide cohort study in men. *Eur Heart J*. (2014) 35:3133–40. doi: 10.1093/eurheartj/ehf527
11. Lang JJ, Phillips EW, Hoffmann MD, Prince SA. Establishing modified Canadian Aerobic Fitness Test (mCAFT) cut-points to detect clustered cardiometabolic risk among Canadian children and youth aged 9 to 17 years. *Appl Physiol Nutr Metab*. (2020) 45:311–7. doi: 10.1139/apnm-2019-0303
12. Ross R, Blair SN, Arena R, Church TS, Després JP, Franklin Ba, et al. Importance of assessing cardiorespiratory fitness in clinical practice: a case for fitness as a clinical vital sign: a scientific statement from the American heart association. *Circulation*. (2016) 134:e653–99. doi: 10.1161/CIR.0000000000000461
13. García-Hermoso A, Ramírez-Vélez R, García-Alonso Y, Alonso-Martínez AM, Izquierdo M, Ramírez-Vélez R. Association of cardiorespiratory fitness levels during youth with health risk later in life: a systematic review and meta-analysis. *JAMA*. (2020) 174:952–60. doi: 10.1001/jamapediatrics.2020.2400
14. Raghuvveer G, Hartz J, Lubans DR, Takken T, Wiltz JL, Miettus-Snyder M, et al. Cardiorespiratory fitness in youth: An important marker of health: a scientific statement from the American heart association. *Circulation*. (2020) 142:e101–18. doi: 10.1161/CIR.0000000000000866
15. Ortega FB, Ruiz JR, Castillo MJ, Sjörström M. Physical fitness in childhood and adolescence: a powerful marker of health. *Int J Obes*. (2008) 32:1–11. doi: 10.1038/sj.ijo.0803774
16. Shephard RJ, Allen C, Benade AJ, Davies CT, Di Prampero PE, Hedman R et al. The maximum oxygen intake. An international reference standard of cardiorespiratory fitness. *Bull World Health Organ*. (1968) 38:757–64.
17. Taylor HL, Buskirk E, Henschel A. Maximal oxygen intake as an objective measure of cardio-respiratory performance. *J Appl Physiol*. (1955) 8:73–80. doi: 10.1152/jap.1955.8.1.73
18. Armstrong N, Welsman J. Twenty-metre shuttle run: (mis)representation, (mis)interpretation and (mis)use. *Br J Sports Med*. (2019) 53:1199. doi: 10.1136/bjsports-2018-100082
19. Armstrong N, Welsman J. Youth cardiorespiratory fitness: evidence, myths and misconceptions. *Bull World Health Organ*. (2019) 97:777–82. doi: 10.2471/BLT.18.227546
20. Tomkinson GR, Lang JJ, Blanchard J, Léger L, Tremblay MS. The 20-m shuttle run : assessment and interpretation of data in relation to youth aerobic fitness and health. *Pediatr Exerc Sci*. (2019) 31:152–63. doi: 10.1123/pes.2018-0179
21. Tomkinson GR, Lang JJ, Tremblay MS. Temporal trends in the cardiorespiratory fitness of children and adolescents representing 19 high-income and upper middle-income countries between 1981 and 2014. *Br J Sports Med*. (2019) 53:478–86. doi: 10.1136/bjsports-2017-097982
22. Olds T, Tomkinson G, Léger L, Cazorla G. Worldwide variation in the performance of children and adolescents: an analysis of 109 studies of the 20-m shuttle run test in 37 countries. *J Sports Sci*. (2006) 24:1025–38. doi: 10.1080/02640410500432193
23. Léger L, Lambert J, Goulet A. Aerobic capacity of 6 to 17-year-old Quebecois 20-meter shuttle run test with 1 minute stages. *Can J Appl Sport Sci*. (1984) 9:64–9.
24. Hoffmann MD, Colley RC, Doyon CY, Wong SL, Tomkinson GR, Lang JJ. Normative-referenced percentile values for physical fitness among Canadians. *Health Rep*. (2019) 30:14–22. doi: 10.25318/82-003-x201901000002-eng
25. Lohman TG, Roche AF, Martorell R. *Anthropometric Standardization Reference Manual*. Champaign, IL: Human Kinetics Books (1988).
26. Cole TJ, Bellizzi MC, Flegal KM, Dietz WH. Establishing a standard definition for child overweight and obesity worldwide: international survey. *BMJ*. (2000) 320:1240–3. doi: 10.1136/bmj.320.7244.1240
27. Box GEP, Cox DR. An analysis of transformations. *J R Stat Soc Series B Stat Methodol*. (1964) 26:211–52. doi: 10.1111/j.2517-6161.1964.tb00553.x
28. Hoaglin DC, Iglewicz B, Tukey JW. Performance of some resistant rules for outlier labeling. *J Am Stat Assoc*. (1986) 81:991–9. doi: 10.1080/01621459.1986.10478363
29. Hoaglin DC, Iglewicz B. Fine tuning some resistant rules for outlier labeling. *J Am Stat Assoc*. (1987) 82:1147–9. doi: 10.1080/01621459.1987.10478551
30. Cole TJ, Green PJ. Smoothing reference centile curves: the LMS method and penalized likelihood. *Stat Med*. (1992) 11:1305–19. doi: 10.1002/sim.4780111005
31. Tarp J, Fagerland MW, Dalene KE, Johannessen JS, Hansen BH, Jefferis BJ, et al. Device-measured physical activity, adiposity and mortality: a harmonised meta-analysis of eight prospective cohort studies. *Br J Sports Med*. (2021) 56:1–9. doi: 10.1136/bjsports-2021-104827
32. Ruiz JR, Cavero-Redondo I, Ortega FB, Welk GJ, Andersen LB, Martínez-Vizcaino V. Cardiorespiratory fitness cut points to avoid cardiovascular disease risk in children and adolescents; what level of fitness should raise a red flag? A systematic review and meta-analysis. *Br J Sports Med*. (2016) 50:1451–8. doi: 10.1136/bjsports-2015-095903
33. Ruiz JR, Huybrechts I, Cuenca-García M. Cardiorespiratory fitness and ideal cardiovascular health in European adolescents. *Heart*. (2015) 101:766–73. doi: 10.1136/heartjnl-2014-306750
34. Dobbins M, Husson H, DeCorby K, LaRocca RL. School-based physical activity programs for promoting physical activity and fitness in children and adolescents aged 6-18 (Review). *Cochrane Database Syst Rev*. (2013) 4:CD007651. doi: 10.1002/14651858.CD007651.pub2
35. Arango CM, Parra DC, Gómez LF, Lema L, Lobelo F, Ekelund U. Screen time, cardiorespiratory fitness and adiposity among school-age children from Montería, Colombia. *J Sci Med Sport*. (2014) 17:491–5. doi: 10.1016/j.jsams.2013.09.010
36. Sigmundová D, El Ansari W, Sigmund E, Frömel K. Secular trends: a ten-year comparison of the amount and type of physical activity and inactivity of random samples of adolescents in the Czech Republic. *BMC Public Health*. (2011) 11:731. doi: 10.1186/1471-2458-11-731
37. Ryu S, Kim H, Kang M, Pedisic Z, Loprinzi PD. Secular Trends in Sedentary Behavior Among High School Students in the United States, 2003 to 2015. *Am J Health Promot*. (2019) 33:1174–81. doi: 10.1177/0890117119854043
38. Guthold R, Stevens GA, Riley LM, Bull FC. Global trends in insufficient physical activity among adolescents: a pooled analysis of 298 population-based surveys with 1.6 million participants. *Lancet Child Adolesc Health*. (2020) 4:23–35. doi: 10.1016/S2352-4642(19)30323-2
39. López-Bueno R, Calatayud J, Andersen LL, Casana J, Ezzatvar Y, Casajus JA, et al. Cardiorespiratory fitness in adolescents before and after the COVID-19 confinement: a prospective cohort study. *Eur J Pediatr*. (2021) 180:2287–93. doi: 10.1007/s00431-021-04029-8
40. Kuczumarski RJ. *2000 CDC Growth Charts for the United States: methods and development (No. 246)*. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Health Statistics. (2002). Available online at: https://www.cdc.gov/nchs/data/series/sr_11/sr11_246.pdf (accessed November 9, 2022).