



OPEN ACCESS

EDITED AND REVIEWED BY

Shuchao Wang,
Chinese Academy of Agricultural
Sciences, China

*CORRESPONDENCE

Oyelola Adegboye
✉ oyelola.adegboye@jcu.edu.au

RECEIVED 10 July 2023

ACCEPTED 13 July 2023

PUBLISHED 24 July 2023

CITATION

Stephen R, Alele F, Olumoh J, Tyndall J,
Okeke MI and Adegboye O (2023)
Corrigendum: The epidemiological trend of
monkeypox and monkeypox-varicella zoster
viruses co-infection in North-Eastern Nigeria.
Front. Public Health 11:1256195.
doi: 10.3389/fpubh.2023.1256195

COPYRIGHT

© 2023 Stephen, Alele, Olumoh, Tyndall, Okeke
and Adegboye. This is an open-access article
distributed under the terms of the [Creative
Commons Attribution License \(CC BY\)](#). The use,
distribution or reproduction in other forums is
permitted, provided the original author(s) and
the copyright owner(s) are credited and that
the original publication in this journal is cited, in
accordance with accepted academic practice.
No use, distribution or reproduction is
permitted which does not comply with these
terms.

Corrigendum: The epidemiological trend of monkeypox and monkeypox-varicella zoster viruses co-infection in North-Eastern Nigeria

Roland Stephen¹, Faith Alele^{2,3}, Jamiu Olumoh ⁴,
Jennifer Tyndall⁵, Malachy Ifeanyi Okeke⁵ and
Oyelola Adegboye ^{2,3,6*}

¹Department of Internal Medicine, Modibbo Adama University Teaching Hospital, Yola, Adamawa, Nigeria, ²Public Health and Tropical Medicine, College of Public Health, Medical and Veterinary Sciences, James Cook University, Townsville, QLD, Australia, ³Australian Institute of Tropical Health and Medicine, James Cook University, Townsville, QLD, Australia, ⁴Department of Mathematics and Statistics, American University of Nigeria, Yola, Adamawa, Nigeria, ⁵Department of Natural and Environmental Sciences, American University of Nigeria, Yola, Adamawa, Nigeria, ⁶World Health Organization Collaborating Center for Vector-Borne and Neglected Tropical Diseases, College of Public Health, Medical and Veterinary Sciences, James Cook University, Townsville, QLD, Australia

KEYWORDS

monkeypox virus, MPX, varicella zoster virus, Nigeria, coinfection

A corrigendum on

[The epidemiological trend of monkeypox and monkeypox-varicella zoster viruses co-infection in North-Eastern Nigeria](#)

by Stephen, R., Alele, F., Olumoh, J., Tyndall, J., Okeke, M. I., and Adegboye, O. (2022). *Front. Public Health* 10:1066589. doi: 10.3389/fpubh.2022.1066589

In the published article, there was an error in [Table 1](#) as published. There were errors regarding the number of patients with negative/inconclusive results by age group and gender. The corrected [Table 1](#) and its caption appear below.

In the published article, there was an error in the second line of the first paragraph of the results section.

A correction has been made to the second line of the first paragraph of the results. This sentence previously stated:

“Males accounted for 79% (28). . .”

The corrected sentence appears below:

“Males accounted for 79% (26). . .”

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated

organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

TABLE 1 Patient demographic and clinical characteristics in all case series.

Characteristics	Total	All MPX	MPX only	All VZ	VZ only	Co-infection	Negative/inconclusive
<i>n</i> (<i>N</i>)	33	11/33	2/33 (6.1%)	22/33 (67%)	13/33 (39%)	9/33 (27%)	9/33 (27%)
Age, median (IQR)	14 (8–30)	14 (10–28)	14 (8–20)	13 (10–28)	12 (7–22)	14 (11–29)	30 (17–36)
Age group, <i>n</i> (%)							
0–9	8 (24)	2 (18)	1 (50)	5 (23)	4 (31)	1 (11)	2 (22.2)
10–19	10 (30)	4 (36)	0 (0)	9 (41)	5 (38)	4 (44)	1 (11.1)
20–29	4 (12)	3 (27)	1 (50)	3 (14)	1 (7.7)	2 (22)	0 (0)
30–39	7 (21)	2 (18)	0 (0)	3 (14)	1 (7.7)	2 (22)	4 (44.4)
40+	2 (6.1)	0 (0)	0 (0)	2 (9.1)	2 (15)	0 (0)	0 (0)
Missing	2 (6.1)						2 (22.2)
Sex, <i>n</i> (%)							
Male	26 (79)	9 (82)	2 (100)	16 (73)	9 (69)	7 (78)	8 (88.9)
Female	7 (21)	2 (18)	0 (0)	6 (27)	4 (31)	2 (22)	1 (11.1)
Presence crust, <i>n</i> (%)	8 (24.2)	NA	NA	NA	7 (53.9)	1 (11.1)	NA
Days, † [mean (SD)]	2.9 (4.1)	NA	NA	NA	5.5 (5.7)	1.2 (1.2)	NA

† Average days between the onset of fever and rashes.

NA, not available.