



## OPEN ACCESS

## EDITED BY

Sabrina Skorski,  
Saarland University, Germany

## REVIEWED BY

Danilo Reis Coimbra,  
Juiz de Fora Federal University, Brazil  
Emma Sweeney,  
Nottingham Trent University, United Kingdom

## \*CORRESPONDENCE

Craig Thomas  
✉ C.Thomas@lboro.ac.uk

<sup>†</sup>These authors share first authorship

## SPECIALTY SECTION

This article was submitted to Elite Sports and Performance Enhancement, a section of the journal Frontiers in Sports and Active Living

RECEIVED 29 June 2022

ACCEPTED 13 December 2022

PUBLISHED 06 January 2023

## CITATION

Thomas C, Langan-Evans C, Germaine M, Artukovic M, Jones H, Whitworth-Turner C, Close GL and Louis J (2023) Case Report: Effect of low energy availability and training load on sleep in a male combat sport athlete. *Front. Sports Act. Living* 4:981755. doi: 10.3389/fspor.2022.981755

## COPYRIGHT

© 2023 Thomas, Langan-Evans, Germaine, Artukovic, Jones, Whitworth-Turner, Close and Louis. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

# Case Report: Effect of low energy availability and training load on sleep in a male combat sport athlete

Craig Thomas<sup>1,2\*†</sup>, Carl Langan-Evans<sup>1†</sup>, Mark Germaine<sup>1,3</sup>, Mario Artukovic<sup>1</sup>, Helen Jones<sup>1</sup>, Craig Whitworth-Turner<sup>4</sup>, Graeme L. Close<sup>1</sup> and Julien Louis<sup>1</sup>

<sup>1</sup>Research Institute for Sport and Exercise Sciences (RISES), Liverpool John Moores University, Liverpool, United Kingdom, <sup>2</sup>School of Sport, Exercise and Health Sciences (SSEHS), Loughborough University, Loughborough, United Kingdom, <sup>3</sup>SFI Centre for Research Training in Machine Learning, Dublin City University, Dublin, Ireland, <sup>4</sup>Department of Athletics, University of Pennsylvania, Philadelphia, PA, United States

**Purpose:** The aim of this case report was to describe the sleep responses in a male combat sport athlete, who was engaging in both chronic (CWL) and acute (AWL) weight loss practices in order to reduce body mass for a national competition.

**Methods:** During the first seven weeks of training (Phases 1 and 2), the athlete adhered to a daily energy intake (EI) equating to their resting metabolic rate (1700 kcal·day<sup>-1</sup>) followed by a reduction in EI (915–300 kcal·day<sup>-1</sup>) in the 5 days before weighing in (Phase 3). Nocturnal sleep was monitored throughout the 8-week training period using wristwatch actigraphy and frequent measurements of body mass/composition, daily exercise energy expenditure and training load (TL) were taken.

**Results:** The athlete was in a state of low energy availability (LEA) during the entire training period. There was a very large decrease in LEA status during phase 3 compared with phases 1 and 2 (3 vs. 20 kcal·kgFFM·day<sup>-1</sup>) and there was a small decrease in TL during phase 3 compared with phase 2 (410 vs. 523 AU). The athlete's sleep efficiency increased throughout the training period, but total sleep time displayed a small to moderate decrease in phase 3 compared with phases 1 and 2 (386 vs. 429 and 430 min). However, correlational analysis demonstrated trivial to small, non-significant relationships between sleep characteristics and the athlete's LEA status and TL.

**Conclusion:** These findings suggest that CWL and AWL practices that cause fluctuations in LEA and TL may be implemented without compromising the sleep of combat sport athletes.

## KEYWORDS

weight making, energy restriction, energy expenditure, training load, actigraphy

## 1. Introduction

Professional and amateur combat sport disciplines are organised into weight categories to promote fair competition between opponents of equal body mass (BM) (1). Despite this, combat sport athletes often aim to compete in the lowest weight category possible *via* a process known as *weight making*, in an attempt to gain competitive advantages over opponents of smaller proportion (2). To that end, there

is a widespread culture of reducing BM within combat sports, with athletes across a range of disciplines typically losing 4–10 kg leading up to competition (3). These losses are achieved by engaging in both acute (AWL—over hours or days) and chronic (CWL—over several weeks or months) weight loss practices, which include a combination of increasing exercise energy expenditure (EEE), restricting energy intake (EI) and techniques to induce specific states of dehydration (4). The AWL approach, when not well considered, has been associated with a number of health risks, as a rapid reduction in BM may affect fluid balance, thermoregulation, and metabolism, as well as cause injury or in extreme cases even death (4). When the goal is to reduce BM gradually, combat sport athletes may utilise a CWL approach by inducing an energy deficit resulting in losses of 0.5–1 kg·week<sup>-1</sup> (1, 5, 6). However, a reduction in EI with concomitant increases in EEE *via* escalations in overall daily training load (TL), is likely to result in a state of low energy availability (LEA), which has been associated with a number of psycho-physiological consequences as characterised within the Athlete Triad (TRIAD) and Relative Energy Deficiency in Sport (RED-S) models (7–9).

Sleep is a key factor in health and wellbeing, which influences a number of psycho-physiological systems (10). Varying sleep parameters have been correlated with a range of performance markers when assessed in an athletic combat sport population (11). To maintain optimal health and daytime functioning, it is recommended adults achieve at least 420 min of total sleep time with ≥85% sleep efficiency per night (12, 13). As previously described, an increase in daily TL is an inherent consequence of adopting a CWL approach. Proliferated training periods may result in greater sleep duration, sleep efficiency and slow wave sleep percentage (14, 15), whilst others have demonstrated either no change or reductions in sleep duration and efficiency (16–18). It has been suggested that sleep may improve with subtle changes in TL but become disturbed with large increases relative to the athlete (19). Whilst the TL of combat sport athletes may not be as high compared to athletes from other sports (20), due to BM elevations induced partly from inactivity (5, 21), these athletes are often exposed to a large increase in TL when making weight. This is supported by data in combat sport athletes, where sleep was rated worse during early preparation training compared with mid preparation training and on training days compared with rest days, indicating sleep may be challenged when commencing training after a rest period (22, 23). Thus, it is plausible that an inflation in TL associated with weight making may negatively impact sleep and subsequently the training and competition readiness of combat sport athletes, concomitant with LEA.

Despite the potential impact of increased TL and LEA, sleep is not considered within the aforementioned TRIAD and RED-S models, yet this may be of concern given a lower sleep duration

may affect appetite, mood, and performance (24–26). Indeed, in male and female dancers who were screened for LEA using the Dance Specific Energy Availability Questionnaire, among those with sleep problems, 37% reported difficulty falling asleep and 21% indicated disrupted sleep patterns (27). Previously, when compared to high level athletes from non-weight making sports (e.g., racquet sports), a greater number of combat sport athletes perceived themselves to have had a sleep problem at some point in their lifetime (28). Furthermore, AWL practices may induce sleep disturbances in combat sport athletes leading into competition, yet current evidence has suggested this may not be the case when utilising techniques such as low fibre/residue diets and water loading across time periods of up to 5 days (29). Such findings suggest that sport specific factors are responsible for sleep differences between sporting demographics and in the case of combat sport athletes, associated periods of LEA may potentially result in sleep disturbances.

Therefore, the aim of this case report was to describe the sleep responses in a male combat sport athlete, who was engaging in both CWL and AWL practices in order to make weight for a national competition. During an 8-week training period the athlete's sleeping patterns were monitored using wristwatch actigraphy and frequent measurements of BM/ composition, daily EEE and TL were assessed, in tandem with controlled restriction of daily EI. This analysis was part of a larger case report, which evaluated the effects of LEA on health and performance related indices associated with the Male TRIAD and RED-S models (5).

## 2. Materials and methods

### 2.1. Athlete overview

The athlete (age: 19 years; stature: 166 cm; BM: 72.5 kg) was a Caucasian male taekwondo competitor with 5 years international competition experience, who would habitually use CWL and AWL practices over time periods of 3–4 weeks to lose 4–5 kg, so they could compete in the <68 kg feather weight category. For this case report, the athlete aimed to compete in the <63 kg bantam weight category at the British University Championships, therefore requiring a reduction of >9.5 kg BM across an 8-week training period (5). Prior to commencement of the training period, the athlete did not use any sleep aids (e.g., medication) or work night shifts, nor had they travelled across different time zones. Additionally, throughout the training period they worked part-time, did not partake in any daytime napping, disclosed that they understood the importance of sleep for optimising performance and wanted to gain an insight into their sleeping patterns. The athlete provided written informed consent and institutional ethical approval was granted by Liverpool John Moores University.

## 2.2. Athlete assessment

### 2.2.1. Measures of energy availability status, daily training load and TRIAD/RED-S consequences

BM was measured each morning, post void of the bladder/bowels and determined to the nearest 0.01 kg on a calibrated digital scale, with measures of stature established to the nearest 0.1 cm using a free-standing stadiometer (Seca 702 & 123; Seca GmbH, Hamburg, Germany). To examine energy availability (EA) status, measures of body composition inclusive of bone mineral content/density, absolute and relative fat mass, and fat free mass (FFM) were assessed using Dual-energy x-ray Absorptiometry (QDR Series Discovery A, software version 12:4:3, Hologic Inc, Bedford, MA, USA), as per best practice guidelines (30). Gross EEE from all training sessions was measured *via* a combined heart rate and actigraphy unit (Actiheart 4, CamNtech Ltd, Cambridgeshire, United Kingdom) and calculated into net EEE as previously described (5). EI was provided for the athlete throughout the entire 8-week training period as highlighted in the following sections. Daily EA was subsequently calculated using the equation: EI minus net EEE divided by FFM with values  $<30 \text{ kcal}\cdot\text{kgFFM}\cdot\text{day}^{-1}$  classified as LEA (31). Daily TL was assessed using the session rating of perceived exertion (s-RPE) method, based on Borg's Category Ratio 10 scale (1–10 arbitrary units) (32). Further details pertaining to the measurement of TRIAD/RED-S health and performance consequences inclusive of resting metabolic rate, cardiac function, maximal dynamic strength and power, cardiorespiratory capacity, psychological state, and blood clinical chemistry (for biomarkers related to endocrine, bone turnover, hydration, electrolyte, renal, liver, and lipid profiles) are highlighted in Langan-Evans et al. (5).

### 2.2.2. Wristwatch actigraphy

To assess nocturnal sleep throughout the 8-week training period, the athlete was provided with a wrist activity monitor (Actiwatch 4, Cambridge Neurotechnology Ltd, Cambridgeshire, United Kingdom) for the first four weeks and then again for the last four weeks, which was set to an epoch length of 1 min at a medium sensitivity. Actigraphy has shown to have a good overall agreement with polysomnography (90%–91%; gold standard measurement of sleep) and is a valid alternative for sleep/wake estimation (33). On each night, the athlete was instructed to wear the activity monitor on their non-dominant wrist 30–60 min before they retired to bed in their home environment. Upon their bedtime, they recorded the clock time in the Core Consensus Sleep Diary (CSD) (34) and pressed the marker button on the activity monitor. The following morning, they pressed the marker button again upon their final awakening and filled in the remainder of the Core CSD. During the monitoring

period, the athlete received no sleep hygiene education. The recorded markers from the activity monitor and the information from the Core CSD were then used to determine bedtime, sleep onset, final awakening time and get up time, so that sleep behaviour could be automatically calculated using the appropriate software (Actiwatch Activity and Sleep Analysis version 5.24, Cambridge Neurotechnology Ltd, Cambridgeshire, United Kingdom). From the analysis, the following sleep characteristics were used: bedtime (BT—h:min), final awakening (FA—h:min), time in bed (TIB—min), sleep onset latency (SOL—minutes between bedtime and sleep onset), total sleep time (TST—min), wake after sleep onset (WASO—min) and sleep efficiency (SE—total sleep time as a percentage of time in bed (sleep quality)).

## 2.3. Dietary and training periodisation

### 2.3.1. Phase 1 (–8 week to –4 week) & phase 2 (–4 week to –1 week)

During the first seven weeks of the training period, the athlete ingested a daily EI that equated to their resting metabolic rate measurement at baseline ( $1700 \text{ kcal}\cdot\text{day}^{-1}$ ) (See **Table 1**), consisting of pre-packaged meals totalling  $3.4 \text{ g}\cdot\text{kgFFM}^{-1}$  carbohydrate (CHO) ( $748 \text{ kcal}\cdot\text{day}^{-1}$ ),  $2.3 \text{ g}\cdot\text{kgFFM}^{-1}$  protein ( $506 \text{ kcal}\cdot\text{day}^{-1}$ ) and  $0.9 \text{ g}\cdot\text{kgFFM}^{-1}$  fat ( $446 \text{ kcal}\cdot\text{day}^{-1}$ ) (5). Alongside restriction in EI, the athlete completed  $12\text{--}15 \text{ h}\cdot\text{week}^{-1}$  of set periodised training across a weekly schedule, consisting of three aerobic, two high intensity interval, two resistance and three taekwondo specific sessions with one sparring-based session. All strength and conditioning sessions were prescribed and delivered by an accredited strength and conditioning coach, with over 10 years' experience. Sport specific sessions were prescribed and delivered by an Olympic licensed coach with over 12 years' experience. During Saturdays no training occurred, and these were classified as recovery periods [see **Figure 1** for schedule and Langan-Evans et al. (5) for further detail on exercise prescription].

### 2.3.2. Phase 3 (5 days prior to weigh in)


In the 5 days prior to the weigh in daily EI was gradually reduced (see **Table 1**), whereby the athlete consumed  $0.5 \text{ g}\cdot\text{kgFFM}^{-1}$  CHO ( $110 \text{ kcal}\cdot\text{day}^{-1}$ ),  $1.8 \text{ g}\cdot\text{kgFFM}^{-1}$  protein ( $385 \text{ kcal}\cdot\text{day}^{-1}$ ) and  $0.9 \text{ g}\cdot\text{kgFFM}^{-1}$  fat ( $420 \text{ kcal}\cdot\text{day}^{-1}$ ) on days –5 and –4, followed by  $0.3 \text{ g}\cdot\text{kgFFM}^{-1}$  CHO ( $67 \text{ kcal}\cdot\text{day}^{-1}$ ),  $1.8 \text{ g}\cdot\text{kgFFM}^{-1}$  protein ( $400 \text{ kcal}\cdot\text{day}^{-1}$ ) and  $0.8 \text{ g}\cdot\text{kgFFM}^{-1}$  fat ( $396 \text{ kcal}\cdot\text{day}^{-1}$ ) on days –3 and –2 (5). On –1 day, the athlete had one meal in the morning, which contained  $1.4 \text{ g}\cdot\text{kgFFM}^{-1}$  protein ( $300 \text{ kcal}\cdot\text{day}^{-1}$ ). This reduction in EI corresponded with a taper to training volume, therefore clamping EEE (see **Table 1**). From 12:00 h on –1 day until 13:00 h at the official competition weigh in the following day, the athlete then abstained from all food and

**TABLE 1** Measurements of energy availability, intake, expenditure, training load and sleep throughout the training period.

Time phase/ characteristic	Monday	Tuesday	Wednesday	Thursday	Friday	Saturday	Sunday	Combined Average	Effect Size
<b>PHASE 1 (−8 WK TO −4 WK)</b>									
EA (kcal·kgFFM·day <sup>−1</sup> )	15 ± 1	20 ± 1	24 ± 0	20 ± 1	9 ± 1	31 ± 0	24 ± 0	20 ± 7	t: VL
EI (kcal·day <sup>−1</sup> )	1700 ± 0	1700 ± 0	1700 ± 0	1700 ± 0	1700 ± 0	1700 ± 0	1700 ± 0	1700 ± 0	t: VL
EEE (kcal·day <sup>−1</sup> )	873 ± 47	584 ± 31	409 ± 22	617 ± 34	1209 ± 66	0 ± 0	376 ± 20	559 ± 370	t: T
TL s-RPE (AU)	728 ± 388	481 ± 212	439 ± 229	304 ± 250	1140 ± 268	0 ± 0	191 ± 21	469 ± 375	t: T
BT (h: min)	23:53 ± 00:17	23:45 ± 00:11	23:25 ± 00:35	23:43 ± 00:12	00:02 ± 00:17	23:34 ± 00:51	23:52 ± 00:07	23:44 ± 00:12	
FA (h: min)	07:22 ± 00:54	06:23 ± 00:42	06:56 ± 01:45	07:23 ± 00:44	08:23 ± 01:27	08:47 ± 02:49	07:54 ± 01:35	07:35 ± 00:49	
TIB (min)	458 ± 56	404 ± 36	472 ± 151	465 ± 38	517 ± 88	558 ± 131	507 ± 128	483 ± 49	s: M
TST (min)	417 ± 30	361 ± 52	418 ± 97	406 ± 44	465 ± 74	490 ± 155	448 ± 92	429 ± 42	t: M
SE (%)	91.6 ± 5.1	89.1 ± 6.7	90.1 ± 8.7	87.1 ± 3.1	90.3 ± 5.3	86.4 ± 10.2	89.0 ± 6.9	89.1 ± 1.8	l: VL
WASO (min)	26 ± 22	28 ± 20	20 ± 16	46 ± 7	23 ± 24	42 ± 34	29 ± 21	31 ± 10	l: VL
SOL (min)	6 ± 2	9 ± 4	13 ± 11	9 ± 9	13 ± 4	22 ± 14	5 ± 4	11 ± 6	t: M
<b>PHASE 2 (−4 WK TO −1 WK)</b>									
EA (kcal·kgFFM·day <sup>−1</sup> )	13 ± 1	22 ± 1	25 ± 0	20 ± 1	7 ± 1	31 ± 0	23 ± 0	20 ± 8	VL
EI (kcal·day <sup>−1</sup> )	1700 ± 0	1700 ± 0	1700 ± 0	1700 ± 0	1700 ± 0	1700 ± 0	1700 ± 0	1700 ± 0	VL
EEE (kcal·day <sup>−1</sup> )	968 ± 52	527 ± 29	340 ± 18	616 ± 34	1333 ± 72	0 ± 0	430 ± 23	579 ± 418	T
TL s-RPE (AU)	710 ± 615	507 ± 78	357 ± 6	600 ± 137	1270 ± 139	0 ± 0	220 ± 9	523 ± 407	S
BT (h: min)	23:38 ± 00:31	23:47 ± 00:06	23:50 ± 00:09	23:57 ± 00:07	00:01 ± 00:06	00:30 ± 00:30	00:10 ± 00:17	23:59 ± 00:17	
FA (h: min)	05:40 ± 00:04	06:16 ± 01:00	05:42 ± 00:15	08:11 ± 00:14	08:18 ± 01:45	10:29 ± 01:06	08:09 ± 01:11	07:32 ± 01:45	
TIB (min)	372 ± 32	399 ± 74	362 ± 16	508 ± 23	510 ± 108	629 ± 97	481 ± 57	466 ± 95	S
TST (min)	340 ± 27	373 ± 64	342 ± 12	462 ± 11	473 ± 105	558 ± 90	459 ± 61	430 ± 81	S
SE (%)	91.4 ± 2.3	93.7 ± 3.5	94.5 ± 1.6	91.0 ± 3.8	92.6 ± 2.1	88.8 ± 5.6	95.3 ± 3.2	92.5 ± 2.3	M
WASO (min)	13 ± 4	5 ± 6	5 ± 8	20 ± 2	16 ± 6	23 ± 9	7 ± 6	13 ± 7	S
SOL (min)	10 ± 5	10 ± 7	5 ± 1	12 ± 7	8 ± 4	17 ± 16	13 ± 9	11 ± 4	M
<b>PHASE 3 (5 DAYS PRIOR TO OCWI)</b>									
	−5 D	−4 D	−3 D	−2 D	−1 D	OCWI	COMP		
EA (kcal·kgFFM·day <sup>−1</sup> )	−4	9	8	7	−3	−	−	3 ± 6	
EI (kcal·day <sup>−1</sup> )	915	915	863	863	300	−	−	771 ± 265	
EEE (kcal·day <sup>−1</sup> )	1120	409	421	467	474	−	−	578 ± 304	
TL s-RPE (AU)	1220	180	160	160	330	−	−	410 ± 458	
BT (h: min)	00:04	00:04	23:56	23:51	00:09	−	−	00:00 ± 00:07	
FA (h: min)	05:40	06:40	05:36	06:44	08:54	−	−	06:42 ± 01:20	
TIB (min)	340	401	342	434	540	−	−	411 ± 82	
TST (min)	313	373	328	408	507	−	−	386 ± 77	
SE (%)	92.1	93.0	95.9	94.0	93.9	−	−	93.8 ± 1.4	
WASO (min)	13	11	6	1	13	−	−	9 ± 5	
SOL (min)	10	12	6	4	5	−	−	7 ± 3	

EA, energy availability; EI, energy intake; EEE, exercise energy expenditure; TL s-RPE, training load session rating of perceived exertion; BT, bedtime; FA, final awakening; TIB, time in bed; TST, total sleep time; SE, sleep efficiency; WASO, wake after sleep onset; SOL, sleep onset latency; WK, week; D, day; OCWI, official competition weigh in; COMP, competition day.

t, trivial; s, small; l, large effect size compared to Phase 2 (−4 WK to −1 WK); T, trivial; S, small; M, moderate; VL, very large effect size compared to Phase 3 (5 days prior to OCWI).



fluids, and participated in self-directed AWL methods as further described in Langan-Evans et al. (5).

## 2.4. Statistical analysis

All phase 1 and 2 data highlighted in **Table 1** are presented as mean ± SD and phase 3 are individual values. To compare between phases, quantifications of daily EA, EI, EEE and TL s-RPE alongside sleep measures of TIB, TST, SE, WASO and SOL were assessed *via* Hedges *g* effect size calculations utilising the following quantitative criteria to explain the practical significance of the findings: trivial <0.2, small 0.2–0.59, moderate 0.6–1.19, large 1.20–1.99, and very large ≥2.0 (35). Given the non-normal distribution of data, sleep measurement variables of TIB, TST, SE, WASO and SOL were also examined in comparison to both daily EA and TL s-RPE *via* a non-parametric Spearman's rank correlation coefficient employing the following criteria to explain the relationship of association: trivial <0.1, small 0.1–0.29, moderate 0.3–0.49, large 0.5–0.69, very large 0.7–0.89 and almost perfect 0.9–1.00 (35). All analyses were performed using Prism version 9.2.0 (GraphPad, San Diego, California, USA) and the alpha level was set at  $p < 0.05$ .

## 3. Results

As highlighted in **Table 1**, the athlete was classified as being in a state of LEA during the entire training period, with no difference (0 kcal·kgFFM·day<sup>-1</sup>; ES = 0.00) during CWL across phases 1 and 2. LEA was further augmented during AWL within phase 3, resulting in very large differences (–17 kcal·kgFFM·day<sup>-1</sup>) compared to both phase 1 (ES = 2.37) and 2 (ES = 2.16). This pattern of LEA was dictated predominantly by changes in EI, which remained unchanged between phases 1 and 2 (0 kcal·day<sup>-1</sup>; ES = 0.00) and was followed by a very large

reduction (–929 kcal·day<sup>-1</sup>; ES = 5.12) during phase 3. To that end, modulations in LEA status were independent of changes in EEE, with minimal variances between 20 to –1 kcal·day<sup>-1</sup> across all three phases, resulting in trivial differences (ES ≤ 0.05). Finally, measures of TL s-RPE also mirrored EEE, with trivial differences between phases 1 and 2 (54 AU; ES = 0.12), 1 and 3 (–59 AU; ES = 0.13) and a small difference between phases 2 and 3 (–113 AU; ES = 0.24).


Also highlighted in **Table 1**, the athlete's TIB gradually reduced across the training period, resulting in a small difference between phases 1 and 2 (–17 min; ES = 0.20), a small difference between phases 2 and 3 (–55 min; ES = 0.56) and a moderate difference between phases 1 and 3 (–72 min; ES = 1.03). Independent of reductions in TIB, TST between phases 1 and 2 remained similar resulting in a trivial effect (1 min; ES = 0.01). However, these differences begin to increase by both small and moderate effects when comparing reductions in TST between phases 2 and 3 (–44 min; ES = 0.51) and 1 and 3 (–43 min; ES = 0.68), respectively. In contrast to TIB and TST, there is a subsequent improvement in SE through each phase, with a large difference between phases 1 and 2 (3.4%; ES = 1.43), very large difference between phases 1 and 3 (4.7%; ES = 2.63) and a moderate difference between phases 2 and 3 (1.3%; ES = 0.60). Measures of WASO also follow a similar pattern to improvements in SE, with continuing reductive improvements of a large effect between phases 1 and 2 (–18 min; ES = 1.81), a very large effect (–22 min; ES = 2.43) between phases 1 and 3 and a small effect (–4 min; ES = 0.59) between phases 2 and 3. Finally, SOL remained stable between phases 1 and 2 (0 min; ES = 0.00), with a decreasing moderate difference between phases 1 and 3 (–4 min; ES = 0.74) and a moderate difference between phases 2 and 3 (–4 min; ES = 1.02), respectively.


**Figure 2** highlights the association between the differing sleep measures and daily EA status, demonstrating small, non-significant, positive relationships with TIB (2A.  $r = 0.23$ ;  $p = 0.10$ ), TST (2B.  $r = 0.19$ ;  $p = 0.18$ ), WASO (2C.  $r = 0.21$ ;

$p=0.13$ ), SOL (2D.  $r = 0.12$ ;  $p=0.40$ ) and a negative relationship with SE (2E.  $r = -0.19$ ;  $p=0.16$ ). Finally, **Figure 3** also showcases the same analysis with a focus on sleep measures and daily TL, yet again highlighting trivial to small, non-significant, negative relationships with TIB (3A.  $r = -0.11$ ;  $p = 0.44$ ), TST (3B.  $r = -0.10$ ;  $p = 0.46$ ), WASO (3C.  $r = -0.01$ ;  $p = 0.94$ ), SOL (3D.  $r = -0.08$ ;  $p = 0.58$ ) and a positive relationship with SE (3E.  $r = 0.01$ ;  $p = 0.92$ ).

### 4. Discussion

The aim of this case report was to describe the sleep responses in a male combat sport athlete, who was engaging in both CWL and AWL practices in order to make weight for a national competition. During the first seven weeks of training, the athlete’s TST and SE were, on average, in line with current sleep recommendations (12, 13). When





comparing between the first two phases, TST was similar, but the athlete did display an increase in SE during phase 2, which was a result of a small reduction in TIB coupled with a large reduction in WASO. The TST and SE values from phases 1 and 2 are comparable to non-athletes [431 min and 88.7%; Leeder et al. (36)] and greater than that of other

individual sport athletes monitored during a typical out of competition training phase [390 min and 85.9%; Lastella et al. (37)]. In phase 3, where the athlete participated in AWL practices, a further reduction in TIB led to a moderate to very large increase in SE compared with the first two phases, however, TST decreased meaning the athlete fell short of the

recommended 420 min per night (12). The TST achieved by the athlete in phase 3 is similar to other combat sport athletes who utilised low fibre/residue diets and water loading across 5-days (386 vs. 373 min); Dunican et al. (29), whilst the SE of the athlete was remarkably higher than those same combat sport athletes (93.8 vs. 79%); Dunican et al. (29). Our data suggests that combat sport athletes may achieve adequate sleep quality during the process of making weight, but that their sleep duration may be suboptimal in the days prior to competition due to changes in daily schedule.

This is the first report to assess nocturnal sleep in relation to daily EA and TL in a combat sport athlete using an objective sleep measurement. As expected, during the first seven weeks of the training period, the athlete was in a state of LEA ( $20 \text{ kcal}\cdot\text{kgFFM}\cdot\text{day}^{-1}$ ). During the AWL in phase 3, there was a further reduction in EA ( $3 \text{ kcal}\cdot\text{kgFFM}\cdot\text{day}^{-1}$ ) compared with the first two phases, due to a very large reduction in daily EI rather than changes in EEE. Despite this, correlational analysis demonstrated small, non-significant, relationships between the athlete's LEA status and sleep characteristics, suggesting that LEA may not be strongly related to the sleep of the athlete in this case report. This opposes the idea that disrupted nocturnal sleep may be used as an indicator of LEA status (27). With there being an inherent increase in TL from a CWL approach, this case report also assessed whether TL may impact sleep, independent of LEA. There was a small reduction in daily TL during phase 3 compared with phase 2, however in a similar manner to LEA, there were trivial to small, non-significant, relationships between this metric and the sleep characteristics of the athlete. This concurs with Caia et al. (16) where a higher daily TL did not result in changes to sleep but contrasts with Hausswirth et al. (17) in which sleep was disturbed in overreached triathletes that undertook longer training sessions. To that end it could be postulated that the athlete's TL in this case report was not large enough to impact their sleep (19). Collectively, our findings indicate that practitioners may implement CWL and AWL practices that induce fluctuations in LEA and TL without compromising the sleep of combat sport athletes.

Our case report also highlights that the athlete altered their sleep schedule during the AWL phase, which likely explains the reduction in TST in the days before competition. In comparison to the first two phases, there was a small to moderate reduction in TIB during phase 3 because of an earlier FA time. Anecdotally, the athlete stated that they had taken on more work-based commitments (due to less scheduled training sessions) that required an earlier time of FA. This is in broad agreement with studies that have shown that early training start times may constrain the time of FA and subsequently impact sleep duration (38, 39). In the case of the athlete in this report, the replacement of training with more work hours can be attributed to them not being at the highest level in their sport, nonetheless, this highlights the need for practitioners to be aware of an athlete's schedule outside of

training. Given the importance of adequate sleep for physiological and psychological restoration (10), it is unknown whether the reduction in TST in the days before competition affected the athlete's competitive readiness. Although it should be noted that the athlete won a total of 4 contests to secure a gold medal placing at the championships. Therefore, more research is warranted to understand the individual sleep need of combat sport athletes and the minimum amount of sleep loss that may result in performance decrements.

## 5. Conclusions & practical implications

In conclusion, sleep duration decreased whilst sleep quality increased across the 8-week training period, though neither daily EA nor TL strongly correlated with overall sleep characteristics, suggesting practitioners may utilise CWL and AWL strategies without compromising sleep. TST fell below sleep guidelines during phase 3 and it is likely this was a product of an earlier FA imposed by the athlete's work schedule; thus, we recommend practitioners are aware of their athlete's schedule outside of training during the competition week in order to optimise the athlete's readiness for competition. The strength of this case report is that daily EI was controlled, and an objective measure of sleep was used, but there are some limitations. Our results reflect only one individual combat sport athlete, and it is possible that LEA and/or TL may impact sleep in other athletes. Nocturnal sleep was monitored across an 8-week training period, however, as sleep was not monitored prior to this, it is unknown whether the athlete's sleep had been further disrupted in comparison to their habitual sleeping patterns. Furthermore, it would have been beneficial to have monitored sleep at certain time points using polysomnography, given this is the gold standard measurement for sleep. Future research should examine the effect of daily EA and TL on sleep in larger combat sport athlete cohorts and across subsequent weight making cycles.

## Data availability statement

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation.

## Ethics statement

This case report involving a human participant was reviewed and approved by Liverpool John Moores University. The participant provided written informed consent to participate in this study.

## Author contributions

CT, CLE, MG, MA, HJ, CWT, GLC and JL were involved in the design of the case report. CT, CLE, MG, and MA were completed data collection and analysis. CT and CLE interpreted the data and wrote the initial manuscript. All authors contributed to the article and approved the submitted version.

## Acknowledgements

The authors would like to thank the athlete for their participation and giving permission to publish this case report.

## References

- Langan-Evans C, Close GL, Morton JP. Making weight in combat sports. *Strength Cond J*. (2011) 33(6):25–39. doi: 10.1519/SSC.0b013e318231bb64
- Langan-Evans C, Reale R, Sullivan J, Martin D. Nutritional considerations for female athletes in weight category sports. *Eur J Sport Sci*. (2021) 22(5):720–32. doi: 10.1080/17461391.2021.1936655
- Barley OR, Chapman DW, Abbiss CR. Weight loss strategies in combat sports and concerning habits in mixed martial arts. *Int J Sports Physiol Perform*. (2018) 13(7):933–9. doi: 10.1123/ijssp.2017-0715
- Burke LM, Slater GJ, Matthews JJ, Langan-Evans C, Horswill CA. ACSM Expert consensus statement on weight loss in weight-category sports. *Curr Sports Med Rep*. (2021) 20(4):199–217. doi: 10.1249/JSR.0000000000000831
- Langan-Evans C, Germaine M, Artukovic M, Oxborough DL, Areta JL, Close GL, et al. The psychological and physiological consequences of low energy availability in a Male combat sport athlete. *Med Sci Sports Exerc*. (2021) 53(4):673–83. doi: 10.1249/MSS.0000000000002519
- Morton JP, Robertson C, Sutton L, MacLaren DPM. Making the weight: a case study from professional boxing. *Int J Sport Nutr Exerc Metab*. (2010) 20(1):80–5. doi: 10.1123/ijsnem.20.1.80
- Mountjoy M, Sundgot-Borgen JK, Burke LM, Ackerman KE, Blauwet C, Constantini N, et al. IOC Consensus statement on relative energy deficiency in sport (RED-S): 2018 update. *Br J Sports Med*. (2018) 52:687–97. doi: 10.1136/bjsports-2018-099193
- Nattiv A, Loucks AB, Manore MM, Sanborn CF, Sundgot-Borgen J, Warren MP. The female athlete triad. *Med Sci Sports Exerc*. (2007) 39(10):1867–82. doi: 10.1249/mss.0b013e318149f111
- Nattiv A, De Souza MJ, Koltun KJ, Misra M, Kussman A, Williams NI, et al. The Male athlete triad – a consensus statement from the female and Male athlete triad coalition part 1: definition and scientific basis. *Clin J Sport Med*. (2021) 31(4):335–48. doi: 10.1097/JSM.0000000000000946
- Watson AM. Sleep and athletic performance. *Curr Sports Med Rep*. (2017) 16(6):413–8. doi: 10.1249/JSR.0000000000000418
- Peacock CA, Mena M, Sanders GJ, Silver TA, Kalman D, Antonio J. Sleep data, physical performance, and injuries in preparation for professional mixed martial arts. *Sports*. (2019) 7(1):1. doi: 10.3390/sports7010001
- Hirshkowitz M, Whiton K, Albert SM, Alessi C, Bruni O, DonCarlos L, et al. National sleep foundation's sleep time duration recommendations: methodology and results summary. *Sleep Health*. (2015) 1(1):40–3. doi: 10.1016/j.sleh.2014.12.010
- Ohayon M, Wickwire EM, Hirshkowitz M, Albert SM, Avidan A, Daly FJ, et al. National sleep foundation's sleep quality recommendations: first report. *Sleep Health*. (2017) 3(1):6–19. doi: 10.1016/j.sleh.2016.11.006
- Taylor SR, Rogers GG, Driver HS. Effects of training volume on sleep, psychological, and selected physiological profiles of elite female swimmers.

## Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

## Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

- Med Sci Sports Exerc*. (1997) 29(5):688–93. doi: 10.1097/00005768-199705000-00016
- Thornton HR, Delaney JA, Duthie GM, Dascombe BJ. Effects of preseason training on the sleep characteristics of professional rugby league players. *Int J Sports Physiol Perform*. (2018) 13(2):176–82. doi: 10.1123/ijssp.2017-0119
- Caia J, Scott TJ, Halson SL, Kelly VG. Do players and staff sleep more during the pre- or competitive season of elite rugby league? *Eur J Sport Sci*. (2017) 17(8):964–72. doi: 10.1080/17461391.2017.1335348
- Hauswirth C, Louis J, Aubry A, Bonnet G, Duffield R, Le Meur Y. Evidence of disturbed sleep and increased illness in overreached endurance athletes. *Med Sci Sports Exerc*. (2014) 46(5):1036–45. doi: 10.1249/MSS.0000000000000177
- Thornton HR, Duthie GM, Pitchford NW, Delaney JA, Benton DT, Dascombe BJ. Effects of a 2-week high intensity training camp on sleep activity of professional rugby league athletes. *Int J Sports Physiol Perform*. (2017) 12(7):928–33. doi: 10.1123/ijssp.2016-0414
- Roberts SSH, Teo WP, Warmington SA. Effects of training and competition on the sleep of elite athletes: a systematic review and meta-analysis. *Br J Sports Med*. (2019) 53(8):513–22. doi: 10.1136/bjsports-2018-099322
- Kirk C, Langan-Evans C, Clark DR, Morton JP. Quantification of training load distribution in mixed martial arts athletes: a lack of periodisation and load management. *PLoS One*. (2021) 16(5):e0251266. doi: 10.1371/journal.pone.0251266
- Morehen JC, Langan-Evans C, Hall ECR, Close GL, Morton JP. A 5-year analysis of weight cycling practices in a Male world champion professional boxer: potential implications for obesity and cardiometabolic disease. *Int J Sport Nutr Exerc Metab*. (2021) 31(6):1–7. doi: 10.1123/ijsnem.2021-0085
- Obminski Z, Mroczko J. Orthostatic trials and perceptual responses to daily loads during 25 days of preparatory period 1-day competition and 5 days of recovery period in elite female judokas. *J Combat Sports Martial Arts*. (2015) 6(1):27–33. doi: 10.5604/20815735.1174228
- Ouergui I, Ardigo LP, Selmi O, Levitt DE, Chtourou H, Bouassida A, et al. Changes in perceived exertion, well-being, and recovery during specific judo training: impact of training period and exercise modality. *Front Physiol*. (2020) 11:931. doi: 10.3389/fphys.2020.00931
- Lastella M, Lovell GP, Sargent C. Athletes precompetitive sleep behaviour and its relationship with subsequent precompetitive mood and performance. *Eur J Sport Sci*. (2014) 14(sup 1):S123–30. doi: 10.1080/17461391.2012.660505
- Mejri MA, Yousfi N, Mhenni T, Tayech A, Hammouda O, Driss T, et al. Does one night of partial sleep deprivation affect the evening performance during intermittent exercise in taekwondo players? *J Exerc Rehabil*. (2016) 12(1):47–53. doi: 10.12965/jer.150256
- Spiegel K, Tasali E, Penev P, Van Cauter E. Brief communication: sleep curtailment in healthy young men is associated with decreased leptin levels,

- elevated ghrelin levels, and increased hunger and appetite. *Ann Intern Med.* (2004) 141(11):846–50. doi: 10.7326/0003-4819-141-11-200412070-00008
27. Keay N, Overseas A, Francis G. Indicators and correlates of low energy availability in male and female dancers. *BMJ Open Sport Exerc Med.* (2020) 6(1):e000906. doi: 10.1136/bmjsem-2020-000906
28. Schaal K, Tafflet M, Nassif H, Thibault V, Pichard C, Alcotte M, et al. Psychological balance in high level athletes: gender-based differences and sport-specific patterns. *PLoS One.* (2011) 6(5):e19007. doi: 10.1371/journal.pone.0019007
29. Dunican IC, Eastwood PR, Murray K, Caldwell JA, Reale R. The effect of water loading for acute weight loss following fluid restriction on sleep quality and quantity in combat sports athletes. *J Exerc Nutr.* (2019) 2(5):S1. ISSN 2640-2572
30. Nana A, Slater GJ, Stewart AD, Burke LM. Methodology review: using dual-energy x-ray absorptiometry (DXA) for the assessment of body composition in athletes and active people. *Int J Sport Nutr Exerc Metab.* (2015) 25(2):198–215. doi: 10.1123/ijsnem.2013-0228
31. Loucks AB, Kiens B, Wright HH. Energy availability in athletes. *J Sports Sci.* (2011) 29(Sup 1):S7–15. doi: 10.1080/02640414.2011.588958
32. Foster C, Florhaug JA, Franklin J, Gottschall L, Hrovatin LA, Parker S, et al. A new approach to monitoring exercise training. *J Strength Cond Res.* (2001) 15(1):109–15. PMID: 11708692
33. Rupp TL, Balkin TJ. Comparison of motionlogger watch and actiwatch actigraphs to polysomnography for sleep/wake estimation in healthy young adults. *Behav Res Methods.* (2011) 43:1152–60. doi: 10.3758/s13428-011-0098-4
34. Carney CE, Buysse DJ, Ancoli-Israel S, Edinger JD, Krystal AD, Lichstein KL, et al. The consensus sleep diary: standardizing prospective sleep self-monitoring. *Sleep.* (2012) 35(2):287–302. doi: 10.5665/sleep.1642
35. Hopkins WG, Marshall SW, Batterham AM, Hanin J. Progressive statistics for studies in sports medicine and exercise science. *Med Sci Sports Exercise.* (2009) 41(1):3–13. doi: 10.1249/MSS.0b013e31818cb278
36. Leeder J, Glaister M, Pizzoferro K, Dawson J, Pedlar C. Sleep duration and quality in elite athletes measured using wristwatch actigraphy. *J Sports Sci.* (2012) 30(6):541–5. doi: 10.1080/02640414.2012.660188
37. Lastella M, Roach GD, Halson SL, Sargent C. Sleep/wake behaviours of elite athletes from individual and team sports. *Eur J Sport Sci.* (2015) 15(2):94–100. doi: 10.1080/17461391.2014.932016
38. Sargent C, Halson S, Roach GD. Sleep or swim? Early-morning training severely restricts the amount of sleep obtained by elite swimmers. *Eur J Sport Sci.* (2014) 14(sup 1):S310–315. doi: 10.1080/17461391.2012.696711
39. Sargent C, Lastella M, Halson SL, Roach GD. The impact of training schedules on the sleep and fatigue of elite athletes. *Chronobiol Int.* (2014) 31(10):1160–8. doi: 10.3109/07420528.2014.957306