



OPEN ACCESS

EDITED BY
Frontiers in Surgery Editorial Office,
Frontiers Media SA, Switzerland

*CORRESPONDENCE
Zhun Xu
xuzhun2012@163.com

[†]These authors have contributed equally to this work.

SPECIALTY SECTION
This article was submitted to Orthopedic Surgery, a section of the journal Frontiers in Surgery

RECEIVED 14 September 2022
ACCEPTED 15 September 2022
PUBLISHED 05 October 2022

CITATION
Zheng B-W, Liu F-S, Zheng B-Y, Niu H, Li J, Lv G-H, Zou M-X and Xu Z (2022)
Corrigendum: Risk factors for tuberculous or nontuberculous spondylitis after percutaneous vertebroplasty or kyphoplasty in patients with osteoporotic vertebral compression fracture: A case-control study.
Front. Surg. 9:1044417.
doi: 10.3389/fsurg.2022.1044417

COPYRIGHT
© 2022 Zheng, Liu, Zheng, Niu, Li, Lv, Zou and Xu. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Corrigendum: Risk factors for tuberculous or nontuberculous spondylitis after percutaneous vertebroplasty or kyphoplasty in patients with osteoporotic vertebral compression fracture: A case-control study

Bo-Wen Zheng^{1,2,3†}, Fu-Sheng Liu^{1,2†}, Bo-Yv Zheng⁴, HuaQing Niu⁴, Jing Li², Guo-Hua Lv², Ming-Xiang Zou¹ and Zhun Xu^{1*}

¹Department of Spine Surgery, The First Affiliated Hospital, Hengyang Medical School, University of South China, Hengyang, China, ²Department of Spine Surgery, The Second Xiangya Hospital, Central South University, Changsha, China, ³Musculoskeletal Tumor Center, Peking University People's Hospital, Peking University, Beijing, China, ⁴Department of Orthopedics Surgery, General Hospital of the Central Theater Command, Wuhan, China

KEYWORDS

percutaneous vertebroplasty, percutaneous kyphoplasty, tuberculous spondylitis, nontuberculous spondylitis, pyogenic spondylitis, risk factors

A corrigendum on

[Risk factors for tuberculous or nontuberculous spondylitis after percutaneous vertebroplasty or kyphoplasty in patients with osteoporotic vertebral compression fracture: A case-control study](#)

by Zheng B-W, Liu F-S, Zheng B-Y, Niu HQ, Li J, Lv G-H, Zou1 M-X and Xu Z. (2022) Front. Surg. 9: 962425. doi: 10.3389/fsurg.2022.962425

Incorrect Funding

In the published article, there was an error in the Funding statement. [This work was supported by the National Natural Science Foundation of China (81871821 to JL and 82002364 to MXZ) and Project for Clinical Research of Hunan Provincial Health Commission (20201956 to MXZ and Natural Science Foundation of Hunan Province)]. The correct Funding statement appears below.

Funding

[This work was supported by the National Natural Science Foundation of China (81871821 to JL and 82002364 to MXZ), Natural Science Foundation of Hunan Province (2022JJ40403 to ZX), and Project for Clinical Research of Hunan Provincial Health Commission (20201956 to MXZ)]

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.