



# Inhibition Controls Asynchronous States of Neuronal Networks

Mario Treviño\*

*Instituto de Neurociencias, Universidad de Guadalajara, Guadalajara, Mexico*

Computations in cortical circuits require action potentials from excitatory and inhibitory neurons. In this mini-review, I first provide a quick overview of findings that indicate that GABAergic neurons play a fundamental role in coordinating spikes and generating synchronized network activity. Next, I argue that these observations helped popularize the notion that network oscillations require a high degree of spike correlations among interneurons which, in turn, produce synchronous inhibition of the local microcircuit. The aim of this text is to discuss some recent experimental and computational findings that support a complementary view: one in which interneurons participate actively in producing *asynchronous* states in cortical networks. This requires a proper mixture of shared excitation and inhibition leading to asynchronous activity between neighboring cells. Such contribution from interneurons would be extremely important because it would tend to reduce the spike correlation between neighboring pyramidal cells, a drop in redundancy that could enhance the information-processing capacity of neural networks.

**Keywords:** oscillations, network, cortex, inhibition, interneuron, synchrony

## OPEN ACCESS

### Edited by:

Martín Cammarota,  
Federal University of Rio Grande do  
Norte, Brazil

### Reviewed by:

M. Gustavo Murer,  
Universidad de Buenos Aires,  
Argentina

Adriano B. L. Tort,  
Federal University of Rio Grande do  
Norte, Brazil

### \*Correspondence:

Mario Treviño  
mariomtv@hotmail.com

**Received:** 09 March 2016

**Accepted:** 29 April 2016

**Published:** 12 May 2016

### Citation:

Treviño M (2016) Inhibition Controls  
Asynchronous States of Neuronal  
Networks.  
*Front. Synaptic Neurosci.* 8:11.  
doi: 10.3389/fnsyn.2016.00011

## INTRODUCTION

Glutamatergic (excitatory) and GABAergic (inhibitory) neurons can be classified on the basis of their morphology and electrophysiological properties, as well as on the neuromodulatory receptors expressed on their membranes. Pyramidal neurons (PyrCs) are the principal type of excitatory cells in cortex. Each one receives approximately  $10^4$  synaptic inputs, of which approximately 75% are excitatory and 25% inhibitory (Amaral and Witter, 1989; Ishizuka et al., 1995). The excitatory synapses on PyrCs are distributed along their dendrites, whereas the inhibitory synapses are located throughout their dendritic tree, the soma, and the initial axon segment. Thus, in PyrCs, excitatory and inhibitory inputs co-exist on the peripheral dendritic tree, while the proximal segments receive only inhibition (Pouille et al., 2013).

Although far less numerous, the GABAergic interneurons (INTs) are exceptionally diverse in terms of their biochemical, biophysical and morphological properties, as well as in their connectivity with other neurons (Freund and Buzsáki, 1996; Kawaguchi and Kubota, 1997; Somogyi and Klausberger, 2005). Probably due to the high metabolic costs involved in building and maintaining long-range axonal projections, the wiring of most INTs is local. An apparent inverse relationship between potential synaptic contacts from interneurons and the distance of their axonal projection supports this idea (Gupta et al., 2000; Buzsáki et al., 2004; Ferreira et al., 2014). One effect of this arrangement is that neighboring PyrCs and INTs share some of their synaptic inputs and process similar information.


Neurons in the brain receive concurrent excitatory and inhibitory inputs. Depending on the driving force, a synaptic response can produce either a depolarization, a hyperpolarization or simply

no change in the membrane potential of the postsynaptic cell. Increasing the net synaptic conductance lowers the input resistance (due to more simultaneous inputs), reducing the voltage gradient generated by additional currents and the membrane time constant of the cell. In consequence, synaptic inhibition can decrease the amount of membrane depolarization in two complementary ways: by producing a hyperpolarizing current (i.e., a “subtractive” effect), or by increasing the global conductance of the cell by acting in such a way as to divide all the synaptic currents (i.e., “shunting inhibition”, also known as “silent inhibition”; Koch et al., 1990; Borg-Graham et al., 1998; Treviño and Gutiérrez, 2005; Silver, 2010; Treviño et al., 2011). Theoretical work supports this notion and predicts that inhibition will have a “divisive” effect on the postsynaptic potentials if conductance change is large and located close to the soma, but will produce a “subtractive” effect if the changes in input conductance are small and spatially-distributed (Silver, 2010). For all these reasons, INTs are key players in controlling the dynamics of recurrent excitatory circuits at various spatial and temporal scales. Perisomatic inhibition tightly regulates the spike timing of postsynaptic cells and is able to synchronize entire neural ensembles (Freund and Buzsáki, 1996; Glickfeld and Scanziani, 2006; Mann and Paulsen, 2007; Silberberg and Markram, 2007).

## BUILDING PRINCIPLES OF CORTICAL CIRCUITS

The complex patterns of excitatory and inhibitory connections in the brain provide the structural basis for a rich repertoire of network activities. Cortical INTs can perform both feed-forward and feed-back inhibitory actions based on their input activity and their sensitivity to neuromodulators (Salgado et al., 2012, 2016). In feed-forward inhibition (FFI), an INT outputs its inhibitory signal to a PyrC that receives shared excitation (Figure 1A). Typical *in vitro* recordings performed from PyrCs responding to stimulation of afferent fibers show a compound synaptic response that consists of an excitatory postsynaptic potential (EPSP) followed by an inhibitory postsynaptic potential (IPSP) mediated by the activation of GABA<sub>A</sub> and, sometimes, GABA<sub>B</sub> receptors (Figure 1B). This built-in structural inequality sharpens the EPSP by narrowing the temporal “window of opportunity” for action potential generation, while also reducing the overall number of action potentials and limiting the temporal summation of EPSPs in the PyrCs (McCormick, 1992; Pouille and Scanziani, 2001; Maccaferri and Dingledine, 2002; Lawrence and McBain, 2003; Gabernet et al., 2005; Higley and Contreras, 2006; Ferrante et al., 2009; Torborg et al., 2010). The nature and strength of inhibition regulate the propagation speed of excitatory activity through cortical circuits (Trevelyan et al., 2007; Moldakarimov et al., 2015).

The generation of an action potential in cortical neurons is the result of complex spatio-temporal interactions between simultaneous synaptic inputs (Treviño et al., 2011). The precise relationship between synaptic excitation and inhibition (i.e., the E/I ratio) determines fundamental cortical operations



such as feature selectivity and gain (Haider and McCormick, 2009; Isaacson and Scanziani, 2011). Interestingly, despite fluctuating network activity levels, PyrCs in the cortex show quite stable averaged E/I ratios over time. This cortical “E/I balance” is achieved because inhibition increases proportionally with excitation, mainly through an enhanced recruitment of INTs (Anderson et al., 2000; Shu et al., 2003; Haider et al., 2006; Okun and Lampl, 2008; Xue et al., 2014). Yet, to complete this concept: some short-term plastic changes in the strength of synaptic transmission can also lead to small but progressive changes in the E/I balance. In the cerebral cortex, for example, both excitation and inhibition on PyrCs undergo short-term depression during repetitive electrical stimulation, but inhibition has a greater depressive effect, increasing the E/I ratio. This, in turn, leads to a progressively longer temporal window for EPSP summation and a decrease in spike-timing precision. Similarly, in the hippocampus, a single action potential in a dentate granule cell evokes a net inhibitory potential in CA3 PyrCs (Figure 1B), but this input signal can lead to an excitatory output if the

stimulation frequency is increased (Gabernet et al., 2005; Treviño et al., 2007, 2011). The process responsible for this switch involves the facilitation of excitatory responses coupled with a rapid depression of inhibitory transmission (Mori et al., 2004; Szabadics and Soltesz, 2009). A last example: in hippocampal CA1, feed-forward excitatory synaptic strength onto fast-spiking basket cells remains quite constant, whereas monosynaptic connections between fast-spiking basket cells and PyrCs are depressed during repetitive stimulation, resulting in an overall depression of the inhibitory drive onto PyrCs. Thus, the E/I balance can be regulated in a frequency-dependent manner by short-term changes in synaptic transmission (Galarreta and Hestrin, 1998; Gabernet et al., 2005; Glickfeld and Scanziani, 2006).

## THE INPUT-OUTPUT TRANSFER FUNCTION OF CORTICAL NEURONS

The probability that a neuron will fire an action potential in response to afferent stimuli is described by its input-output (I/O) function. The I/O function depends on the passive and active electrical properties of the cell, but also on the dendritic morphology and on the E/I balance. Mathematically, this function can be represented by a logistic curve: the neuron is nearly silent at low input rates, but increases its firing output with a given slope (also referred to as the “gain control”) before finally reaching a plateau at saturation input rates (Llinás, 1988; Bartesaghi et al., 2006; Campanac and Debanne, 2008; Pouille et al., 2013).

Typically, the all-or-none properties of axonal spiking, together with the presence of active dendritic properties, yield a steep I/O sigmoid with a very limited operative range for input stimuli. However, because neurons are regularly subjected to an intense ongoing synaptic bombardment *in vivo*, this leads to an actual I/O function with a smaller gain. In other words, the combination of variable “background” synaptic noise with the intrinsic properties of the cells renders an I/O relationship with a broader dynamic range (for input signals), mixing single-spike and burst responses at multiple membrane potentials (Borg-Graham et al., 1998; Paré et al., 1998; Wolfart et al., 2005; Monier et al., 2008). Mechanistically, transformations of the I/O relationship can occur because the voltage fluctuations allow synaptic inputs to cross the action potential threshold more frequently, even when the average membrane potential is well below the spike threshold. Interestingly, such high levels of voltage noise fluctuations can only be achieved during uncorrelated (asynchronous) inputs, when excitatory and inhibitory synaptic conductances are, on average, well-balanced (Okun and Lampl, 2008; Petersen and Crochet, 2013).


## INHIBITION CONTROLS ASYNCHRONOUS STATES OF NEURONAL NETWORKS

Evidence from *in vitro* and *in vivo* recordings indicates that the activity of INTs plays a major role in coordinating

spikes and generating oscillatory network activity (Freund and Buzsáki, 1996; Bartos et al., 2007; Treviño et al., 2007). By counterbalancing excitatory inputs in a temporally precise way, correlated inhibition can orchestrate action potential timing in multiple postsynaptic neurons. This is crucial for general cortical functioning and also for inducing synaptic plasticity (Kwag and Paulsen, 2009). Contrary to this view, however, recent experimental evidence reveals that neural networks, both *in vitro* and *in vivo*, generally display asynchronous states with low spiking correlations among cortical neurons including INTs (Okun and Lampl, 2008; Petersen and Crochet, 2013; Sippy and Yuste, 2013). One candidate mechanism to explain the low correlated spiking output is that presynaptic excitatory and inhibitory (phasic) inputs are themselves correlated, shunting the membranes of postsynaptic cells. Some authors have interpreted the high correlations in IPSCs observed in neighboring PyrCs as an indication of a high correlation in the spiking activity of presynaptic INTs (Hasenstaub et al., 2005). However, recent experiments indicate that correlations in IPSCs can be easily explained by the fact that nearby cells share some of their inhibitory input (**Figure 2A**). In other words, this means that inhibition can indeed be locally coherent, but not due to synchrony among INTs, but simply because PyrCs share common presynaptic inputs from INTs. Here, importantly, inhibition is globally no more correlated than excitation (Okun and Lampl, 2008; Sippy and Yuste, 2013).

In their theoretical model of de-correlated networks, Renart et al. (2010) proposed that shared excitatory and inhibitory input correlations could indeed explain the low levels of spiking (output) correlation between pairs of PyrCs, as observed in dozens of *in vivo* experiments (**Figure 2A**). To test this model, Sippy and Yuste (2013) performed dual voltage-clamp recordings in PyrCs from brain slices during thalamically-triggered UP states at different holding potentials. This strategy changed the driving force of inhibition and excitation (at least in the perisomatic region) and provided control of their relative synaptic efficacy. In agreement with the model, the intracellular recordings revealed a V-shaped relationship between the correlation of membrane currents of nearby PyrCs and their membrane potential (**Figure 2B**). That is, the correlation between synaptic currents was highest at  $-70$  mV and  $+0$  mV, where excitation and inhibition are relatively isolated, whereas at intermediate potentials, the net effect of the synchronous excitatory and inhibitory currents lead to a lower output correlation between the membranes of nearby PyrCs (Sippy and Yuste, 2013; see also Hasenstaub et al., 2005). Thus, a proper mixture of excitation and inhibition leads to a state of relative asynchrony between neighboring cells (**Figures 2A,B**). Certainly, in this explanatory framework, inhibition and excitation must be correlated in postsynaptic membranes (Renart et al., 2010; Sippy and Yuste, 2013).

Further support to these ideas: *in vivo* experiments by Okun and Lampl (2008) have shown that, during spontaneous and sensory-evoked activities, neighboring cortical cells display



highly-correlated: (i) EPSPs/EPSPs (**Figure 2C**, upper panel); (ii) IPSPs/IPSPs; and (iii) EPSPs/IPSPs (**Figure 2C**, lower panel). Moreover, the cross-correlations of the EPSPs and IPSPs from neighboring cells peaked at a positive delay of  $\sim 3$  ms. This indicates that, on average, inhibition lags behind excitation

by several milliseconds. Such tight coupling is consistent with the hypothesis that inhibition controls the integration time window of excitation, enabling neurons to operate as coincidence detectors. In other words, because nearby cortical neurons receive highly-similar excitatory inputs, FFI might

indeed constitute a robust mechanism for de-correlating network activity (Figure 2D).

Assuming that FFI accounts for the strong correlations between excitatory and inhibitory inputs into PyrCs, it would be natural to ask whether the correlation state of a network could be sensitive to (or manipulated by) changes in the inhibitory “tone” in cortical networks. This has been confirmed experimentally: substantial increases in network synchrony occur both *in vitro* and *in vivo* when using pharmacological agonists that reduce the probability of GABA release, and also with antagonists that block GABAergic conductances (Cohen and Miles, 2000; Treviño et al., 2007; Petersen and Crochet, 2013). Similarly, a partial blockage of GABAergic receptors can produce an increase in pair-wise correlations of EPSCs between PyrCs, confirming that IPSCs participate in de-correlating output spikes (Sippy and Yuste, 2013). Therefore, reducing the GABA function increases network synchrony *in vitro*, whereas adding INTs into simulated networks *in silico* decreases their output correlation (Figure 2D).

## POSSIBLE COMPUTATIONAL BENEFIT FROM ASYNCHRONOUS ACTIVITY

Neural activity can represent information as a firing rate, as correlations in spike-timing, or as a complex combination of both (Silver, 2010; Luczak et al., 2015). Many researchers believe that networks can display a rich repertoire of synchronous and asynchronous states and that they can switch between them by adjusting the E/I balance. Experimental measurements *in vivo* corroborate this view, and theoretical studies suggest that networks can self-organize to reach such balanced states in which different neurons emit action potentials asynchronously (van Vreeswijk and Sompolinsky, 1996; Brunel, 2000; Renart et al., 2010). Two fundamentally different types of asynchronous activity have been distinguished (Ostojic, 2014). When synaptic couplings between neurons are weak, the network at rest can operate in an “homogenous asynchronous state”, a regime well-suited for transmitting information about the firing rate of external inputs. This is because neurons in the network will change their mean firing rate in proportion to the input. In contrast, for strong couplings, a network at rest displays rich internal dynamics, in which the firing rates of individual neurons fluctuate strongly, both in time and across neurons. In this scenario, the internal state of the network is such that repeated presentations of the same external stimulus leads to very different responses (i.e., the input is dynamically transformed). Certainly, this variability in the population degrades the transmission of information but provides a rich substrate for a nonlinear processing of the stimuli (Ostojic, 2014).

Neuronal responses are typically variable in the sense that the number and timing of the spikes in response to the same stimulus is never identical among trials (Lee et al., 1998; Kilgard and Merzenich, 1999; Ringach et al., 2002; Chelaru and Dragoi, 2008; Moreno-Bote et al., 2014). The responses of nearby neurons located within the same “functional

column” in the cortex, which are thought to encode the “same stimulus property”, exhibit a high degree of heterogeneity. Yet, interestingly, the trial-by-trial variability in neuronal responses is not independent, but exhibits correlations (Zohary et al., 1994; Lee et al., 1998; Ringach et al., 2002; Chelaru and Dragoi, 2008). One possibility could be that population coding depends not only on the response properties of the cells but also on the distribution of neuronal correlations across the network (Abbott and Dayan, 1999; Pouget et al., 2000; Sompolinsky et al., 2001; Luczak et al., 2015). And in this respect, the high variability of intrinsic response properties of individual cells may change the structure of neuronal correlations thereby improving the information encoded in the population activity (Chelaru and Dragoi, 2008). This proposal predicts that the most energy-efficient codes for representing information would be sparse, as observed in multiple *in vivo* preparations (Petersen and Crochet, 2013). Under a sparse coding regime, a neuron could act as a coincidence detector of temporally-correlated inputs. Furthermore, computational strategies used by the brain should depend on the amount of information that can be stored in population activity. *In vivo*, noise correlations tend to be positive and proportional to the similarity in tuning properties, and are thought to limit information. It would be thus natural to think that de-correlating neuronal activity could increase the information capacity of the network (Abbott and Dayan, 1999; Pouget et al., 2000; Sompolinsky et al., 2001; Seriès et al., 2004). In other words, because information usually decreases as correlations increase, it could be advantageous if the brain were to possess a mechanism to de-correlate neural activity, either through a passive process such as balancing or synchronizing excitation and inhibition, or an active one such as attention (Cohen and Maunsell, 2009; Mitchell et al., 2009; Renart et al., 2010).

## CONCLUDING REMARKS

Inhibition is generally conceptualized as a mechanism that restricts the probability of action potential generation by reorganizing spikes in multiple cells and promoting network oscillations (Bartos et al., 2007). Here, I have discussed how inhibition can also promote and control the generation of asynchronous states. A proposal is that depending on the spatiotemporal dynamics of synaptic inputs, inhibition could act as a correlating or de-correlating agent that controls network synchrony. Signal integration of postsynaptic cells is transformed through shared inhibition, affecting their I/O transfer function and de-correlating spikes in neighboring cells which might increase the network’s storage-capacity (but see Moreno-Bote et al., 2014).

Spontaneous spiking by most cortical neurons *in vivo* and *in vitro* is generally asynchronous and infrequent (usually <1 Hz; Renart et al., 2010; Petersen and Crochet, 2013) but, intuitively, such sporadic activity should eventually reach a point of complete inactivity because it cannot sufficiently depolarize postsynaptic neurons to

reach their spike threshold. How cortical networks are able to maintain asynchronous and infrequent activity is unclear. Maybe spikes result from cell-intrinsic—i.e., non-synaptically driven—processes: some studies suggest that certain brain areas can generate discharges in the absence of (sensory) evoked synaptic inputs (e.g., Cohen and Miles, 2000). Different levels of neuromodulators, or different attentional or arousal levels could also regulate the intrinsic conductances and the firing and excitability of neurons as well as their transition into desynchronized UP states *in vivo* (Destexhe et al., 2003; Wladyka and Kunze, 2006; Constantinople and Bruno, 2011). Perhaps both intrinsic cellular properties and shared synaptic interactions contribute to the asynchronous state of cortical networks. If so, fine-tuning of the “E/I balance” may be a key mechanism for modulating spike output correlations and transitions among brain states.

## REFERENCES

- Abbott, L. F., and Dayan, P. (1999). The effect of correlated variability on the accuracy of a population code. *Neural Comput.* 11, 91–101. doi: 10.1162/089976699300016827
- Amaral, D. G., and Witter, M. P. (1989). The three-dimensional organization of the hippocampal formation: a review of anatomical data. *Neuroscience* 31, 571–591. doi: 10.1016/0306-4522(89)90424-7
- Anderson, J. S., Carandini, M., and Ferster, D. (2000). Orientation tuning of input conductance, excitation and inhibition in cat primary visual cortex. *J. Neurophysiol.* 84, 909–926.
- Bartese, R., Migliore, M., and Gessi, T. (2006). Input-output relations in the entorhinal cortex-dentate-hippocampal system: evidence for a non-linear transfer of signals. *Neuroscience* 142, 247–265. doi: 10.1016/j.neuroscience.2006.06.001
- Bartos, M., Vida, I., and Jonas, P. (2007). Synaptic mechanisms of synchronized  $\gamma$  oscillations in inhibitory interneuron networks. *Nat. Rev. Neurosci.* 8, 45–56. doi: 10.1038/nrn2044
- Borg-Graham, L. J., Monier, C., and Frégnac, Y. (1998). Visual input evokes transient and strong shunting inhibition in visual cortical neurons. *Nature* 393, 369–373. doi: 10.1038/30735
- Brunel, N. (2000). Dynamics of sparsely connected networks of excitatory and inhibitory spiking neurons. *J. Comput. Neurosci.* 8, 183–208. doi: 10.1023/A:1008925309027
- Buzsáki, G., Geisler, C., Henze, D. A., and Wang, X. J. (2004). *Interneuron Diversity series: circuit complexity and axon wiring economy of cortical interneurons.* *Trends Neurosci.* 27, 186–193. doi: 10.1016/j.tins.2004.02.007
- Campanac, E., and Debanne, D. (2008). Spike timing-dependent plasticity: a learning rule for dendritic integration in rat CA1 pyramidal neurons. *J. Physiol.* 586, 779–793. doi: 10.1113/jphysiol.2007.147017
- Chelaru, M. I., and Dragoi, V. (2008). Efficient coding in heterogeneous neuronal populations. *Proc. Natl. Acad. Sci. U S A* 105, 16344–16349. doi: 10.1073/pnas.0807744105
- Cohen, M. R., and Maunsell, J. H. (2009). Attention improves performance primarily by reducing interneuronal correlations. *Nat. Neurosci.* 12, 1594–1600. doi: 10.1038/nn.2439
- Cohen, I., and Miles, R. (2000). Contributions of intrinsic and synaptic activities to the generation of neuronal discharges in *in vitro* hippocampus. *J. Physiol.* 524, 485–502. doi: 10.1111/j.1469-7793.2000.00485.x
- Constantinople, C. M., and Bruno, R. M. (2011). Effects and mechanisms of wakefulness on local cortical networks. *Neuron* 69, 1061–1068. doi: 10.1016/j.neuron.2011.02.040
- Destexhe, A., Rudolph, M., and Paré, D. (2003). The high-conductance state of neocortical neurons *in vivo*. *Nat. Rev. Neurosci.* 4, 739–751. doi: 10.1038/nrn1198

## AUTHOR CONTRIBUTIONS

MT conceived and wrote the article.

## FUNDING

Research in my laboratory was funded by grants from the Consejo Nacional de Ciencia y Tecnología (CONACyT; 220862, 251406).

## ACKNOWLEDGMENTS

I thank Y. Elyada (The Hebrew University of Jerusalem) for useful comments during an earlier version of the manuscript; J. Bargas (IFC-UNAM, México) and P.R. Killeen (Arizona State University) for exciting discussions and E. Matute for constant support.

- Ferrante, M., Migliore, M., and Ascoli, G. A. (2009). Feed-forward inhibition as a buffer of the neuronal input-output relation. *Proc. Natl. Acad. Sci. U S A* 106, 18004–18009. doi: 10.1073/pnas.0904784106
- Ferreira, T. A., Blackman, A. V., Oyrer, J., Jayabal, S., Chung, A. J., Watt, A. J., et al. (2014). Neuronal morphometry directly from bitmap images. *Nat. Methods* 11, 982–984. doi: 10.1038/nmeth.3125
- Freund, T. F., and Buzsáki, G. (1996). Interneurons of the hippocampus. *Hippocampus* 6, 347–470. doi: 10.1002/(SICI)1098-1063(1996)6:4<347::AID-HIPO1>3.0.CO;2-I
- Gabernet, L., Jadhav, S. P., Feldman, D. E., Carandini, M., and Scanziani, M. (2005). Somatosensory integration controlled by dynamic thalamocortical feed-forward inhibition. *Neuron* 48, 315–327. doi: 10.1016/j.neuron.2005.09.022
- Galarreta, M., and Hestrin, S. (1998). Frequency-dependent synaptic depression and the balance of excitation and inhibition in the neocortex. *Nat. Neurosci.* 1, 587–594. doi: 10.1038/2822
- Glickfeld, L. L., and Scanziani, M. (2006). Distinct timing in the activity of cannabinoid-sensitive and cannabinoid-insensitive basket cells. *Nat. Neurosci.* 9, 807–815. doi: 10.1038/nn1688
- Gupta, A., Wang, Y., and Markram, H. (2000). Organizing principles for a diversity of GABAergic interneurons and synapses in the neocortex. *Science* 287, 273–278. doi: 10.1126/science.287.5451.273
- Haider, B., Duque, A., Hasenstaub, A. R., and McCormick, D. A. (2006). Neocortical network activity *in vivo* is generated through a dynamic balance of excitation and inhibition. *J. Neurosci.* 26, 4535–4545. doi: 10.1523/jneurosci.5297-05.2006
- Haider, B., and McCormick, D. A. (2009). Rapid neocortical dynamics: cellular and network mechanisms. *Neuron* 62, 171–189. doi: 10.1016/j.neuron.2009.04.008
- Hasenstaub, A., Shu, Y., Haider, B., Kraushaar, U., Duque, A., and McCormick, D. A. (2005). Inhibitory postsynaptic potentials carry synchronized frequency information in active cortical networks. *Neuron* 47, 423–435. doi: 10.1016/j.neuron.2005.06.016
- Higley, M. J., and Contreras, D. (2006). Balanced excitation and inhibition determine spike timing during frequency adaptation. *J. Neurosci.* 26, 448–457. doi: 10.1523/jneurosci.3506-05.2006
- Isaacson, J. S., and Scanziani, M. (2011). How inhibition shapes cortical activity. *Neuron* 72, 231–243. doi: 10.1016/j.neuron.2011.09.027
- Ishizuka, N., Cowan, W. M., and Amaral, D. G. (1995). A quantitative analysis of the dendritic organization of pyramidal cells in the rat hippocampus. *J. Comp. Neurol.* 362, 17–45. doi: 10.1002/cne.903620103
- Kawaguchi, Y., and Kubota, Y. (1997). GABAergic cell subtypes and their synaptic connections in rat frontal cortex. *Cereb. Cortex* 7, 476–486. doi: 10.1093/cercor/7.6.476

- Kilgard, M. P., and Merzenich, M. M. (1999). Distributed representation of spectral and temporal information in rat primary auditory cortex. *Hear. Res.* 134, 16–28. doi: 10.1016/s0378-5955(99)00061-1
- Koch, C., Douglas, R., and Wehmer, U. (1990). Visibility of synaptically induced conductance changes: theory and simulations of anatomically characterized cortical pyramidal cells. *J. Neurosci.* 10, 1728–1744.
- Kwag, J., and Paulsen, O. (2009). The timing of external input controls the sign of plasticity at local synapses. *Nat. Neurosci.* 12, 1219–1221. doi: 10.1038/nn.2388
- Lawrence, J. J., and McBain, C. J. (2003). *Interneuron Diversity series*: containing the detonation–feedforward inhibition in the CA3 hippocampus. *Trends Neurosci.* 26, 631–640. doi: 10.1016/j.tins.2003.09.007
- Lee, D., Port, N. L., Kruse, W., and Georgopoulos, A. P. (1998). Variability and correlated noise in the discharge of neurons in motor and parietal areas of the primate cortex. *J. Neurosci.* 18, 1161–1170.
- Llinás, R. R. (1988). The intrinsic electrophysiological properties of mammalian neurons: insights into central nervous system function. *Science* 242, 1654–1664. doi: 10.1126/science.3059497
- Luczak, A., McNaughton, B. L., and Harris, K. D. (2015). Packet-based communication in the cortex. *Nat. Rev. Neurosci.* 16, 745–755. doi: 10.1038/nrn4026
- Maccaferri, G., and Dingledine, R. (2002). Control of feedforward dendritic inhibition by NMDA receptor-dependent spike timing in hippocampal interneurons. *J. Neurosci.* 22, 5462–5472.
- Mann, E. O., and Paulsen, O. (2007). Role of GABAergic inhibition in hippocampal network oscillations. *Trends Neurosci.* 30, 343–349. doi: 10.1016/j.tins.2007.05.003
- McCormick, D. A. (1992). Neurotransmitter actions in the thalamus and cerebral cortex and their role in neuromodulation of thalamocortical activity. *Prog. Neurobiol.* 39, 337–388. doi: 10.1016/0301-0082(92)90012-4
- Mitchell, J. F., Sundberg, K. A., and Reynolds, J. H. (2009). Spatial attention decorrelates intrinsic activity fluctuations in macaque area V4. *Neuron* 63, 879–888. doi: 10.1016/j.neuron.2009.09.013
- Moldakarimov, S., Bazhenov, M., and Sejnowski, T. J. (2015). Feedback stabilizes propagation of synchronous spiking in cortical neural networks. *Proc. Natl. Acad. Sci. U S A* 112, 2545–2550. doi: 10.1073/pnas.1500643112
- Monier, C., Fournier, J., and Fregnac, Y. (2008). *In vitro* and *in vivo* measures of evoked excitatory and inhibitory conductance dynamics in sensory cortices. *J. Neurosci. Methods* 169, 323–365. doi: 10.1016/j.jneumeth.2007.11.008
- Moreno-Bote, R., Beck, J., Kanitscheider, I., Pitkow, X., Latham, P., and Pouget, A. (2014). Information-limiting correlations. *Nat. Neurosci.* 17, 1410–1417. doi: 10.1038/nn.3807
- Mori, M., Abegg, M. H., Gähwiler, B. H., and Gerber, U. (2004). A frequency-dependent switch from inhibition to excitation in a hippocampal unitary circuit. *Nature* 431, 453–456. doi: 10.1038/nature02854
- Okun, M., and Lampl, I. (2008). Instantaneous correlation of excitation and inhibition during ongoing and sensory-evoked activities. *Nat. Neurosci.* 11, 535–537. doi: 10.1038/nn.2105
- Ostojic, S. (2014). Two types of asynchronous activity in networks of excitatory and inhibitory spiking neurons. *Nat. Neurosci.* 17, 594–600. doi: 10.1038/nn.3658
- Paré, D., Shink, E., Gaudreau, H., Destexhe, A., and Lang, E. J. (1998). Impact of spontaneous synaptic activity on the resting properties of neocortical pyramidal neurons *in vivo*. *J. Neurophysiol.* 79, 1450–1460.
- Petersen, C. C., and Crochet, S. (2013). Synaptic computation and sensory processing in neocortical layer 2/3. *Neuron* 78, 28–48. doi: 10.1016/j.neuron.2013.03.020
- Pouget, A., Dayan, P., and Zemel, R. (2000). Information processing with population codes. *Nat. Rev. Neurosci.* 1, 125–132. doi: 10.1038/35039062
- Pouille, F., and Scanziani, M. (2001). Enforcement of temporal fidelity in pyramidal cells by somatic feed-forward inhibition. *Science* 293, 1159–1163. doi: 10.1126/science.1060342
- Pouille, F., Watkinson, O., Scanziani, M., and Trevelyan, A. J. (2013). The contribution of synaptic location to inhibitory gain control in pyramidal cells. *Physiol. Rep.* 1:e00067. doi: 10.1002/phy2.67
- Renart, A., de la Rocha, J., Bartho, P., Hollender, L., Parga, N., Reyes, A., et al. (2010). The asynchronous state in cortical circuits. *Science* 327, 587–590. doi: 10.1126/science.1179850
- Ringach, D. L., Shapley, R. M., and Hawken, M. J. (2002). Orientation selectivity in macaque V1: diversity and laminar dependence. *J. Neurosci.* 22, 5639–5651.
- Salgado, H., Köhr, G., and Treviño, M. (2012). Noradrenergic ‘tone’ determines dichotomous control of cortical spike-timing-dependent plasticity. *Sci. Rep.* 2:417. doi: 10.1038/srep00417
- Salgado, H., Treviño, M., and Atzori, M. (2016). Layer- and area-specific actions of norepinephrine on cortical synaptic transmission. *Brain Res.* doi: 10.1016/j.brainres.2016.01.033 [Epub ahead of print].
- Seriès, P., Latham, P. E., and Pouget, A. (2004). Tuning curve sharpening for orientation selectivity: coding efficiency and the impact of correlations. *Nat. Neurosci.* 7, 1129–1135. doi: 10.1038/nn1321
- Shu, Y., Hasenstaub, A., and McCormick, D. A. (2003). Turning on and off recurrent balanced cortical activity. *Nature* 423, 288–293. doi: 10.1038/nature01616
- Silberberg, G., and Markram, H. (2007). Disynaptic inhibition between neocortical pyramidal cells mediated by Martinotti cells. *Neuron* 53, 735–746. doi: 10.1016/j.neuron.2007.02.012
- Silver, R. A. (2010). Neuronal arithmetic. *Nat. Rev. Neurosci.* 11, 474–489. doi: 10.1038/nrn2864
- Sippy, T., and Yuste, R. (2013). Decorrelating action of inhibition in neocortical networks. *J. Neurosci.* 33, 9813–9830. doi: 10.1523/JNEUROSCI.4579-12.2013
- Somogyi, P., and Klausberger, T. (2005). Defined types of cortical interneurone structure space and spike timing in the hippocampus. *J. Physiol.* 562, 9–26. doi: 10.1113/jphysiol.2004.078915
- Sompolinsky, H., Yoon, H., Kang, K., and Shamir, M. (2001). Population coding in neuronal systems with correlated noise. *Phys. Rev. E Stat. Nonlin. Soft Matter Phys.* 64:051904. doi: 10.1103/physreve.64.051904
- Szabadics, J., and Soltesz, I. (2009). Functional specificity of mossy fiber innervation of GABAergic cells in the hippocampus. *J. Neurosci.* 29, 4239–4251. doi: 10.1523/JNEUROSCI.5390-08.2009
- Torborg, C. L., Nakashiba, T., Tonegawa, S., and McBain, C. J. (2010). Control of CA3 output by feedforward inhibition despite developmental changes in the excitation-inhibition balance. *J. Neurosci.* 30, 15628–15637. doi: 10.1523/JNEUROSCI.3099-10.2010
- Trevelyan, A. J., Sussillo, D., and Yuste, R. (2007). Feedforward inhibition contributes to the control of epileptiform propagation speed. *J. Neurosci.* 27, 3383–3387. doi: 10.1523/jneurosci.0145-07.2007
- Treviño, M., and Gutiérrez, R. (2005). The GABAergic projection of the dentate gyrus to hippocampal area CA3 of the rat: pre- and postsynaptic actions after seizures. *J. Physiol.* 567, 939–949. doi: 10.1113/jphysiol.2005.092064
- Treviño, M., Vivar, C., and Gutiérrez, R. (2007).  $\beta/\gamma$  oscillatory activity in the CA3 hippocampal area is depressed by aberrant GABAergic transmission from the dentate gyrus after seizures. *J. Neurosci.* 27, 251–259. doi: 10.1523/jneurosci.3815-06.2007
- Treviño, M., Vivar, C., and Gutiérrez, R. (2011). Excitation-inhibition balance in the CA3 network–neuronal specificity and activity-dependent plasticity. *Eur. J. Neurosci.* 33, 1771–1785. doi: 10.1111/j.1460-9568.2011.07670.x
- van Vreeswijk, C., and Sompolinsky, H. (1996). Chaos in neuronal networks with balanced excitatory and inhibitory activity. *Science* 274, 1724–1726. doi: 10.1126/science.274.5293.1724
- Wladyka, C. L., and Kunze, D. L. (2006). KCNQ/M-currents contribute to the resting membrane potential in rat visceral sensory neurons. *J. Physiol.* 575, 175–189. doi: 10.1113/jphysiol.2006.113308
- Wolfart, J., Debay, D., Le Masson, G., Destexhe, A., and Bal, T. (2005). Synaptic background activity controls spike transfer from thalamus to cortex. *Nat. Neurosci.* 8, 1760–1767. doi: 10.1038/nn1591

- Xue, M., Atallah, B. V., and Scanziani, M. (2014). Equalizing excitation-inhibition ratios across visual cortical neurons. *Nature* 511, 596–600. doi: 10.1038/nature13321
- Zohary, E., Shadlen, M. N., and Newsome, W. T. (1994). Correlated neuronal discharge rate and its implications for psychophysical performance. *Nature* 370, 140–143. doi: 10.1038/370140a0

**Conflict of Interest Statement:** The author declares that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

The reviewer AT and handling Editor MC declared their shared affiliation, and the handling Editor states that the process nevertheless met the standards of a fair and objective review.

*Copyright © 2016 Treviño. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution and reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.*