



OPEN ACCESS

EDITED AND REVIEWED BY
Leonid A. Dombrovsky,
Joint Institute for High Temperatures
(RAS), Russia

*CORRESPONDENCE
Yatao Ren,
✉ renyt@hit.edu.cn

RECEIVED 06 July 2023
ACCEPTED 26 July 2023
PUBLISHED 03 August 2023

CITATION
Ren Y, Singh S and Soni S (2023), Editorial:
Computational modeling of various
procedures in thermal therapy of
human tumors.
Front. Front. Therm. Eng. 3:1253718.
doi: 10.3389/fther.2023.1253718

COPYRIGHT
© 2023 Ren, Singh and Soni. This is an
open-access article distributed under the
terms of the [Creative Commons
Attribution License \(CC BY\)](#). The use,
distribution or reproduction in other
forums is permitted, provided the original
author(s) and the copyright owner(s) are
credited and that the original publication
in this journal is cited, in accordance with
accepted academic practice. No use,
distribution or reproduction is permitted
which does not comply with these terms.

Editorial: Computational modeling of various procedures in thermal therapy of human tumors

Yatao Ren^{1*}, Sundeep Singh² and Sanjeev Soni³

¹School of Energy Science and Engineering, Harbin Institute of Technology, Harbin, China, ²Faculty of Sustainable Design Engineering, University of Prince Edward Island, Charlottetown, PE, Canada, ³Biomedical Applications Group, CSIR-Central Scientific Instruments Organization, Chandigarh, India

KEYWORDS

thermal therapy, computational modeling, bioheat and mass transfer, tumor, photothermal therapy (PPT)

Editorial on the Research Topic

Computational modeling of various procedures in thermal therapy of human tumors

Cancer is the leading cause in many countries, which lead to almost 10 million deaths in 2020 (Sung et al., 2021). In the meantime, the traditional cancer therapeutics including radiotherapy, chemotherapy, and surgery, all suffer from different side effects or limitations (Ren et al., 2022). Thermal therapy, commonly known as hyperthermia has been used to treat cancer and other diseases since at least 4,000 years ago (Hornback, 1989; Glazer and Curley, 2010). Hyperthermia cancer therapy involves subjecting a tumor (locally or regionally) to a temperature within 42°C–48°C for a duration of an hour to a few minutes depending on the temperature used. Due to its advantages of minimal- or non-invasive nature, it has attracted the interest of researchers worldwide. Thermal therapy can be characterized as photothermal therapy, ultrasound thermal therapy, radiofrequency thermal therapy, magnetic hyperthermia, etc., according to the heat generation methods (Khurana et al., 2022; Singh et al., 2022). In the context of photothermal therapy, an indirect heating strategy has been proposed in which the tissue around the tumor is heated instead of the tumor itself, which will reduce or even stop the oxygen supply of the tumor cells (Dombrovsky et al., 2012; Dombrovsky, 2022). Due to the development of nanotechnology, plasmonic nanoparticles are employed to enhance energy absorption during photothermal therapy, termed as plasmonic photothermal therapy, which is very popular in recent years. The most important Research Topic for thermal therapy is precise temperature control to kill the tumor cell selectively. Accurate numerical modeling of thermal therapy, especially the heat transfer process, is very important for the optimization and real time adjustment of thermal therapy procedure or pre-treatment planning before thermal therapy (Ren et al., 2022). This Research Topic aims to report the important development in the field of computational modeling of different kinds of procedures involved in thermal therapy, including but not limited to heat transfer, light propagation, blood flow, thermal damage, etc., which is very important to quantify the optimal thermal dosage for patient-specific settings.

In this Research Topic, Mariappan et al. has proposed a point source model to predict the power generated by the needle which does not require to obtain the complex electric field during radiofrequency ablation. The proposed model was applied to both two-dimensional

and three-dimensional simulations and the obtained results were in good agreement with those predicted with the Joule heat model. The proposed point source model approximation would provide the source term to be utilized in the Pennes bioheat transfer equation without the need to actually solve (or coupling) the electric field model, leading to a significant reduction in the computational time for predicting the temperature distribution and ablation volume. It should be mentioned that blood flow plays an extremely important role in the heat transfer process within living tissue. However, this effect has been considered as a simplified blood perfusion rate term in the Pennes bioheat transfer equation, which accounts for volumetric heat transfer in organs containing large vessels. To get information about the local blood temperature, another model, named two coupled energy equations, has been proposed to take into account the spatial and time variations in arterial blood temperature (Xuan and Roetzel, 1997). It has also been extended and applied to solve the heat transfer problem during photothermal therapy (Dombrovsky et al., 2011; Dombrovsky et al., 2012).

In another article of this Research Topic, Tenorio et al. presented an *in silico* photothermal therapy system aimed at enhancing the efficacy of therapy in attaining complete tumor necrosis while restricting damage to the surrounding healthy tissue. An external temperature-tracking system, a real-time thermal damage predictor, a fuzzy logic or a PI controller, and a continuous wave laser is included in this system. A closed-loop feedback system was designed that would be able to modulate the input power delivered to the tissue on the basis of predicted thermal response in both target (tumorous) and healthy regions of the tissue, thus, providing a safe and effective thermal therapy.

Besides the above numerical work, this Research Topic also includes an experimental study by Li et al. related to the application of moxibustion in treating cancer-related fatigue in colorectal patients. Governor Vessel Moxibustion trials were conducted on 80 patients for a duration of 3 weeks. The presented results provide

experimental evidence showing greater reductions in interleukin-6 and tumor necrosis factor- α levels, with no serious adverse events. Thus, highlighting the safety and effectiveness of Governor Vessel Moxibustion for alleviating cancer-related fatigue in colorectal patients.

The three research papers in this Research Topic cover a wide range of current state-of-the-art developments in the field of thermal therapy, from numerical to experimental works. In view of the rising interest in exploring safe and effective alternative therapies for cancer, more exciting work regarding numerical models and treatment strategies of thermal therapy is still expected.

Author contributions

YR: Writing–original draft, Writing–review and editing, SuS: Writing–original draft, Writing–review and editing, SaS: Writing–original draft, Writing–review and editing.

Conflict of interest

The authors YR, SuS, and SaS declared that they were an editorial board member of Frontiers, at the time of submission. This had no impact on the peer review process and the final decision.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

References

- Dombrovsky, L. A., Timchenko, V., Jackson, M., and Yeoh, G. H. (2011). A combined transient thermal model for laser hyperthermia of tumors with embedded gold nanoshells. *Int. J. Heat Mass Transf.* 54 (25), 5459–5469. doi:10.1016/j.ijheatmasstransfer.2011.07.045
- Dombrovsky, L. A., Timchenko, V., and Jackson, M. (2012). Indirect heating strategy for laser induced hyperthermia: An advanced thermal model. *Int. J. Heat Mass Transf.* 55 (17–18), 4688–4700. doi:10.1016/j.ijheatmasstransfer.2012.04.029
- Dombrovsky, L. A. (2022). Laser-induced thermal treatment of superficial human tumors: An advanced heating strategy and non-Arrhenius law for living tissues. *Front. Therm. Eng.* 1, 807083. doi:10.3389/fther.2021.807083
- Glazer, E. S., and Curley, S. A. (2010). The ongoing history of thermal therapy for cancer. *Surg. Oncol. Clin. N. Am.* 20 (2), 229–235. doi:10.1016/j.soc.2010.11.001
- Hornback, N. B. (1989). Historical aspects of hyperthermia in cancer therapy. *Radiologic Clin. N. Am.* 27 (3), 481–488. doi:10.1016/s0033-8389(22)02133-9
- Khurana, D., Dudi, R., Shukla, S. K., Singh, D., Mondhe, D. M., and Soni, S. (2022). Gold nanoblockbodies mediated plasmonic photothermal cancer therapy for melanoma. *Nanomedicine* 17 (19), 1323–1338. doi:10.2217/nmm-2022-0052
- Ren, Y., Yan, Y., and Qi, H. (2022). Photothermal conversion and transfer in photothermal therapy: From macroscale to nanoscale. *Adv. Colloid Interface Sci.* 308, 102753. doi:10.1016/j.cis.2022.102753
- Singh, S., Saccomandi, P., and Melnik, R. (2022). Three-phase-lag bio-heat transfer model of cardiac ablation. *Fluids* 7 (5), 180. doi:10.3390/fluids7050180
- Sung, H., Ferlay, J., Siegel, R. L., Laversanne, M., Soerjomataram, I., Jemal, A., et al. (2021). Global cancer statistics 2020: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA A Cancer J. Clin.* 71 (3), 209–249. doi:10.3322/caac.21660
- Xuan, Y., and Roetzel, W. (1997). Bioheat equation of the human thermal system. *Chem. Eng. Technol.* 20 (4), 268–276. doi:10.1002/ceat.270200407