



## OPEN ACCESS

## EDITED BY

Orhan Sahin,  
Iowa State University, United States

## REVIEWED BY

Marwa Ibrahim Abd El-Hamid,  
Zagazig University, Egypt  
Issmat I. Kassem,  
University of Georgia, United States

## \*CORRESPONDENCE

Beatrix Stessl  
✉ Beatrix.Stessl@vetmeduni.ac.at

RECEIVED 14 July 2023

ACCEPTED 07 November 2023

PUBLISHED 22 November 2023

## CITATION

Mustedanagic A, Matt M, Weyermair K,  
Schrattenecker A, Kubitzka I, Firth CL, Loncaric I,  
Wagner M and Stessl B (2023) Assessment of  
microbial quality in poultry drinking water on  
farms in Austria.  
*Front. Vet. Sci.* 10:1254442.  
doi: 10.3389/fvets.2023.1254442

## COPYRIGHT

© 2023 Mustedanagic, Matt, Weyermair,  
Schrattenecker, Kubitzka, Firth, Loncaric,  
Wagner and Stessl. This is an open-access  
article distributed under the terms of the  
[Creative Commons Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/).  
The use, distribution or reproduction in other  
forums is permitted, provided the original  
author(s) and the copyright owner(s) are  
credited and that the original publication in this  
journal is cited, in accordance with accepted  
academic practice. No use, distribution or  
reproduction is permitted which does not  
comply with these terms.

# Assessment of microbial quality in poultry drinking water on farms in Austria

Azra Mustedanagic<sup>1,2</sup>, Monika Matt<sup>3</sup>, Karin Weyermair<sup>4</sup>,  
Anna Schrattenecker<sup>2</sup>, Isabella Kubitzka<sup>2</sup>, Clair L. Firth<sup>5</sup>,  
Igor Loncaric<sup>6</sup>, Martin Wagner<sup>1,2</sup> and Beatrix Stessl<sup>2\*</sup>

<sup>1</sup>FFoQSI GmbH-Austrian Competence Centre for Feed and Food Quality, Safety and Innovation, Tulln, Austria, <sup>2</sup>Unit of Food Microbiology, Institute of Food Safety, Food Technology and Veterinary Public Health, University of Veterinary Medicine Vienna, Vienna, Austria, <sup>3</sup>Department of Statistics and Analytical Epidemiology, Austrian Agency for Health and Food Safety (AGES), Innsbruck, Austria, <sup>4</sup>Department of Statistics and Analytical Epidemiology, Austrian Agency for Health and Food Safety (AGES), Graz, Austria, <sup>5</sup>Unit of Veterinary Public Health and Epidemiology, Institute of Food Safety, Food Technology and Veterinary Public Health, University of Veterinary Medicine Vienna, Vienna, Austria, <sup>6</sup>Institute of Microbiology, Department of Pathobiology, University of Veterinary Medicine Vienna, Vienna, Austria

The quality of poultry drinking water has a significant effect on broiler health and performance. This study conducted an analysis of aerobic mesophilic counts (AMC), *Enterobacteriaceae* (EB), *Pseudomonadaceae* (PS), and screened for the presence of *Campylobacter* spp. in water samples collected from a total of 14 farms in Austria, with either a public or private water source. The efficacy of two water line treatment methods was evaluated: a chemical treatment of the water lines with 4.0 ppm ClO<sub>2</sub> (T1) and a combined chemical (4.0 ppm active ClO<sub>2</sub> and 3.0% peracetic acid) and mechanical treatment (purging of the water lines with a high-pressure air pump; T2). However, both the T1 and T2 treatments failed to reduce the AMC counts below the maximum acceptable microbial limit of 4.0 log<sub>10</sub> CFU/ml in water samples. In addition, no significant reduction in EB and PS counts was observed in water samples after either T1 or T2 water line treatment. The water samples showed a high level of microbial diversity with 18 to 26 different genera. The genus *Pseudomonas* was most frequently isolated across all poultry farms, while *Campylobacter jejuni* was identified in a single sample collected before water line treatment. Isolate analysis revealed the presence of opportunistic pathogens in water samples both before (T1 43.1%, T2 30.9%) and after (T1 36.3%, T2 33.3%) water line treatment. Opportunistic pathogens belonging to genera including *Pseudomonas* spp., *Stenotrophomonas* spp., and *Ochrobactrum* spp., were most frequently isolated from poultry drinking water. These isolates exhibited multidrug resistance and resistance phenotypes to antimicrobials commonly used in Austrian poultry farms. The findings of this study emphasize the potential risk of exposure to opportunistic pathogens for poultry and personnel, underscoring the importance of efficient water line management.

## KEYWORDS

water line treatment, opportunistic pathogens, poultry health, *Pseudomonas*, antimicrobial susceptibility

## 1 Introduction

Poultry is one of the main sources of meat production worldwide (1). In 2020, more than 97 million chickens were processed in Austria, representing 124,000 tons of processed poultry meat (2). Drinking water is a vital nutrient for commercial poultry and has a significant impact on poultry health, liveweight, feed conversion ratios, and overall performance (3, 4). The water consumption of poultry is approximately twice the amount of feed intake (5). Poultry health and water intake are directly influenced by microbial water quality (4, 6, 7).

In Europe, the water quality standards for poultry drinking water have been adapted from water quality regulations intended for human drinking water consumption (8), EC Directive 98/83/EC (Drinking Water Directive [DWD] 9). According to the Austrian Poultry Hygiene Regulation (10) drinking water used for poultry production must not exceed a total aerobic mesophilic count (AMC) of  $2.0 \log_{10}$  and  $1.3 \log_{10}$  colony forming units (CFU/ml) at 22° and 37°C, respectively. Currently, there is no legal requirement to examine microbial contamination inside the drinking water lines (11). Hence, maintenance of water line hygiene is primarily the responsibility of the poultry producer, and it is typically conducted between the production cycles (12). The standard water line practices involve mechanical cleaning by flushing the water lines, followed by oxidative disinfection, primarily using chlorination or acidifiers (7, 12–14).

While water line treatment is a crucial component of an effective biosecurity program, its effectiveness does not ensure the complete elimination of the microorganisms within the water lines (15–17). *Escherichia coli*, *Salmonella* spp., and *Campylobacter* spp. have been detected in poultry drinking water (7, 18). Elevated temperatures and low water flow rates in enclosed water line systems have been found to adversely affect water quality, as indicated by previous studies (4, 12). These conditions are favorable for the accumulation of dissolved organic substances, minerals, and solid particles, which facilitate growth and promote the formation of biofilms. Among biofilm-forming bacteria, primarily *Pseudomonas* and *Stenotrophomonas* are responsible for biofilm formation on surfaces of poultry drinking lines (12). Biofilms may provide a favorable surface for attachment of opportunistic pathogens (OP), such as *Acinetobacter*, *Aeromonas*, *Citrobacter*, *Enterobacter*, and *Klebsiella* whose members are natural inhabitants of plumbing systems and adapted to survival in drinking water (19). Although these bacteria are generally not pathogenic, some have the potential to cause infections in susceptible poultry and farm workers (20). Hence, the detachment of pathogen and OP rich biofilms and their contamination of the water system present a significant risk for waterborne transmission of these bacteria, posing a potential threat to both poultry and human health. Moreover, the administration of medication to poultry through drinking water, which is a preferred route, has been linked to presence of multidrug-resistant (MDR) bacteria (21, 22).

Microbial water quality is frequently evaluated at its source, but assessments at the end of the drinking lines are infrequent, despite the potential for substantial variations in microbial quality between the source and endpoint (12). Thus, the objective of this study was to evaluate the microbial quality of water samples collected at the end of a production cycle of five to six weeks and shortly before restocking for the subsequent production cycle, following the water line treatment. Previous studies have demonstrated the presence of pathogens such as *Campylobacter* spp. in poultry water on farms with

private water supplies compared to those with a public supply (23, 24). This highlights the critical role of poultry drinking water as a potential source of *Campylobacter* spp. infection on the farm (25, 26). The presence of *Campylobacter* spp. in drinking water on poultry farms may indicate lapses in biosecurity, contaminated water source, ineffective and/or incorrectly applied water line cleaning procedures (11, 18). Therefore, one of our objectives was to assess the microbial quality of poultry drinking water in farms with either public or private water supply. We applied ISO-based reference methods to assess bacterial load and presence of *Campylobacter* spp. in poultry drinking water, followed by partial 16S rRNA sequencing of bacterial isolates. Antibiotic susceptibility patterns of commonly isolated OP were then determined.


## 2 Materials and methods

### 2.1 Water line treatment and sample collection

Twenty-eight poultry farms producing broilers for local slaughterhouses in Austria voluntarily participated in the study between May 2019 and August 2020, some of which had private ( $n=11$ ) and others public ( $n=17$ ) water supplies. The fattening period at the participating poultry farms in Austria was five to six weeks. The poultry farms were divided into two distinct groups based on whether the farms employed solely chemical (T1) or a combination of chemical and mechanical (T2) water line treatment methods. An overview of the poultry farms included in the study is presented in Figure 1. Cleaning and water line treatment at the poultry farms was performed by the farmer. Since the participation of poultry farms in the study was voluntary, poultry farms 6, 8, 9, 12, and 13 withdrew their participation after T1 and were substituted by the poultry farms 15–19 during T2. The study was conducted in collaboration with a private laboratory (HYGIENICUM GmbH, Graz, Austria), which provided training on the water line cleaning procedures to be implemented at the poultry farms to the participating farmers.

During T1 water line treatment, water lines were drained and filled with a commercially-available solution of which the main disinfecting component contained 4.0 ppm active chlorine dioxide ( $\text{ClO}_2$ ) solution (Calgonit CD-K1/K2, Calvatis GmbH, Ladenburg, Germany). The commercial solution was retained in the water lines for 24 h. Measurements of free  $\text{ClO}_2$  inside the waterlines were not obtained. Subsequently, the water lines were washed with the supply water by continuous flushing for 10 min. Under normal operating conditions. The T2 water line was performed by continuous pumping of acidic cleaner containing 3.0% peroxyacetic acid (PAA) and hydrogen peroxide (Calgonit DS 625, Calvatis GmbH, Ladenburg, Germany) continuously for 30 min using high-pressure air pump. The water lines were then washed with the supply water and purged using a high-pressure air pump until no inorganic and organic debris were visible in the water. Subsequently, the water line disinfection was performed using a commercial disinfection solution containing 4.0 ppm active  $\text{ClO}_2$  solution (Calgonit CD-K1/K2) which was retained in the water lines for 24 h. Subsequently, the water lines were washed with supply water by flushing for 10 min. Under normal operating conditions.

Water samples were collected by employees from the private laboratory, samples were taken from the end nipple of the drinking



water line inside the vacant poultry house (HYGIENICUM GmbH, Graz, Austria). One water line was sampled at four and five poultry farms, while two water lines (line 1 and 2) were sampled at ten and nine poultry farms during T1 and T2 water line treatments (Figure 1). Two sampling timepoints were chosen, namely before treatment (BT) at the end of fattening period of 5–6 weeks, and after the water line treatment (AT) before restocking of the subsequent production cycle. As shown in Figure 1 and Supplementary Table S1, in six poultry farms during the T1 and T2 water line treatment, water samples were collected at two different sampling intervals, while other poultry farms were sampled only once. Additionally, at some poultry farms from some water lines the duplicate samples were collected, while from other poultry farms only a single sample was collected. Therefore, in total 36 (T1) and 33 (T2) BT and corresponding AT samples were collected for the microbial analysis in the present study. The water samples were collected in sterile 500 mL bottles by the private laboratory and immediately transported to the laboratory at 4°C for microbial analysis.

## 2.2 Sample processing and microbial analysis

Prior to analysis, 500 mL of water samples were centrifuged at 8000 rpm for 30 min at 4°C (Thermo Scientific, Sorvall Lynx 4000 centrifuge). All but 10 mL of the supernatant was discarded, the remainder was then resuspended using a serological 10 mL pipette (Greiner Bio One, Frickenhausen, Germany) and vortexed for 30 s.

*Campylobacter* selective enrichment and isolation were performed according to the ISO 10272-1:2006 standard for the detection of *Campylobacter* spp. in foodstuff (27). Five milliliters of the supernatant were transferred to 45 mL of Bolton broth (Thermo Fisher Scientific Ltd., Hampshire, United Kingdom) supplemented with 5% hemolyzed horse blood (Oxoid Ltd., Hampshire, United Kingdom). The Bolton broth enrichment was incubated for up to 48 h at 42°C under microaerobic conditions (10% CO<sub>2</sub>, 3% O<sub>2</sub>, 87% N<sub>2</sub>). After incubation modified charcoal cefoperazone deoxycholate agar (mCCDA) (Oxoid Ltd) was inoculated by fractionated loop inoculation (10 µL) and incubated at 42°C for 48 h under microaerobic conditions. Quantification of aerobic mesophilic count (AMC), *Enterobacteriaceae* (EB), and *Pseudomonadaceae* (PS) counts was carried out according to ISO reference methods (28, 29). For enumeration of AMC, EB, and PS, 5 mL of the re-suspended supernatant was transferred to 45 mL buffered peptone water (BPW) (Biokar Solabia diagnostics, Pantin Cedex, France). Subsequently, serial ten-fold dilutions were prepared up to dilution 10<sup>-5</sup> in BPW (Biokar Solabia diagnostics, Pantin Cedex, France). The AMC were enumerated on trypto-caseine soy agar with 0.6% yeast extract (TSAYE) (Biokar Solabia diagnostics), while EB and PS were enumerated on red bile glucose agar (VRBG) (Merck KGaA, Darmstadt, Germany). Each dilution step (100 µL) was plated on selective agar media for the enumeration of AMC, EB, and PS counts. For dilution 10<sup>-1</sup> the volume of 1 mL was divided (333 µL) on three agar plates per selective medium. Agar plates were incubated at 30°C (AMC) and 37°C (EB, PS) aerobically for up to 48 h. The EB and PS counts on VRBG agar were differentiated by their ability to ferment

glucose, leading to pink colonies with or without precipitation and pale colonies for PS. Presumptive EB and PS isolates were confirmed using oxidase reaction (BioMerieux, Marcy l'Etoile, France). The minimum and maximum limits for the determination of the AMC, EB, and PS in the samples ranged between 10 and 300 CFU.

Microbial quality of water samples before (BT) and after (AT) sanitation were categorized according to AMC, EB, and PS load in two contamination levels,  $<4.0 \log_{10}$  CFU/ml and  $\geq 4.0 \log_{10}$  CFU/ml based on existing studies (4, 7, 12).

## 2.3 Isolation and identification of bacterial and *Campylobacter* spp. isolates

The predominant bacterial colony morphologies were collected from each water sample for further confirmation. Specifically, 1–5 colonies were selected from TSAYE ( $n=224$ ), VRBG ( $n=206$ ) and mCCDA agar ( $n=41$ ) and then subcultured on the respective medium. The isolate list is provided in the [Supplementary Table S1](#). The purified colonies, comprising isolates from T1 BT samples ( $n=123$ ), T1 AT samples ( $n=113$ ), T2 BT samples ( $n=139$ ), T2 AT samples ( $n=96$ ) were stored at  $-80^{\circ}\text{C}$  in brain heart infusion broth (Biokar Solabia diagnostics) supplemented with 25% (v/v) glycerol (Merck KgaA).

For DNA extraction of *Campylobacter* spp. isolates 10  $\mu\text{L}$  loop of bacterial material was resuspended in 100  $\mu\text{L}$  of 0.1 M Tris–HCl buffer pH 7 (Sigma Aldrich, St. Louis, MO, United States) and mixed with 400  $\mu\text{L}$  Chelex<sup>®</sup> 100-Resin (BioRad, Hercules, CA, United States) (30). The bacterial Chelex<sup>®</sup> 100Resin suspension was heated at  $100^{\circ}\text{C}$  for 10 min on a block heater (Thermo Fischer Scientific Inc.), followed by short centrifugation step at 15,000  $\times g$  (Eppendorf Centrifuge 5,425) for 5 s. The supernatant (100  $\mu\text{L}$ ) was transferred to a maximum recovery tube (Corning Incorporated Life Sciences, Reynosa, Mexico) and stored at  $-20^{\circ}\text{C}$  until analysis. *Campylobacter* spp. were identified using multiplex PCR targeting genes including the conserved genus-specific 23S rRNA gene, the *Campylobacter jejuni* hippuricase gene (*hipO*) and the *Campylobacter coli* serine hydroxymethyltransferase (*glyA*) gene, as previously described (31). Briefly, a single reaction mixture (20  $\mu\text{L}$ ) contained diethylpyrocarbonate (DEPC) treated water (Sigma Aldrich), 1 $\times$  PCR buffer, 2 mM  $\text{MgCl}_2$ , 500 nm *hipO* forward and reverse primer, 1,000 *glyA* forward and reverse primer, 200 nm 23S forward and reverse primer, 200  $\mu\text{M}$  dNTP mix, 1.5 U of Platinum Taq DNA polymerase (Platinum<sup>™</sup> Taq DNA Polymerase, DNAfree, Thermo Fisher Scientific Inc., Waltham, MA, United States), and 5  $\mu\text{L}$  template genomic DNA. The amplification was performed in T100<sup>™</sup> Thermal Cycler (Bio-Rad, Hercules, CA, United States). The PCR cycling conditions included initial denaturation at  $94^{\circ}\text{C}$  for 2 min, 30 cycles of denaturation ( $94^{\circ}\text{C}$  for 30 s), primer annealing ( $59^{\circ}\text{C}$  for 30 s), elongation ( $72^{\circ}\text{C}$  for 30 s) and final elongation ( $72^{\circ}\text{C}$  for 7 min). The gel electrophoresis of PCR amplicons was performed in a 1.5% agarose gel containing 0.5 $\times$  TrisBorateEDTA (TBE) buffer (Sigma Aldrich, St. Louis, MO, United States) and 3.5  $\mu\text{L}$  peqGREEN DNA gel stain (VWR International, Radnor, United States), at 120 V for 30 min. The DNA standard Thermo Scientific<sup>™</sup> GeneRuler<sup>™</sup> 100 bp (Thermo Fisher Scientific Inc., Waltham, United States) was applied for fragment length comparison. We utilized the following control isolates for the DNA extraction and multiplex PCR: *C. jejuni* strain DSM 4688 and *C. coli* strain DSM 4689, obtained from Deutsche Sammlung von

Mikroorganismen und Zellkulturen (DSMZ), Braunschweig, Germany.

For DNA extraction of isolates from TSAYE and VRBG, bacterial cells were lysed by boiling the suspension. A 10  $\mu\text{L}$  loop of bacterial material was re-suspended in 100  $\mu\text{L}$  0.1 M Tris–HCl pH 7 buffer (Sigma Aldrich, St. Louis, MO, United States), briefly vortexed and heated at  $100^{\circ}\text{C}$  for 15 min (Thermo Scientific<sup>™</sup> block heater, Thermo Fischer Scientific Inc.). The suspension was then centrifuged for 5 s at 15,000  $\times g$  (Eppendorf Centrifuge 5,425, Hamburg, Germany) and the supernatant (70  $\mu\text{L}$ ) was transferred into maximum recovery tubes (Corning Incorporated Life Sciences, Reynosa, Mexico) and stored at  $-20^{\circ}\text{C}$  until analysis. For identification of bacteria isolates ( $n=471$ ) the partial amplification of 16S rRNA gene was performed following the methods of (32, 33), using universal primer pairs 616F (5'AGAGTTTGATYMTGGCTC3') and 1492R (5'GGYTACCTTGTTACGACTT3') (both Microsynth AG, Blagach, Switzerland). A single PCR reaction (45  $\mu\text{L}$ ) contained 1 $\times$  PCR buffer, 2 mM  $\text{MgCl}_2$ , 200 nM forward and reverse primer, 250  $\mu\text{M}$  dNTP mix, 2 U of Platinum Taq DNA polymerase (Platinum<sup>™</sup> Taq DNA Polymerase, DNAfree, Thermo Fisher Scientific Inc.) and 5  $\mu\text{L}$  template genomic DNA. The DNA amplification was performed in T100<sup>™</sup> Thermal Cycler (BioRad, Hercules, CA, United States). The PCR cycling conditions included initial denaturation at  $95^{\circ}\text{C}$  for 5 min, 35 cycles of denaturation ( $94^{\circ}\text{C}$  for 30 s), primer annealing ( $52^{\circ}\text{C}$  for 30 s), elongation ( $72^{\circ}\text{C}$  for 60 s) and final elongation ( $72^{\circ}\text{C}$  for 7 min). Subsequently, the PCR amplicons were sent for purification and sanger sequencing to LGC Genomics (LGC Genomics GmbH, Berlin, Germany). The gel electrophoresis of PCR-amplicons was performed in a 1.5% agarose gel containing 0.5 $\times$  TrisBorateEDTA (TBE) buffer (Sigma Aldrich, St. Louis, MO, United States) and 3.5  $\mu\text{L}$  peqGREEN DNA gel stain (VWR International, Radnor, United States), at 120 V for 30 min. The DNA standard Thermo Scientific<sup>™</sup> GeneRuler<sup>™</sup> 100 bp (Thermo Fisher Scientific Inc., Waltham, United States) was applied for fragment length comparison. The PCR amplicons were sequenced using a 1492R (5'GGYTACCTTGTTACGACTT3') primer. The nucleotide sequences were qualityevaluated by using Finch TV 1.4.0 (34) and MEGA X (35). The bacterial nucleotide Basic Local Alignment Search Tool (BLAST) algorithm from the National Centre for Biotechnology Information (NCBI)<sup>1</sup> was used for taxonomy assignment. Sequences were assigned to genus or species level according to best matches and highest similarities (1,040 to 1,120 bp fragment length, similarity cutoff  $\geq 97.0\%$ ). The partial rRNA gene sequence data from the isolates were deposited in the GenBank database under accession numbers MZ642358 to MZ643011.<sup>2</sup> Subsequent identification of opportunistic pathogens among identified isolates was performed using the bacterial metadata base BacDive (36) and List of Prokaryotic names with Standing in Nomenclature (LPSN) (37).

## 2.4 Antimicrobial susceptibility testing

Opportunistic pathogens with clinical relevance isolated from water samples during T1 and T2 water line treatment were subjected

1 <https://blast.ncbi.nlm.nih.gov/Blast.cgi>

2 <https://www.ncbi.nlm.nih.gov/genbank>

to antimicrobial susceptibility testing (AST). The set of isolates included most frequently isolated OP, such as *Pseudomonas* spp. ( $n=17$ ), *Ochrobactrum* spp. ( $n=4$ ), *Stenotrophomonas* spp. ( $n=3$ ), and human relevant opportunistic pathogens including *Citrobacter* spp. ( $n=2$ ), *Enterobacter* spp. ( $n=2$ ), *Klebsiella* spp. ( $n=1$ ), and *Aeromonas* spp. ( $n=1$ ).

AST was performed for a total of 30 bacterial isolates using Sensititre™ Avian AVIAN1F Vet AST Plate (ThermoFischer Scientific Inc., Waltham, MA, United States), according to the manufacturer's instructions. Briefly, single colonies were picked from fresh cultures grown on TSAYE for 24 h at 30°C, suspended in sterile water to an optical density of a 0.5 McFarland standard ( $\sim 10^8$  CFU/mL). 50  $\mu$ l volumes of the bacterial suspension were transferred to wells containing different concentrations of lyophilized antimicrobials. Plates were sealed and incubated at 30°C for 24 to 48 h, after which minimum inhibitory concentrations (MIC) were read visually and defined as the lowest concentration of a given antibiotic at which no growth of the test organism was observed. *E. coli* strain ATCC 25922 was used as the internal quality control isolate. The minimum inhibitory concentration (MIC) breakpoints and definitions for multi-drug resistance (MDR; resistance to two or more antibiotic classes) (38) were determined following the standards provided by the Clinical and Laboratory Standards Institute (CLSI) manuals (39–41).

## 2.5 Data analysis

A descriptive analysis was carried out (mean, median, and standard deviation) for AMC, EB, and PS counts. The normal distribution of each data set (T1 and T2) was investigated using the Shapiro–Wilks test. Due to nonnormal distribution of data, the median values of AMC, EB, and PS counts were calculated. The Wilcoxon–Mann–Whitney rank sum test performed as a twosided test was applied to identify whether there was a significant difference between median AMC, EB and PS counts of BT and AT samples. Median AMC, EB, and PS counts in AT samples were compared for different water supplies (public vs. private), water line treatments (T1 vs. T2), following  $\log_{10}$  transformation, using Wilcoxon–Mann–Whitney rank sum test. Values of  $p < 0.05$  were considered as statistically significant. Statistical analyses were carried out using the R software package for statistical computing.<sup>3</sup>

## 3 Results

### 3.1 Aerobic mesophilic count, Enterobacteriaceae, and Pseudomonadaceae count in poultry drinking water

Ninety-nine BT samples and their corresponding AT water samples were microbiologically assessed, with a maximum acceptable microbial limit of 4.0  $\log_{10}$  CFU/ml for AMC, EB, and PS counts (Table 1). Due to non-normal distribution of the data, we used the

Wilcoxon–Mann–Whitney twosided rank sum test to assess the median values for AMC, EB, and PS counts. No significant differences ( $p \geq 0.05$ ) were observed between the median AMC, EB, and PS counts of the BT and AT samples after T1 water line treatment (Table 1). Furthermore, we did not observe any significant difference between median AMC, EB, and PS counts in poultry farms with private and public water supply. Among the water samples, the highest median AMC counts were observed in BT ( $5.9 \pm 1.02 \log_{10}$  CFU/ml, median  $\pm$  MAD; MAD: median absolute deviation) and AT ( $6.0 \pm 1.17 \log_{10}$  CFU/ml) samples. Higher median AMC counts in BT and AT samples were observed in poultry farms with a private well than those with a public water supply (Table 1). The lowest median counts were observed for EB in both BT ( $3.6 \pm 2.13 \log_{10}$  CFU/ml) and AT ( $2.3 \pm 1.52 \log_{10}$  CFU/ml) samples. In AT samples higher median EB counts were observed in poultry farms with public water supply. The PS resulted in the second highest median counts, which remained unchanged in BT ( $4.7 \pm 1.44 \log_{10}$  CFU/ml) and AT ( $4.7 \pm 2.48 \log_{10}$  CFU/ml) samples. Higher median PS counts were detected in both BT and AT samples in poultry farms with public water supply.

After T1 water line treatment, high ( $>4.0 \log_{10}$  CFU/ml) AMC, EB, and PS counts from BT samples decreased below the maximum acceptable microbial limit in 8/36, 7/36, and 9/36 AT samples, respectively (Supplementary Table S2). The AMC, EB, and PS below the microbial limit were observed in 1/36, 18/36, and 7/36 BT and AT samples, respectively. The AMC, EB, and PS counts above the maximum acceptable microbial limit were observed in 27/36, 11/36, and 20/36 AT samples, respectively, after T1 treatment.

During T2 water line sampling, no significant differences ( $p \geq 0.05$ ) were observed in the median AMC, EB, and PS counts between the BT and AT samples (Table 1). No significant difference was observed between median AMC, EB, and PS count in poultry farms with private and public water supply. The highest median counts were for AMC counts in both BT ( $4.6 \pm 1.55 \log_{10}$  CFU/ml) and AT ( $4.7 \pm 1.85 \log_{10}$  CFU/ml) samples, followed by the PS counts in BT ( $3.5 \pm 1.62 \log_{10}$  CFU/ml) and AT ( $3.1 \pm 2.05 \log_{10}$  CFU/ml) samples. The lowest counts were observed in the median EB counts of BT ( $2.4 \pm 1.63 \log_{10}$  CFU/ml) and AT ( $1.6 \pm 0.42 \log_{10}$  CFU/ml) samples. Higher median AMC, EB, and PS counts were detected in AT samples in poultry farms with public water supply.

After T2 water line treatment, high ( $>4.0 \log_{10}$  CFU/ml) AMC, EB, and PS counts from BT samples decreased below the maximum acceptable microbial limit in 8/33, 5/33, and 14/33 AT samples, respectively (Supplementary Table S3). The AMC, EB, and PS counts below the microbial limit were detected in 4/33, 25/33, and 10/33 samples in both BT and AT, respectively. The AMC, EB, and PS counts remained above the maximum acceptable microbial limit in 21/33, 3/33, and 9/33 AT samples, respectively, after T2 water line treatment.

The impact of T1 and T2 water line treatment on private and public water supply was evaluated by calculating the  $\log_{10}$  ratio from CFU  $\log_{10}$  counts detected in BT and AT water samples (Table 1). No significant differences ( $p \geq 0.05$ ) in  $\log_{10}$  ratios were observed for AMC, EB, and PS counts after T1 and T2 water line treatment. The  $\log_{10}$  ratio was not significantly different ( $p \geq 0.05$ ) between private and public supplied poultry farms after T1 and T2 water line treatment. The median AMC, EB, and PS ratios after T1 waterline treatment were  $-0.2 \pm 2.13$ ,  $-0.6 \pm 1.79$ , and  $0.0 \pm 2.26$ , respectively. The analysis of  $\log_{10}$  ratios after T2 waterline treatment resulted in median values of  $-1.1 \pm 2.13$  for AMC,  $0.0 \pm 2.94$  for EB, and  $0.0 \pm 3.12$  for PS counts. Although  $\log_{10}$

<sup>3</sup> www.r-project.org

TABLE 1 The median aerobic mesophilic count (AMC), *Enterobacteriaceae* (EB), and *Pseudomonadaceae* (PS) in poultry drinking water samples were determined before (BT) and after waterline treatment (AT) during T1 and T2 waterline treatment using culture-dependent methods.

| Treatment (T)  | Water supply | Median AMC               |            | Median AMC log <sub>10</sub> ratio | Median EB                |            | Median EB log <sub>10</sub> ratio | Median PS                |            | Median PS log <sub>10</sub> ratio | Campylobacter spp. |      |
|----------------|--------------|--------------------------|------------|------------------------------------|--------------------------|------------|-----------------------------------|--------------------------|------------|-----------------------------------|--------------------|------|
|                |              | log <sub>10</sub> CFU/ml |            |                                    | log <sub>10</sub> CFU/ml |            |                                   | log <sub>10</sub> CFU/ml |            |                                   |                    |      |
|                |              | BT                       | AT         | BT                                 | AT                       | BT         | AT                                | BT                       | AT         | BT                                | AT                 |      |
| 1              | Private      | 5.8 ± 1.30               | 5.4 ± 1.99 | -0.5 ± 2.37                        | 3.6 ± 2.13               | 1.6 ± 0.53 | -1.1 ± 1.85                       | 4.9 ± 1.30               | 3.7 ± 2.01 | -0.7 ± 2.71                       | 0/15               | 0/15 |
| 1              | Public       | 5.9 ± 0.81               | 6.4 ± 0.88 | -0.2 ± 1.68                        | 3.5 ± 2.13               | 3.2 ± 2.77 | -0.6 ± 1.01                       | 4.6 ± 1.63               | 5.3 ± 1.95 | 0.3 ± 2.14                        | 1/21               | 0/21 |
| 2              | Private      | 5.0 ± 1.47               | 4.1 ± 0.82 | -1.1 ± 1.68                        | 1.3 ± 0.00               | 1.3 ± 0.00 | 0.0 ± 0.00                        | 3.1 ± 2.11               | 2.5 ± 1.84 | -1.8 ± 2.00                       | 0/9                | 0/24 |
| 2              | Public       | 4.5 ± 1.54               | 4.8 ± 2.15 | -1.2 ± 2.88                        | 2.6 ± 1.36               | 1.9 ± 0.83 | -0.3 ± 3.74                       | 3.7 ± 1.47               | 3.4 ± 1.54 | 0.8 ± 3.83                        | 0/9                | 0/24 |
| Total after T1 |              | 5.9 ± 1.02               | 6.0 ± 1.17 | -0.2 ± 2.13                        | 3.6 ± 2.13               | 2.3 ± 1.52 | -0.6 ± 1.79                       | 4.7 ± 1.44               | 4.7 ± 1.44 | 0.0 ± 2.26                        | 1/36               | 0/36 |
| Total after T2 |              | 4.6 ± 1.55               | 4.7 ± 1.85 | -1.1 ± 2.13                        | 2.4 ± 1.63               | 1.6 ± 0.42 | 0.0 ± 2.94                        | 3.5 ± 1.62               | 3.1 ± 2.05 | 0.0 ± 3.12                        | 0/33               | 0/33 |

The AMC, EB, and PS values are provided as median values (log<sub>10</sub> CFU/g) and standard deviations. The presence (+) or absence (-) of *Campylobacter* spp. in water samples identified by the multiplex PCR assay. MAD, median absolute deviation.

ratios between poultry farms with private and public water supplies were not significantly different, we observed higher median log<sub>10</sub> reduction of AMC, EB, and PS counts at poultry farms with private water supply. During T2 water line treatment higher median log<sub>10</sub> reduction was observed for AMC and EB counts at poultry farms with public water supply, while higher median log<sub>10</sub> reduction for PS counts was observed in poultry farms with private water supply.

Out of the 14 poultry farms assessed, five farms exhibited microbial counts below the acceptable microbial limit (<4.0 log<sub>10</sub> CFU/ml) subsequent to the T1 water line treatment (Figures 2A–C). Among these farms, three had a private water supply, while the remaining two had public water supplies. Notably, poultry farm 7, which had a public water supply, exhibited an AMC count below the maximum acceptable microbial limit in both BT and corresponding AT water sample. Furthermore, 11 poultry farms exhibited EB counts below the maximum acceptable microbial limit. Of these, nine poultry farms demonstrated EB counts below the microbial limit in both BT and corresponding AT samples. Additionally, among 14 poultry farms examined, a total of eight poultry farms exhibited PS counts below the microbial limit. Out of these, four poultry farms demonstrated PS counts below the microbial limit in both BT and corresponding AT samples. Among the poultry farms that underwent two samplings, poultry farms 12 and 13 exhibited AMC and PS counts exceeding the microbial limit in one of the sampling events. Furthermore, poultry farm 12 demonstrated EB counts above the microbial limit on one of two sampling occasions.


During T2 waterline treatment AMC counts below the microbial limit were observed in six out of 14 poultry farms (Figures 3A–C). Of these, two poultry farms demonstrated AMC count below the microbial limit in both BT and corresponding AT samples (Figure 3B). EB counts below the microbial limit were observed in 12 out of 14 poultry farms, and among them, nine poultry farms had EB counts below the microbial limit in both the BT and corresponding AT samples. Similarly, PS counts below the microbial limit were observed in ten from 14 poultry farms, and among them, three poultry farms demonstrated PS counts below the microbial limit in BT and corresponding AT samples. Among the poultry farms subjected to two samplings, poultry farm 18 demonstrated AMC, PS and EB counts below the microbial limit during one of the sampling occasions. However, after second sampling, the AMC load in water samples exceeded the microbial limit. Notably, the PS and EB counts remained below the microbial limit during both sampling occasions.

### 3.2 Bacterial isolate identification in poultry drinking water


Isolate taxonomic assignment was performed using partial sequencing of 16S rRNA gene. In the present study, isolate sequences showed ≥97.0% similarity to the reference sequence in the NCBI database. In BT samples, 123 isolates corresponded to 24 genera and 55 species, while in AT samples, the 113 isolates corresponded to 22 genera and 40 species. Further analysis of bacterial isolates revealed that in BT and AT samples, 43.1% ( $n=41$  isolates) and 36.3% ( $n=53$  isolates) of sequenced isolates were assigned to OP, found in 29/36 BT and 17/36 AT samples (Table 2). The isolates from BT samples contained OP represented by 16 genera and 19 species, while isolates from AT samples contained OP represented by 12 genera and 12 species OP. Furthermore, *C. jejuni* was detected using multiplex PCR in a single BT water sample from a poultry farm with a public water supply.

During the T2 water line treatment, 139 isolates in the BT corresponded to 26 genera and 46 species, whereas 96 isolates in AT samples corresponded to 21 genera and 33 species (Table 2). Among the sequenced isolates, 30.9% ( $n=43$  isolates) and 33.3% ( $n=33$  isolates) corresponded to OP, isolated from 20/33 BT and 14/33 AT samples, respectively. The OP in the BT samples comprised 10 genera, and 14 species, while the OP in the AT samples comprised 11 genera and 14 species. No *Campylobacter* spp. were detected in poultry drinking water samples during the T2 water line treatment.

Figures 4A,B represents the taxonomic classification of assigned isolate sequences at phylum, and genus level. The predominant phyla in BT and AT samples were *Pseudomonadota*, followed by *Bacillota*, *Actinomycetota*, and *Bacteroidota* (Figure 4A). The frequently isolated genera during both T1 and T2 water line treatment in BT and AT samples were *Aeromonas*, *Bacillus*, *Citrobacter*, *Enterobacter*, *Pseudomonas* and *Stenotrophomonas* (Figure 4B). Among these, *Pseudomonas* (BT 38.2%; AT 32.7%) and *Bacillus* (BT, 13.0%, AT, 11.5%) were most commonly observed genera during T1 water line treatment. Similarly, during T2 water line treatment, *Pseudomonas* (BT, 31.7%; AT, 33.3%) and *Bacillus* (BT, 10.1%; AT, 11.5%) were predominant genera in BT and AT samples. The Figure 4B depicts the percentage identification of other observed genera during T1 and T2 water line treatments. The majority of sequenced isolates classified as OP in BT and AT samples during T1 and T2 water belonged to the *Pseudomonas* spp., followed by *Stenotrophomonas* spp., *Citrobacter* spp., *Ochrobactrum* spp., and *Acinetobacter* spp. (Figure 4C).



**FIGURE 2**  
 Log<sub>10</sub> transformed average fold changes (before/after waterline treatment) obtained from aerobic mesophilic counts (AMC) (A), *Enterobacteriaceae* (EB) (B), and *Pseudomonadaceae* (PS) (C) in poultry drinking water. The x-axis indicates the comparison between poultry farms with private and public water supply (WS) after T1 waterline treatment. The y-axis shows the log<sub>10</sub> AMC, EB, and PS count ratio. The log<sub>10</sub> AMC, EB, and PS ratio was not significantly different between poultry farms with private and public water supply. No significant differences were observed in the AMC, EB, and PS log<sub>10</sub> ratio after T1 waterline treatment between poultry farms with private and public WS.



**FIGURE 3** Log<sub>10</sub> transformed average fold changes (before/after waterline treatment) obtained from aerobic mesophilic counts (AMC) (A), *Enterobacteriaceae* (EB) (B), and *Pseudomonadaceae* (PS) (C) in poultry drinking water. The x-axis indicates the comparison between poultry farms with private and public water supply (WS) after T2 waterline treatment. The y-axis shows the log<sub>10</sub> AMC, EB, and PS count ratio. The log<sub>10</sub> AMC, EB, and PS ratio was not significantly different between poultry farms with private and public water supply. No significant differences were observed in the AMC, EB, and PS log<sub>10</sub> ratio after T2 waterline treatment between poultry farms with private and public WS.

**TABLE 2** An overview of the number of isolate sequences assigned to the different phyla and genera using similarity cut-off of ≥97.0% after partial sequencing of 16S rRNA gene.

| Sampling timepoint and isolate number                | T1                |              | T2                |             |
|--|-------------------|--------------|-------------------|-------------|
|  | Isolate diversity |              | Isolate diversity |             |
|  | BT (n = 123)      | AT (n = 113) | BT (n = 139)      | AT (n = 93) |
|  | n                 | n            | n                 | n           |
| Phylum   | 4                 | 3            | 4                 | 3           |
| Genus  | 24                | 18           | 26                | 21          |
| Opportunistic pathogens (≥97.0% sequence similarity) | n=53              | 53           | 43                | 33          |

The assigned bacterial isolate sequences encompass the classification of opportunistic pathogens present in water samples collected before (BT) and after (AT) the T1 and T2 water line treatment.


Furthermore, isolates of *Enterobacter* spp. and *Klebsiella* spp. genera were isolated during T1 and T2 sampling. The *Pseudomonas* spp. isolates identified as OP were most frequently observed bacteria sequences during both T1 (BT, 22/123 isolates; AT, 10/113 isolates) and T2 (BT, 13/139 isolates; AT, 10/96 isolates) sampling.

Before and after the T1 water line treatment, *Pseudomonas* spp. was isolated from BT and AT samples in 12/14 and 9/14 poultry farms, respectively (Table 3). Isolate sequences of OPs were detected in BT samples of 11 out of 14 poultry farms and in AT samples of 9 out of 14 poultry farms. Among the frequently observed genera before and after T2 treatment, the genus *Pseudomonas* was isolated from the BT and AT samples in 12 out of 14 poultry farms and 9 out of 14 poultry farms, respectively (Table 4). The OP were observed in 10 out of 14 poultry farms in BT samples and in 9 out of 14 poultry farms in AT samples after T2 water line treatment.

### 3.3 Antibiotic susceptibility patterns of bacterial isolates obtained from poultry drinking water

The susceptibility of bacterial isolates recovered from BT (n = 14) and AT (n = 16) water samples during T1 and T2 water line treatments to 18 antibiotic agents commonly used in poultry production was evaluated using Avian AVIAN1F Vet AST susceptibility plates (Table 5). The goal was to investigate AMR in the most frequently isolated OP isolates, including isolates belonging to *Pseudomonas* spp., *Stenotrophomonas* spp., *Ochrobactrum* spp., as well as AMR in specific waterborne OP important to human health, such as *Aeromonas* spp., *Citrobacter* spp., *Enterobacter* spp., and *Klebsiella* spp.





**FIGURE 4** Taxonomic classification of isolates based on partial sequencing of 16S rRNA gene on phylum (A), genus (B), and opportunistic pathogens (C) level in water samples during T1 and T2 waterline treatments. Sequence similarity cut-off of  $\geq 97.0\%$  was applied for assignment of isolate sequences (1,040 to 1,120 bp fragment) to type strain was applied. (C) The bacterial sequences that were isolated from water samples one to two times are indicated by the grey color.

The highest level of AMR was observed against spectinomycin and sulfadimethoxin (90.0%; 27/30 isolates each), followed by ceftiofur (83.3%; 25/30 isolates), florfenicol (66.6%; 20/30 isolates), and neomycin (53.5%, 16/30 isolates). Further, some isolates were resistant to enrofloxacin (23.3%; 13/30 isolates), trimethoprim-sulfamethoxazole (23.1%; 3/13 isolates), sulfathiazole (20.0%; 6/30

isolates), streptomycin (16.7%, 5/30 isolates), gentamicin (13.3%; 4/30 isolates), and trimethoprim-sulfamethoxazole (10.0%, 3/30 isolates).

The MDR was exhibited among the isolates of *Pseudomonas* spp. (17/17 isolates), and *Stenotrophomonas* spp. (1/3 isolates), *Ochrobactrum* spp. (4/4 isolates), *Citrobacter* spp. (2/2 isolates), and *Enterobacter* spp. (1/2 isolates). All *Pseudomonas* spp. isolates showed

**TABLE 3** The isolate diversity in poultry drinking water samples was assessed using partial sequencing of 16S rRNA gene of cultured isolates collected during chemical waterline treatment with 4.0 ppm active ClO<sub>2</sub> waterline treatment (T1) at poultry farms.

|                                     | Water supply            | Waterline treatment (T1) |         |         |         |         |         |        |        |        |        |        |                  |        |        | Per sample isolation |             |
|-------------------------------------|-------------------------|--------------------------|---------|---------|---------|---------|---------|--------|--------|--------|--------|--------|------------------|--------|--------|----------------------|-------------|
|                                     |                         | Private                  | Private | Private | Private | Private | Private | Public | Public | Public | Public | Public | Public           | Public | Public | BT<br>(n = 36)       | AT (n = 36) |
|                                     |                         | 4                        | 6       | 8       | 10      | 11      | 12      | 1      | 2      | 3      | 5      | 7      | 9                | 13     | 14     |                      |             |
| Poultry farm                        | BT/AT                   | BT/AT                    | BT/AT   | BT/AT   | BT/AT   | BT/AT   | BT/AT   | BT/AT  | BT/AT  | BT/AT  | BT/AT  | BT/AT  | BT/AT            | BT/AT  | BT/AT  | BT/AT                |             |
| Phylum (n = 4)                      | Genus (n = 29)          |                          |         |         |         |         |         |        |        |        |        |        |                  |        |        |                      |             |
| <i>Pseudomonadota</i> (n = 20)      | <i>Achromobacter</i>    |                          |         |         |         |         |         |        |        |        |        |        | 1/1 <sup>1</sup> |        |        | 1                    | 1           |
|                                     | <i>Acinetobacter</i>    | 1/0                      | 1/0     | 2/0     |         |         |         |        | 4/0    |        |        |        |                  |        |        | 5                    | 0           |
|                                     | <i>Aeromonas</i>        | 0/1                      |         |         | 0/1     |         |         | 1/0    | 2/9    | 1/0    | 0/2    |        |                  | 2/0    |        | 5                    | 5           |
|                                     | <i>Atlantibacter</i>    |                          |         |         |         |         |         |        |        |        |        |        |                  | 0/1    |        | 0                    | 1           |
|                                     | <i>Campylobacter</i>    |                          |         |         |         |         |         |        |        |        |        |        |                  | 2/0    |        | 1                    | 0           |
|                                     | <i>Citrobacter</i>      |                          |         |         |         |         |         |        |        | 3/5    | 1/0    |        |                  | 2/0    |        | 2                    | 3           |
|                                     | <i>Comamonas</i>        |                          | 0/2     |         |         |         | 1/0     |        |        |        |        |        |                  | 0/1    |        | 1                    | 2           |
|                                     | <i>Enterobacter</i>     |                          | 2/0     |         |         |         | 1/1     |        |        | 0/1    |        |        |                  | 2/10   | 1/0    | 4                    | 6           |
|                                     | <i>Escherichia</i>      |                          | 0/1     |         |         |         |         |        |        |        |        |        |                  |        |        | 0                    | 1           |
|                                     | <i>Klebsiella</i>       |                          |         |         |         |         |         |        |        | 2/1    | 1/0    |        |                  |        |        | 2                    | 1           |
|                                     | <i>Kluyvera</i>         |                          |         |         | 1/0     |         |         |        |        |        |        |        |                  |        |        | 1                    | 0           |
|                                     | <i>Lecleercia</i>       |                          |         | 1/0     |         |         | 1/3     |        |        |        |        |        |                  |        |        | 2                    | 1           |
|                                     | <i>Ochrobactrum</i>     |                          |         |         | 0/1     |         |         |        | 0/5    |        |        |        |                  |        | 1/0    | 1                    | 3           |
|                                     | <i>Pantonea</i>         |                          |         |         | 0/1     |         |         |        |        |        |        |        |                  |        |        | 0                    | 1           |
|                                     | <i>Phytobacter</i>      |                          |         |         |         |         |         |        |        |        |        |        |                  | 0/1    | 1/0    | 1                    | 1           |
|                                     | <i>Pigmentiphaga</i>    |                          |         |         |         |         |         | 1/0    |        |        |        |        |                  |        |        | 1                    | 0           |
|                                     | <i>Pseudomonas</i>      | 3/3                      | 10/7    |         | 2/2     | 3/0     | 3/7     | 2/2    | 3/0    | 2/6    | 7/6    | 2/0    | 7/1              | 3/3    |        | 26                   | 18          |
|                                     | <i>Raoultella</i>       |                          | 1/0     |         |         |         |         |        |        |        |        |        |                  |        |        | 1                    | 0           |
| <i>Rhizobium</i>                    |                         |                          |         |         |         |         |         |        |        |        |        |        | 0/1              |        | 0      | 1                    |             |
| <i>Stenotrophomonas</i>             |                         |                          | 1/0     |         |         | 3/1     |         | 0/1    |        | 0/5    |        | 3/2    | 1/1              | 1/0    | 5      | 10                   |             |
| <i>Actinomycetota</i> (n = 2)       | <i>Brachybacterium</i>  |                          |         |         |         |         |         |        |        |        |        |        | 1/0              |        | 1      | 0                    |             |
|                                     | <i>Microbacterium</i>   |                          |         | 1/0     |         |         |         |        |        |        |        |        |                  |        | 1      | 0                    |             |
| <i>Bacillota</i> (n = 5)            | <i>Aerococcus</i>       | 1/0                      |         |         |         |         |         |        |        |        |        |        |                  |        | 1      | 0                    |             |
|                                     | <i>Bacillus</i>         |                          | 1/0     | 1/0     | 2/2     |         |         | 4/6    | 4/1    | 4/3    |        |        |                  | 0/1    | 10     | 7                    |             |
|                                     | <i>Lysinibacillus</i>   |                          |         |         |         | 1/0     |         |        |        |        |        |        |                  |        | 1      | 0                    |             |
|                                     | <i>Planococcus</i>      |                          |         |         |         |         |         | 1/0    |        |        |        |        |                  |        | 1      | 0                    |             |
|                                     | <i>Staphylococcus</i>   |                          | 1/0     |         | 1/1     | 3/0     |         | 1/0    |        |        |        |        | 3/0              | 2/0    | 10     | 1                    |             |
| <i>Bacteroidota</i> (n = 2)         | <i>Chryseobacterium</i> |                          |         |         | 0/1     |         |         |        |        |        |        |        |                  |        | 0      | 1                    |             |
|                                     | <i>Sphingobacterium</i> |                          | 1/0     |         |         |         |         |        |        |        |        |        |                  |        | 1      | 0                    |             |
| Bacterial diversity on poultry farm |                         | 3/2                      | 6/3     | 5/0     | 4/6     | 2/1     | 7/4     | 3/1    | 6/4    | 5/5    | 4/4    | 1/0    | 5/6              | 6/4    | 5/1    |                      |             |
| Identified opportunistic pathogens  |                         | 1/0                      | 4/1     | 0/0     | 2/1     | 0/0     | 4/3     | 1/1    | 2/2    | 3/2    | 4/3    | 0/0    | 4/3              | 1/1    | 0/0    | 29/36                | 17/36       |

The occurrence of each genus in sample collected before treatment (BT, n = 36) and after treatment (AT, n = 36) was determined. The percentage of isolate occurrence was calculated based on cultured isolates from BT (n = 123) and AT (n = 113) samples. The isolate diversity at each poultry farm was evaluated in both BT and AT samples, and the presence of opportunistic pathogens was also determined. <sup>1</sup>Number of bacterial isolates isolated from BT and AT samples.

TABLE 4 The isolate diversity in poultry drinking water samples was assessed using partial sequencing of 16S rRNA gene of cultured isolates collected during combined chemical (3.0% peroxyacetic acid [PAA] and 4.0 ppm active ClO<sub>2</sub>) with mechanical (purging of waterlines with a high-pressure air pump) waterline treatment (T2) in poultry farms.

|                                   | Water supply             | Waterline treatment (T2) |         |         |         |                  |        |        |        |        |        |        |        |        |        | Per sample isolation |                |
|-----------------------------------|--------------------------|--------------------------|---------|---------|---------|------------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|----------------------|----------------|
|                                   |                          | Private                  | Private | Private | Private | Private          | Public | Public | Public | Public | Public | Public | Public | Public | Public | BS<br>(n = 33)       | AS<br>(n = 33) |
|                                   |                          | Poultry farms            | 4       | 10      | 11      | 15               | 19     | 1      | 2      | 3      | 5      | 7      | 14     | 16     | 17     |                      |                |
| Water sample                      | BT/AT                    | BT/AT                    | BT/AT   | BT/AT   | BT/AT   | BT/AT            | BT/AT  | BT/AT  | BT/AT  | BT/AT  | BT/AT  | BT/AT  | BT/AT  | BT/AT  | BT/AT  |                      |                |
| Pylum (n = 4)                     | Genus (n = 33)           |                          |         |         |         |                  |        |        |        |        |        |        |        |        |        |                      |                |
| <i>Pseudomonadota</i><br>(n = 24) | <i>Acidovorax</i>        |                          |         |         |         | 0/2 <sup>1</sup> |        |        |        |        |        | 2/0    |        |        | 0/1    | 5                    | 4              |
|                                   | <i>Acinetobacter</i>     |                          |         | 1/0     |         |                  | 0/1    | 1/0    |        |        | 0/2    |        |        |        |        | 2                    | 2              |
|                                   | <i>Aeromonas</i>         | 0/1                      |         |         |         |                  |        |        | 4/1    |        |        |        | 1/0    |        | 6/0    | 7                    | 2              |
|                                   | <i>Atlantibacter</i>     |                          |         |         |         |                  | 0/1    |        |        |        |        |        |        |        |        | 0                    | 1              |
|                                   | <i>Brevundomonas</i>     |                          |         |         | 0/1     |                  |        |        |        |        |        |        |        |        |        | 0                    | 1              |
|                                   | <i>Buttiauxella</i>      |                          |         |         |         |                  |        | 1/0    |        |        |        |        |        |        |        | 1                    | 0              |
|                                   | <i>Chromobacterium</i>   |                          |         |         |         |                  |        |        |        |        |        |        | 2/0    |        |        | 2                    | 0              |
|                                   | <i>Citrobacter</i>       | 0/2                      |         |         | 0/3     |                  | 1/0    | 0/1    | 3/0    | 4/0    | 1/1    |        |        |        | 1/0    | 8                    | 4              |
|                                   | <i>Comamonas</i>         |                          |         |         |         |                  |        |        | 0/4    |        |        |        |        |        |        | 0                    | 3              |
|                                   | <i>Cupriavidus</i>       |                          |         | 1/0     | 1/0     |                  |        |        |        |        |        |        | 1/0    |        |        | 3                    | 0              |
|                                   | <i>Enterobacter</i>      |                          |         |         |         | 1/0              |        |        |        | 1/3    |        |        |        |        | 0/4    | 2                    | 3              |
|                                   | <i>Janthinobacterium</i> |                          |         |         |         |                  |        | 0/1    | 0/3    |        |        |        |        |        |        | 0                    | 3              |
|                                   | <i>Klebsiella</i>        |                          |         |         | 0/1     |                  | 3/0    |        |        |        | 2/0    |        |        |        |        | 2                    | 1              |
|                                   | <i>Kluyvera</i>          |                          |         |         |         |                  | 0/1    |        |        |        |        |        |        |        |        | 0                    | 1              |
|                                   | <i>Moraxella</i>         |                          |         |         |         |                  |        |        | 0/2    |        |        |        |        |        |        | 0                    | 1              |
|                                   | <i>Ochrobactrum</i>      |                          |         |         |         |                  |        | 2/2    |        |        |        |        |        |        | 2/0    | 3                    | 1              |
|                                   | <i>Pantonea</i>          |                          |         |         |         |                  |        |        | 0/2    |        |        |        |        |        |        | 0                    | 1              |
|                                   | <i>Phytobacter</i>       |                          |         |         |         |                  | 1/0    |        |        |        |        |        |        |        |        | 1                    | 0              |
|                                   | <i>Pigmentiphaga</i>     |                          |         |         |         |                  |        |        |        |        |        |        |        |        | 1/0    | 1                    | 0              |
|                                   | <i>Pseudacromonas</i>    |                          |         |         |         |                  |        |        |        |        |        |        | 1/0    |        |        | 1                    | 0              |
| <i>Pseudomonas</i>                | 1/1                      | 5/0                      | 1/0     | 6/0     | 4/1     | 0/2              | 2/1    | 0/8    | 5/6    | 3/3    | 5/0    | 1/0    | 4/3    | 7/7    | 19     | 16                   |                |
| <i>Raoultella</i>                 |                          |                          |         |         |         |                  | 1/0    |        |        |        |        | 4/0    |        |        | 3      | 0                    |                |
| <i>Stenotrophomonas</i>           |                          |                          | 2/0     | 1/0     | 2/1     |                  |        | 6/0    | 0/5    |        |        |        |        |        | 7      | 3                    |                |
| <i>Variovorax</i>                 | 2/0                      |                          | 2/0     | 1/0     | 0/1     |                  |        |        |        |        |        |        |        | 1/2    | 5      | 2                    |                |
| <i>Actinomycetota</i><br>(n = 2)  | <i>Brachybacterium</i>   |                          |         |         |         |                  |        |        |        |        |        |        | 2/0    |        | 1      | 0                    |                |
|                                   | <i>Microbacterium</i>    | 1/0                      |         | 1/0     |         |                  |        |        |        |        |        |        |        | 1/0    | 3      | 0                    |                |
| <i>Bacillota</i> (n = 4)          | <i>Bacillus</i>          | 1/1                      | 3/0     |         | 1/0     |                  | 6/5    | 2/0    | 1/5    |        |        |        |        |        | 10     | 8                    |                |
|                                   | <i>Jeotgalicoccus</i>    |                          |         |         |         |                  |        |        |        |        |        |        |        | 1/1    | 1      | 1                    |                |
|                                   | <i>Staphylococcus</i>    |                          |         |         |         |                  | 1/0    |        |        |        |        |        | 2/0    | 1/1    | 3      | 1                    |                |
|                                   | <i>Trichococcus</i>      |                          |         |         |         |                  |        |        | 1/0    |        |        |        |        |        | 1      | 0                    |                |

(Continued)

TABLE 4 (Continued)

|                                     | Waterline treatment (T2) |         |         |         |         |         |        |        |        |        |        |        | Per sample isolation |                |                |
|-------------------------------------|--------------------------|---------|---------|---------|---------|---------|--------|--------|--------|--------|--------|--------|----------------------|----------------|----------------|
|                                     | Private                  | Private | Private | Private | Private | Private | Public | Public | Public | Public | Public | Public | Public               | BS<br>(n = 33) | AS<br>(n = 33) |
| Water supply                        | 4                        | 10      | 11      | 15      | 19      | 1       | 2      | 3      | 5      | 7      | 14     | 16     | 17                   | 18             |                |
| Poultry farms                       |                          |         |         |         |         |         |        |        |        |        |        |        |                      |                |                |
| Water sample                        | BT/AT                    | BT/AT   | BT/AT   | BT/AT   | BT/AT   | BT/AT   | BT/AT  | BT/AT  | BT/AT  | BT/AT  | BT/AT  | BT/AT  | BT/AT                | BT/AT          | BT/AT          |
| Genus (n = 33)                      |                          |         |         |         |         |         |        |        |        |        |        |        |                      |                |                |
| Bacteroidata (n = 3)                | <i>Chryseobacterium</i>  |         | 1/0     |         |         |         |        |        |        |        |        |        |                      | 1              | 0              |
|                                     | <i>Flavobacterium</i>    |         |         |         |         | 0/1     |        | 0/2    |        |        | 1/0    |        |                      | 1              | 3              |
|                                     | <i>Pedobacter</i>        |         |         |         |         |         |        |        | 1/0    |        |        |        |                      | 1              | 0              |
| Bacterial diversity on poultry farm | 4/4                      | 1/0     | 9/0     | 4/3     | 4/4     | 3/5     | 8/8    | 4/5    | 6/4    | 2/2    | 4/0    | 5/0    | 3/1                  | 9/6            |                |
| Identified opportunistic pathogens  | 0/1                      | 0/0     | 2/0     | 1/1     | 2/0     | 1/1     | 2/2    | 3/2    | 3/2    | 1/1    | 1/0    | 0/0    | 1/1                  | 2/3            | 14/33          |

The occurrence of each genus in sample collected before treatment (BT, n = 33) and after treatment (AT, n = 33) was determined. The percentage of isolate occurrence was calculated based on cultured isolates from BT (n = 139) and AT (n = 96) samples. The isolate diversity at each poultry farm was evaluated in both BT and AT samples, and the presence of opportunistic pathogens was also determined. Number of bacterial isolates isolated from BT and AT samples.

resistance patterns exhibiting resistance to a minimum of four and a maximum of eight antibiotics. Tested *Stenotrophomonas* spp. isolates also demonstrated resistance patterns to a minimum of four and a maximum of six antibiotics. All tested *Ochrobactrum* spp. were resistant to four antibiotics. The isolates of *Citrobacter* spp. were resistant to six antimicrobial classes and nine different antibiotics. The isolates of *Enterobacter* spp. showed resistance patterns to a minimum of two and a maximum of four antibiotics. The isolates of *Klebsiella* spp. were resistant to five antibiotics, while *Aeromonas* spp. isolate was susceptible to all tested antibiotic agents.

### 4 Discussion

Providing poultry with water that meets the highest quality standards is essential to ensure the safety and quality of the products derived from these animals. The presence of high microbial loads and biofilms in the drinking water lines can have a negative effect on poultry health and performance (14). Moreover, when health issues arise within a poultry flock, antibiotics are often administered through drinking water. This practice increases the risk of antibiotic resistance within poultry farms, presenting a potential threat to both animal and human health (12).

We assessed microbial quality of poultry drinking water at the end of the drinking line based on established limits from previous studies, where AMC, EB, and PS counts below 4.0 log<sub>10</sub> CFU/ml were deemed acceptable (4, 7, 12). At the end of the fattening period, AMC exceeded acceptable limits in most poultry farms tested, with similar trends observed for PS counts. However, EB remained within acceptable levels in the majority of farms. Environmental factors, such as ambient temperatures (±25°C), low water flow rates, pipeline installation type, and feed additives (often mixed with glucose) provided ample nutrients for bacteria, contributing to a high microbial load at the end of the fattening period (42). Poultry farms opt to chlorinate and/or acidify their drinking water systems due to the easy application, cost-effectiveness, and broad antimicrobial properties of these treatment systems (12). Additionally, mechanical cleaning helps remove biofilm from surfaces inside the drinking water system. Surprisingly, plate count analysis did not show a significant reduction of microbial load (AMC, EB, and PS counts) in AT samples after chemical water line treatment (T1) or combined chemical with mechanical treatment (T2). Unlike previous reports associating poultry farms with a private water supply with elevated microbial loads, we did not observe significant differences in microbial load between poultry farms with private or public water supplies (43). The microbial counts observed in our study were similar to those found on surfaces inside poultry house drinking water systems, which were typically above 6.0 log<sub>10</sub> CFU (12). This suggests a limited disinfection effectiveness likely due to low concentration of applied disinfectant. Despite mechanical cleaning and subsequent disinfection, high microorganism levels persisted in the water lines, indicating that the disinfectant concentration post-mechanical treatment was insufficient to eliminate the majority of microorganisms. However, our study focused solely on microbiological parameters, overlooking vital factors such as water hardness, pH, temperature, and free ClO<sub>2</sub> residues within the water lines. This limited our ability to comprehensively evaluate the

TABLE 5 Antimicrobial resistance among bacterial isolates before (BT) and after (AT) waterline treatment to a panel of veterinary antimicrobials commonly used in the poultry production.

| Opportunistic pathogens <sup>2</sup> | Treatment <sup>3</sup> | Time-point <sup>4</sup> | Isolates (n) | Antimicrobial class <sup>1</sup> (in µg/ml): |            |            |               |                  |                |                   |           |              |      |                                |
|--------------------------------------|------------------------|-------------------------|--------------|--|------------|------------|---------------|------------------|----------------|-------------------|-----------|--------------|------|--------------------------------|
|                                      |                        |                         |              | Aminoglycosides                              |            |            |               | Fluoroquinolones | Cephalosporins | Tetracyclines     | Phenicols | Sulfonamides |      | Diaminopyrimidine/sulfonamides |
|                                      |                        |                         |              | GEN<br>≥8                                    | SPE<br>≥64 | NEO<br>≥32 | STR<br>≥1,024 | ENR<br>≥2/1      | XNL<br>≥4      | TET and OXY<br>≥8 | FFN<br>≥8 | SDM<br>≥256  | STZ  | SXT<br>≥2/38                   |
| <i>Citrobacter</i> spp.              | 1                      | BT                      | 1            | 0/1  | 1/1        | 1/1        | 0/1           | 1/1              | 0/1            | 1/1               | 1/1       | 1/1          | 1/1  | 1/1                            |
|                                      | 2                      | AT                      | 1            | 0/1  | 1/1        | 1/1        | 0/1           | 1/1              | 0/1            | 1/1               | 1/1       | 1/1          | 1/1  | 1/1                            |
| <i>Enterobacter</i> spp.             | 1                      | AT                      | 1            | 0/1  | 0/1        | 0/1        | 0/1           | 0/1              | 1/1            | 1/1               | 1/1       | 1/1          | 0/1  | 0/1                            |
|                                      | 2                      | AT                      | 1            | 0/1  | 0/1        | 0/1        | 0/1           | 0/1              | 0/1            | 1/1               | 0/1       | 1/1          | 0/1  | 0/1                            |
| <i>Klebsiella</i> spp.               | 2                      | BT                      | 1            | 1/1  | 1/1        | 1/1        | 0/1           | 0/1              | 0/1            | 0/1               | 0/1       | 1/1          | 1/1  | 0/1                            |
| <i>Ochrobactrum</i> spp.             | 1                      | AT                      | 1            | 0/1  | 1/1        | 1/1        | 0/1           | 0/1              | 1/1            | 0/1               | 0/1       | 1/1          | 0/1  | 0/1                            |
|                                      | 2                      | BT                      | 2            | 0/2  | 2/2        | 2/2        | 0/2           | 0/2              | 2/2            | 0/2               | 0/2       | 2/2          | 0/2  | 0/2                            |
|                                      | 2                      | AT                      | 1            | 0/1  | 1/1        | 1/1        | 0/1           | 0/1              | 1/1            | 0/1               | 0/1       | 1/1          | 0/1  | 0/1                            |
| <i>Pseudomonas</i> spp.              | 1                      | BT                      | 8            | 0/8  | 8/8        | 0/8        | 1/8           | 3/7              | 8/8            | NA <sup>5</sup>   | 8/8       | 8/8          | 1/8  | NA                             |
|                                      | 1                      | AT                      | 6            | 0/6  | 6/6        | 6/6        | 2/6           | 2/6              | 6/6            |                   | 6/6       | 1/6          |      |                                |
|                                      | 2                      | BT                      | 1            | 1/1  | 1/1        | 0/1        | 0/1           | 0/1              | 1/1            |                   | 1/1       | 0/1          |      |                                |
|                                      | 2                      | AT                      | 2            | 0/2  | 2/2        | 0/2        | 1/2           | 0/2              | 2/2            |                   | 2/2       | 0/2          |      |                                |
| <i>Aeromonas</i> spp.                | 2                      | AT                      | 1            | 0/1  | 0/1        | 0/1        | 0/1           | 0/1              | 0/1            | 0/1               | 0/1       | 0/1          | 0/1  | 0/1                            |
| <i>Stenotrophomonas</i> spp.         | 1                      | AT                      | 1            | 1/1  | 1/1        | 1/1        | 1/1           | 0/1              | 1/1            | NA                | 0/1       | 0/1          | 0/1  | 0/1                            |
|                                      | 2                      | BT                      | 1            | 0/1  | 1/1        | 1/1        | 0/1           | 0/1              | 1/1            |                   | 0/1       | 1/1          | 1/1  | 1/1                            |
|                                      | 2                      | AT                      | 1            | 1/1  | 1/1        | 1/1        | 0/1           | 0/1              | 1/1            |                   | 0/1       | 0/1          | 0/1  | 0/1                            |
| BT (n/N) <sup>6</sup>                |                        |                         | 14           | 2/14   | 14/14      | 5/14       | 1/14          | 4/14             | 12/14          | 1/14              | 10/14     | 14/14        | 4/14 | 2/14                           |
| AT (n/N)                             |                        |                         | 16           | 2/16   | 13/16      | 11/16      | 4/16          | 4/16             | 13/16          | 3/16              | 10/16     | 13/16        | 2/16 | 1/16                           |
| Total (n/N)                          |                        |                         | 30           | 4/30   | 27/30      | 16/30      | 5/30          | 7/30             | 25/30          | 4/10              | 20/30     | 27/30        | 6/30 | 3/30                           |

<sup>1</sup>The resistance breakpoints for selected antimicrobial classes represented by the antimicrobial agents in µg/ml for ≥8 gentamicin (GEN); ≥64 spectinomycin (SPE); ≥32 neomycin (NEO); ≥1,024 streptomycin (STR); ≥2/1 enrofloxacin (ENR); ≥4 ceftiofur (XNL); ≥8 tetracycline (TET) and oxytetracycline (OXY); ≥8 florfenicol (FFN); ≥256 sulfadimethoxine (SDM) and sulfathiazole (STZ); ≥2/38 trimethoprim-sulfamethoxazole (STX). <sup>2</sup>Bacteria species identified by partial sequencing of 16S rRNA gene. <sup>3</sup>Waterline treatment type.

<sup>4</sup>Isolate identification in water sample before treatment (BT) and after treatment. <sup>5</sup>NA: not applicable, bacteria have intrinsic resistance against the antimicrobial agent. <sup>6</sup>n/N: number of isolates resistant to particular antimicrobial agent/total isolates tested.

efficiency of the 4 ppm active ClO<sub>2</sub> and 3% PAA during water line treatments. Previous studies have highlighted the limited effectiveness of water line disinfection practices using oxidizing agents such as chlorine or hydrogen peroxide (12). This limitation primarily arises from applied concentrations being lower than recommended by suppliers, which is in alignment with our observations of high microbial load in AT samples. In addition, inconsistencies were noted in AT water samples among poultry farms that were sampled twice, emphasizing the need for frequent water quality checks in a closed system. Even with the addition of typical concentrations of hydrogen peroxide (25–50 ppm) and free chlorine (2–5 ppm) to poultry drinking water during fattening, biofilm formation was observed in minimally contaminated water (7). Therefore, regular monitoring of microbial water quality, combined with consistent water line treatment during the fattening period, is a crucial aspect of robust biosecurity programs at poultry farms. Moreover, specialized contractors have been noted to achieve more effective water line treatment compared to farmers (42, 44). Finally, Zou et al. (45) demonstrated a significant reduction of *E. coli*, *Salmonella*, *Staphylococcus aureus*, and mold in poultry drinking water after treatment with sodium dichloroisocyanurate, correlating positively with poultry health.

The presence of high microbial load in water samples led to a wide taxonomic variety among isolates in both BT and AT samples, ranging between 18 and 26 genera. While definitive taxonomic conclusions require further extensive studies, the frequent presence of genera such as *Aeromonas*, *Bacillus*, *Citrobacter*, *Enterobacter*, *Pseudomonas* and *Stenotrophomonas*, commonly associated with waste and surface waters, underscores an increased risk to both poultry and human health in this study (19, 46). Identification of genera, including *Pseudomonas*, *Stenotrophomonas*, and *Ochrobactrum*, were in line with the isolates found on surfaces in poultry drinking water system (12). The majority of the identified bacteria found at poultry farms independent of their water supply were OP, specifically those belonging to *Pseudomonas* spp., *Stenotrophomonas* spp., and *Ochrobactrum* spp. The OP belonging to *Pseudomonas* spp. are linked to secondary infections in both poultry and humans. In poultry, these infections can manifest as septicemia, skin lesion infections, and hemorrhagic pneumonia (47). In immunocompromised humans, they can lead to septicemia, pneumonia, and urinary tract infections (48). Previous studies have also emphasized an increased mortality rate in poultry following *P. aeruginosa* OP infection (49, 50). A previous study demonstrated enhanced adhesion to abiotic surfaces, tissue invasion through cytotoxic effects, resistance to 0.2 mg/mL chlorine, and increased AMR among *P. aeruginosa* isolates from water (51). Moreover, *Stenotrophomonas maltophilia* and *Ochrobactrum intermedium* are emerging human environmental pathogens causing infections, primarily in immunocompromised patients (52). *S. maltophilia* and *P. aeruginosa* are often co-isolated from the lungs of cystic fibrosis patients, and previous research findings suggest that *S. maltophilia* modulates the virulence of *P. aeruginosa* in a multispecies biofilm (53). While *S. maltophilia* and *O. intermedium* have been recognized to cause infections in immunocompromised humans, no established link between water quality and disease development in poultry production involving these bacterial species has been reported yet. Nevertheless, notable characteristics of these bacteria, such as resistance to disinfection

and heat, slow growth, and biofilm formation, emphasize the potential risk of poultry and farmer infection through direct contact with drinking water, along with the risk of cross-contamination of chicken meat products during post-slaughter processing.

During T1 water line treatment, *C. jejuni* was detected in one water sample collected before water line treatment at a poultry farm with a public water supply, while other analyzed samples tested negative. The detection of *Campylobacter* spp. in water depends on factors such as sample volume, sample number, and bacterial concentration (54, 55). Furthermore, *Campylobacter* spp. can enter a viable but nonculturable state (VBNC) under environmental stress, potentially hindering growth on conventional culture media due to limited metabolic activity (56). Consequently, *Campylobacter* spp. might have been overlooked in other analyzed water samples due to limitations in the processing method. These limitations include a small sample volume, the absence of water sample filtration, and the potential presence of *Campylobacter* spp. in the VBNC state, which cannot be detected using the ISO-based methods used in the current study. While this approach may have led to missing *Campylobacter* spp., our assessment of bacterial load and diversity in the water samples examined provided a comprehensive insight into both quantitative and qualitative microbial content in poultry drinking water. Notably, previous research emphasizes that a significant presence of *Pseudomonas* spp. in poultry drinking water heightens the risk of *Campylobacter* spp. infection, as *Campylobacter* sp. isolates from poultry can persist for extended periods within *P. aeruginosa* biofilms in drinking water (57–59).

Previous studies have established poultry farms as significant reservoirs of antimicrobial resistance genes, contributing to the emergence of AMR and transmission dynamics of MDR bacteria at the human-animal-environment interface (60–62). Our findings align with these observations, revealing MDR patterns in all tested isolates of both *Pseudomonas* spp. and *Ochrobactrum* spp. isolates from BT and AT water samples. Furthermore, a single *Stenotrophomonas* spp. from BT water sample exhibited MDR pattern. The consistent AMR patterns observed in both BT and AT water samples align with our observations of ineffective water line treatment characterized by limited disinfectant concentrations that allow for the survival and persistence of AMR bacteria within the water lines. The antimicrobials permitted for poultry treatment in Austria at the time of this study include enrofloxacin, doxycycline, trimethoprim-sulfamethoxazole, amoxicillin-clavulanic acid, colistin sulfate, tetracycline, and gentamicin (63–67). For the isolates we utilized in the AST, information or protocols regarding the current or past treatment of poultry on these farms were not available to the authors; therefore a detailed analysis of the potential causes of AMR in these isolates was not possible. The isolates from both BT and AT water samples exhibited increased resistance patterns to spectinomycin, sulfadimethoxin, ceftiofur, florfenicol, and neomycin, likely attributed to their widespread use in poultry health management on farms. This raises concerns, as elevated streptomycin resistance in *E. coli* isolates from broilers in several countries in Europe, including Poland, Germany, Great Britain, France and Spain was previously reported (68). Additionally, resistance to streptomycin and sulfadimethoxin was previously reported in *Salmonella* spp. isolates from poultry farms in Canada and the United States (69–72). Furthermore, these isolates exhibited resistance to ceftiofur and enrofloxacin, both of which are

recognized as top priority critically important antimicrobials by the World Health Organization (73). This antimicrobial resistance raises concerns, as it can be indirectly transmitted through horizontal gene transfer to *E. coli*, *Salmonella* spp., *Campylobacter* spp. and other potential poultry and human pathogens. Heinemann et al. (42) reported isolation of extended-spectrum beta-lactamase-producing bacteria (ESBL) such as *P. aeruginosa*, *Enterobacter* spp., *Klebsiella* spp., and *Acinetobacter baumannii* from poultry drinking water lines and sprinkler systems. ESBL bacteria can hydrolyze extended-spectrum cephalosporins, monobactams, and penicillins and thus lead to elevated morbidity and mortality, further complicating therapeutic choices, particularly among elderly and immunocompromised individuals (74–76). The observed AMR resistance patterns in poultry drinking water isolates highlight the potential for acquiring antimicrobial resistance through water-administered medication, posing a risk and limiting treatment options in both veterinary and human medicine (1, 42, 77–79).

The study emphasizes the persistent challenge of maintaining microbial quality in poultry drinking water. The high microbial load observed is attributed to established microbiota in the water system, resistant to suboptimal disinfectant concentrations used during cleaning. Furthermore, our findings suggest that current poultry treatment and antibiotic usage may elevate the presence of AMR bacteria in drinking water due to inefficient management. Addressing this issue necessitates regular water monitoring, consistent water line treatment, and improved farmer education. Enhancing understanding of biological processes in drinking water systems and microorganism viability can lead to better guidance on herd health and farm productivity. Identifying and mitigating on-farm water quality risks, including assessing waterline technologies affecting microbiota in drinking water and water lines, is essential for controlling pathogen and antibiotic transmission in poultry production.

## 5 Conclusion

In conclusion, the majority of poultry farms in Austria exhibited high microbial loads in drinking water, largely attributed to inadequate water line management practices, including the use of suboptimal disinfectant concentrations and inconsistent treatment. Notably, there were no significant differences observed between chemical and combined chemical and mechanical water line treatments. The prevalent microbiota in poultry included *Pseudomonas* spp., *Stenotrophomonas* spp., and *Ochrobactrum* spp. Moreover, these isolates from both before and after water line treatment samples displayed increased resistance patterns to commonly used antimicrobials to treat bacterial infections in poultry. Our results underscore the need for future studies to consider appropriate water supply management on poultry farms in terms of the One Health approach, to protect public health, and to raise awareness among farmers and veterinarians.

## Data availability statement

The datasets presented in this study can be found in online repositories. The names of the repository/repositories and accession number(s) can be found in the article/Supplementary material.

## Author contributions

AM: Data curation, Writing – original draft, Investigation, Methodology, Software, Visualization. MM: Data curation, Methodology, Writing – review & editing. KW: Data curation, Methodology, Writing – review & editing. AS: Methodology, Writing – review & editing, Investigation. IK: Investigation, Methodology, Writing – review & editing. CF: Writing – review & editing. IL: Formal analysis, Methodology, Writing – review & editing. MW: Conceptualization, Writing – review & editing. BS: Conceptualization, Data curation, Funding acquisition, Project administration, Resources, Supervision, Validation, Writing – original draft, Writing – review & editing.

## Funding

The author(s) declare financial support was received for the research, authorship, and/or publication of this article. The competence center FFOQSI is funded by the Austrian ministries BMVIT, BMDW, and the Austrian provinces Lower Austria, Upper Austria and Vienna within the scope of COMET – Competence Centers for Excellent Technologies. The COMET program is handled by the Austrian Research Promotion Agency FFG.

## Acknowledgments

We would like to thank Egon Singer and the team at the private laboratory HYGIENICUM GmbH Graz for the organization of the sampling and good cooperation. The authors wish to acknowledge and thank Sonja Muri-Klinger for her valuable help in the laboratory. Additionally, we are grateful for the careful input and review of the manuscript text by Patrick-Julian Mester.

## Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

## Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

## Supplementary material

The Supplementary material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fvets.2023.1254442/full#supplementary-material>

## References

- Savin M, Alexander J, Bierbaum G, Hammerl JA, Hembach N, Schwartz T, et al. Antibiotic-resistant bacteria, antibiotic resistance genes, and antibiotic residues in wastewater from a poultry slaughterhouse after conventional and advanced treatments. *Sci Rep.* (2021) 11:16622–11. doi: 10.1038/s41598-021-96169-y
- Statistics Austria. Hühnerproduktion 2021: Zahl der Bruteier Gestiegen, Mehr Schlachtungen; (2020). Available at: <https://www.statistik.at/en/statistics/agriculture-and-forestry/animals-animal-production> (Accessed March 21, 2023)
- King AJ. Water quality and poultry production. *Poult Sci.* (1996) 75:852–3. doi: 10.3382/PS.0750852
- Maharjan P, Clark T, Kuenzel C, Foy MK, Watkins S. On farm monitoring of the impact of water system sanitation on microbial levels in broiler house water supplies. *J Appl Poult Res.* (2016) 25:266–71. doi: 10.3382/JAPR/PFW010
- Pesti GM, Amato SV, Minear LR. Water consumption of broiler chickens under commercial conditions. *Poult Sci.* (1985) 64:803–8. doi: 10.3382/ps.0640803
- Lilliard HS, Thomson JE. Efficacy of hydrogen peroxide as a bactericide in poultry chiller water. *J Food Sci.* (1983) 48:125–6. doi: 10.1111/j.1365-2621.1983.tb14804.x
- Maharjan P, Huff G, Zhang W, Watkins S. Effects of chlorine and hydrogen peroxide sanitation in low bacterial content water on biofilm formation model of poultry brooding house waterlines. *Poult Sci.* (2017) 96:2145–50. doi: 10.3382/PS/PEX009
- Regulation (EC) No 178/2002 of the European Parliament and of the Council. 28 January 2002 Laying Down the General Principles and Requirements of Food Law, Establishing the European Food Safety Authority and Laying Down Procedures in Matters of Food Safety; (2002). Available at: <https://eur-lex.europa.eu/legal-content/EN/ALL/?uri=celex%3A32002R0178> (Accessed March 10, 2021)
- Council Directive 98/83/EC. On the Quality of Water Intended for Human Consumption; (1998). Available at: <https://leap.unep.org/countries/eu/national-legislation/council-directive-9883ec-quality-water-intended-human-consumption> (Accessed March 12, 2021)
- BGBI. II Nr. 100/2007. Gesamte Rechtsvorschrift für Geflügelhygieneverordnung; (2007). Available at: <https://www.ris.bka.gv.at/GeltendeFassung.wxe?Abfrage=Bundesnormen&Gesetzesnummer=20005323> (Accessed June 19, 2023)
- Di Martino G, Piccirillo A, Giacomelli M, Comin D, Gallina A, Capello K, et al. Microbiological, chemical and physical quality of drinking water for commercial turkeys: a cross-sectional study. *Poult Sci.* (2018) 97:2880–6. doi: 10.3382/ps/pey130
- Maes S, Vackier T, Nguyen Huu S, Heyndrickx M, Steenackers H, Sampers I, et al. Occurrence and characterisation of biofilms in drinking water systems of broiler houses. *BMC Microbiol.* (2019) 19:77–15. doi: 10.1186/s12866-019-1451-5
- Mateus-Vargas RH, Kemper N, Volkmann N, Kietzmann M, Meissner J, Schulz J. Low-frequency electromagnetic fields as an alternative to sanitize water of drinking systems in poultry production? *PLoS One.* (2019) 14:e0220302–17. doi: 10.1371/journal.pone.0220302
- Sparks NHC. The role of the water supply system in the infection and control of *Campylobacter* in chicken. *Worlds Poult Sci J.* (2009) 65:459–74. doi: 10.1017/S0043933909000324
- Lehtola MJ, Miettinen IT, Lampola T, Hirvonen A, Vartiainen T, Martikainen PJ. Pipeline materials modify the effectiveness of disinfectants in drinking water distribution systems. *Water Res.* (2005) 39:1962–71. doi: 10.1016/j.watres.2005.03.009
- Mathieu L, Francius G, El Zein R, Angel E, Block JC. Bacterial repopulation of drinking water pipe walls after chlorination. *Biofouling.* (2016) 32:925–34. doi: 10.1080/08927014.2016.1212989
- Trachoo N, Frank JF. Effectiveness of chemical sanitizers against *Campylobacter jejuni*-containing biofilms. *J Food Prot.* (2002) 65:1117–21. doi: 10.4315/0362-028X-65.7.1117
- Mantzios T, Tsiouris V, Kiskinis K, Economou V, Petridou E, Tsiotsos A, et al. In vitro investigation of the antibacterial activity of nine commercial water disinfectants, acidifiers, and glyceride blends against the most important poultry zoonotic bacteria. *Pathogens.* (2023) 12:381. doi: 10.3390/pathogens12030381
- Huang K, Mao Y, Zhao F, Zhang XX, Ju F, Ye L, et al. Free-living bacteria and potential bacterial pathogens in sewage treatment plants. *Appl Microbiol Biotechnol.* (2018) 102:2455–64. doi: 10.1007/s00253-018-8796-9
- Falkinham JO, Pruden A, Edwards M. Opportunistic premise plumbing pathogens: increasingly important pathogens in drinking water. *Pathogens.* (2015) 4:373–86. doi: 10.3390/pathogens4020373
- Raciewicz P, Majewski M, Biesiada H, Nowaczewski S, Wilczyński J, Wyslaska D, et al. Prevalence and characterisation of antimicrobial resistance genes and class I and 2 integrons in multiresistant *Escherichia coli* isolated from poultry production. *Sci Rep.* (2022) 12:6062. doi: 10.1038/s41598-022-09996-y
- Vermeulen B, De Backer P, Remon JP. Drug administration to poultry. *Adv Drug Deliv Rev.* (2002) 54:795–803. doi: 10.1016/S0169-409X(02)00069-8
- Jonsson ME, Chriél M, Norström M, Hofshagen M. Effect of climate and farm environment on *Campylobacter* spp. colonisation in Norwegian broiler flocks. *Prev Vet Med.* (2012) 107:95–104. doi: 10.1016/j.prevetmed.2012.05.002
- Lyngstad TM, Jonsson ME, Hofshagen M, Heier BT. Risk factors associated with the presence of *Campylobacter* species in Norwegian broiler flocks. *Poult Sci.* (2008) 87:1987–94. doi: 10.3382/ps.2008-00132
- Sahin O, Morishita TY, Zhang Q. *Campylobacter* colonization in poultry: sources of infection and modes of transmission. *Anim Health Res Rev.* (2002) 3:95–105. doi: 10.1079/ahr.2002244
- Shane SM. *Campylobacter* infection of commercial poultry. *OIE Rev Sci Tech.* (2000) 19:376–95. doi: 10.20506/rst.19.2.1224
- ISO 10272-1:2006. *Microbiology of food and Animal Feeding Stuffs—Horizontal Method for Detection and Enumeration of Campylobacter spp. — Part 1: Detection method.* Geneva, Switzerland: International Organisation for Standardisation (2006).
- ISO 4833-2:2013. *Microbiology of the Food Chain: Horizontal Method for the Enumeration of Microorganisms—Part 1: Colony Count at 30° C by the Pour Plate Technique.* Geneva, Switzerland: International Organisation for Standardisation (2013).
- ISO 21528-2:2017. *Microbiology of the Food Chain—Horizontal Method for the Detection and Enumeration of Enterobacteriaceae—Part 2: Colony-Count Technique.* Geneva, Switzerland: International Organisation for Standardisation (2017).
- Walsh PS, Metzger DA, Higuchi R. Chelex 100 as a medium for simple extraction of DNA for PCR-based typing from forensic material. *BioTechniques.* (1991) 54:134–9. doi: 10.2144/000114018
- Wang G, Clark CG, Taylor TM, Pucknell C, Barton C, Price L, et al. Colony multiplex PCR assay for identification and differentiation of *Campylobacter jejuni*, *C. coli*, *C. lari*, *C. upsaliensis*, and *C. fetus* subsp. *fetus*. *Society.* (2002) 40:4744–7. doi: 10.1128/JCM.40.12.4744-4747.2002
- Juretschko S, Timmermann G, Schmid M, Schleifer KH, Pommerening-Röser A, Koops HP, et al. Combined molecular and conventional analyses of nitrifying bacterium diversity in activated sludge: Nitrosococcus mobilis and Nitrospira-like bacteria as dominant populations. *Appl Environ Microbiol.* (1998) 64:3042–51. doi: 10.1128/AEM.64.8.3042-3051.1998
- Weisburg WG, Barns SM, Pelletier DA, Lane DJ. 16S ribosomal DNA amplification for phylogenetic study. *J Bacteriol.* (1991) 173:697–703. doi: 10.1128/jb.173.2.697-703.1991
- Geospiza Inc. FinchTV 1.4.0 Chromatogram Viewer; (2004). Available at: <https://digitalworldbiology.com/FinchTV> (Accessed December 18, 2021)
- Kumar S, Stecher G, Li M, Knyaz C, Tamura K. MEGA X: molecular evolutionary genetics analysis across computing platforms. *Mol Biol Evol.* (2018) 35:1547–9. doi: 10.1093/molbev/msy096
- Reimer LC, Sardá Carbasse J, Koblitz J, Ebeling C, Podstawka A, Overmann J. BacDive in 2022: the knowledge base for standardized bacterial and archaeal data. *Nucleic Acids Res.* (2022) 50:D741–6. doi: 10.1093/nar/gkab961
- Parte AC, Carbasse JS, Meier-Kolthoff JP, Reimer LC, Göker M. List of prokaryotic names with standing in nomenclature (LPSN) moves to the DSMZ. *Int J Syst Evol Microbiol.* (2020) 70:5607–12. doi: 10.1099/ijsem.0.004332
- Magiorakos AP, Srinivasan A, Carey RB, Carmeli Y, Falagas ME, Giske CG, et al. Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: an international expert proposal for interim standard definitions for acquired resistance. *Clin Microbiol Infect.* (2012) 18:268–81. doi: 10.1111/j.1469-0691.2011.03570.x
- Clinical and Laboratory Standards Institute (CLSI). *Performance Standards for Antimicrobial Disk and Dilution Susceptibility Tests for Bacteria Isolated from Animals. CLSI Supplement VET01S. 5th ed* Wayne, PA: Clinical and Laboratory Standards Institute (2020). Available at: <http://clsiweb.org/Login.aspx>
- Clinical and Laboratory Standards Institute (CLSI). *VET01S Performance Standards for Antimicrobial Disk and Dilution Susceptibility Tests for Bacteria Isolated From Animals.* An Informational Supplement for Global Application Developed through the Clinical and Laboratory Standards Institute Wayne, PA: Clinical and Laboratory Standards Institute (2015). Available at: <https://clsi.org/>
- Clinical and Laboratory Standards Institute (CLSI). *M100 Performance Standards for Antimicrobial Susceptibility Testing* Wayne, PA: Clinical and Laboratory Standards Institute (2021). Available at: <https://clsi.org/standards/products/microbiology/documents/m100/>
- Heinemann C, Leubner CD, Savin M, Sib E, Schmuthausen RM, Steinhoff-Wagner J. Research note: tracing pathways of entry and persistence of facultative pathogenic and antibiotic-resistant bacteria in a commercial broiler farm with substantial health problems. *Poult Sci.* (2020) 99:5481–6. doi: 10.1016/J.PS.2020.08.050
- Hynds PD, Misstear BD, Gill LW. Development of a microbial contamination susceptibility model for private domestic groundwater sources. *Water Resour Res.* (2012) 48:1–13. doi: 10.1029/2012WR012492
- Maertens H, De Reu K, Van Weyenberg S, Van Coillie E, Meyer E, Meirhaeghe V, et al. Evaluation of the hygienogram scores and related data obtained after cleaning and disinfection of poultry houses in Flanders during the period 2007 to 2014. *Poult Sci.* (2018) 97:620–7. doi: 10.3382/ps/pex327
- Zou Q, Meng W, Wang C, Wang T, Liu X, Li D. Sodium dichloroisocyanurate: improving broiler health by reducing harmful microbial levels in the waterline. *Front Vet Sci.* (2023) 10:1234949. doi: 10.3389/fvets.2023.1234949



46. Wang P, Chen B, Yuan R, Li C, Li Y. Characteristics of aquatic bacterial community and the influencing factors in an urban river. *Sci Total Environ.* (2016) 569–570:382–9. doi: 10.1016/j.scitotenv.2016.06.130
47. Gong Q, Ruan MD, Niu MF, Qin CL, Hou Y, Guo JZ. Immune efficacy of DNA vaccines based on *oprL* and *oprF* genes of *Pseudomonas aeruginosa* in chickens. *Poult Sci.* (2018) 97:4219–27. doi: 10.3382/PS/PEY307
48. Lyczak JB, Cannon CL, Pier GB. Establishment of *Pseudomonas aeruginosa* infection: lessons from a versatile opportunist. *Microbes Infect.* (2000) 2:1051–60. doi: 10.1016/S1286-4579(00)01259-8
49. Medts GD, Viaene NJ. *Pseudomonas aeruginosa* infection on a broiler farm. *Avian Pathol.* (1975) 4:233–7. doi: 10.1080/03079457509353870
50. Walker SE, Sander JE, Cline JL, Helton JS. Characterization of *Pseudomonas aeruginosa* isolates associated with mortality in broiler chicks. *Avian Dis.* (2002) 46:1045–50. doi: 10.1637/0005-2086(2002)046[1045:COPAIA]2.0.CO;2
51. Zamberlan da Silva ME, Filho IC, Endo EH, Nakamura CV, Ueda-Nakamura T, Filho BPD. Characterisation of potential virulence markers in *Pseudomonas aeruginosa* isolated from drinking water. *Anton Leeuw Int J Gen Mol Microbiol.* (2008) 93:323–34. doi: 10.1007/s10482-007-9209-8
52. Teyssier C, Marchandin H, Jean-Pierre H, Diego I, Darbas H, Jeannot JL, et al. Molecular and phenotypic features for identification of the opportunistic pathogens *Ochrobactrum* spp. *J Med Microbiol.* (2005) 54:945–53. doi: 10.1099/jmm.0.46116-0
53. Pompilio A, Crocetta V, De Nicola S, Verginelli F, Fiscarelli E, Bonaventura D, et al. Cooperative pathogenicity in cystic fibrosis: *Stenotrophomonas maltophilia* modulates *Pseudomonas aeruginosa* virulence in mixed biofilm. *Front Microbiol.* (2015) 6:1–13. doi: 10.3389/fmicb.2015.00951
54. Magajna BA, Schraft H. *Campylobacter jejuni* biofilm cells become viable but non-culturable (VBNC) in low nutrient conditions at 4°C more quickly than their planktonic counterparts. *Food Control.* (2015) 50:45–50. doi: 10.1016/j.foodcont.2014.08.022
55. Strakova N, Korena K, Gelbicova T, Kulich P, Karpiskova R. A rapid culture method for the detection of *Campylobacter* from water environments. *Int J Environ Res Public Health.* (2021) 18:6098. doi: 10.3390/ijerph18116098
56. Lv R, Wang K, Feng J, Heeney DD, Liu D, Lu X. Detection and quantification of viable but non-culturable *Campylobacter jejuni*. *Front Microbiol.* (2020) 10:1–8. doi: 10.3389/fmicb.2019.02920
57. Cools I, Uyttendaele M, Caro C, D'Haese E, Nelis HJ, Debevere J. Survival of *Campylobacter jejuni* strains of different origin in drinking water. *J Appl Microbiol.* (2003) 94:886–92. doi: 10.1046/j.1365-2672.2003.01916.x
58. Culotti A, Packman AI. *Pseudomonas aeruginosa* facilitates *Campylobacter jejuni* growth in biofilms under oxic flow conditions. *FEMS Microbiol Ecol.* (2015) 91:136. doi: 10.1093/femsec/fiv136
59. Hanning I, Jarquin R, Slavik M. *Campylobacter jejuni* as a secondary colonizer of poultry biofilms. *J Appl Microbiol.* (2008) 105:1199–208. doi: 10.1111/j.1365-2672.2008.03853.x
60. Aslam B, Khurshid M, Arshad MI, Muzammil S, Rasool M, Yasmeen N, et al. Antibiotic resistance: one health one world outlook. *Front Cell Infect Microbiol.* (2021) 11:771510. doi: 10.3389/fcimb.2021.771510
61. Huijbers PMC, Blaak H, De Jong MCM, Graat EAM, Vandenbroucke-Grauls CMJE, De Roda Husman AM. Role of the environment in the transmission of antimicrobial resistance to humans: a review. *Environ Sci Technol.* (2015) 49:11993–2004. doi: 10.1021/acs.est.5b02566
62. Yang Y, Ashworth AJ, Willett C, Cook K, Upadhyay A, Owens PR, et al. Review of antibiotic resistance, ecology, dissemination, and mitigation in U.S. *Broiler Poultry Syst.* *Front. Microbiol.* (2019) 10:1–10. doi: 10.3389/fmicb.2019.02639
63. Austrian Agency for Health and Food Safety GmbH. Antibiotika-Vertriebsmengen in der Veterinärmedizin; (2022). Available at: <https://www.ages.at/tier/tierarzneimittel-hormone/antibiotika-vertriebsmengen-in-der-veterinaermedizin> (Accessed September 27, 2023)
64. BGBl. I Nr. 36/2008. Bundesgesetz, mit dem das Tierseuchengesetz, das Arzneiwareneinfuhrgesetz das Tierarzneimittelkontrollgesetz und das Rezeptpflichtgesetz Geändert Werden (Veterinärrechtsänderungsgesetz 2007); (2002). Available at: <https://www.ris.bka.gv.at/eli/bgbli/1/2008/36/20080111> (Accessed April 17, 2023)
65. Griesbacher A, Schließnig H, Weber S, Fuchs K. Antimicrobial use in poultry flocks in Austria between 2013 and 2019. *Vet Rec.* (2021) 189:e508. doi: 10.1002/vetr.508
66. BMASGK-74330/0008-IX/B/15/2018, AVN Nr. 2018/11a. Leitlinien für den Sorgfältigen Umgang mit Antibakteriell Wirksamen Tierarzneimitteln; (2018). Available at: [https://www.ris.bka.gv.at/Dokumente/Avn/AVN\\_20181129\\_AVN\\_2018\\_11a\\_2/201811a\\_AntibiotikaLL\\_Beilage.pdf](https://www.ris.bka.gv.at/Dokumente/Avn/AVN_20181129_AVN_2018_11a_2/201811a_AntibiotikaLL_Beilage.pdf) (Accessed March 05, 2023)
67. Qualitätsgeflügelvereinigung (QGV). Antibiotikareport 2022 der Österreichischen Qualitätsgeflügelvereinigung; (2022). Available at: <https://www.qgv.at/allgemein/antibiotikareport-2022> (Accessed September 27, 2023)
68. Roth N, Käsbohrer A, Mayrhofer S, Zitz U, Hofacre C, Domig KJ. The application of antibiotics in broiler production and the resulting antibiotic resistance in *Escherichia coli*: a global overview. *Poult Sci.* (2019) 98:1791–804. doi: 10.3382/ps/pey539
69. Carson C, Li XZ, Agunos A, Loest D, Chapman B, Finley R, et al. Ceftiofur-resistant *Salmonella enterica* serovar Heidelberg of poultry origin - a risk profile using the codex framework. *Epidemiol Infect.* (2019) 147:e296. doi: 10.1017/S0950268819001778
70. Liljebjelke KA, Hofacre CL, White DG, Ayers S, Lee MD, Maurer JJ. Diversity of antimicrobial resistance phenotypes in *Salmonella* isolated from commercial poultry farms. *Front Vet Sci.* (2017) 4:96. doi: 10.3389/fvets.2017.00096
71. Velasquez CG, Macklin KS, Kumar S, Bailey M, Ebner PE, Oliver HF, et al. Prevalence and antimicrobial resistance patterns of *Salmonella* isolated from poultry farms in southeastern United States. *Poult Sci.* (2018) 97:2144–52. doi: 10.3382/ps/pex449
72. World Health Organization (WHO). *Critically Important Antimicrobials for Human Medicine. 6th Revision 2018.* Geneva, Switzerland: World Health Organization (2019).
73. Aldeyab MA, Harbarth S, Vernaz N, Kearney MP, Scott MG, Darwish Elhaggi FW, et al. The impact of antibiotic use on the incidence and resistance pattern of extended-spectrum beta-lactamase-producing bacteria in primary and secondary healthcare settings. *Br J Clin Pharmacol.* (2012) 74:171–9. doi: 10.1111/j.1365-2125.2011.04161.x
74. Lu SY, Zhang YL, Geng SN, Li TY, Ye ZM, Zhang DS, et al. High diversity of extended-spectrum beta-lactamase-producing bacteria in an urban river sediment habitat. *Appl Environ Microbiol.* (2010) 76:5972–6. doi: 10.1128/AEM.00711-10
75. Slama TG. Gram-negative antibiotic resistance: there is a price to pay. *Crit Care.* (2008) 12:1–7. doi: 10.1186/ccXXXX
76. Ashbolt NJ. Microbial contamination of drinking water and human health from community water systems. *Curr Environ Health Rep.* (2015) 2:95–106. doi: 10.1007/s40572-014-0037-5
77. Rilstone V, Vignale L, Craddock J, Cushing A, Filion Y, Champagne P. The role of antibiotics and heavy metals on the development, promotion, and dissemination of antimicrobial resistance in drinking water biofilms. *Chemosphere.* (2021) 282:131048. doi: 10.1016/j.chemosphere.2021.131048
78. Agunos A, Léger D, Carson C. Review of antimicrobial therapy of selected bacterial diseases in broiler chickens in Canada. *Can Vet J.* (2012) 53:1289–300.
79. Alfifi A, Christensen JP, Hounmanou YMG, Sandberg M, Dalsgaard A. Characterization of *Escherichia coli* and other bacteria isolated from condemned broilers at a Danish abattoir. *Front Microbiol.* (2022) 13:1–13. doi: 10.3389/fmicb.2022.1020586