



OPEN ACCESS

EDITED BY
Massimo Castagnaro,
University of Padua, Italy

REVIEWED BY
Barbara Brunetti,
University of Bologna, Italy

*CORRESPONDENCE
Carlos Rosales
✉ carosal@unam.mx

RECEIVED 04 March 2024
ACCEPTED 11 April 2024
PUBLISHED 02 May 2024

CITATION
Uribe-Querol E, Romero-Romero L,
Govezensky T and Rosales C (2024)
Corrigendum: Neutrophil to lymphocyte ratio
and principal component analysis offer
prognostic advantage for dogs with mammary
tumors. *Front. Vet. Sci.* 11:1395930.
doi: 10.3389/fvets.2024.1395930

COPYRIGHT
© 2024 Uribe-Querol, Romero-Romero,
Govezensky and Rosales. This is an
open-access article distributed under the
terms of the [Creative Commons Attribution
License \(CC BY\)](#). The use, distribution or
reproduction in other forums is permitted,
provided the original author(s) and the
copyright owner(s) are credited and that the
original publication in this journal is cited, in
accordance with accepted academic practice.
No use, distribution or reproduction is
permitted which does not comply with these
terms.

Corrigendum: Neutrophil to lymphocyte ratio and principal component analysis offer prognostic advantage for dogs with mammary tumors

Eileen Uribe-Querol¹, Laura Romero-Romero²,
Tzipe Govezensky³ and Carlos Rosales^{4*}

¹Laboratorio de Biología del Desarrollo, División de Estudios de Posgrado e Investigación, Facultad de Odontología, Universidad Nacional Autónoma de México, Mexico City, Mexico, ²Departamento de Patología, Facultad de Medicina Veterinaria y Zootecnia, Universidad Nacional Autónoma de México, Mexico City, Mexico, ³Apoyo de estadística, Instituto de Investigaciones Biomédicas, Universidad Nacional Autónoma de México, Mexico City, Mexico, ⁴Departamento de Inmunología, Instituto de Investigaciones Biomédicas, Universidad Nacional Autónoma de México, Mexico City, Mexico

KEYWORDS

neutrophil, lymphocyte, breast cancer, canine mammary tumor, inflammation, cancer, albumin, globulin

A corrigendum on

Neutrophil to lymphocyte ratio and principal component analysis offer prognostic advantage for dogs with mammary tumors

by Uribe-Querol, E., Romero-Romero, L., Govezensky, T., and Rosales, C. (2023). *Front. Vet. Sci.* 10:1187271. doi: 10.3389/fvets.2023.1187271

In the published article, there was an error. In section 3.3, data on separation of low-grade from high-grade tumors was referred instead of the correct data on separation of surviving from not-surviving dogs.

A correction has been made to Section 3.3 High neutrophil to lymphocyte ratio associates with more aggressive tumors; Second paragraph. The sentence previously stated: “ROC curve analysis identified a cutoff value of NLR = 5 (sensitivity 72.7%; specificity 74.6%) as a threshold separating low-grade from high-grade tumors (Figure 2).”

The corrected sentence appears below:

“ROC curve analysis also identified a cutoff value of NLR = 5 (sensitivity 72.7%; specificity 74.6%) as a threshold separating surviving from not-surviving dogs (Figure 3).”

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.